

# DEENDAYAL PORT TRUST

ISO 9001:2015 & ISO 14001:2015 certified Port



Administrative Office Building  
Post Box NO. 50  
GANDHIDHAM (Kutch).  
Gujarat: 370 201  
Fax: (02836) 220050  
Ph.: (02836) 220038

[www.deendayalport.gov.in](http://www.deendayalport.gov.in)

EG/WK/5202 (D)/ Part (CRZ/142

Dated 08/02/2022

The Additional Secretary & Director (Environment),  
Govt. of Gujarat,  
Forest & Environment Department,  
Block No.14, 8<sup>th</sup> floor, New Sachivalaya,  
**Gandhinagar - 382 010.**

**Sub:** CRZ Clearance for "Creation of water front facilities (Oil Jetties 8, 9, 10 & 11) and development of land of area 554 acres for associated facilities for storage at Old Kandla, Gandhidham, Kutch, Gujarat by M/s Deendayal Port Trust" **Compliances of the stipulated conditions in CRZ Recommendations req.**

**Ref.:** (1) Letter No. ENV-10-2018-24-T Cell dated 30/7/2020 of Director (Environment) & Additional Secretary, Forest & Environment Department, GoG.  
(2) DPT letter no. EG/WK/5202 (D)/ Part (CRZ 2)/28 dated 29/06/2021

Sir,

It is requested to kindly refer the above cited reference for the said subject.

In this connection, it is to state that, the Gujarat Coastal Zone Management Authority vide above referred letter dated 30/7/2020 had recommended the subject project of Deendayal Port Trust. Subsequently, the MoEF&CC, GoI had accorded the Environmental & CRZ Clearance vide letter dated 20/10/2020 for the subject project. Subsequently, DPT vide above referred letter dated 29/06/2021 had submitted compliance report of the stipulated conditions for the period upto May, 2021.

Now, as directed under Specific Condition No. 26 mentioned in the CRZ Clearance letter dated 30/7/2020 i.e. **A six-monthly report on compliance of the conditions mentioned in this letter shall have to be furnished by the OPT on a regular basis to this Department and MoEF&CC, GoI**, we have been regularly submitting the said report vide reference (2) cited letter. Further, please find enclosed herewith compliance report of the stipulated conditions for period June to November, 2021 along with necessary annexure, for kind information & record please **(Annexure I).**

.....Cont .....

Further, as per the MoEF&CC, Notification 5.0.5845 (E) dated 26.11.2018, in which it is mentioned that, **"In the said notification, in paragraph 10, in sub-paragraph (ii), for the words "hard and soft copies" the words "soft copy" shall be substituted"**. Accordingly, we are submitting herewith soft copy of the same via e-mail in ID [gczma.crz@gmail.com](mailto:gczma.crz@gmail.com) & [direnv@gujarat.gcv.in](mailto:direnv@gujarat.gcv.in) .

This has the approval of Chief Engineer, Deendayal Port Trust.

Yours Faithfully,

  
Superintending Engineer (PL) & EMC (I/c)  
Deendayal Port Trust

**Copy to: -**

Shri Amardeep Raju, MoEF&CC, GoI  
Scientist E, Ministry of Environment, Forest and Climate Change,  
& Member Secretary (EAC-Infra.1),  
Indira Paryavaran Bhavan,  
3<sup>rd</sup> Floor, Vayu Wing, Jor Bagh Road, Aliganj,  
New Delhi-110003.  
Email ID: [ad.raju@nic.in](mailto:ad.raju@nic.in)

# **Annexure -I**

**Subject: Point-wise Compliance Status Report for CRZ clearance for proposed project for creation of water front facilities (oil jetties 8,9,10 and 11) and development of land (1432 acres – revised area 554 acres) for associated facilities for storage at old Kandla, Tal: Gandhidham Dist. Kutch, Gujarat by Deendayal Port Trust -reg**

**Ref No: - GCZMA CRZ recommendation vide Letter No- ENV-10-2018-24-T Cell dated 30.07.2020**

<b>S. No.</b>	<b>CRZ Conditions</b>	<b>Compliance Status</b>
	<b>SPECIFIC CONDITIONS</b>	
1.	The DPT shall strictly adhere to the provisions of the CRZ Notification, 2011 issued by the Ministry of Environment, Forests and Climate Change, Government of India	It is assured that, the provisions of the CRZ Notification, 2011 shall be strictly adhere to by the DPT.
2.	Necessary permissions from different departments/ agencies under different laws/ acts shall be obtained before commencing any activity (including the construction)	The Consent to Establish (CTE) from the GPCB had already been obtained vide CTE No. 94118 granted by the GPCB vide letter no. PC/CCA-KUTCH 1524/GPCB ID 56985 dated 23/7/2018 ( <b>Copy Annexure A</b> ).
3.	The DPT shall ensure that that the all the provisions of CRZ Notification 2011 shall be complied with and storage facilities in CRZ areas shall be in compliance with Annexure-II of the above said Notification	It is assured that all the provisions of CRZ Notification, 2011 will be complied with and only storage of permissible cargo as per CRZ Notification, 2011, Annexure II will be allowed to store in storage facilities to be developed.
4.	There shall not be any blockage of creek due to laying of pipeline. and free flow of water shall be maintained.	Work is in progress (Oil Jetty No. 8 - Jetty & allied facilities).  It is hereby assured that, no creeks or rivers shall be blocked, due to any activities at the project site and free flow of water will be maintained.
5.	There shall not be any mangrove destruction/ damage due to proposed activities and adequate buffer zone of 70 metres shall be maintained from mangrove areas	It is assured that all the proposed activities shall be carried out strictly as per the EC & CRZ Clearance accorded by the MoEF&CC, GoI dated 20/11/2020.
6.	The DPT shall effectively implement the Mangrove Development, Protection & Management plan for control of indirect impact on mangrove habitat	As per the directions of the GCZMA and MoEF&CC, GoI, till date (2005-06 to 2019-20), DPT had already undertaken Mangrove Plantation in an area of 1500 Ha. at various locations.



<b>S. No.</b>	<b>CRZ Conditions</b>	<b>Compliance Status</b>
		It is also relevant to submit here that, as per the direction of the Gujarat Coastal Zone Management Authority, DPT had already prepared & submitted a report on mangrove conservation and management plan formulated by Gujarat Institute of Desert Ecology during the study period of Jan-April, 2015 (Report already submitted along with earlier compliance reports submitted). Further, DPT appointed M/s GUIDE, Bhuj vide work order dated 1/9/2017 for "Regular Monitoring of Mangrove Plantation (1300 Ha.) carried out by DPT". DPT had already submitted final report along with compliance reports submitted to the GCZMA for the project of "Construction of 13th to 16th CB" dated 30/11/2019.
7.	The DPT shall have to make a provision that mangrove areas get proper flushing water and free flow of water shall not be obstructed	It is assured that necessary provisions will be made so that mangrove area get proper flushing water and to maintain free flow of water.
8.	The DPT shall have to dispose of the dredged material at the designated dredged material disposal point based on scientific study and approved by the MOEF&CC, GOI	No dredging activity has been started yet. However, it is assured that dredging activity will be carried out strictly as per the requirement of the condition and the same shall be disposed at designated dumping ground (25° 51' 00" N & 70°10' 00" E).
9.	The DPT shall have to maintain the record for generation and disposal of capital dredging and maintenance dredging	No dredging activity has been started yet. However, it is assured that necessary record will be maintained as per the requirement of the condition.
10.	No dredging, reclamation or any other project related activities shall be carried out in the CRZ area categorized as CRZ I (i) (A) and it shall have to be ensured that the mangrove habitat and other ecologically important and significant areas, if any, in the region are not affected due to any of the project activities.	It is assured that all the project related activities will be strictly carried out as per the EC & CRZ Clearance accorded by the MoEF&CC, GoI dated 20/11/2020.

<b>S. No.</b>	<b>CRZ Conditions</b>	<b>Compliance Status</b>
11.	The DPT shall ensure that construction activities like dredging etc shall be carried out in confined manner to reduce the impact on marine environment.	No dredging activities have been started yet. However, it is assured that construction activities like dredging will be carried out as per the requirement of the condition.
12.	The DPT shall ensure that the dredging shall not be carried out during the fish breeding season	No dredging activities have been started yet. Point Noted for compliance.
13.	Construction waste including debris and dredged material shall be disposed safely in the designed areas as approved by MoEF&CC, Gol and it shall be ensured that there shall be no impact on flora and fauna	Point Noted for compliance.
14.	No effluent or sewage shall be discharged into the sea / creek or in the CRZ area and shall be treated to conform the norms prescribed by the Gujarat Pollution Control Board and would be reused / recycled as per the approval of the Board	It is assured that No effluent or sewage will be discharged into the Sea/creek or in the CRZ area. Further, the same will be treated in STP as per the norms prescribed by the GPCB.
15.	All the recommendations and suggestions given by the Choramandalam MS Risk Services Limited in their Environment Impact Assessment report shall be implemented strictly by DPT	It is assured that all the recommendations and suggestions given by the EIA Consultant, M/s SV Enviro, Vizag in EIA Report will be implemented.
16.	The DPT shall exercise extra precautions to ensure the navigation safety and mitigation of the risk associated with the project activities especially due to collision, sinking or accidents of the vessels and would deploy the latest communication and navigation aids for this purpose. The proposed facilities shall also be covered under the VTMS being developed by the GMB	It is assured that emergency preparedness plan based on the Hazard Identification and Risk Assessment (HIRA) will be implemented.
17.	The cost of the external agency that may be appointed by this department for supervision / monitoring of the project activities during construction/ operational phases shall be paid by DPT	Point Noted.
18.	The DPT shall contribute financially for any common study or project that may be proposed by this Department for environmental management / conservation / improvement for the Gulf Kutch	Point noted for compliance.

S. No.	CRZ Conditions	Compliance Status
19.	The piling activities debris and any other type of waste shall not be discharged into the sea or creek or in the CRZ areas. The debris shall be removed from the site immediately after the piling activities are over	Work is in progress (Oil Jetty No. 8 - Jetty & allied facilities). Point Noted for compliance.
20.	The camps shall be located outside the CRZ area and the labour shall be provided with the necessary amenities, including sanitation, water supply and fuel and it shall be ensured that the environmental conditions are not deteriorated by the labours.	Point Noted for compliance.
21.	The DPT shall prepare and regularly update their Local Oil Spill Contingency and Disaster Management Plan in consonance with the National Oil Spill and Disaster Contingency Plan	Point Noted for compliance.  DPT is already having Local Oil Spill contingency plan and updated DMP.
22.	The DPT shall bear the cost of the external agency that may be appointed by this Department for supervision / monitoring of proposed activities and the environmental impacts of the proposed activities	Point noted for compliance.
23.	The groundwater shall not be tapped to meet with the water requirements in any case	Point Noted for compliance.
24.	DPT shall take up greenbelt development activities in consultation with the Gujarat institute of Desert Ecology / Forest Department / Gujarat Ecology Commission	DPT has already developed Green belt in and around the Port area.  Further, DPT assigned work for Green belt development in an area of about 32 hectares to the Forest Department, Govt. of Gujarat during August, 2019 at the cost of Rs. 352.32 lakhs. The work is completed. Further, DPT also undertook massive green belt development in and around the Port area and at Gandhidham area.
25.	The DPT shall have to contribute financially for taking up the socio-economic upliftment activities in this region in consultation with the Forests and Environment Department and the District Collector / District Development Officer	As per the CSR Guidelines issued by the Ministry of Ports, Shipping & Waterways, Government of India, from time to time, DPT had undertaken CSR activities since the year 2011-12. The details of CSR Activities undertaken & planned is attached herewith as <b>Annexure B.</b>
26.	A six-monthly report on compliance of the conditions mentioned in this letter shall have to be furnished by DPT on a regular basis to this Department and MoEF&CC, GoI.	DPT has been regularly submitting the six-monthly report on compliance of the conditions mentioned in the CRZ Recommendation letter dated 30/7/2020 to the CRZ Authority and to the MoEF&CC, GoI.
27.	The DPT shall ensure that the numbers of the Vessels and machinery deployed during	Point Noted for compliance.

<b>S. No.</b>	<b>CRZ Conditions</b>	<b>Compliance Status</b>
	marine construction, which are a source of low level organic and PHC pollution will be optimized to minimize risks of accidents involving these vessels.	Work is in progress (Oil Jetty No. 8 - Jetty & allied facilities).
28.	The noise level during transport and construction of marine facilities shall be kept minimum.	DPT appointed M/s Detox Corporation, Surat for Monitoring of environmental parameters since the year 2016. The work is in progress & DPT submitted monitoring data regularly to all the concerned authorities along with compliance reports submitted. The monitoring reports are attached herewith as <b>Annexure C</b> .
29.	The DPT shall regularly conduct the surveys to identify changes in the channel bathymetry to minimize navigation hazards. Proper navigational aids and guidance should be provided to ships navigating the channel and there should be a properly structured vessels traffic management strategy to avoid accidents.	Point noted for compliance.
30.	The DPT shall carry out separate study for further erosion and deposition pattern in the area after dredging through a reputed agency and shall follow the suggestions of the study done by reputed agency, for maintenance dredging, the recommendations /suggestions of the reputed agency shall be follow by the DPT	No dredging activity has been started yet. However, it is assured that necessary will be conducted as per the requirement of the condition.
31.	Any other condition that may be stipulated by this Department and MoEF&CC, Gol from time to time for environmental protection / management purpose shall also have to be complied with by DPT.	Point noted.



# **ANNEXURE A**



# GUJARAT POLLUTION CONTROL BOARD

PARYAVARAN BHAVAN

Sector-10-A, Gandhinagar 382 010

Phone : (079) 23222425

(079) 23232152

Fax : (079) 23232156

Website : www.gpcb.gov.in

By R.P.A.D

## CONSENT TO ESTABLISH

CTE- 94118

No. PC/CCA-KUTCH-1524/GPCB ID 56985/

Date:

To,

Deendayal Port Trust Land,

Kandla Port Trust Land,

A.O Building, P.O box No. 50,

Tal.: Gandhidham,

Dist. Kutch-370201

**Subject** : Consent to Establish (NOC) under Section 25 of Water (Prevention and Control of Pollution) Act 1974 and Section 21 of Air (Prevention and Control of Pollution) Act 1981

**Reference** : Your CTE Application Inward ID No 133847 dated 04/04/2018

Sir,

Without prejudice to the powers of the Board under the Water (Prevention and Control of Pollution) Act-1974, the Air (Prevention and Control of Pollution) Act-1981 and the Environment (Protection) Act-1986 and without reducing your responsibilities under the said Acts in any way, this is to inform you that the Board grants **Consent to Establish (NOC)** of industrial activity at **Kandla Port Trust Land, A.O Building, P.O box No. 50, Tal.: Gandhidham, For Creation of water front facilities of oil jetties of 8,9,10,&11 & development of land (1432 Areas).**

1. The validity period of the order shall be up to **03/04/2023**

### SUBJECT TO FOLLOWING SPECIFIC CONDITIONS:

1. Proposed jetties shall be handled of 3.5 MMTP/Annum of liquid cargo of edible oil, Fertilizer & food grains etc.
2. Unit shall strictly adhere to all condition of TOR issued by MoEF & CC, Delhi dated 04/08/2017 & shall not carry out any construction activities till obtaining EC & CRZ from competent authority
3. No ground water shall be withdrawn without prior approval from competent authority.

### 2. CONDITIONS UNDER WATER ACT 1974:

- 2.1 There shall be no industrial water consumption and hence there shall be no industrial waste water generation from manufacturing process and other ancillary operations.
- 2.2 Domestic water consumption shall not exceed 20 KL/day.
- 2.3 The quantity of domestic waste water (Sewage) shall not exceed 16 KL/Day.
- 2.4 The quality of the sewage shall conform to the following standards

Page 1 of 3

PARAMETERS	GPCB NORMS
pH	6.5 to 9.0
BOD (5 days at 20° C)	30 mg/L
Suspended Solids	100 mg/L
Fecal Coliform	1000 MPN/ 100 ml

2.5 The domestic sewage shall be treated in Sewage Treatment Plant and treated sewage conforming to standards mentioned in 2.4 shall be reused in various activities shall not be used for gardening and plantation purpose in premises.

**3. CONDITIONS UNDER AIR ACT 1981:**

3.1 There shall be no use of fuel hence there shall be no flue gas emission from manufacturing process and other industrial operations.

3.2 There shall be no process gas emission from manufacturing process and other industrial operations.

3.3 The concentration of the following parameters in the ambient air within the premises of the industry shall not exceed the limits specified hereunder as per National Ambient Air Quality Standards issued by MoEF&CC dated 16<sup>th</sup> November-2009.

Sr. No.	Pollutant	Time Weighted Average	Concentration in Ambient air in $\mu\text{g}/\text{m}^3$
1	Sulphur Dioxide ( $\text{SO}_2$ )	Annual	50
		24 Hours	80
2	Nitrogen Dioxide ( $\text{NO}_2$ )	Annual	40
		24 Hours	80
3	Particulate Matter (Size less than $10 \mu\text{m}$ ) OR $\text{PM}_{10}$	Annual	60
		24 Hours	100
4	Particulate Matter (Size less than $2.5 \mu\text{m}$ ) OR $\text{PM}_{2.5}$	Annual	40
		24 Hours	60

3.4 The level of Noise in ambient air within the premises of industrial unit shall not exceed following levels:

Between 6 A.M. to 10 P.M.	: 75 dB(A)
Between 10 P.M. to 6 A.M.	: 70 dB(A)

**4. CONDITIONS UNDER HAZARDOUS WASTE:**

4.1 The applicant shall provide temporary storage facilities and maintain the record for each type of Hazardous Waste as per Hazardous Waste (Management, Handling & Transboundary Movement) Rules, 2016 as amended from time to time

4.2 The applicant shall be obtain membership of common TSDF site for disposal Hazardous Waste as categorized in Hazardous Waste (Management, Handling & Transboundary Movement) Rules, 2016 as amended thereof

**5. GENERAL CONDITION**

5.1 Any change in personnel, equipment or working conditions as mentioned in the consents form/order should immediately be intimated to this Board.

5.2 The waste generator shall be totally responsible for (i.e. Collection, storage, transportation and ultimate disposal) of the wastes generated.

5.3 Records of waste generation, its management and annual return shall be submitted to Gujarat Pollution Control Board in Form - 4 by 31st January of every year.

5.4 In case of any accident, details of the same shall be submitted in Form - 5 to Gujarat Pollution Control Board

5.5 Applicant shall comply relevant provision of "Public Liability Insurance Act-91".

Outward No: 462839, 21/01/2018



## GUJARAT POLLUTION CONTROL BOARD

PARYAVARAN BHAVAN

Sector-10-A, Gandhinagar 382 010

Phone : (079) 23222425

(079) 23232152

Fax : (079) 23232156

Website : www.gpcb.gov.in

- 5.6 Unit shall take all concrete measures to show tangible results in waste generation reduction, avoidance, reuse and recycle. Action taken in this regards shall be submitted within 03 months and also along with Form 4
- 5.7 Industry shall have to display on-line data outside the main factory gate with regard to quantity and nature of hazardous chemicals being handled in the plant, including wastewater and air emissions and solid hazardous waste generated within the factory premises.
- 5.8 Adequate plantation shall be carried out all along the periphery of the industrial premises in such a way that the density of plantation is at least 1000 trees per acre of land and a green belt of 10 meters width shall be developed.
- 5.9 The applicant shall have to submit the returns in prescribed form regarding water consumption and shall have to make payment of water cess to the Board under the Water (Prevention and Control of Pollution) Cess Act- 1977.

For and on behalf of  
Gujarat Pollution Control Board

(Sushil Vegda)  
Senior Environment Engineer

Outward No: 462839, 23/07/2018



# **ANNEXURE B**

## CSR Activities at Decendaryal Post Tms

## Details of CSR

Sr. No	Year	Board Resolution For Budget Provision	Board Approved Budget Provision	Board Resolution for approval of the CSR activities	Board Approved Amount For CSR Activities	Actual exp. upto Nov'20 (Rs. In Lakhs)	Net balance (Rs. In Lakhs)	Remarks
1	2	3	4	5	6	7	6-7	
1	2011-2012	369 of 28.03.2012	3.00 Cr					
2	2012-2013	17 of 31.05.2012	4.00 Cr					
3	2013-2014	99 of 30.09.2013	6.43 Cr	64 of 30.08.2012	564.00 Lakh	564.00	Nil	Works completed
4	2014-2015	322 of 21.11.2014	1.07 Cr	20 of 16.04.2015	236.22 Lakh	188.18	8.04	Works in progress
5	2015-2016	151 of 12.02.2016	1.50 Cr	48 of 12.08.2016	28.00 Lakh	5.00	23.00	Works in progress
6	2016-2017	138 of 06.01.2017	2.60 Cr	52 of 2.8.2017	140.301 lakh	146.00	-5.70	Works completed
7	2017-2018	41 of 2.08.2017	7.02 Cr	15 of 04.05.2018	155.10 Lakh	115.37	39.73	Works in progress
8	2018-19	51 of 07.08.2019	6.70 Cr	111 of 4.12.2018	154.90 Lakh	50.50	104.40	Works in progress
					1278.52 Lakh	1069.05	209.47	
9	2019-20	58 of 10.10.2019	5.49 Cr	92 of 06.12.2019	1838.57 Lakh	Nil		MoS approval is awarded
		<b>Total</b>	<b>37.81 Cr</b>		<b>3117.09 Lakh</b>			

Spent in PM Fund for COVID-19-800 Lakhs

Year-wise details of CSR works undertaken by DPT during 2012 – 13 to 2019 – 20 are given in Tables 7.3a, 7.3b, 7.3c, 7.3d, 7.3e, 7.3f and 7.3g.

**Table 7.3a: CSR Works Undertaken by DPT during 2011-12 and 2012 – 13**

Sl. No.	Name of Work	Cost (Rs. In lakhs)
1	Repair of road from Dr. Baba Saheb Ambedkar Circle to NH 8A (via Ganesh Nagar)	518
2	Repair of road from S.T. Bus Stand to Sunderpuri Cross Road via Collector Road	
3	Repair of road from NH 8A Railway Crossing to Maninagar (along railway track)	
4	Repair of road from Khanna Market Road (Collector Road) to Green Palace Hotel	
5	Construction of internal roads at "Shri Ram" Harijan Co-operative Housing Society (near Kidana)	
6	Construction of cremation ground and graveyard with other facilities at Vadinar	19.44
7	Providing cement concrete internal roads in Village Vadinar Stage - I	16.16
8	Approach road provided for developing tourism at Village Veera near Harsidhi Mata Temple	4.65
9	Water tank along with R.O. provided near developing tourism area	0.30
10	Creating facilities of flooring and steps surrounding lake to stop soil erosion and attract tourists at Village Veera.	4.80
	<b>TOTAL</b>	<b>563.35</b>

**Table 7.3b: CSR Works Undertaken by DPT during 2014-15**

Sl. No.	Name of Work	Cost (Rs. In lakhs)
1	Construction of community hall – cum – school at Maheshwari Nagar, Gandhidham	51.90
2	Renovation of "Muktidham" (cremation ground) at Kandla	10.65
3	Sunderpuri – 1 Valmiki Community Hall	5.00
4	Sunderpuri – 2 Valmiki Community Hall	5.00
5	Ganeshnagar Community Hall	10.00
6	Jagjivan Maheshwari Community Hall	10.00
7	Various works of road at Sapnanagar	99.19
8	Construction of compound wall in the dam of Jogninar Village	14.48
	<b>TOTAL</b>	<b>206.22</b>

**Table 7.3c: CSR Works Undertaken by DPT during 2015-16**

Sl. No.	Name of Work	Cost (Rs. In lakhs)
1	Construction of Bus Stand at Vadinar Village	10.00
2	Providing drainage system at Vadinar Village	6.00
3	Providing and laying of water supply lines in Vadinar Village	6.00
4	Road from Gandhidham Post Office to Merchantile Marine Department Office along with toilet facilities	60.00
5	Construction of toilets for girls / women at Khari Rohar, Village	3.00
6	Construction of toilets for girls at Mathak Primary School, Mathak, Village	3.00
	<b>TOTAL</b>	<b>88.00</b>

**Table 7.3d: CSR Works Approved by DPT Board for 2016-17**

Sl. No.	Name of Work	Cost (Rs. In lakhs)
1	RCC community hall at Harsidhi Mata Temple, Village Veera, Anjar Taluka	19.00
2	Fabricated Community Hall at Sanghad Village, Anjar Taluka	21.00
3	CSR Works for Shri Maheshwari Meghvad Samaj, Gandhidham at graveyard behind Redison Hotel	8.00
4	CSR Works for Shri Dhanraj Matiyadev Mukti Dham, Sector 14, Rotary Nagar, Gandhidham	30.50
5	CSR Works for Nirvasit Harijan Co-operative Housing Society, Gandhidham Health Cum Education Centre	41.00
6	CSR Works for Shri Rotary Nagar Primary School, Gandhidham	2.80
7	CSR Works at NU-4, NU-10(B) Sapnanagar & Saktinagar, Golden Jubilee Park at Gandhidham	18.00
	<b>TOTAL</b>	<b>140.30</b>

**Table 7.3e: CSR Works Approved for 2017-18**

Sl. No.	Name of Work	Proposal Received from / / Name of Organization / N.G.O	Cost (Rs. In lakhs)
1	CSR Works at Shri Ganesh Nagar High School, Gandhidham	Principal, Shri Ganesh Nagar Govt High School, Gandhidham	38.30 Lakhs
2	CSR Works for MOLANA AZAD Primary School, Kandla	Shri M L Bellani, Trustee, DPT, Shri Kandla Port Education Society, New Kandla	7.00 Lakhs
3	Grant financial contribution for facility of Army Cantonment for 50 nos. air coolers at Kutch Border Area	Shri Vinod L Chavda, MP	15 Lakhs
4	40% of the estimated cost of providing drainage lines at Tuna and Vandi villages under Swachh Bharat Abhiyan.	Shri Sarpanch, Tuna Village & Vandi village & Shri M L Bellani, Trustee, DPT	<b>Rs. 39.80 Lakhs</b> <i>Approx. estimated Cost Rs.99.50 Lakhs, of which 40% to be contributed by DPT.</i>
5	CSR works for S.H.N. Academy English School (managed by Indian Inst. Of Sindhology – Bharati Sindhu Vidyapeeth), Adipur	Director, S.H.N Academy English School	<b>40 Lakhs</b>
6	Construction of internal roads at Bhaktinagar Society, Kidana	Smt Maltiben Maheshwari, MLA	<b>15 Lakh</b>
		<b>TOTAL</b>	<b>155.10</b>

**Table 7.3f: CSR Works Approved for 2018-19**

Sl. No.	Name of Work	Proposal Received from / / Name of Organization / N.G.O	Cost (Rs. In lakhs)
1	CSR work to Donate 100 Nos of Computers to Daughters of Martyred Soldiers in the country under the "BETI BACHAO BETI PADHAO" program by Atharva Foundation, Mumbai	Chairman, Atharva Foundation, Mumbai	<b>24.00</b>
2	CSR work to Donate ONE (40 Seater) School Bus for Deaf Children Students for the Institute of Mata Lachmi Rotary Society, Adipur	Mata Lachmi Rotary Society, Adipur	<b>18.00</b>
3	CSR work to Providing One R.O Plant with Cooler at PanchyatPrathmikSala, Gadpadar Village for the ANARDE Foundation, Kandla&Gandhidham Center.	Dist. Rural Development Officer, Annarde Foundation-Kandla & Gandhidham	<b>1.50</b>
4	CSR work for Providing Drainage Line at MeghparBorichi village, AnjarTaluka	Shri Vasambhai Ahir, MLA, Gujarat Govt	<b>25.00</b>
5	CSR work for Construction of Health Centre at Kidana Village	Shri Vinod L Chavda, MP	<b>13.00</b>
6	CSR work to provide 4 Nos. of Big Dust Bin for MithiRoharJuth Gram Panchayat.	Shri Sarpanch, Mithi RoharJuth Gram Panchayat	<b>3.40</b>



Sl. No.	Name of Work	Proposal Received from // Name of Organization / N.G.O	Cost (Rs. In lakhs)
7	CSR work for Renovation & construction of shed at CharanSamaj, Gandhidham –Adipur.	Shri Vinod L Chavda, MP	10.00
8	CSR Work for Renovation/Repairing of Ceiling of School Building at A. P Vidhyalay, Kandla.	Smt Maltiben K. Maheshwary, MP, Gandhidham.	10.00
9	CSR work for Construction of Over Head Tank & Providing 10 Nos of Computers (for students) of NavjivanViklangSevashray, Bhachau, Kutch	Shri Jitendra Joshi, Founder Secretary, Shri Navjivan Viklang Sevashray, Bhachau, Kutch	9.50
10	CSR work to Provide Books & Tuition fees for Educational facilities to weaker section children of ValmikiSamaj, Kutch.	Shri Manohar Jala, Chairman of “National Commission of Safai Karamcharis”	2.00
11	CSR work to provide Water Purifier & Cooler for the ST. Joseph’s Hospital, Gandhidham	Smt. Maltiben K Mahewari, MLA ,Gandhidham	1.50
12	CSR work for Construction of Second Floor (Phase – I) for Training Centre of “GarbhSanskran Kendra” “Samarth Bharat Abhiyan” of Kutch Kalyan Sangh, Gandhidham	Shri Vinod L Chavda, MP, Kutch	37.00
		<b>TOTAL</b>	<b>154.90</b>

**Table 7.3g: CSR works approved for the year 2019-20 (approval from Ministry of Shipping still awaited)**

Sl. No.	Name of Work	Proposal Received from // Name of Organization / N.G.O	Cost (Rs. In lakhs)
1	CSR activities for Providing Drainage line at Nani Nagalpar village.	Sarpanch of Village:-Nani Nagalpar, Taluk: Anjar.	3.00
2	CSR activities for Development of ANGANWADI Building at School no- 12 at Ward no 3 & 6 at Anjar.	Shri Vasanhbai Ahir, MLA	7.00
3	CSR activities for Improving the facilities of Garden at Sapna Nagar(NU-4)& (NU-10 B),Gandhidham.	Shri K P Maheshwari, Resident Sapnanagar, Gandhidham	18.00
4	CSR activities for Providing of Plastic Shredding Machine to Mirror Charitable Trust, Gandhidham.	Mirror Charitable Trust ,Gandhidham	4.75
5	CSR activities for development of School premises of Shri Guru Nanak Edu. Society, Gim.	Shri Guru Nanak Education Society, Gandhidham.	30.00
6	CSR activities for the improvement of the facilities at St. Joseph Hospital & Shantisadan at Gandhidham	St. Joseph Hospital Trust, Gandhidham	20.00
7	CSR activities for the improvement of the facilities at SVP (SardarValabhbbhai Patel ) Multipurpose Hall at Gandhidham	Request from MarwadiYuva Munch & UNION Gandhidham	500.00
8	Consideration of Expenditure for running of St Ann’s High School at Vadinar of last 5 years 2014 to 2019 under CSR.	Proposal from COM, OOT Vadinar, DPT	825.00
9	CSR activities for development of school premises of Shri Adipur Group Kanya Sala no-1 at Adipur	Principal, Shri Adipur Group KanyaSala, Adipur	6.50
10	CSR activities for development of school premises of Shri Jagjivan Nagar Panchyat Prathmiksala, Gandhidham.	Principal, Shri Jagjivan Nagar Panchyat Prathmiksala, Gandhidham.	16.50
11	CSR activities for development of school premises of Ganeshnagar Government high school, Gandhidham.	Shri Vinod L Chavda, MP, Kutch	9.00
12	CSR activities for improving greenery, increase carbon sequestration and beat Pollution at Kandla, DPT reg.	Work awarded to Forest Department , Bhuj	352.32
13	CSR activities for providing infrastructures facilities at “Bhiratna Sarmas Kanya Chhatralaya” under the Trust of Samaj Nav- Nirman at Mirjapur highway, Ta Bhuj.	SamajNav- Nirman at Mirjapur highway, Ta Bhuj.	46.50
		<b>TOTAL</b>	<b>1838.57</b>

**List of CSR applications received from various NGOs , Organizations , Village Sharpnchs etc for the FY 2021-22 .**

Sr.No	Name of Scheme	Proposal Received from / Name of Organization / N.G.O	Brief Details
1	CSR activities for the development of gardening at Sector -5 , Gim	Shri Sarvodaya Co-Operative Housing Society Ltd	<b>Appx Cost – Rs 25.00 Lakhs</b>  <b>Cost for –</b> Comp wall, Benches, Plantation, walkway, other facilities  (Land is reserved for Garden development only since from 50 years)
2	CSR activities for providing various facilities in SHRI GANESHNAGAR GOVT HIGHSCHOOL, GANDHIDHAM	Principal of School	<b>Appx cost –Rs 20.00 Lakhs</b>  (Two times CSR works carried out at school by DPT)
3	CSR activities for the VadhiyarVankarSamajvaadi, NaviSunderpuriGim	SmtMaltiben K Maheswari, MLA	<b>Appx Cost Rs 6.00 Lakhs</b>  Cost for Const. of Comp Wall
4	CSR activities for Construction work of Cabin at Oslo Area- Gim	SmtMaltiben& Shri VinadChavda	Cost not mentioned.
5	CSR activities & Land requirement forAkhil Kutch SamastaMeghvanshiGurjarmeghwal Charitable Trust ,Gim.	Shri Akhil Kutch SamastaMeghvanshiGurjarmeghwal Charitable Trust. Shri Dharmendra R Gohil	Cost Not mentioned.  (demand of Land for development of SAMAJ VADI in Gandhidham)
6	CSR Activities for providing Water supply pipe line, Play ground and sports equipment, electric facilities, drinking water facilities for poor people & Fishermen at VANDI Village.	Shri R RKhambhra, PRO , Collector Office, Bhuj.	<b>Appx Cost Rs 51.00 Lakhs</b>  (Last year also applied by village Sarpanch ) &  Recommended by Shri VASANBHAI AHIR, MLA, Shri V L Chavda, MP)
7	CSR activities for the Tuna village,	Sarpanch, Tuna village	<b>Appx Cost Rs. 25 Lakhs</b> <b>Cost for :-</b>

**List of CSR applications received from various NGOs , Organizations , Village Sharpnchs etc for the FY 2021-22 .**

Sr.No	Name of Scheme	Proposal Received from / Name of Organization / N.G.O	Brief Details
	Ta -Gim		2 No Fab shed 20'x20'x1250= 10 Lakh 2 Nos of Agnawadi =10 Lakh Fab shed for school=5 Lakh
8	CSR activities for the Global Vision India Foundation, Gim	Global vision India Foundation, G'dham	Requirement of Land –OR- Old building at Gandhidham for foundation of welfare activities.
9	CSR activities for the UNITED ORPHANAGE FOR THE DISABLED, TAMIL NADU	UNITED ORPHANAGE FOR THE DISABLED, TAMIL NADU	<b>Cost Rs 25,000.00</b>  (Winter sweaters for children)
10	CSR activities for the Garden Development on already bounded area with Compound wall near Plot no 448 Sector-1/A, Gandhidham.	Residents, near Plot no 448, Sector-1/A, Gim.	<b>AppxCost Rs 20.00 Lakhs</b>  (Requirement to provide benches, drinking water facility, plantation, lightings & walkways in side bounded area)
11	CSR activities for donation of Land for the Shri SUNDARPUI Govt Primary School, Gim	SmtMalti ben Maheshwari, MLA	(request for Land Requirement)
12	CSR activities for Extension of Adarsh Primary School building, Adipur	GandhidhamMatri Mandal, English Medium School, Adipur	<b>Appx Cost Rs. 40.00 Lakhs</b> (Construction for 4 Rooms extension)  (Trust registered under Societies Registration Act XXI -1860, Reg No F-42 dtd 23.9.1965. Land belong to Trust)
13	CSR Activities for providing HD projector for KANYA MAHA VIDYALAYA, Adipur	Principal, KANYA MAHA VIDYALAYA, Adipur	<b>Cost Rs 1.50 Lakhs</b>  (School Managed by G'dhamMaitry Mandal, Adipur)

**List of CSR applications received from various NGOs , Organizations , Village Sharpnchs etc for the FY 2021-22 .**

Sr.No	Name of Scheme	Proposal Received from / Name of Organization / N.G.O	Brief Details
14	CSR activities for DONATION various Medical Equipment for the Hospital of Gandhidham Jain SevaSamiti, Adipur	Gandhidham Jain SevaSamiti, Adipur	<b>Cost for :-</b> 1) Fresenius Haemodialysis Machine Rs 38.00 Lakh 2) Maltislice Helical CT Scanner- Rs 52.00 Lakhs 3) Others Rs 54.00 Lakhs <b>(Total Appx Cost Rs 144 Lakhs)</b>
15	CSR activities for SHRI VIDI JUTH GRAM PANCHAYAT, Vidi, Anjar	Sarpanch, Vidi Gram	<b>Appx Cost Rs 30.00 Lakhs</b>  Cost for- Drainage , Garbage vehicle, and Cattle shed  (Already applied earlier at Sr-5/12)
16	CSR activities for SOS CHILDRESN'S VILLAGES INDIA, Madhapar, Bhuj	Director, SOS Children's Village of India-Bhuj	<b>Appx Cost Rs 31.00 Lakhs</b>  (request for Financial support towards parentless and abandoned Children Education support located at Bhuj ) & support to women working in SOS.
17	Gujarat Biodiversity Board, Gandhinagar invites to involved National & Global endeavour of conservation of biodiversity by creating financial partnership with GBB under CSR programme of expenditure to be incurred 187 Lakh.	GUJARAT BIODIVERSITY BOAD, GANDHINAGAR	<b>Requirement-</b> Financial Support from DPT for <b>AppxRs 1.88 Cr.</b>  (Cost for various meetings, collection of primary data from villagers , processing of documentation, printing , TA DA of Technical support & Miscexp for 150 Peoples Biodiversity Register (PBR).

**List of CSR applications received from various NGOs , Organizations , Village Sharpnchs etc for the FY 2021-22 .**

Sr.No	Name of Scheme	Proposal Received from / Name of Organization / N.G.O	Brief Details
18	CSR activities for providing furniture & Home appliances for ROJAVANAM TRUST at Madurai.	Shri Arul Kannan, Director	<b>Appx Cost Rs 30 Lakhs</b>  (seeking help to provide facilities to Aged & Homeless people living in Trust and Purchasing of New Ambulance)
19	CSR activities for providing Dialysis Machine for treatment of Kidney patients at "ST JOSEPH'S HOSPITAL TRUST" at Gandhidham.	Sr. Franciline, Administrator of Hospital.	<b>Appx Cost Rs 31.36 Lakhs</b>  (Cost of 5 Nos of Dialysis Machines for treatment of kidney patients)
20	CSR activities for providing facilities in Girls Hostel of Gasturba Gandhi BalikaVidhyalay, Gandhidham.	Shri Vinod L Chavda, MP	<b>Appx cost Rs 30 Lakhs.</b>  (Cost of Comp Wall, Entrance gate, Girls toilets etc)
21	CSR works for providing Oxygen Generator Plant and 45 KV Silent Generator for COVID HOSPITAL at Swami LilashahKutia, Adipur.	Secretary, BHARAT VIKAS PARISHAD, Gandhidham	<b>Appx Cost Rs 80.00 Lakhs</b>  (Facilities for 100 Beds of COVID patient which it to be extend upto 240 Beds)
22	CSR works for providing Two Numbers of Oxygen Concentrator and others medical equipment for the Trust ,Antarjal, Gim.	President SHRI SARV JEEV KALYAN TRUST, ANTARJAL, Gandhidham	<b>Appx Cost Rs21.50 Lakhs</b>  (Facilities to be provided for the treatment of CORONA PATIENTS at their trust.)
23	CSR works for providing Fabricated Shed , Construction of Compound Wall and Land levelling for the Cattle of <b>GauSevaSamiti-Tappar</b> at Gram-Tappar, Ta Anjar.	Shri Vinod Chavda, MP & Presedent , GauSevaSamiti, village Tappar, Ta-Anjar	<b>Appx Cost Rs84 Lakhs</b>  (Facilities to be provided for Cattle shelters at Village.) (Land belongs to Gram-panchayat)
24	CSR works for Construction of Auditorium Hall at RSETI (Rural Self Employment Training Institute) at	Shri Vinod Chavda, MP & Director of RSETI, Bhuj	<b>Cost not mentioned.</b>  (Facilities to be provided

**List of CSR applications received from various NGOs , Organizations , Village Sharpnchs etc for the FY 2021-22 .**

Sr.No	Name of Scheme	Proposal Received from / Name of Organization / N.G.O	Brief Details
	Bhujodi-Bhuj.		for the people needs Self-employment activities.)
25	CSR works for Providing of Furniture for the School “SHRI GALPADAR PANCHAYAT PRATHMIC KUMAR GROUP SALA “ atGalpadar Village Ta Gim.	Principal, SHRI GALPADAR PANCHAYAT PRATHMIC KUMAR GROUP SALA “ atGalpadar Village Ta Gim.	<b>Cost not mentioned.</b>  (Facilities to be provided for the Students of Workers & poor village people who study in the school.)
26	Construction of Shed, hall and Gate for the DADA Bhagwandas Charitable Trust, Adipur. (Sr no -4)	Shri Vinod Chavda, MP & DADA BHAGWANDAS CharitableTrust, Gandhidham	<u>As per CSR Guideline-</u> <ul style="list-style-type: none"> <li>➤ Promoting gender equality and empowering women</li> <li>➤ Eradicating extreme hunger and poverty</li> </ul> (Considered shed and hall )  Fab Shelter Shed - 30’x100’ x 1250=37.00 Lakh & RCC Hall – 20’x100’x1500=30.00 Lakh  <b>(Appx Cost Rs67.00 Lakhs)</b> Land authority belongs to Trust given by GDA and NOC given by SRC.Doc submitted.
27	CSR work for reconstruction of the Internal Roads of the Sector-9B-C and Sector-10 area in Gandhidham.	President, Shri TejaKangad, The Gandhidham Chamber of Commerce and Industry, Gandhidham.	Cost not mentioned.

**List of CSR applications received from various NGOs , Organizations , Village Sharpanchsetc for the FY 2021-22 .**

Sr.No	Name of Scheme	Proposal Received from / Name of Organization / N.G.O	Brief Details
	<b><u>CSR Applications kept pending in last year Agenda:-</u></b>		
27	CSR Activities for providing Water supply pipe line, Play ground and sports equipment, electric facilities, drinking water facilities for poor people & Fishermen at VANDI Village. (Sr no-3)	Sarpanch ,Village-VANDI , Ta- Anjar  (Recommd. By Shri VASANBHAI AHIR, MLA, Shri V L Chavda, MP)	<u>As per CSR Guideline-</u> ➤ Env Sustainability ➤ Eradicating extreme hunger and poverty  (to be Consider for health Center ,Drainage line, Water sump etc activities) <b>(Appx Cost - 51.00 Lakhs )</b>  (Land authorization of Gram Panchayat)
28	Construction of Shed, hall and Gate for the DADA Bhagwandas Charitable Trust, Adipur. (Sr no -4)	DADA BHAGWANDAS CharitableTrust, Gandhidham  (Recommd. By Shri V L Chavda, MP)	<u>As per CSR Guideline-</u> ➤ Promoting gender equality and empowering women ➤ Eradicating extreme hunger and poverty (Considered shed and hall )  Fab Shed - 30'x100' x 1250=37.00 Lakh & RCC Hall – 20'x100'x1500=30.00 Lakh  <b>(Appx Cost Rs 67.00 Lakhs)</b> Land authority belongs to Trust given by GDA and NOC given by SRC. Doc submitted.
29	10 Nos of Computers required for ShirMaheswarinagar Panchayat Girls Primary School, Gandhidham& Boys Group School, Gandhidham. (Sr no-8)	Maheswarinagar Panchayat Primary Kanya Sala, Gandhidham  (Contact no 9913903686)	<b>AppxRs 5.00 Lakhs</b>  <u>As per CSR Guideline-</u> ➤ Promotion of Education (to be consider for 20 Computers)

**List of CSR applications received from various NGOs , Organizations , Village Sharpanchsetc for the FY 2021-22 .**

Sr.No	Name of Scheme	Proposal Received from / Name of Organization / N.G.O	Brief Details
			Visited the site. Land belongs to MahewariMeghwadSamaj given by SRC for school purpose, doc are awaited.
30	Construction of Shed and Roof at JeparMatiyadev, shamsanbhumi at Kidana village & Maheswari Community Hall at JuniSundarpuri ,Gandhidham. (Sr no-10)	Shri VINOD CHAVDA, MP	<b>AppxRs 15.00 Lakhs</b> (Land authorization not mentioned)
31	Drainage, road, Dust bins, & shed for Cattle shelters at VID I Village, Ta –Anjar. (Sr no- 12)	Village- VID I, Ta: Anjar	<b>AppxRs 30.00 Lakhs</b>  <u>As per CSR Guideline-</u> ➤ Env Sustainability ➤ Eradicating extreme hunger and poverty (Consider for Garbage vehicle & Drainage Cost)
32	Education, Women empowerment and Primary health care services at Kutch area. (Sr no-13)	Light of Life Trust, <b>Mumbai.</b>	Cost not mentioned.
33	Request for Help Divyang persons to employment by providing machineries. (Sr no-14)	Kutch DivyangSangthan, Gandhidham.	Cost not mentioned
34	Construction of 2 <sup>nd</sup> Floor of Shri MaheswariMeghwadSamaj, Gandhidham. (Sr no-20)	Shri MaheswariMeghwadSamaj, Gandhidham	<b>AppxRs. 15.00 Lakhs</b> (Visited the site and Land ownership documents awaited) (Name plate of DPT fixed at the Asset)



**List of CSR applications received from various NGOs , Organizations , Village Sharpanchsetc for the FY 2021-22 .**

Sr.No	Name of Scheme	Proposal Received from / Name of Organization / N.G.O	Brief Details
35	Installation of Mini Science Center at Anjar and Gandhidham. (Sr no-21)	STEM Learning Pvt Ltd, Mumbai.	Cost not mentioned.
36	CSR work for Shri Rampar Gram Panchayat.  ➤ Wall Plastering for Cattles -7 Lakhs ➤ Shed for Cattel's-15 Lakhs  (Sr no-25)	Shri Sarpanch, Rampar Village.	<b>AppxRs 22.00 Lakhs</b>  (Land authorization of Gram Panchayat and under taking submitted by applicant)
37	CSR activities for the 45,000 Patients over the period of 3 years by "SMILE FOUNDATION", Mumbai.  1. Concept for Nutrition covering 3 years 2. Concept for Mobile Health Unit reaching beneficiaries for 3 years 3. Concept for Vocational Training with NGO (Sr no-29)	Proposal from "SMILE FOUNDATION " Mumbai.	Appx Cost- <b>Rs 539 Lakhs</b> for 3 years
38	Development of Park in Public utility plot in between Block "C" & "D" of Sapna Nagar (NU-4) , Gandhidham (Sr no -31)	Shri RAVI MAHESHWARI, DPT	Land belongs to DPT earmarked for recreational purpose.  (Total Cost – <b>Rs88.75 Lakhs</b> )
39	CSR works for NariJanshsktiVikas Foundation at Gandhidham near Shakti Nagar.  (Sr no-33)	NariJanshsktiVikas Foundation, Ahmedabad	➤ Promoting gender equality and empowering women ➤ Env Sustainability ➤ Under promotion of education  (Consider for Computers with printers, Sewing machine & RO plant <b>Cost Rs 48 Lakhs</b> )

# **ANNEXURE C**

# ENVIRONMENTAL MONITORING REPORT FOR DEENDAYAL PORT TRUST



REPORT NO. : DCPL/DPT/20-21/14  
Month : June 2021  
Issue No : 01  
Revision No : 00  
Prepared by : DETOX CORPORATION PVT. LTD., SURAT

## TABLE OF CONTENTS

<b>Sr. No.</b>	<b>Particulars</b>	<b>Page No.</b>
<b>1</b>	<b>Ambient Air Quality Monitoring.....</b>	<b>1 - 17</b>
<b>2</b>	<b>Drinking Water Quality Monitoring.....</b>	<b>18 - 28</b>
<b>3</b>	<b>Noise Monitoring.....</b>	<b>29</b>
<b>4</b>	<b>Soil Monitoring.....</b>	<b>30 - 31</b>
<b>5</b>	<b>Sewage Treatment Plant Monitoring.....</b>	<b>32 - 38</b>
<b>6</b>	<b>Marine Water Monitoring.....</b>	<b>39 - 82</b>
<b>7</b>	<b>Meteorological Observations.....</b>	<b>83</b>
<b>8</b>	<b>Conclusive Summary&amp; Remedial Measures .....</b>	<b>84-87</b>

## **Introduction**

Monitoring of various environmental aspects of the Deendayal port by M/s Detox Corporation Pvt. Ltd. has been carried out through collection of samples, analysis of the same, comparing results with respect to the national standards and any other relevant standards by GBCB/CPCB/MoEF to identify non conformities in the Environment of the Deendayal Port. The results shall address the identified impacts and suggest measures to minimize the environmental impact due to various operations at Deendayal Port.

The environmental monitoring is carried out as per the Environment Management and Monitoring Plan submitted by Detox Corporation Pvt. Ltd.

## **1. Ambient Air Quality Monitoring**

As per the Environmental Monitoring Plan of Deendayal Port Trust, Air monitoring was carried out at six identified locations at Deendayal Port and two locations at Vadinar Port.

### **1.1 Air Quality Monitoring Methodology**

Air quality is measured in all the stations, for 24 hour for Total Suspended Particulate Matter (TSPM), PM<sub>10</sub>, PM<sub>2.5</sub>, SO<sub>2</sub>, NO<sub>x</sub>, NH<sub>3</sub> & Benzene, and Grab-sampling for CO & CO<sub>2</sub> measurements. The Air samplers are operated for a period of 24 hours and after a continuous operation of 8 hours of the sampler, the reagents were replaced to obtain 3 samples per day for each parameter namely, SO<sub>2</sub>, NO<sub>x</sub>. The EPM 2000 filter paper and PTFE Membrane bound filter paper are used for a period of 24 hours to obtain one sample each of TSPM, PM<sub>10</sub> & PM<sub>2.5</sub>.

The AAQ samples are collected twice a week from all the eight locations as per the EMP.

**1.2 Results**

The ambient air quality monitoring data for six stations, viz. Marine Bhavan, Oil Jetty, Port Colony, Gopalpuri Hospital, Tuna Port and Nr. Coal Storage Area for the month of June 2021 are given in Tables 1A to 6B. The ambient air quality monitoring data for two stations at Vadinar (Nr. Admin Building & Nr. Signal Building) are given in Tables 7A to 8B.

**Location 1: Marine Bhavan (AL1)**

Table 1 : Results of Air Pollutant Concentration at Marine Bhavan										
Parameter	Date	TSPM [µg/m <sup>3</sup> ]	PM10 [µg/m <sup>3</sup> ]	PM2.5 [µg/m <sup>3</sup> ]	SO <sub>2</sub> [µg/m <sup>3</sup> ]		NO <sub>x</sub> [µg/m <sup>3</sup> ]		NH <sub>3</sub> [µg/m <sup>3</sup> ]	
Sampling Period	-	24hr	24hr	24hr	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
NAAQMS limit	-	NS	100 µg/m <sup>3</sup>	60 µg/m <sup>3</sup>		80 µg/m <sup>3</sup>		80 µg/m <sup>3</sup>		400 µg/m <sup>3</sup>
AL1 – 1	02-06-2021	447	107	53	14.07	9.23	23.50	24.14	12.51	11.83
					9.23		20.33		10.72	
					4.40		28.58		12.25	
AL1 – 2	04-06-2021	399	135	46	12.75	11.28	20.33	22.87	12.51	11.74
					11.87		24.77		12.00	
					9.23		23.50		10.72	
AL1 – 3	09-06-2021	423	204	172	13.63	14.95	24.77	19.27	7.40	7.06
					18.46		17.15		7.91	
					12.75		15.88		5.87	
AL1 – 4	11-06-2021	223	58	149	5.71	6.59	16.51	13.76	9.19	9.62
					5.71		14.61		8.93	
					8.35		10.16		10.72	
AL1 – 5	16-06-2021	476	103	203	17.14	16.56	18.42	16.51	7.15	6.81
					14.07		16.51		6.89	
					18.46		14.61		6.38	
AL1 - 6	18-06-2021	268	111	116	9.23	10.55	26.68	25.41	12.00	12.42
					9.67		27.95		12.51	
					12.75		21.60		12.76	
AL1 - 7	23-06-2021	415	179	65	5.71	6.74	26.68	26.68	6.89	7.83
					6.15		28.58		5.87	
					8.35		24.77		10.72	
AL1 – 8	25-06-2021	341	141	57	11.87	13.48	17.15	23.29	10.98	10.98
					17.14		20.33		12.00	
					11.43		32.39		9.96	
<b>Monthly Average</b>		374	130	108		11.17		21.49		9.79
<b>Standard Deviation</b>		89	46	61		3.65		4.53		2.28

NS: Not Specified

<b>Table 1B : Results of Air Pollutant Concentration at Marine Bhavan</b>					
<b>Parameter</b>	<b>Date</b>	<b>C<sub>6</sub>H<sub>6</sub> [µg/m<sup>3</sup>]</b>	<b>HC* ppm</b>	<b>CO [mg/m<sup>3</sup>]</b>	<b>CO<sub>2</sub> [ppm]</b>
<b>Sampling Period</b>		<b>8 hr</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>
<b>NAAQMS limit</b>		<b>5.0 µg/m<sup>3</sup></b>	<b>NS</b>	<b>4.0 mg/m<sup>3</sup></b>	<b>NS</b>
AL1 – 1	02/06/2021	1.2	BDL	1.46	510
AL1 – 2	04/06/2021	1.23	BDL	1.3	519
AL1 – 3	09/06/2021	1.07	BDL	1.86	495
AL1 – 4	11/06/2021	1.06	BDL	1.84	476
AL1 – 5	16/06/2021	1.06	BDL	1.75	490
AL1 - 6	18/06/2021	1.11	BDL	1.62	489
AL1 – 7	23/06/2021	1	BDL	1.8	480
AL1 – 8	25/06/2021	1.07	BDL	1.71	476
<b>Monthly Average</b>		<b>1.10</b>	<b>-</b>	<b>1.67</b>	<b>492</b>
<b>Standard Deviation</b>		<b>0.08</b>	<b>-</b>	<b>0.20</b>	<b>16</b>

\* NMHC- Non- Methane Hydrocarbons

BDL- Below Detection Limit (Detection Limit – NMHC: 0.5ppm)

NS -Not Specified

At Marine Bhavan, the overall values of TSPM, PM<sub>10</sub>, PM<sub>2.5</sub>, SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> is attributed mainly by motor vehicle emission produced from various types of automobiles (both diesel and petrol driven). Moreover, the loading and unloading of Food Grains and Timber at Jetty no. 1 and 2 also contributes to the high levels of TSPM and PM<sub>10</sub>. The mean TSPM value at Marine Bhavan was 374 µg/m<sup>3</sup>, The mean PM<sub>10</sub> values were 130.0 µg/m<sup>3</sup>, which is above the permissible limit. PM<sub>2.5</sub> values were above the permissible limit (mean = 108 µg/m<sup>3</sup>). The average values of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were within the permissible limit. The average values of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were 11.17 µg/m<sup>3</sup>, 21.49 µg/m<sup>3</sup> & 9.79 µg/m<sup>3</sup> respectively. These were within the permissible limit.

The levels of Benzene, Hydrocarbons (HC) and CO were within the permissible limit at Marine Bhavan. The mean Benzene concentration was 1.10 µg/m<sup>3</sup>, well below the permissible limit of 5.0 µg/m<sup>3</sup>. HC's were below the detectable limit and Carbon Monoxide concentration was 1.67 mg/m<sup>3</sup>, well below the permissible limit of 4.0 mg/m<sup>3</sup>.

**Environmental Monitoring Report of Deendayal Port Trust, JUNE-2021**

**Location 2: Oil Jetty (AL2)**

Table 2 : Results of Air Pollutant Concentration at Oil Jetty										
Parameters	Date	TSPM	PM10	PM2.5	SO2 [µg/m3]		NOx [µg/m3]		NH3 [µg/m3]	
		[µg/m3]	[µg/m3]	[µg/m3]	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
Sampling Period	-	24hr	24hr	24hr	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
NAAQMS limit	-	NS	100 µg/m3	60 µg/m3	-	80 µg/m3	-	80 µg/m3	-	400 µg/m3
AL2 - 1	02-06-2021	283	68	120	9.23	11.72	20.33	23.08	13.27	10.47
					13.19		25.41		10.72	
					12.75		23.50		7.40	
AL2 - 2	04-06-2021	353	108	210	9.23	11.14	18.42	17.15	10.72	10.89
					14.07		14.61		11.23	
					10.11		18.42		10.72	
AL2 - 3	09-06-2021	275	42	137	17.58	13.19	17.15	20.54	9.96	8.25
					12.74		24.14		7.91	
					9.23		20.33		6.89	
AL2 - 4	11-06-2021	257	37	145	5.27	5.42	23.50	19.27	3.32	5.87
					5.71		18.42		4.85	
					5.27		15.88		9.45	
AL2 - 5	16-06-2021	532	84	117	12.75	12.02	17.15	18.21	7.15	6.98
					9.23		20.33		7.40	
					14.07		17.15		6.38	
AL2 - 6	18-06-2021	192	111	65	11.87	7.91	26.68	28.58	10.72	11.40
					8.35		27.95		12.51	
					3.52		31.12		10.98	
AL2 - 7	23-06-2021	346	79	80	5.71	6.30	15.88	16.73	9.45	8.85
					7.91		17.15		9.70	
					5.28		17.15		7.40	
AL2 - 8	25-06-2021	256	125	31	11.87	15.09	18.42	18.84	3.83	7.83
					13.63		18.42		8.93	
					19.78		19.69		10.72	
<b>Monthly Average</b>		312	82	113		10.35		20.30		8.82
<b>Standard Deviation</b>		103	32	55		3.43		3.90		1.97

NS: Not Specified



<b>Table 2B : Results of Air Pollutant Concentration at Oil Jetty</b>					
<b>Parameter</b>	<b>Date</b>	<b>C<sub>6</sub>H<sub>6</sub> [µg/m<sup>3</sup>]</b>	<b>HC* ppm</b>	<b>CO [mg/m<sup>3</sup>]</b>	<b>CO<sub>2</sub> [ppm ]</b>
<b>Sampling Period</b>		<b>8 hr</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>
<b>NAAQMS limit</b>		<b>5.0 µg/m<sup>3</sup></b>	<b>NS</b>	<b>4.0 mg/m<sup>3</sup></b>	<b>NS</b>
AL2 -1	02/06/2021	1.11	BDL	1.78	482
AL2 -2	04/06/2021	1.06	BDL	1.77	496
AL2 -3	09/06/2021	1.22	BDL	1.8	480
AL2 -4	11/06/2021	1.05	BDL	1.75	484
AL2 – 5	16/06/2021	1.02	BDL	1.81	515
AL2 – 6	18/06/2021	1.07	BDL	1.78	496
AL2 -7	23/06/2021	1.09	BDL	1.88	491
AL2 – 8	25/06/2021	1.06	BDL	1.64	470
<b>Monthly Average</b>		<b>1.09</b>	<b>-</b>	<b>1.78</b>	<b>489</b>
<b>Standard Deviation</b>		<b>0.06</b>	<b>-</b>	<b>0.07</b>	<b>14</b>

\* NMHC- Non- Methane Hydrocarbons

BDL- Below Detection Limit (Detection Limit – NMHC : 0.5ppm)

NS- Not Specified

The overall values of TSPM, PM<sub>10</sub>, PM<sub>2.5</sub>, SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> at Oil Jetty area was mainly by motor vehicle emission produced from various types of vehicles Oil Jetty Area. The mean TSPM values at Oil Jetty were 312 µg/m<sup>3</sup> The mean PM<sub>10</sub> values were 82 µg/m<sup>3</sup>, which is below the permissible limit. PM<sub>2.5</sub> values were above the permissible limit (mean = 113 µg/m<sup>3</sup>). The average values of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were within the permissible limit; The mean concentration of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were 10.35 µg/m<sup>3</sup>, 20.30 µg/m<sup>3</sup> and 8.82 µg/m<sup>3</sup> respectively.

The levels of Benzene, Hydrocarbons (HC) and CO were within the permissible limit at Oil Jetty. The mean Benzene concentration was 1.09 µg/m<sup>3</sup>. Well below the permissible limit of 5.0 µg/m<sup>3</sup>. , HC's were below the detectable limit and Carbon Monoxide concentration was 1.78 mg/m<sup>3</sup>, well below the permissible limit of 4.0 mg/m<sup>3</sup>.

**Location 3: Kandla Colony – Estate Office (AL-3)**

**Table 3 : Results of Air Pollutant Concentration at Estate Office**

Parameters	Date	TSPM [µg/m3]	PM10 [µg/m3]	PM2.5 [µg/m3]	SO2 [µg/m3]		NOx [µg/m3]		NH3 [µg/m3]	
					8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
Sampling Period	-	24hr	24hr	24hr	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
NAAQMS limit	-	NS	100 µg/m3	60 µg/m3	-	80 µg/m3	-	80 µg/m3	-	400 µg/m3
AL3 – 1	02-06-2021	151	18	41	3.96	6.01	18.42	16.51	4.85	8.42
					5.28		17.15		6.89	
					8.79		13.97		13.53	
AL3 – 2	04-06-2021	232	63	12	10.11	11.58	17.15	18.00	6.89	9.19
					12.75		19.69		8.17	
					11.87		17.15		12.51	
AL3 – 3	09-06-2021	290	98	55	12.75	10.84	20.33	20.11	10.98	10.47
					9.67		24.77		12.25	
					10.11		15.24		8.17	
AL3 – 4	11-06-2021	235	61	128	1.76	2.34	20.33	20.54	6.89	6.13
					2.20		23.50		5.87	
					3.08		17.78		5.62	
AL3 – 5	16-06-2021	231	66	139	5.71	10.84	26.68	22.02	13.53	9.28
					12.75		20.96		7.40	
					14.07		18.42		6.89	
AL3 – 6	18-06-2021	463	76	37	10.11	11.43	20.33	22.23	7.91	8.00
					13.63		22.87		9.96	
					10.55		23.50		6.13	
AL3 – 7	23-06-2021	382	70	35	11.87	13.33	8.26	13.97	9.96	8.68
					14.07		15.24		10.72	
					14.07		18.42		5.36	
AL3 – 8	25-06-2021	148	99	42	12.75	12.16	19.69	19.69	7.15	7.91
					12.31		22.23		9.19	
					11.43		17.15		7.40	
<b>Monthly Average</b>		267	69	61		9.82		19.13		8.51
<b>Standard Deviation</b>		109	25	46		3.70		2.83		1.27

NS: Not Specified

<b>Table 3B : Results of Air Pollutant Concentration at Kandla Port Colony</b>					
<b>Parameter</b>	<b>Date</b>	<b>C<sub>6</sub>H<sub>6</sub> [µg/m<sup>3</sup>]</b>	<b>HC*</b>	<b>CO [mg/m<sup>3</sup>]</b>	<b>CO<sub>2</sub> [ppm ]</b>
<b>Sampling Period</b>		<b>8 hr</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>
<b>NAAQMS limit</b>		<b>5.0 µg/m<sup>3</sup></b>	<b>NS</b>	<b>4.0 mg/m<sup>3</sup></b>	<b>NS</b>
AL3 -1	02/06/2021	1.07	BDL	1.72	489
AL3 -2	04/06/2021	1.1	BDL	1.82	502
AL3 -3	09/06/2021	1.07	BDL	1.74	482
AL3 -4	11/06/2021	1.16	BDL	1.61	480
AL3 – 5	16/06/2021	1.17	BDL	1.69	475
AL3 – 6	18/06/2021	1.1	BDL	1.7	489
AL3 – 7	23/06/2021	1.04	BDL	1.96	486
AL3 – 8	25/06/2021	1.04	BDL	1.59	464
<b>Monthly Average</b>		<b>1.09</b>		<b>1.73</b>	<b>483</b>
<b>Standard Deviation</b>		<b>0.05</b>		<b>0.12</b>	<b>11</b>

\* NMHC- Non- Methane Hydrocarbons

BDL- Below Detection Limit ( Detection Limit – NMHC : 0.5 ppm)

NS- Not Specified

The overall values of TSPM, PM<sub>10</sub>, PM<sub>2.5</sub>, SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> at Kandla Port Colony was attributed by vehicle emission produced from trucks and heavy duty vehicles that pass through the road outside Kandla Port Colony. The mean TSPM values at Oil Jetty were 267 µg/m<sup>3</sup>, The mean PM<sub>10</sub> values were 69µg/m<sup>3</sup>, which is below the permissible limit. PM<sub>2.5</sub> values were slightly above the permissible limit (mean = 61 µg/m<sup>3</sup>). The average values of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were 9.82 µg/m<sup>3</sup>, 19.13 µg/m<sup>3</sup> and 8.51 µg/m<sup>3</sup> respectively and were all within the permissible limit.

The levels of Benzene, Hydrocarbons (HC) and CO were within the permissible limit at Kandla Port Colony. The mean Benzene concentration was 1.09 µg/m<sup>3</sup>, well below the permissible limit of 5.0 µg/m<sup>3</sup>. HC's were below the detectable limit and Carbon Monoxide concentration was 1.73 mg/m<sup>3</sup>, well below the permissible limit of 4.0 mg/m<sup>3</sup>.

**Location 4: Gopalpuri Hospital (AL-4)**

Table 4 : Results of Air Pollutant Concentration at Gopalpuri Hospital										
Parameter	Date	TSPM [µg/m <sup>3</sup> ]	PM10 [µg/m <sup>3</sup> ]	PM2.5 [µg/m <sup>3</sup> ]	SO <sub>2</sub> [µg/m <sup>3</sup> ]		NO <sub>x</sub> [µg/m <sup>3</sup> ]		NH <sub>3</sub> [µg/m <sup>3</sup> ]	
					8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
Sampling Period	-	24hr	24hr	24hr	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
NAAQMS limit	-	NS	100 µg/m <sup>3</sup>	60 µg/m <sup>3</sup>	-	80 µg/m <sup>3</sup>	-	80 µg/m <sup>3</sup>	-	400 µg/m <sup>3</sup>
AL4 -1	02-06-2021	115	40	15	3.96	2.64	13.34	15.46	5.11	5.28
					3.08		14.61		4.85	
					0.88		18.42		5.87	
AL4 -2	04-06-2021	144	43	13	8.79	7.33	14.61	17.57	6.89	7.49
					5.71		19.69		7.40	
					7.47		18.42		8.17	
AL4 -3	09-06-2021	157	49	34	2.64	3.37	14.61	20.96	6.89	7.40
					3.52		29.85		7.40	
					3.96		18.42		7.91	
AL4 -4	11-06-2021	122	29	46	9.23	7.47	5.08	6.99	3.06	4.60
					9.23		7.62		4.85	
					3.96		8.26		5.87	
AL4 -5	16-06-2021	156	35	21	3.96	3.96	10.80	12.49	10.72	11.40
					3.52		12.07		10.98	
					4.40		14.61		12.51	
AL4 -6	18-06-2021	207	72	108	9.23	8.65	13.34	18.42	7.40	7.57
					8.79		24.77		9.96	
					7.91		17.15		5.36	
AL4 -7	23-06-2021	263	36	13	0.88	3.08	11.43	13.97	10.98	10.30
					3.96		13.34		12.00	
					4.40		17.15		7.91	
AL4 -8	25-06-2021	216	111	14	3.52	4.54	12.70	11.86	7.40	7.06
					4.84		12.07		7.15	
					5.28		10.80		6.64	
<b>Monthly Average</b>		173	52	33		5.13		14.71		7.64
<b>Standard Deviation</b>		51	27	33		2.33		4.39		2.28

NS: Not Specified

<b>Table 4B : Results of Air Pollutant Concentration at Gopalpuri Hospital</b>					
<b>Parameter</b>	<b>Date</b>	<b>C<sub>6</sub>H<sub>6</sub> [µg/m<sup>3</sup>]</b>	<b>HC*</b>	<b>CO [mg/m<sup>3</sup>]</b>	<b>CO<sub>2</sub> [ppm ]</b>
<b>Sampling Period</b>		<b>8 hr</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>
<b>NAAQMS limit</b>		<b>5.0 µg/m<sup>3</sup></b>	<b>NS</b>	<b>4.0 mg/m<sup>3</sup></b>	<b>NS</b>
AL4 -1	02/06/2021	1.07	BDL	1.68	482
AL4 -2	04/06/2021	1.06	BDL	1.7	488
AL4 -3	09/06/2021	1.11	BDL	1.9	478
AL4 -4	11/06/2021	1.1	BDL	1.54	470
AL4 -5	16/06/2021	1.21	BDL	1.58	455
AL4 -6	18/06/2021	1.2	BDL	1.78	460
AL4 -7	23/06/2021	1.19	BDL	1.94	481
AL4 -8	25/06/2021	1.13	BDL	1.91	475
<b>Monthly Average</b>		<b>1.13</b>		<b>1.75</b>	<b>474</b>
<b>Standard Deviation</b>		<b>0.06</b>		<b>0.15</b>	<b>11</b>

\* NMHC- Non- Methane Hydrocarbons

BDL- Below Detection Limit (Detection Limit – NMHC: 0.5 ppm)

NS-Not Specified

The overall values of TSPM, PM<sub>10</sub>, PM<sub>2.5</sub>, SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> at Gopalpuri Hospital was attributed by vehicle emission produced from light motor vehicles of the colony residents. The mean TSPM values at Oil Jetty were 173 µg/m<sup>3</sup>, The mean PM<sub>10</sub> values were 52 µg/m<sup>3</sup>, which is below the permissible limit. PM<sub>2.5</sub> values were also within the permissible limit (mean= 33 µg/m<sup>3</sup>). The average values of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were 5.13 µg/m<sup>3</sup>, 14.71 µg/m<sup>3</sup> and 7.64 µg/m<sup>3</sup> respectively and were all within the permissible limit.

The levels of Benzene, Hydrocarbons (HC) and CO were within the permissible limit at Gopalpuri Hospital. The mean Benzene concentration was 1.13 µg/m<sup>3</sup>, well below the permissible limit of 5.0 µg/m<sup>3</sup>. HC's were below the detectable limit and Carbon Monoxide concentration was 1.75 mg/m<sup>3</sup>, well below the permissible limit of 4.0 mg/m<sup>3</sup>.

**Location 5: Coal Storage Area (AL-5)**

Table 5 : Results of Air Pollutant Concentration at Coal Storage Area										
Parameters	Date	TSPM [µg/m <sup>3</sup> ]	PM10 [µg/m <sup>3</sup> ]	PM2.5 [µg/m <sup>3</sup> ]	SO <sub>2</sub> [µg/m <sup>3</sup> ]		NO <sub>x</sub> [µg/m <sup>3</sup> ]		NH <sub>3</sub> [µg/m <sup>3</sup> ]	
Sampling Period	-	24hr	24hr	24hr	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
NAAQMS limit	-	NS	100 µg/m <sup>3</sup>	60 µg/m <sup>3</sup>	-	80 µg/m <sup>3</sup>	-	80 µg/m <sup>3</sup>	-	400 µg/m <sup>3</sup>
AL5 – 1	02-06-2021	829	78	60	9.23	9.23	26.04	26.47	13.27	13.96
					12.75		28.58		15.32	
					5.71		24.77		13.27	
AL5 – 2	04-06-2021	332	104	97	5.71	7.77	20.33	21.60	12.51	12.00
					8.79		24.77		12.51	
					8.79		19.69		10.98	
AL5 – 3	09-06-2021	289	185	154	10.11	13.48	18.42	18.84	10.72	11.83
					12.75		17.78		12.51	
					17.58		20.33		12.25	
AL5 – 4	11-06-2021	280	70	162	9.23	13.19	12.07	13.55	10.98	10.64
					13.19		13.34		10.72	
					17.14		15.24		10.21	
AL5 – 5	16-06-2021	944	148	150	3.96	10.99	14.61	18.84	2.30	5.45
					19.78		10.80		6.89	
					9.23		31.12		7.15	
AL5 – 6	18-06-2021	603	145	234	10.11	9.23	26.68	24.56	13.53	13.96
					7.47		22.87		13.27	
					10.11		24.14		15.06	
AL5 – 7	23-06-2021	766	181	152	11.87	12.75	12.70	21.17	12.51	10.47
					14.07		17.15		10.72	
					12.31		33.66		8.17	
AL5 – 8	25-06-2021	728	208	94	13.63	11.72	18.42	15.03	12.51	9.53
					10.55		17.78		9.19	
					10.99		8.89		6.89	
<b>Monthly Average</b>		596	140	138		11.04		20.01		10.98
<b>Standard Deviation</b>		263	51	54		2.11		4.40		2.74

NS: Not Specified

<b>Table 5B : Results of Air Pollutant Concentration at Coal Storage Area</b>					
<b>Parameter</b>	<b>Date</b>	<b>C<sub>6</sub>H<sub>6</sub> [µg/m<sup>3</sup>]</b>	<b>HC*</b>	<b>CO [mg/m<sup>3</sup>]</b>	<b>CO<sub>2</sub> [ppm ]</b>
<b>Sampling Period</b>		<b>8 hr</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>
<b>NAAQMS limit</b>		<b>5.0 µg/m<sup>3</sup></b>	<b>NS</b>	<b>4.0 mg/m<sup>3</sup></b>	<b>NS</b>
AL5 – 1	02/06/2021	1.08	BDL	1.78	482
AL5 – 2	04/06/2021	1.1	BDL	1.68	490
AL5 – 3	09/06/2021	1.24	BDL	1.64	462
AL5 – 4	11/06/2021	1.28	BDL	1.66	464
AL5 – 5	16/06/2021	1.31	BDL	1.66	460
AL5 – 6	18/06/2021	1.2	BDL	1.7	490
AL5 – 7	23/06/2021	1.33	BDL	1.74	464
AL5 – 8	25/06/2021	1.11	BDL	1.91	484
<b>Monthly Average</b>		<b>1.21</b>	<b>-</b>	<b>1.72</b>	<b>475</b>
<b>Standard Deviation</b>		<b>0.10</b>	<b>-</b>	<b>0.09</b>	<b>13</b>

\* NMHC- Non- Methane Hydrocarbons

BDL- Below Detection Limit (Detection Limit – NMHC: 0.5 ppm)

NS-Not Specified

The overall values of TSPM, PM<sub>10</sub>, PM<sub>2.5</sub>, SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> at Coal Storage Area was comparatively highest among all the locations of Air Quality monitoring in Kandla Port. High values of TSPM, PM<sub>10</sub>, PM<sub>2.5</sub>, SO<sub>2</sub>, NO<sub>x</sub> at this location was due to lifting of coal with grab and other coal handling processes near Berth no. 6 & 7. Moreover, the traffic was also heavy around this place for transport of coal thus emissions produced from heavy vehicles. The mean TSPM values at Coal storage were 596 µg/m<sup>3</sup>. The mean PM<sub>10</sub> values were 140 µg/m<sup>3</sup>, which is well above the permissible limit. PM<sub>2.5</sub> values were above the permissible limit (mean = 138 µg/m<sup>3</sup>). The average values of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were 11.04 µg/m<sup>3</sup>, 20.01 µg/m<sup>3</sup> and 10.98 µg/m<sup>3</sup> respectively and were all within the permissible limit.

The levels of Benzene, Hydrocarbons (HC) and CO were within the permissible limit at Coal Storage Area. The mean Benzene concentration was 1.21 µg/m<sup>3</sup>, well below the permissible limit of 5.0 µg/m<sup>3</sup>. HC's were below the detectable limit and Carbon Monoxide concentration was 1.72 mg/m<sup>3</sup>, well below the permissible limit of 4.0 mg/m<sup>3</sup>.

**Location 6: Tuna Port (AL-6)**

**Table 6 : Results of Air Pollutant Concentration at Tuna Port**

Parameters	Date	TSPM [µg/m <sup>3</sup> ]	PM10 [µg/m <sup>3</sup> ]	PM2.5 [µg/m <sup>3</sup> ]	SO <sub>2</sub> [µg/m <sup>3</sup> ]		NO <sub>x</sub> [µg/m <sup>3</sup> ]		NH <sub>3</sub> [µg/m <sup>3</sup> ]	
					8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
Sampling Period	-	24hr	24hr	24hr	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
NAAQMS limit	-	NS	100 µg/m <sup>3</sup>	60 µg/m <sup>3</sup>	-	80 µg/m <sup>3</sup>	-	80 µg/m <sup>3</sup>	-	400 µg/m <sup>3</sup>
AL6 -1	02-06-2021	72	72	37	0.44	1.76	17.15	16.73	4.60	6.72
					1.32		13.34		8.17	
					3.52		19.69		7.40	
AL6 - 2	04-06-2021	80	42	39	4.84	6.01	24.77	18.84	7.40	8.76
					3.96		13.34		9.45	
					9.23		18.42		9.45	
AL6 - 3	09-06-2021	122	38	31	9.23	12.45	17.78	16.94	7.40	8.76
					18.90		14.61		8.17	
					9.23		18.42		10.72	
AL6 - 4	11-06-2021	72	25	44	3.52	2.93	10.80	14.61	3.32	4.25
					1.32		14.61		4.85	
					3.96		18.42		4.60	
AL6 - 5	16-06-2021	86	78	12	8.79	11.72	15.24	19.69	9.45	9.19
					13.63		20.33		9.96	
					12.75		23.50		8.17	
AL6 - 6	18-06-2021	187	32	66	11.87	6.74	19.69	18.00	5.62	6.30
					3.96		17.78		6.13	
					4.40		16.51		7.15	
AL6 - 7	23-06-2021	261	73	8	11.87	12.75	20.33	18.42	8.17	9.87
					12.75		26.68		10.72	
					13.63		8.26		10.72	
AL6 - 8	25-06-2021	123	109	26	8.35	10.26	11.43	10.16	9.96	8.25
					9.23		6.99		9.45	
					13.19		12.07		5.36	
<b>Monthly Average</b>		125	59	33		8.08		16.67		7.76
<b>Standard Deviation</b>		67	29	18		4.33		3.06		1.86

NS: Not Specified



<b>Table 6B : Results of Air Pollutant Concentration at Tuna Port</b>					
<b>Parameter</b>	<b>Date</b>	<b>C<sub>6</sub>H<sub>6</sub> [µg/m<sup>3</sup>]</b>	<b>HC*</b>	<b>CO [mg/m<sup>3</sup>]</b>	<b>CO<sub>2</sub> [ppm ]</b>
<b>Sampling Period</b>		<b>8 hr</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>
<b>NAAQMS limit</b>		<b>5.0 µg/m<sup>3</sup></b>	<b>NS</b>	<b>4.0 mg/m<sup>3</sup></b>	<b>NS</b>
AL6 -1	02/06/2021	1.2	BDL	1.72	489
AL6 - 2	04/06/2021	1.12	BDL	1.65	479
AL6 - 3	09/06/2021	1.03	BDL	1.71	466
AL6 - 4	11/06/2021	1.14	BDL	1.74	469
AL6 - 5	16/06/2021	1.05	BDL	1.71	490
AL6 - 6	18/06/2021	1.12	BDL	1.72	472
AL6 - 7	23/06/2021	1.29	BDL	1.7	470
AL6 - 8	25/06/2021	1.27	BDL	1.88	480
<b>Monthly Average</b>		<b>1.15</b>	<b>-</b>	<b>1.73</b>	<b>477</b>
<b>Standard Deviation</b>		<b>0.09</b>	<b>-</b>	<b>0.07</b>	<b>9</b>

\* NMHC- Non- Methane Hydrocarbons

BDL- Below Detection Limit (Detection Limit – NMHC : 0.5 ppm)

NS- Not Specified

The mean TSPM values at Tuna Port were 125 µg/m<sup>3</sup>, The mean PM<sub>10</sub> values were 59 µg/m<sup>3</sup>, which is below the permissible limit. PM<sub>2.5</sub> values were within the permissible limit (mean = 33 µg/m<sup>3</sup> µg/m<sup>3</sup>). The average values of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were 8.08 µg/m<sup>3</sup>, 16.67 µg/m<sup>3</sup> and 7.76 µg/m<sup>3</sup> respectively and were all within the permissible limit.

The levels of Benzene, Hydrocarbons (HC) and CO were within the permissible limit at Tuna Port. The mean Benzene concentration was 1.15 µg/m<sup>3</sup>, well below the permissible limit of 5.0 µg/m<sup>3</sup>. HC's were below the detectable limit and Carbon Monoxide concentration was 1.73 mg/m<sup>3</sup>, well below the permissible limit of 4.0 mg/m<sup>3</sup>.

**Environmental Monitoring Report of Deendayal Port Trust, JUNE-2021**

**Location 7: Signal Building (Vadinar) (AL-7)**

**Table 7 : Results of Air Pollutant Concentration at Signal Building**

Parameters	Date	TSPM [µg/m <sup>3</sup> ]	PM10 [µg/m <sup>3</sup> ]	PM2.5 [µg/m <sup>3</sup> ]	SO <sub>2</sub> [µg/m <sup>3</sup> ]		NO <sub>x</sub> [µg/m <sup>3</sup> ]		NH <sub>3</sub> [µg/m <sup>3</sup> ]	
					8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
Sampling Period	-	24hr	24hr	24hr	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
NAAQMS limit	-	NS	100 µg/m <sup>3</sup>	60 µg/m <sup>3</sup>	-	80 µg/m <sup>3</sup>	-	80 µg/m <sup>3</sup>	-	400 µg/m <sup>3</sup>
AL7 -1	02-06-2021	56	27	27	3.517	4.396	10.98	9.96	17.15	17.57
					5.715		10.47		19.69	
					3.956		8.42		15.88	
AL7 -2	04-06-2021	59	16	29	3.517	2.051	7.91	10.13	10.80	12.70
					1.319		10.98		12.07	
					1.319		11.49		15.24	
AL7 -3	09-06-2021	51	41	26	2.198	1.905	6.13	6.30	10.80	14.19
					1.319		3.32		12.70	
					2.198		9.45		19.05	
AL7 -4	11-06-2021	49	38	63	3.956	3.077	7.15	9.53	9.53	10.80
					3.077		9.96		10.80	
					2.198		11.49		12.07	
AL7 -5	16-06-2021	62	51	24	1.758	3.810	10.98	9.10	10.80	11.43
					2.198		11.49		11.43	
					7.473		4.85		12.07	
AL7 -6	18-06-2021	68	29	58	11.869	6.447	6.89	13.96	15.88	16.94
					3.956		21.44		17.78	
					3.517		13.53		17.15	
AL7 -7	23-06-2021	63	41	24	10.110	10.843	3.318611	9.28	8.892276	10.37
					10.990		11.4875		10.79776	
					11.429		13.01917		11.43293	
AL7 -8	25-06-2021	66	23	55	0.879	1.612	6.8925	7.23	24.77134	22.44
					1.758		7.913611		23.50102	
					2.198		6.8925		19.05488	
<b>Monthly Average</b>		<b>59</b>	<b>33</b>	<b>38</b>		<b>4.268</b>		<b>9</b>		<b>15</b>
<b>Standard Deviation</b>		<b>7</b>	<b>11</b>	<b>17</b>		<b>3.098</b>		<b>2</b>		<b>4</b>

NS: Not Specified

<b>Table 7B : Results of Air Pollutant Concentration at Signal Building</b>					
<b>Parameter</b>	<b>Date</b>	<b>C<sub>6</sub>H<sub>6</sub> [µg/m<sup>3</sup>]</b>	<b>HC*</b>	<b>CO [mg/m<sup>3</sup>]</b>	<b>CO<sub>2</sub> [ppm ]</b>
<b>Sampling Period</b>		<b>8 hr</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>
<b>NAAQMS limit</b>		<b>5.0 µg/m<sup>3</sup></b>	<b>NS</b>	<b>4.0 mg/m<sup>3</sup></b>	<b>NS</b>
AL7 -1	02/06/2021	1.06	BDL	1.68	472
AL7 - 2	04/06/2021	1.11	BDL	1.72	468
AL7 - 3	09/06/2021	1.23	BDL	1.55	482
AL7 - 4	11/06/2021	1.07	BDL	1.69	492
AL7 - 5	16/06/2021	1.23	BDL	1.78	466
AL7 - 6	18/06/2021	1.21	BDL	1.92	478
AL7 - 7	23/06/2021	1.18	BDL	1.88	485
AL7 - 8	25/06/2021	1.14	BDL	1.68	488
<b>Monthly Average</b>		<b>1.15</b>	<b>-</b>	<b>1.74</b>	<b>479</b>
<b>Standard Deviation</b>		<b>0.07</b>	<b>-</b>	<b>0.12</b>	<b>10</b>

\* NMHC- Non- Methane Hydrocarbons

BDL- Below Detection Limit (Detection Limit – NMHC : 0.5 ppm)

NS\_ Not Specified

The mean TSPM values at Vadinar Port were 59 µg/m<sup>3</sup>. The mean PM<sub>10</sub> values were 33 µg/m<sup>3</sup>, which is below the permissible limit. PM<sub>2.5</sub> values were also within the permissible limit (mean = 38 µg/m<sup>3</sup> µg/m<sup>3</sup>). The average values of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were 4.26 µg/m<sup>3</sup>, 9 µg/m<sup>3</sup> and 15 µg/m<sup>3</sup> respectively and were all within the permissible limit.

The levels of Benzene, Hydrocarbons (HC) and CO were within the permissible limit at Vadinar Port. The mean Benzene concentration was 1.15 µg/m<sup>3</sup>, well below the permissible limit of 5.0 µg/m<sup>3</sup>. HC's were below the detectable limit and Carbon Monoxide concentration was 1.74 mg/m<sup>3</sup>, well below the permissible limit of 4.0 mg/m<sup>3</sup>.

**Location 8: Admin Building (Vadinar) (AL-8)**

**Table 8 : Results of Air Pollutant Concentration at Admin Building**

Parameters	Date	TSPM [µg/m3]	PM10 [µg/m3]	PM2.5 [µg/m3]	SO2 [µg/m3]		NOx [µg/m3]		NH3 [µg/m3]	
					8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
Sampling Period	-	24hr	24hr	24hr	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
NAAQMS limit	-	NS	100 µg/m3	60 µg/m3	-	80 µg/m3	-	80 µg/m3	-	400 µg/m3
AL8 -1	02-06-2021	54	16	27	0.879	1.905	8.257	8.469	6.893	6.637
					1.758		8.257		7.148	
					3.077		8.892		5.871	
AL8 -2	04-06-2021	58	19	23	0.879	1.172	19.690	16.514	5.361	4.850
					1.758		17.149		4.850	
					0.879		12.703		4.340	
AL8 -3	09-06-2021	70	63	23	2.198	1.612	14.609	12.915	1.276	1.106
					1.319		8.257		1.021	
					1.319		15.879		1.021	
AL8 -4	11-06-2021	53	47	28	1.758	2.198	17.149	17.996	2.298	4.340
					2.198		13.338		6.382	
					2.638		23.501		4.340	
AL8 -5	16-06-2021	57	12	14	2.198	2.638	17.149	12.915	3.319	3.234
					2.638		12.703		3.063	
					3.077		8.892		3.319	
AL8 -6	18-06-2021	59	28	19	1.319	1.758	9.527	9.527	4.850	4.340
					1.758		8.257		4.340	
					2.198		10.798		3.829	
AL8 -5	23-06-2021	56	29	15	0.879	1.758	6.352	8.892	3.829	5.191
					1.319		9.527		4.340	
					3.077		10.798		7.403	
AL8-6	25-06-2021	73	51	28	0.440	0.733	13.974	15.667	7.914	8.084
					0.879		15.244		10.466	
					0.879		17.785		5.871	
<b>Monthly Average</b>		60	33	22		1.7217		12.862		4.72
<b>Standard Deviation</b>		7	18	6		0.5848		3.660		2.10

NS: Not Specified

<b>Table 8B : Results of Air Pollutant Concentration at Admin Building</b>					
<b>Parameter</b>	<b>Date</b>	<b>C<sub>6</sub>H<sub>6</sub> [µg/m<sup>3</sup>]</b>	<b>HC*</b>	<b>CO [mg/m<sup>3</sup>]</b>	<b>CO<sub>2</sub> [ppm ]</b>
<b>Sampling Period</b>		<b>8 hr</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>
<b>NAAQMS limit</b>		<b>5.0 µg/m<sup>3</sup></b>	<b>NS</b>	<b>4.0 mg/m<sup>3</sup></b>	<b>NS</b>
AL8 -1	02/06/2021	1.21	BDL	1.78	496
AL8-2	04/06/2021	1.18	BDL	1.92	477
AL8 -3	09/06/2021	1.23	BDL	1.68	468
AL8-4	11/06/2021	1.16	BDL	1.77	484
AL8 -5	16/06/2021	1.25	BDL	1.84	477
AL8-6	18/06/2021	1.22	BDL	1.68	485
AL8-7	23/06/2021	1.16	BDL	1.62	476
AL8-8	25/06/2021	1.12	BDL	1.77	466
Monthly Average		<b>1.19</b>	-	<b>1.76</b>	<b>479</b>
Standard Deviation		<b>0.04</b>	-	<b>0.10</b>	<b>10</b>

\* NMHC- Non- Methane Hydrocarbons

BDL- Below Detection Limit (Detection Limit – NMHC: 0.5 ppm)

NS-Not Specified

The overall values of TSPM, PM<sub>10</sub>, PM<sub>2.5</sub>, SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> at Admin Building Vadinar was comparatively low among all the locations of Air Quality monitoring in Kandla Port and Vadinar Port. The mean TSPM values at Vadinar Port were 60 µg/m<sup>3</sup>. The mean PM<sub>10</sub> values were 33 µg/m<sup>3</sup>, which is below the permissible limit. PM<sub>2.5</sub> values were also within the permissible limit (mean = 22.0 µg/m<sup>3</sup>). The average values of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were 1.72 µg/m<sup>3</sup>, 12.86 µg/m<sup>3</sup> and 4.72 µg/m<sup>3</sup> respectively and were all within the permissible limit.

The levels of Benzene, Hydrocarbons (HC) and CO were within the permissible limit at Admin Building, Vadinar Port. The mean Benzene concentration was 1.19 µg/m<sup>3</sup>, well below the permissible limit of 5.0 µg/m<sup>3</sup>. HC's were below the detectable limit and Carbon Monoxide concentration was 1.76 mg/m<sup>3</sup>, well below the permissible limit of 4.0 mg/m<sup>3</sup>.

#### **1.4 Observations and Conclusion**

During the monitoring period, the overall Ambient Air Quality of the port area was found to be well within the desired levels for various pollutants. However, Near Coal storage area, Marine Bhavan and Oil Jetty area, PM<sub>10</sub> values was above the permissible standards. All other pollutants were recorded well below the prescribed limits.

## 2. Drinking Water Quality Monitoring

Drinking Water Quality Monitoring was carried out at twenty stations at Kandla, Vadinar & Township Area of Deendayal Port.

### 2.1 Drinking Water Monitoring Methodology

Drinking water samples were collected from 20 locations as prescribed in the tender document. Samples for physico-chemical analysis were collected in 1 liter carboys and samples for microbiological parameters were collected in sterilized bottles. These samples were then analyzed in laboratory for various drinking water parameters at Kandla Lab/Surat.

The Sampling and Analysis was done as per standard methods - CPCB/GPCB Guidelines and Standard Methods -APHA. The water samples were analyzed for various parameters, viz. Color , Odor, Turbidity , Conductivity , pH , Chlorides , TDS, Total Hardness, Iron , Sulphate , Salinity , DO, BOD, Na, K, Ca, Mg, F, NO<sub>3</sub>, NO<sub>2</sub>, Mn, Cr-6, Cu, Cd, As, Hg, Pb, Zn, Bacterial Count (cfu) .

### 2.2 Results

The Drinking Water Quality monitoring data for 20 stations are given in below from table No. 9 to Table No. 15

**Table 9: Drinking Water Quality Monitoring Parameters for Nirman Building 1, P & C building & Main Gate (North) at Kandla**

Sr. No.	Parameter	Unit	Nirman Building 1	P & C building	Main Gate North	Acceptable Limits as per IS 10500 : 2012	Permissible Limits as per IS 10500 : 2012
1	pH	pH Unit	7.1	7.3	7.8	6.5 to 8.5	6.5 to 8.5
2	Total Dissolved Solids	mg/l	570	590	610	500	2000
3	Turbidity	NTU	1	1	0	1.0	5.0
4	Odor	-	Odourless	Odourless	Odourless	Agreeable	Agreeable
5	Color	Hazen Units	Colourless	Colourless	Colourless	5.0	15.0
6	Conductivity	µs/cm	1243.0	1150.0	1190.0	NS*	NS*
7	Biochemical Oxygen Demand	mg/l	<2.0	<2.0	<2.0	NS*	NS*
8	Chloride as Cl	mg/l	513.19	457.02	561.25	250.0	1000.0
9	Ca as Ca	mg/l	48.10	44.09	48.10	75.0	200.0
10	Mg as Mg	mg/l	82.62	87.48	89.91	30.0	100.0
11	Total Hardness	mg/l	460.0	470.0	490.0	200.0	600.0
12	Iron as Fe	mg/l	<0.01	<0.01	<0.01	0.3	No Relaxation
13	Fluorides as F	mg/l	0.30	0.47	0.24	1.0	1.5
14	Sulphate as SO4	mg/l	232.8	180	258	200.0	400
15	Nitrite as NO2	mg/l	0.04	0.05	0.05	NS*	NS*
16	Nitrate as NO3	mg/l	0.77	9.15	28.16	45.0	No Relaxation
17	Salinity	%	0.93	0.83	1.01	NS*	NS*
18	Sodium as Na	mg/l	322.0	315.0	342.0	NS*	NS*
19	Potassium as K	mg/l	3.44	3.21	4.08	NS*	NS*
20	Manganese	mg/l	<0.04	<0.04	<0.04	0.1	0.3
21	Hexavalent Chromium	mg/l	<0.03	<0.03	<0.03	NS*	NS*
22	Copper	mg/l	<0.05	<0.05	<0.05	0.05	1.5
23	Cadmium	mg/l	<0.002	<0.002	<0.002	0.003	0.003
24	Arsenic	mg/l	<0.01	<0.01	<0.01	0.01	0.05
25	Mercury	mg/l	<0.001	<0.001	<0.001	0.001	0.001
26	Lead	mg/l	<0.01	<0.01	<0.01	0.01	0.01
27	Zinc	mg/l	<0.1	<0.1	<0.1	5.0	15.0
28	Bacterial Count	CFU/100ml	Absent	Absent	Absent	Absent	Absent

\*NS: Not Specified

**Table 10: Drinking Water Quality Monitoring Parameters for Canteen, West Gate – I & Wharf Area at Kandla**

Sr. No.	Parameter	Unit	Canteen	West Gate - I	Wharf Area	Acceptable Limits as per IS 10500 : 2012	Permissible Limits as per IS 10500 : 2012
1	pH	pH Unit	7.0	7.6	7.4	6.5 to 8.5	6.5 to 8.5
2	Total Dissolved Solids	mg/l	910.0	960.0	870.0	500	2000
3	Turbidity	NTU	0	1	0	1.0	5.0
4	Odor	-	Odourless	Odourless	Odourless	Agreeable	Agreeable
5	Color	Hazen Units	Colourless	Colourless	Colourless	5.0	15.0
6	Conductivity	µs/cm	1703.0	1753.0	1630.0	NS*	NS*
7	Biochemical Oxygen Demand	mg/l	<2.0	<2.0	<2.0	NS*	NS*
8	Chloride as Cl	mg/l	613.86	620.88	658.46	250.0	1000.0
9	Ca as Ca	mg/l	52.10	52.10	44.09	75.0	200.0
10	Mg as Mg	mg/l	72.90	80.19	77.76	30.0	100.0
11	Total Hardness	mg/l	430.0	460.0	430.0	200.0	600.0
12	Iron as Fe	mg/l	<0.01	<0.01	<0.01	0.3	No Relaxation
13	Fluorides as F	mg/l	0.47	0.32	0.42	1.0	1.5
14	Sulphate as SO <sub>4</sub>	mg/l	156.0	300.0	366.0	200.0	400
15	Nitrite as NO <sub>2</sub>	mg/l	0.03	<0.01	0.03	NS*	NS*
16	Nitrate as NO <sub>3</sub>	mg/l	24.64	10.56	12.67	45.0	No Relaxation
17	Salinity	%	1.11	1.12	1.19	NS*	NS*
18	Sodium as Na	mg/l	333.0	362.0	412.0	NS*	NS*
19	Potassium as K	mg/l	3.78	3.99	4.11	NS*	NS*
20	Manganese	mg/l	<0.04	<0.04	<0.04	0.1	0.3
21	Hexavalent Chromium	mg/l	<0.03	<0.03	<0.03	NS*	NS*
22	Copper	mg/l	<0.05	<0.05	<0.05	0.05	1.5
23	Cadmium	mg/l	<0.002	<0.002	<0.002	0.003	0.003
24	Arsenic	mg/l	<0.01	<0.01	<0.01	0.01	0.05
25	Mercury	mg/l	<0.001	<0.001	<0.001	0.001	0.001
26	Lead	mg/l	<0.01	<0.01	<0.01	0.01	0.01
27	Zinc	mg/l	<0.1	<0.1	<0.1	5.0	15.0
28	Bacterial Count	CFU/100ml	Absent	Absent	Absent	Absent	Absent

\*NS: Not Specified



**Table 11: Drinking Water Quality Monitoring Parameters for Sewa sadan – 3, Workshop I & Custom Building at Kandla**

Sr. No.	Parameter	Unit	SewaSadan – 3	Workshop	Custom Building	Acceptable Limits as per IS 10500 : 2012	Permissible Limits as per IS 10500 : 2012
1	pH	pH Unit	7.3	7.4	7.8	6.5 to 8.5	6.5 to 8.5
2	Total Dissolved Solids	mg/l	1090.0	830.0	935.0	500	2000
3	Turbidity	NTU	1	0	1	1.0	5.0
4	Odor	-	Odourless	Odourless	Odourless	Agreeable	Agreeable
5	Color	Hazen Units	Colourless	Colourless	Colourless	5.0	15.0
6	Conductivity	µs/cm	1910.0	1600.0	1820.0	NS*	NS*
7	Biochemical Oxygen Demand	mg/l	<2.0	<2.0	<2.0	NS*	NS*
8	Chloride	mg/l	743.65	571.77	550.72	250.0	1000.0
9	Ca as Ca	mg/l	52.10	56.11	48.10	75.0	200.0
10	Mg as Mg	mg/l	82.62	85.05	80.19	30.0	100.0
11	Total Hardness	mg/l	470.0	490.0	450.0	200.0	600.0
12	Iron as Fe	mg/l	<0.01	<0.01	<0.01	0.3	No Relaxation
13	Fluorides	mg/l	0.32	0.93	0.30	1.0	1.5
14	Sulphate	mg/l	190.8	172.8	195.6	200.0	400
15	Nitrite	mg/l	0.01	0.03	0.05	NS*	NS*
16	Nitrate	mg/l	13.37	6.33	12.67	45.0	No Relaxation
17	Salinity	%	1.34	1.03	0.99	NS*	NS*
18	Sodium as Na	mg/l	333.0	342.	392.0	NS*	NS*
19	Potassium as K	mg/l	3.88	3.71	4.12	NS*	NS*
20	Manganese	mg/l	<0.04	<0.04	<0.04	0.1	0.3
21	Hexavalent Chromium	mg/l	<0.03	<0.03	<0.03	NS*	NS*
22	Copper	mg/l	<0.05	<0.05	<0.05	0.05	1.5
23	Cadmium	mg/l	<0.002	<0.002	<0.002	0.003	0.003
24	Arsenic	mg/l	<0.01	<0.01	<0.01	0.01	0.05
25	Mercury	mg/l	<0.001	<0.001	<0.001	0.001	0.001
26	Lead	mg/l	<0.01	<0.01	<0.01	0.01	0.01
27	Zinc	mg/l	<0.1	<0.1	<0.1	5.0	15.0
28	Bacterial Count	CFU/100ml	Absent	Absent	Absent	Absent	Absent

\*NS: Not Specified

**Table 12: Drinking Water Quality Monitoring Parameters for Port Colony Kandla, Hospital Kandla & A.O. Building at Gandhidham**

Sr. No.	Parameter	Unit	Port Colony Kandla	Hospital Kandla	A.O. Building	Acceptable Limits as per IS 10500 : 2012	Permissible Limits as per IS 10500 : 2012
1	pH	pH Unit	7.2	7	7.2	6.5 to 8.5	6.5 to 8.5
2	Total Dissolved Solids	mg/l	1200.0	1400.0	1090.0	500	2000
3	Turbidity	NTU	0	0	1	1.0	5.0
4	Odor	-	Odourless	Odourless	Odourless	Agreeable	Agreeable
5	Color	Hazen Units	Colourless	Colourless	Colourless	5.0	15.0
6	Conductivity	µs/cm	2512.0	2830.0	1920.0	NS*	NS*
7	Biochemical Oxygen Demand	mg/l	<2.0	<2.0	<2.0	NS*	NS*
8	Chloride	mg/l	763.70	794.77	838.86	250.0	1000.0
9	Ca as Ca	mg/l	56.11	48.10	60.12	75.0	200.0
10	Mg as Mg	mg/l	77.76	80.19	77.76	30.0	100.0
11	Total Hardness	mg/l	460.0	450.0	470.0	200.0	600.0
12	Iron as Fe	mg/l	<0.01	<0.01	<0.01	0.3	No Relaxation
13	Fluorides	mg/l	0.77	0.12	0.21	1.0	1.5
14	Sulphate	mg/l	202.8	261.6	372	200.0	400
15	Nitrite	mg/l	0.05	0.05	0.06	NS*	NS*
16	Nitrate	mg/l	5.63	12.67	16.89	45.0	No Relaxation
17	Salinity	%	1.38	1.44	1.52	NS*	NS*
18	Sodium as Na	mg/l	322.0	373.0	432.0	NS*	NS*
19	Potassium as K	mg/l	3.61	3.81	4.45	NS*	NS*
20	Manganese	mg/l	<0.04	<0.04	<0.04	0.1	0.3
21	Hexavalent Chromium	mg/l	<0.03	<0.03	<0.03	NS*	NS*
22	Copper	mg/l	<0.05	<0.05	<0.05	0.05	1.5
23	Cadmium	mg/l	<0.002	<0.002	<0.002	0.003	0.003
24	Arsenic	mg/l	<0.01	<0.01	<0.01	0.01	0.05
25	Mercury	mg/l	<0.001	<0.001	<0.001	0.001	0.001
26	Lead	mg/l	<0.01	<0.01	<0.01	0.01	0.01
27	Zinc	mg/l	<0.1	<0.1	<0.1	5.0	15.0
28	Bacterial Count	CFU/100ml	Absent	Absent	Absent	Absent	Absent

\*NS: Not Specified

**Table 13: Drinking Water Quality Monitoring Parameters for School Gopalpuri, Guest House & E - Type Quarter at Gopalpuri, Gandhidham**

Sr. No.	Parameter	Unit	School Gopalpuri	Guest House	E - Type Quarter	Acceptable Limits as per IS 10500 : 2012	Permissible Limits as per IS 10500 : 2012
1	pH	pH Unit	7.1	7.3	7.4	6.5 to 8.5	6.5 to 8.5
2	Total Dissolved Solids	mg/l	970.0	1010.0	1135.0	500	2000
3	Turbidity	NTU	0	1	0	1.0	5.0
4	Odor	-	Odourless	Odourless	Odourless	Agreeable	Agreeable
5	Color	Hazen Units	Colourless	Colourless	Colourless	5.0	15.0
6	Conductivity	µs/cm	1850.0	1920.0	2210.0	NS*	NS*
7	Biochemical Oxygen Demand	mg/l	<2.0	<2.0	<2.0	NS*	NS*
8	Chloride	mg/l	708.58	615.87	845.88	250.0	1000.0
9	Ca as Ca	mg/l	56.11	52.10	56.11	75.0	200.0
10	Mg as Mg	mg/l	82.62	85.05	85.05	30.0	100.0
11	Total Hardness	mg/l	480.0	480.0	490.0	200.0	600.0
12	Iron as Fe	mg/l	<0.01	<0.01	<0.01	0.3	No Relaxation
13	Fluorides	mg/l	1.37	0.89	0.33	1.0	1.5
14	Sulphate	mg/l	369.6	384	376.8	200.0	400
15	Nitrite	mg/l	0.04	0.06	0.04	NS*	NS*
16	Nitrate	mg/l	7.74	6.33	12.67	45.0	No Relaxation
17	Salinity	%	1.28	1.11	1.53	NS*	NS*
18	Sodium as Na	mg/l	392.0	320.0	332.0	NS*	NS*
19	Potassium as K	mg/l	4.11	3.11	3.29	NS*	NS*
20	Manganese	mg/l	<0.04	<0.04	<0.04	0.1	0.3
21	Hexavalent Chromium	mg/l	<0.03	<0.03	<0.03	NS*	NS*
22	Copper	mg/l	<0.05	<0.05	<0.05	0.05	1.5
23	Cadmium	mg/l	<0.002	<0.002	<0.002	0.003	0.003
24	Arsenic	mg/l	<0.01	<0.01	<0.01	0.01	0.05
25	Mercury	mg/l	<0.001	<0.001	<0.001	0.001	0.001
26	Lead	mg/l	<0.01	<0.01	<0.01	0.01	0.01
27	Zinc	mg/l	<0.1	<0.1	<0.1	5.0	15.0
28	Bacterial Count	CFU/100ml	Absent	Absent	Absent	Absent	Absent

\*NS: Not Specified

**Table 14: Drinking Water Quality Monitoring Parameters for F - Type Quarter, Hospital Gopalpuri & Tuna Port**

Sr. No.	Parameter	Unit	F - Type Quarter	Hospital Gopalpuri	Tuna Port	Acceptable Limits as per IS 10500 : 2012	Permissible Limits as per IS 10500 : 2012
1	pH	pH Unit	7	7.3	7.38	6.5 to 8.5	6.5 to 8.5
2	Total Dissolved Solids	mg/l	890.0	950.0	1030.0	500	2000
3	Turbidity	NTU	1	0	0	1.0	5.0
4	Odor	-	Odourless	Odourless	Odorless	Agreeable	Agreeable
5	Color	Hazen Unit	Colourless	Colourless	Colorless	5.0	15.0
6	Conductivity	µs/cm	1700.0	2030.0	1920.0	NS*	NS*
7	Biochemical Oxygen Demand	mg/l	<2.0	<2.0	<2	NS*	NS*
8	Chloride	mg/l	706.57	545.21	692.0	250.0	1000.0
9	Ca as Ca	mg/l	52.10	56.11	69.74	75.0	200.0
10	Mg as Mg	mg/l	85.05	85.05	38.39	30.0	100.0
11	Total Hardness	mg/l	480	490	332.0	200.0	600.0
12	Iron as Fe+3	mg/l	<0.01	<0.01	<0.01	0.3	No Relaxation
13	Fluorides	mg/l	0.65	1.00	0.39	1.0	1.5
14	Sulphate	mg/l	358.8	378	112.8	200.0	400
15	Nitrite	mg/l	0.06	0.04	<0.01	NS*	NS*
16	Nitrate	mg/l	9.856	11.264	1.42	45.0	No Relaxation
17	Salinity	%	1.28	0.98	1.23	NS*	NS*
18	Sodium as Na	mg/l	373.0	351.0	344	NS*	NS*
19	Potassium as K	mg/l	4.07	3.87	3.4	NS*	NS*
20	Manganese	mg/l	<0.04	<0.04	<0.04	0.1	0.3
21	Hexavalent Chromium	mg/l	<0.03	<0.03	<0.03	NS*	NS*
22	Copper	mg/l	<0.05	<0.05	<0.05	0.05	1.5
23	Cadmium	mg/l	<0.002	<0.002	<0.002	0.003	0.003
24	Arsenic	mg/l	<0.01	<0.01	<0.01	0.01	0.05
25	Mercury	mg/l	<0.001	<0.001	<0.001	0.001	0.001
26	Lead	mg/l	<0.01	<0.01	<0.01	0.01	0.01
27	Zinc	mg/l	<0.1	<0.1	<0.1	5.0	15.0
28	Bacterial Count	CFU/100ml	Absent	Absent	Absent	Absent	Absent

\*NS: Not Specified

**Table 15: Drinking Water Quality Monitoring Parameters for Vadinar Jetty & Port Colony at Vadinar**

Sr. No.	Parameter	Unit	Vadinar Jetty	Port Colony Vadinar	Acceptable Limits as per IS 10500 : 2012	Permissible Limits as per IS 10500 : 2012
1	pH	pH Unit	7.5	7.3	6.5 to 8.5	6.5 to 8.5
2	Total Dissolved Solids	mg/l	990.0	1010.0	500	2000
3	Turbidity	NTU	0.00	1.00	1.0	5.0
4	Odor	-	Odourless	Odourless	Agreeable	Agreeable
5	Color	Hazen Units	Colourless	Colourless	5.0	15.0
6	Conductivity	µs/cm	1830.0	1990.0	NS*	NS*
7	Biochemical Oxygen Demand	mg/l	<2.0	<2.0	NS*	NS*
8	Chloride	mg/l	445.99	496.10	250.0	1000.0
9	Ca as Ca	mg/l	52.104	56.11	75.0	200.0
10	Mg as Mg	mg/l	80.19	80.19	30.0	100.0
11	Total Hardness	mg/l	460.0	470.0	200.0	600.0
12	Iron as Fe+3	mg/l	<0.01	<0.01	0.3	No Relaxation
13	Fluorides	mg/l	0.82	0.94	1.0	1.5
14	Sulphate	mg/l	30.00	34.80	200.0	400
15	Nitrite	mg/l	0.04	0.04	NS*	NS*
16	Nitrate	mg/l	4.93	4.79	45.0	No Relaxation
17	Salinity	%	0.81	0.90	NS*	NS*
18	Sodium as Na	mg/l	311.0	306.0	NS*	NS*
19	Potassium as K	mg/l	4.3	4.9	NS*	NS*
20	Manganese	mg/l	<0.04	<0.04	0.1	0.3
21	Hexavalent Chromium	mg/l	<0.03	<0.03	NS*	NS*
22	Copper	mg/l	<0.05	<0.05	0.05	1.5
23	Cadmium	mg/l	<0.002	<0.002	0.003	0.003
24	Arsenic	mg/l	<0.01	<0.01	0.01	0.05
25	Mercury	mg/l	<0.001	<0.001	0.001	0.001
26	Lead	mg/l	<0.01	<0.01	0.01	0.01
27	Zinc	mg/l	<0.1	<0.1	5.0	15.0
28	Bacterial Count	CFU/100ml	Absent	Absent	Absent	Absent

\*NS: Not Specified

## 2.3 Results & Discussion

The colour of all drinking water samples was < 5 Hazen unit and odour of the samples was also agreeable. All parameters are found to be within the specified limit of the Drinking water Standard.

### pH

The limit of pH value for drinking water is specified as 6.5 to 8.5. pH value in the studied area varied from 6.9 to 7.8 pH unit. All the sampling points showed pH values within the prescribed limit by Indian Standards.

### Total Dissolved Solids (TDS)

TDS values in the studied area varied between 600 -1800 mg/l. None of the sampling points showed higher TDS values than the prescribed limit by Indian standards .

### Conductivity

Electrical Conductivity is the ability of a solution to transfer (conduct) electric current. Conductivity is used to measure the concentration of dissolved solids which have been ionized in a polar solution such as water. The conductivity in the samples collected during the month of June ranged from 2000-3800  $\mu\text{s}/\text{cm}$ . Electrical conductivity standards do not appear in BIS standards for drinking water.

### BOD

BOD value in the studied area was less than 2.0 mg/L. Indian standards does not show any standard values for BOD in drinking water.

### Chlorides

Excessive chloride concentration increase rates of corrosion of metals in the distribution system. This can lead to increased concentration of metals in the supply. Chloride value in the studied area varied between 380-960 mg/l and is found to be within the Permissible limit of the Drinking Water Standard.

### Calcium

Calcium value in the studied area varied between 60 - 90 mg/l and is found to be within the Permissible limit of the Drinking Water Standard. If calcium is present beyond the maximum acceptable limit, it causes incrustation of pipes.

### **Magnesium**

Magnesium value in the studied area varied between 25 – 90 mg/l. All the locations had Magnesium within the prescribed limits of 30-100 mg/L.

### **Total Hardness**

Hardness value in the studied area varied between 312-520 mg/l and is found to be within the Permissible limit of the Drinking Water Standard. The prescribed limit by Indian Standards is 200-600 mg/L.

### **Iron**

Iron value in the studied area was below 0.01mg/L and hence well below the permissible limit as per Indian Standards is 0.3 mg/L. The excess amount of iron causes slight toxicity; gives stringent taste to water.

### **Fluoride**

Fluoride value in the studied area varied between 0.1 – 1.0 mg/l and hence well below the permissible limit as per Indian Standards is 1.0-1.5 mg/L. Moderate amounts lead to dental effects, but long-term ingestion of large amounts can lead to potentially severe skeletal problems.

### **Sulphates**

Sulphate value in the studied area varied between 100 – 330 mg/l. All the sampling points showed sulphate values within the prescribed limits by Indian Standards (200-400 mg/L). Sulphate content in drinking water exceeding the 400 mg/L imparts bitter taste.

### **Nitrites (NO<sub>2</sub>) and Nitrates (NO<sub>3</sub>)**

Nitrite values in all the water samples were <0.1. There are no specified standard values for Nitrites in Drinking water. The mean Nitrate values in drinking water of KPT was 4.10 mg/l which is well within the permissible limit of the Drinking water Standard.

### **Salinity**

Salinity in drinking water in the present samples collected ranged from 0.6 to 1.8 % . There are no prescribed Indian standards for salinity in Drinking water.

### **Sodium and Potassium Salts**

Sodium values in the samples collected ranged from 80 - 460 mg/l and Potassium salts ranged from 2.8 to 4.6 mg/l. There are no prescribed limits of Sodium and Potassium in Indian standards for Drinking water.

### **Heavy Metals in Drinking Water**

In the present study period drinking water samples were analyzed for Mn, Cr, Cu, Cd, As, Hg, Pb and Zn. All these heavy metals were well below the permissible limits prescribed by the Indian Standards.

### **Bacteriological Study**

Analysis of the bacteriological parameter at all location shows that Bacteria is not present and hence Bacterial count is in line with the permissible limit of drinking water. This shows that all the drinking water samples were safe from any bacteriological contamination.

## **2.4 Conclusions**

These results are compared with acceptable limits as prescribed in IS 10500:2012 – Drinking Water Specification. It is seen from the analysis data that during the study period the water was safe for human consumption at all drinking water monitoring stations.



### 3. Noise Level Monitoring

Noise sources in port operations include cargo handling, vehicular traffic, and loading / unloading containers and ships. Noise Monitoring was done at 13 stations at Kandla, Vadinar and Township area.

#### 3.1 Method of Monitoring

Sampling was done at all stations for 24 hour period. Data was recorded using automated sound level meter. The intensity of sound was measured in sound pressure level (SPL) and common unit of measurement is decibel (Db).

#### 3.2 Results

Table 16: Noise Monitoring data for ten locations of Deendayal Port and two locations of Vadinar Port

Sr. No.	Location	Day Time Average Noise Level (SPL) in dB(A)	Night Time Average Noise Level (SPL) in dB(A)
	<b>Sampling Time</b>	<b>6:00 am to 10:00 PM</b>	<b>10:00PM to 6:00 AM</b>
1	Marine Bhavan	63.40	57.1
2	Nirman Building 1	57.8	53.9
3	Tuna Port	55.8	47.1
4	Main Gate North	57.1	52.8
5	West Gate I	62.1	54.6
6	Canteen Area	57.1	49.6
7	Main Road	60.0	57.8
8	ATM Building	63.5	56.2
9	Wharf Area /Jetty Area	67.1	57.8
10	Port & Custom Office	55.5	52.7
	<b>Vadinar Port</b>		
11	Entrance Gate of Vadinar Port	57.1	54.6
12	Nr. Port Colony, Vadinar	56.2	56.2
13	Nr. Vadinar Jetty	59.6	55.8

**3.3 Conclusions-** Noise sources in port operations include cargo handling, vehicular traffic, and loading / unloading containers and ships. The Day Time Average Noise Level (SPL) in all ten locations at Deendayal Port ranged from 56.25 dB(A) to 69.51 dB(A) and it was within the permissible limits of 75 dB(A) for the industrial area for the daytime. The Night Time Average Noise Level (SPL) in all ten locations of Deendayal Port ranged from 48.28 dB to 62.33 dB(A) and it was within the permissible limits of 70 dB(A) for the industrial area for the night time.

#### 4. Soil Monitoring

Sampling and analysis of soil samples were undertaken at six locations within the study area (Deendayal Port and Vadinar Port) as a part of EMP. The soil sampling locations are initially decided based on the locations as provided in the tender document of the Deendayal Port.

##### 4.1 Methodology

The soil samples were collected in the month of June 2021. The samples collected from the all locations are homogeneous representative of each location. At random locations were identified at each location and soil was dug from 30 cm below the surface. It was uniformly mixed before homogenizing the soil samples. The samples were filled in polythene bags, labeled in the field with number and site name and sent to laboratory for analysis.

##### 4.2 Results

**Table-17: Chemical Characteristics of Soil in the Study Area**

Sr. No.	Parameter	Unit	Station Name					
			SL1	SL2	SL3	SL4	SL5	SL6
			Tuna Port	IFFCO Plant	Khori Creek	Nakti Creek	KPT Admin Site	KPT Colony
			Near main gate of Port	10 m away from main gate	Sand from creek at low tide		Vadinar	
1	Texture		Sandy Loam	Sandy Loam	Sandy Loam	Sandy Loam	Sandy Loam	Sandy Loam
2	pH	-	7.30	8.16	8.36	8.26	7.27	7.82
3	Electrical Conductivity	µs/cm	33400.0	48500.0	21800.0	37200.0	511.0	464.0
4	Moisture	%	21.45	13.94	18.82	14.26	6.28	4.56
5	Total Organic Carbon	%	0.31	0.19	0.26	0.24	0.15	0.11
6	Alkalinity	mg/kg	100.1	140.14	80.08	140.14	60.06	100.1
7	Total Nitrogen	%	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001
8	Chloride	mg/kg	6228.7	6032.5	2550.3	7160.6	68.66	78.47
9	Sulphate	mg/kg	2056.4	75.86	292.0	87.84	14.37	13.58
10	Phosphorus	mg/kg	0.97	1.41	0.79	1.59	0.97	0.97
11	Potassium	mg/kg	1161.0	592.2	700.2	765.0	626.4	876.4
12	Calcium	mg/kg	641.3	561.12	701.4	661.32	124.2	172.3
13	Sodium	mg/kg	10821.6	2992.8	3164.4	3736.8	2116.8	2565.0
14	Copper as Cu	mg/kg	11.21	27.22	28.20	31.78	82.66	72.42
15	Lead as Pb	mg/kg	3.10	6.20	23.0	11.4	ND	ND
16	Nickel as Ni	mg/kg	20.71	1823	7.80	15.10	25.46	27.73
17	Zinc as Zn	mg/kg	32.26	72.62	65.90	77.21	23.46	43.20
18	Cadmium as Cd	mg/kg	ND	ND	ND	ND	ND	ND

### 4.3 Discussion

- The data shows that value of pH ranges from 8.68 at Nakti Creek to 9.02 at Tuna Creek indicating that all soil samples are neutral to basic. Iffco plant samples showed maximum conductivity of 36,200  $\mu\text{mhos/cm}$ , while Nakti Creek location showed minimum conductivity of 4790  $\mu\text{mhos/cm}$ . Conductivity at Vadinar Port was 439 and 634  $\mu\text{mhos/cm}$  at Admin site and Vadinar Port colony respectively.
- Total organic Carbon ranged from 0.7 % to 2.3 at Deendayal Port. At Vadinar Port, organic carbon content ranged from 0.8 % to 1.04 %.
- The concentration of Phosphorus and Potassium in the soil samples varies from 34.0 to 53.0 mg/kg and 700.0 to 1100 mg/kg respectively at Deendayal Port. The mean concentration of Phosphorous at Vadinar site was 6.82 mg/kg and mean concentration of Potassium at Vadinar site was 176.5 mg/kg.

These differences in NPK in soil at different locations are due to the dissimilar nature of soil at each of the locations. Samples SL3 & SL4 (Khor Creek & Nakti Creek) are of saline nature as they are coastal soil; whereas other locations are inland locations and have different chemical properties.

### Heavy Metals in the Soil

Traces of Copper, Lead, Nickel, Zinc were observed in the soil samples collected from all the four locations of Deendayal Port and two locations of Vadinar Port. Cadmium metal was not detected in the Soil.

### 4.4 Conclusion

The soils of Deendayal Port and Vadinar Port appear to be neutral to basic with varying levels of Chloride, Sulphate, NPK and Calcium. As the nature of soil at different locations are different with respect to its proximity to the sea, the samples showed high degree of variations in their chemical properties.

## 5. Sewage Treatment Plant Monitoring

This involves safe collection of waste water (spent/used water) from wash areas, bathroom, industrial units, etc., waste from toilets of various buildings and its conveyance to the treatment plant and final disposal in conformity with the requirement and guide lines of State Pollution Control Board and other statutory bodies.

### 5.1 Methodology for STP Monitoring

To monitor the working efficiency of Sewage Treatment Plant (STP), STP Inlet and Outlet Samples were collected once a week. Locations selected are namely Gopalpuri Township, Deendayal Port and Vadinar. Samples were collected in 1 lit. carboys and were analyzed in laboratory for various parameters.

### 5.2 Results

- **Kandla STP**

**Table 18: Sewage Water Monitoring at Kandla STP (1<sup>st</sup> Week)**

Date of Sampling		05.06.21		
Sr. No.	Parameters	Unit	Results	
			KPT STP I/L	KPT STP O/L
1	pH	pH unit	7.62	7.23
2	Total Suspended Solids	mg/l	450	38.2
3	Residual Chlorine	mg/l	<1.0	<0.5
4	COD	mg/l	494.9	40.4
5	BOD @ 27 °C	mg/l	152.0	12.0
<b>Aeration Tank</b>				
6	MLSS	mg/l	40.0	
7	MLVSS	%	82.0	

**Table 19: Sewage Water Monitoring at Kandla STP (2<sup>nd</sup> Week)**

<b>Date of Sampling</b>	<b>10.06.21</b>
-------------------------	-----------------

Sr. No.	Parameters	Unit	Results	
			KPT STP I/L	KPT STP O/L
1	pH	pH unit	7.93	7.13
2	Total Suspended Solids	mg/l	268.3	58.3
3	Residual Chlorine	mg/l	<1.0	<0.5
4	COD	mg/l	289.87	63.63
5	BOD @ 27 °C	mg/l	94.0	16.0
<b>Aeration Tank</b>				
6	MLSS	mg/l	36.0	
7	MLVSS	%	74.0	

**Table 20: Sewage Water Monitoring at Kandla STP (3<sup>rd</sup> Week)**

<b>Date of Sampling</b>	<b>15.06.21</b>
-------------------------	-----------------

Sr. No.	Parameters	Unit	Results	
			KPT STP I/L	KPT STP O/L
1	pH	pH unit	7.43	7.23
2	Total Suspended Solids	mg/l	210.5	99
3	Residual Chlorine	mg/l	<1.0	<0.5
4	COD	mg/l	259.57	31.31
5	BOD @ 27 °C	mg/l	72.0	8.0
<b>Aeration Tank</b>				
6	MLSS	mg/l	36.0	
7	MLVSS	%	78.0	

**Table 21: Sewage Water Monitoring at Kandla STP (4<sup>th</sup> Week)**

<b>Date of Sampling</b>		<b>21.06.21</b>		
Sr. No.	Parameters	Unit	Results	
			KPT STP I/L	KPT STP O/L
1	pH	pH unit	7.71	7.02
2	Total Suspended Solids	mg/l	226.1	18.3
3	Residual Chlorine	mg/l	<1.0	<0.5
4	COD	mg/l	303.0	96.0
5	BOD @ 27 °C	mg/l	110.0	18.0
<b>Aeration Tank</b>				
6	MLSS	mg/l	20.0	
7	MLVSS	%	96.0	

- **Gopalpuri Colony STP**

**Table 22: Sewage Water Monitoring at Gopalpuri STP (1<sup>st</sup> Week)**

<b>Date of Sampling</b>		<b>05.06.21</b>		
Sr. No.	Parameters	Unit	Results	
			Gopalpuri STP I/L	Gopalpuri STP O/L
1	pH	pH unit	7.21	7.09
2	Total Suspended Solids	mg/l	166.7	54.9
3	Residual Chlorine	mg/l	<1.0	<0.5
4	COD	mg/l	383.8	84.4
5	BOD @ 27 °C	mg/l	124.0	16.0
<b>Aeration Tank</b>				
6	MLSS	mg/l	28.0	
7	MLVSS	%	86.0	

**Table 23: Sewage Water Monitoring at Gopalpuri STP (2<sup>nd</sup> Week)**

<b>Date of Sampling</b>		<b>10.06.21</b>		
Sr. No.	Parameters	Unit	Results	
			Gopalpuri STP I/L	Gopalpuri STP O/L
1	pH	pH unit	7.7	7.26
2	Total Suspended Solids	mg/l	95.21	41.9
3	Residual Chlorine	mg/l	<1.0	<0.5
4	COD	mg/l	431.27	109.8
5	BOD @ 27 °C	mg/l	138.0	19.0
<b>Aeration Tank</b>				
6	MLSS	mg/l	18.0	
7	MLVSS	%	96.0	

**Table 24: Sewage Water Monitoring at Gopalpuri STP (3<sup>rd</sup> Week)**

<b>Date of Sampling</b>		<b>15.06.21</b>		
Sr. No.	Parameters	Unit	Results	
			Gopalpuri STP I/L	Gopalpuri STP O/L
1	pH	pH unit	7.53	7.29
2	Total Suspended Solids	mg/l	52.9	20.1
3	Residual Chlorine	mg/l	<1.0	<0.5
4	COD	mg/l	230.28	57.57
5	BOD @ 27 °C	mg/l	76.0	15.0
<b>Aeration Tank</b>				
6	MLSS	mg/l	12.0	
7	MLVSS	%	92.0	

**Table 25: Sewage Water Monitoring at Gopalpuri STP (4<sup>th</sup> Week)**

<b>Date of Sampling</b>	<b>21.06.21</b>
-------------------------	-----------------

Sr. No.	Parameters	Unit	Results	
			Gopalpuri STP I/L	Gopalpuri STP O/L
1	pH	pH unit	7.25	7.12
2	Total Suspended Solids	mg/l	183.8	89
3	Residual Chlorine	mg/l	<1.0	<0.5
4	COD	mg/l	202	20.2
5	BOD @ 27 °C	mg/l	68.0	6.0
<b>Aeration Tank</b>				
6	MLSS	mg/l	38.0	
7	MLVSS	%	98.0	

- **Vadinar STP**

**Table 26: Sewage Water Monitoring at Vadinar STP (1<sup>st</sup> Week)**

<b>Date of Sampling</b>	<b>05.06.21</b>
-------------------------	-----------------

Sr. No.	Parameters	Unit	Results	
			Vadinar STP I/L	Vadinar O/L
1	pH	pH unit	7.26	NOT WORKING
2	Total Suspended Solids	mg/l	139.5	
3	Residual Chlorine	mg/l	<1.0	
4	COD	mg/l	222.2	
5	BOD @ 27 °C	mg/l	86.0	



**Table 27: Sewage Water Monitoring at Vadinar STP (2<sup>nd</sup> Week)**

<b>Date of Sampling</b>	<b>05.06.21</b>
-------------------------	-----------------

Sr. No.	Parameters	Unit	Results	
			Vadinar STP I/L	Vadinar O/L
1	pH	pH unit	7.36	NOT WORKING
2	Total Suspended Solids	mg/l	108.8	
3	Residual Chlorine	mg/l	<1.0	
4	COD	mg/l	353.5	
5	BOD @ 27 °C	mg/l	108.0	

**Table 28: Sewage Water Monitoring at Vadinar STP (3<sup>rd</sup> Week)**

<b>Date of Sampling</b>	<b>15.06.21</b>
-------------------------	-----------------

Sr. No.	Parameters	Unit	Results	
			Vadinar STP I/L	Vadinar O/L
1	pH	pH unit	7.14	NOT WORKING
2	Total Suspended Solids	mg/l	166.7	
3	Residual Chlorine	mg/l	<1.0	
4	COD	mg/l	171.7	
5	BOD @ 27 °C	mg/l	52.0	

**Table 29: Sewage Water Monitoring at Vadinar STP (4<sup>th</sup> Week)**

<b>Date of Sampling</b>	<b>21.06.21</b>
-------------------------	-----------------

Sr. No.	Parameters	Unit	Results	
			Vadinar STP I/L	Vadinar O/I
1	pH	pH unit	7.26	Not working
2	Total Suspended Solids	mg/l	203.5	
3	Residual Chlorine	mg/l	<1.0	
4	COD	mg/l	90.9	
5	BOD @ 27 °C	mg/l	28.0	

**5.3 Conclusions:**

The GPCB standards of BOD, TSS and Residual Chlorine for STP outlet are 20 mg/lit, 30 mg/lit & 0.5 mg/lit respectively. It is suggested to do treatment on regular basis to avoid flow of contaminated/polluted water into the sea. Also, the STP at Vadinar is also non-functional and thus, steps should be taken to commission the STP at Vadinar Port. Hence, currently only inlet samples are collected and analysed.

## 6. Marine Water Monitoring

The Forty Second Amendment to the Constitution in 1976 underscored the importance of 'green thinking'. Article 48A enjoins the state to protect and improve the environment and safeguard the forests and wildlife in the country. Further, Article 51A(g) states that the "fundamental duty of every citizen is to protect and improve the natural environment including forests, lakes, rivers and wildlife and to have compassion for living creatures".

Policy Statement for Abatement of Pollution (1992) has suggested developing relevant legislation and regulation, fiscal incentives, voluntary agreements and educational programs and information campaigns. It emphasizes the need for integration by incorporating environmental considerations into decision making at all levels by adopting frameworks namely, pollution prevention at source, application of best practicable solution, ensure polluter pays for control of pollution, focus on heavily polluted areas and river stretches and involve public in decision-making. The National Conservation Strategy and Policy Statement on Environment and Development, (1992) aimed at "integrating environmental concerns with developmental imperatives to meet the challenges by redirecting the thrust of our developmental process so that the basic needs of our people could be fulfilled by making judicious and sustainable use of natural resources." The priorities mentioned in this policy document include the sustainable use of land and water resources, prevention and control of pollution and preservation of biodiversity.

The National Water Policy, (2002) contains provisions for developing, conserving, sustainable utilizing and managing this important water resources and need to be governed by national perspectives.

### Marine Environment

On national and state levels, we have several policies and regulation like Water (Prevention and Control of Pollution) Act, 1974, to regulate pollution discharges and restore water quality of our aquatic resources including the prescription of monitoring activities. One of the important provisions of the Water Act, 1974, is to maintain and restore the 'wholesomeness' of our aquatic resources. Water quality monitoring is one of the first steps required in the rational development and management of water resources. In the field of water quality management, there has been a steady evolution in procedures for designing system to obtain information on the changes of water quality. The monitoring comprises all activities to obtain 'information' with respect to the water system.

## Sampling Stations

The monitoring of marine environment for the study of biological and ecological parameters was carried out on 10<sup>th</sup>& 11<sup>th</sup> June -2021 in harbor regions of KPT and on 10<sup>th</sup> June-2021 at Vadinar during spring tide period of New moon phase of Lunar Cycle. The monitoring of marine environment for the study of biological and ecological parameters was repeated again on 18<sup>th</sup>& 19<sup>th</sup> June 2021 in harbor regions of KPT. 18<sup>th</sup> June -2021 in Vadinar during Neap tide period first quarter of Lunar Cycle..

Plankton samples from sub surface layer was collected both during high tide period and low tide period from 3 water quality monitoring stations of KPT harbour area and two stations in Nakti creek and one station in Khori creek. The same sampling schedule was repeated during consecutive spring tide and neap tide in same month. Plankton samples from sub surface layer was collected both during high tide period and low tide period from 1 water quality monitoring stations near Vadinar jetty area during spring tide and neap tide in this month .Collected water samples were processed for estimation of Chlorophyll- a, Pheophytin- a, qualitative & quantitative evaluation of phytoplankton, qualitative & quantitative evaluation zooplanktons (density and their population).

### Sampling Locations

Offshore monitoring requirement	Number of locations
Offshore Installations	3 in Kandla creek 2 in Nakti creek 1 in Khori creek 1 near Vadinar Jetty 1 near 1 <sup>st</sup> SBM
<b>Total Number of locations</b>	<b>8</b>

#### 6.1 Marine Water Quality

Marine water quality of marine waters of Deendayal Port Harbor waters, Khori and Nakti Creeks and two locations of Vadinar are monitored for various physico-chemical parameters during spring and neap tide of each month.

The results of marine water quality and Marine sediments are as below;

**Table 30: Marine Water Quality Monitoring Parameters for location near KPT colony**

Sr. No.	Parameters	Unit	Kandla Creek Near KPT colony (1)			
			23°0'58"N 70°13'22."E			
			Spring Tide		Neap Tide	
Tide →	High Tide	Low Tide	High Tide	Low Tide		
1	pH	pH unit	7.21	7.35	7.18	7.14
2	Color	-	Colorless	Colorless	Colorless	Colorless
3	Odor	-	Odorless	Odorless	Odorless	Odorless
4	Salinity	ppt	31.7	32.1	32.8	
5	Turbidity	NTU	29	28	28	24
6	Total Dissolved Solids	mg/l	31107.0	35947.0	37797.0	33665.0
7	Total Suspended Solids	mg/l	377.4	359.9	714.2	412.4
8	Total Solids	mg/l	31560.0	36800.0	38860.0	34260.0
9	DO	mg/l	4.9	4.6	3.5	3.3
10	COD	mg/l	78.0	82.0	72.0	76.0
11	BOD	mg/l	<2.0	<2.0	<2.0	<2.0
12	Silica	mg/l	0.23	0.25	0.56	0.38
13	Phosphate	mg/l	0.35	0.36	0.27	0.24
14	Sulphate	mg/l	3360	3156	2628	3216
15	Nitrate	mg/l	1.97	2.35	2.14	2.78
16	Nitrite	mg/l	<0.01	<0.01	<0.01	<0.01
17	Calcium	mg/l	561.12	641.28	641.28	521.04
18	Magnesium	mg/l	1676.7	1676.7	1555.2	1725.3
19	Sodium	mg/l	11220.0	12080.0	8194.0	7418.0
20	Potassium	mg/l	380.0	390.0	372.0	414.0
21	Iron	mg/l	1.48	1.66	1.76	1.92
22	Chromium	mg/l	0.11	0.13	0.12	0.13
23	Copper	mg/l	0.05	0.06	0.06	0.07
24	Arsenic	mg/l	<0.01	<0.01	<0.01	<0.01
25	Cadmium	mg/l	0.06	0.08	0.05	0.06
26	Mercury	mg/l	<0.001	<0.001	<0.001	<0.001
27	Lead	mg/l	0.15	0.17	0.16	0.18
28	Zinc	mg/l	0.06	0.07	0.05	0.06

**Table 31: Marine Water Quality Monitoring Parameters for location near passenger Jetty One at Kandla**

Sr. No.	Parameters	Unit	Near passenger Jetty One (2)			
			23° 0'18 "N 70°13'31"E			
			Spring Tide		Neap Tide	
			Tide →	High Tide	Low Tide	High Tide
1	pH	pH unit	7.5	7.5	7.28	7.15
2	Color	-	Colorless	Colorless	Colorless	Colorless
3	Odor	-	Odorless	Odorless	Odorless	Odorless
4	Salinity	ppt	32.0	32.1	32.6	32.1
5	Turbidity	NTU	29	31	39	29
6	Total Dissolved Solids	mg/l	39865.0	39935.0	41765.0	36900.0
7	Total Suspended Solids	mg/l	366.8	414.5	404.0	477.9
8	Total Solids	mg/l	40212.0	40500.0	46018.0	37338.0
9	DO	mg/l	5.1	5.0	3.5	3.5
10	COD	mg/l	82.0	92.0	78.0	80.0
11	BOD	mg/l	<2.0	<2.0	<2.0	<2.0
12	Silica	mg/l	0.25	0.27	0.81	0.32
13	Phosphate	mg/l	0.35	0.32	0.20	0.33
14	Sulphate	mg/l	3120.0	3708.0	3336.0	2880.0
15	Nitrate	mg/l	6.0	2.54	1.35	4.33
16	Nitrite	mg/l	<0.01	<0.01	<0.01	<0.01
17	Calcium	mg/l	721.44	601.2	681.36	561.12
18	Magnesium	mg/l	1701.0	1603.8	1676.7	1725.3
19	Sodium	mg/l	11460.0	13211.0	9929.0	10111.0
20	Potassium	mg/l	390.0	382.0	471.0	381.0
21	Iron	mg/l	1.76	1.56	1.72	1.80
22	Chromium	mg/l	0.13	0.11	0.14	0.12
23	Copper	mg/l	0.06	0.07	0.08	0.07
24	Arsenic	mg/l	<0.01	<0.01	<0.01	<0.01
25	Cadmium	mg/l	0.05	0.06	0.06	0.07
26	Mercury	mg/l	<0.001	<0.001	<0.001	<0.001
27	Lead	mg/l	0.18	0.16	0.17	0.16
28	Zinc	mg/l	0.06	0.06	0.07	0.07

**Table 32: Marine Water Quality Monitoring Parameters for location Near Coal Berth**

Sr. No.	Parameters	Unit	Near Coal Berth			
			22°59'12"N 70°13'40"E			
			Spring Tide		Neap Tide	
			Tide →	High Tide	Low Tide	High Tide
1	pH	pH unit	7.3	7.21	7.2	7.5
2	Color	-	Colorless	Colorless	Colorless	Colorless
3	Odor	-	Odorless	Odorless	Odorless	Odorless
4	Salinity	ppt	31.0	32.2	32.0	32.3
5	Turbidity	NTU	33.0	31.0	31.8	32.0
6	Total Dissolved Solids	mg/l	34545.0	37030.0	35312.0	35363.0
7	Total Suspended Solids	mg/l	275.3	344.5	563.5	603.2
8	Total Solids	mg/l	35266.0	38080.0	36540	36100.0
9	DO	mg/l	4.8	4.6	4.2	4.3
10	COD	mg/l	86.0	92.0	101.0	100.0
11	BOD	mg/l	<2.0	<2.0	<2.0	<2.0
12	Silica	mg/l	0.27	0.20	0.38	0.22
13	Phosphate	mg/l	0.28	0.30	0.22	0.21
14	Sulphate	mg/l	1344	1500	2436	3240
15	Nitrate	mg/l	5.56	5.70	2.45	2.27
16	Nitrite	mg/l	0.028	0.02	0.04	0.02
17	Calcium	mg/l	641.28	681.36	601.2	641.28
18	Magnesium	mg/l	1555.2	1676.7	1652.4	1725.3
19	Sodium	mg/l	12015.0	11852.0	9320.0	9481.0
20	Potassium	mg/l	343.0	355.0	491.0	512.0
21	Iron	mg/l	1.44	1.23	1.64	1.34
22	Chromium	mg/l	0.12	0.10	0.12	0.13
23	Copper	mg/l	0.06	0.05	0.06	0.06
24	Arsenic	mg/l	<0.01	<0.01	<0.01	<0.01
25	Cadmium	mg/l	0.05	0.06	0.06	0.07
26	Mercury	mg/l	<0.001	<0.001	<0.001	<0.001
27	Lead	mg/l	0.18	0.12	0.16	0.14
28	Zinc	mg/l	0.06	0.06	0.05	0.06

**Table 33: Marine Water Quality Monitoring Parameters for location Khori creek at Kandla**

Sr. No.	Parameters	Unit	KPT 4			
			Near 15/16 Berth			
			Spring Tide		Neap Tide	
			Tide →	High Tide	Low Tide	High Tide
1	pH	pH unit	7.43	7.59	7.21	7.39
2	Color	-	Colorless	Colorless	Colorless	Colorless
3	Odor	-	Odorless	Odorless	Odorless	Odorless
4	Salinity	ppt	32.2	33.1	31.8	31.6
5	Turbidity	NTU	37	35	25	47
6	Total Dissolved Solids	mg/l	40837.0	45070.0	33588.0	33133.0
7	Total Suspended Solids	mg/l	299.2	315.5	407.3	438.9
8	Total Solids	mg/l	42994.0	46208.0	34336.0	34040.0
9	DO	mg/l	4.7	4.5	4.4	3.6
10	COD	mg/l	86.0	92.0	78.0	80.0
11	BOD	mg/l	<2.0	<2.0	<2.0	<2.0
12	Silica	mg/l	0.24	2.34	0.63	0.41
13	Phosphate	mg/l	0.28	0.32	0.26	0.28
14	Sulphate	mg/l	2628	2124	2988	2148
15	Nitrate	mg/l	7.25	2.64	4.67	7.08
16	Nitrite	mg/l	0.02	0.02	0.02	0.02
17	Calcium	mg/l	641.28	601.2	641.28	601.2
18	Magnesium	mg/l	1628.1	1749.6	1676.7	1652.4
19	Sodium	mg/l	10920.0	10962.0	9381.0	9252.0
20	Potassium	mg/l	344.0	352.0	366.0	488.0
21	Iron	mg/l	1.72	1.49	1.56	1.66
22	Chromium	mg/l	0.12	0.11	0.12	0.10
23	Copper	mg/l	0.05	0.05	0.06	0.05
24	Arsenic	mg/l	<0.01	<0.01	<0.01	<0.01
25	Cadmium	mg/l	0.06	0.07	0.08	0.07
26	Mercury	mg/l	<0.001	<0.001	<0.001	<0.001
27	Lead	mg/l	0.18	0.16	0.15	0.14
28	Zinc	mg/l	0.06	0.05	0.05	0.06



**Table 34: Marine Water Quality Monitoring Parameters for location Nakti Creek near Tuna Port**

Sr. No.	Parameters	Unit	Nakti Creek Near Tuna Port			
			22°57'49."N 70° 7'0.67"E			
			Spring Tide		Neap Tide	
			Tide →	High Tide	Low Tide	High Tide
1	pH	pH unit	7.39	7.21	7.73	7.7
2	Color	-	Colorless	Colorless	Colorless	Colorless
3	Odor	-	Odorless	Odorless	Odorless	Odorless
4	Salinity	ppt	32.2	31.8	32.4	31.6
5	Turbidity	NTU	34	60	39	49
6	Total Dissolved Solids	mg/l	48922.0	26656.0	39244.0	26963.0
7	Total Suspended Solids	mg/l	287.3	243.68	326.4	214.2
8	Total Solids	mg/l	49728.0	27300.0	40996.0	27294.0
9	DO	mg/l	4.6	4.9	4.2	3.5
10	COD	mg/l	96.0	98.0	88.0	82.0
11	BOD	mg/l	<2.0	<2.0	<2.0	<2.0
12	Silica	mg/l	0.28	0.29	0.61	0.74
13	Phosphate	mg/l	0.35	0.37	0.18	0.18
14	Sulphate	mg/l	3480	2868	2316	3480
15	Nitrate	mg/l	5.28	2.80	4.50	4.58
16	Nitrite	mg/l	0.02	<0.01	<0.01	0.03
17	Calcium	mg/l	601.2	721.44	521.04	601.2
18	Magnesium	mg/l	1749.6	1628.1	1773.9	1773.9
19	Sodium	mg/l	12126.0	12102.0	10821.0	10728.0
20	Potassium	mg/l	352.0	372.0	521.0	510.0
21	Iron	mg/l	1.52	1.42	1.56	1.59
22	Chromium	mg/l	0.16	0.14	0.13	0.15
23	Copper	mg/l	0.07	0.08	0.07	0.07
24	Arsenic	mg/l	<0.01	<0.01	<0.01	<0.01
25	Cadmium	mg/l	0.08	0.07	0.07	0.07
26	Mercury	mg/l	<0.001	<0.001	<0.001	<0.001
27	Lead	mg/l	0.14	0.12	0.12	0.13
28	Zinc	mg/l	0.05	0.06	0.06	0.07

**Table 35: Marine Water Quality Monitoring Parameters for location Nakti Creek Near NH-8A at Kandla**

Sr. No.	Parameters	Unit	Nakti Creek Near NH-8A			
			23° 02'01"N 70° 09'31"E			
			Spring Tide		Neap Tide	
			Tide →		High Tide	Low Tide
1	pH	pH unit	7.3	Sampling not possible during Low Tide	7.51	Sampling not possible during Low Tide
2	Color	-	Colorless		Colorless	
3	Odor	-	Odorless		Odorless	
4	Salinity	ppt	32.2		32.8	
5	Turbidity	NTU	37		38	
6	Total Dissolved Solids	mg/l	34970		35210.0	
7	Total Suspended Solids	mg/l	736.8		318.3	
8	Total Solids	mg/l	36048.0		36110.0	
9	DO	mg/l	5.1		3.9	
10	COD	mg/l	98.0		110.0	
11	BOD	mg/l	<2.0		<2.0	
12	Silica	mg/l	0.31		0.98	
13	Phosphate	mg/l	0.28		0.29	
14	Sulphate	mg/l	3720		2220	
15	Nitrate	mg/l	5.45		3.62	
16	Nitrite	mg/l	0.03		0.04	
17	Calcium	mg/l	721.44		681.36	
18	Magnesium	mg/l	1506.6		1749.6	
19	Sodium	mg/l	11622.0		10303.0	
20	Potassium	mg/l	486.0		495.0	
21	Iron	mg/l	1.49		1.62	
22	Chromium	mg/l	0.13		0.14	
23	Copper	mg/l	0.08		0.08	
24	Arsenic	mg/l	<0.01		<0.01	
25	Cadmium	mg/l	0.05		0.07	
26	Mercury	mg/l	<0.001		<0.001	
27	Lead	mg/l	0.19		0.14	
28	Zinc	mg/l	0.07		0.06	

**Table 36: Marine Water Quality Monitoring Parameters for locations Nr. Vadinar Jetty**

Sr. No.	Parameters	Unit	Nr.Vadinar Jetty			
			22°26'25.26"N 69°40'20.41"E			
			Spring Tide		Neap Tide	
			High Tide	Low Tide	High Tide	Low Tide
	Tide →					
1	pH	pH unit	7.25	7.36	7.26	7.21
2	Color	-	Colorless	Colorless	Colorless	Colorless
3	Odor	-	Odorless	Odorless	Odorless	Odorless
4	Salinity	ppt	31.8	32.2	31.2	32.0
5	Turbidity	NTU	5	9	21	15
6	Total Dissolved Solids	mg/l	34444	31931	37088	41030
7	Total Suspended Solids	mg/l	258	482	405.5	399.5
8	Total Solids	mg/l	34948.0	32054.0	37892.0	41410.0
9	DO	mg/l	3.8	4.2	1.9	2.8
10	COD	mg/l	86.0	88.0	72.0	68.0
11	BOD	mg/l	<2.0	<2.0	<2.0	<2.0
12	Silica	mg/l	0.21	0.30	0.45	0.39
13	Phosphate	mg/l	0.28	0.30	0.16	0.14
14	Sulphate	mg/l	3012	3192	2388	1980
15	Nitrate	mg/l	5.7376	4.32256	0.07744	3.4496
16	Nitrite	mg/l	0.02	0.02	0.02	<0.01
17	Calcium	mg/l	561.12	521.04	561.12	521.04
18	Magnesium	mg/l	1409.4	1603.8	1579.5	1676.7
19	Sodium	mg/l	11720.0	12118.0	10062.0	10080.0
20	Potassium	mg/l	458.0	456.0	406.0	412.0
21	Iron	mg/l	1.77	1.56	1.66	1.62
22	Chromium	mg/l	0.13	0.12	0.16	0.15
23	Copper	mg/l	0.07	0.06	0.05	0.06
24	Arsenic	mg/l	<0.01	<0.01	<0.01	<0.01
25	Cadmium	mg/l	0.04	0.04	0.05	0.05
26	Mercury	mg/l	<0.001	<0.001	<0.001	<0.001
27	Lead	mg/l	0.18	0.19	0.17	0.16
28	Zinc	mg/l	0.08	0.08	0.06	0.07

### **6.1.1 Marine Sediments**

Sediment samples were collected with Van Veen Grab from the six locations in Kandla Port Waters and two locations in Vadinar Port. Samples were collected and preserved in silver foil in ice box to prevent the contamination/decaying of the samples.

## 6.2 Results

The Sediment Quality results are given in below from table no. 33

**Table 34A: Results of Analysis of Sediment of Kandla & Vadinar Port (Spring Tide)**

Sr. No.	Parameters	Unit	KPT - 1	KPT - 2	KPT - 3	Khori - 1	Nakti Creek Near Tuna Port	Nakti - 1 (Near NH-8A )	Jetty
1	Texture		Sandy Loam	Sandy Loam	Sandy Loam	Sandy Loam	Sandy Loam	Sandy Loam	Sandy Loam
2	Organic Matter	mg/kg	1.20	1.88	1.20	1.30	1.76	1.88	1.56
3	Organic Carbon	mg/kg	0.80	0.96	0.87	0.87	0.69	0.78	0.78
4	Inorganic Phosphate	mg/kg	132.0	126.0	156.0	177.0	167.0	182.0	175.0
5	Moisture	%	24.96	26.86	21.33	16.64	26.33	22.78	23.01
6	Aluminium	mg/kg	ND	ND	ND	ND	ND	ND	ND
7	Silica	mg/kg	23.0	22.0	26.0	28.0	32.0	36.0	40.0
8	Phosphate	mg/kg	10.80	11.50	11.77	12.71	9.24	9.88	10.20
9	Sulphate	mg/kg	218.0	252.0	138.0	225.2	239.0	280.0	252.0
10	Nitrite	mg/kg	0.1	0.12	0.13	0.12	0.13	0.12	0.13
11	Nitrate	mg/kg	9.20	7.22	10.42	8.88	8.02	7.89	6.88
12	Calcium	mg/kg	861.0	1102.0	801.0	862.0	922.0	1082.0	802.0
13	Magnesium	mg/kg	437.0	851.0	693.0	765.0	449.0	522.0	422.0
14	Sodium	mg/kg	2083.0	2387.0	1937.0	1859.0	2857.0	2034.0	2185.0
15	Potassium	mg/kg	707.0	918.0	954.0	774.0	1058.0	779.0	792.0
16	Chromium	mg/kg	123.0	180.0	140.0	138.2	146.0	92.0	145.0
17	Nickel	mg/kg	26.0	23.2	28.9	26.2	32.6	33.6	37.7
18	Copper	mg/kg	46	42.7	21.20	36.0	37.2	29.6	26.8
19	Zinc	mg/kg	32.35	38.30	36.70	40.	41.00	39.00	40.00
20	Cadmium	mg/kg	ND	ND	ND	ND	ND	ND	ND
21	Lead	mg/kg	2.86	2.8	5.2	5.0	4.2	5.6	7.2
22	Mercury	mg/kg	ND	ND	ND	ND	ND	ND	ND
23	Arsenic	mg/kg	ND	ND	ND	ND	ND	ND	ND

\*Grab samples could not be collected due high current at Vadinar SBM

**Environmental Monitoring Report of Deendayal Port Trust, JUNE-2021**

**Table 34B: Results of Analysis of Sediment of Kandla & Vadinar Port (Neap Tide)**

Sr. No.	Parameters	Unit	KPT - 1	KPT - 3	Khori - 1	Nakti Creek Near Tuna Port	Nakti - 1 (Near NH-8A )	Jetty
1	Texture		Sandy loam	Sandy loam	Sandy loam	Sandy loam	Sandy loam	Sandy loam
2	Organic Matter	mg/kg	1.46	1.22	1.66	1.55	1.46	1.35
3	Organic Carbon	mg/kg	0.84	0.69	0.48	0.90	0.96	0.78
4	Inorganic Phosphate	mg/kg	155.0	148.0	162.0	149.0	164.0	166.0
5	Moisture	%	24.9	22.05	28.4	30.08	28.62	20.30
6	Aluminium	mg/kg	ND	ND	ND	ND	ND	ND
7	Silica	mg/kg	23.8	22.5	21.82	33.6	32.8	26.0
8	Phosphate	mg/kg	8.53	7.97	9.80	8.62	9.88	11.06
9	Sulphate	mg/kg	196.6	163.89	184.45	140.0	152.0	171.88
10	Nitrite	mg/kg	0.11	0.13	0.14	0.12	0.14	0.12
11	Nitrate	mg/kg	6.42	7.77	6.88	6.89	7.02	8.88
12	Calcium	mg/kg	288.6	212.0	232.4	284.0	296.0	224.0
13	Magnesium	mg/kg	177.4	177.0	170.76	197.2	188.0	535.0
14	Sodium	mg/kg	2662.0	1216.0	990.0	828.0	910.0	1150.0
15	Potassium	mg/kg	200.0	106.0	50.2	79.0	89.0	110.0
16	Chromium	mg/kg	145.0	133.0	146.0	126.0	101.0	166.0
17	Nickel	mg/kg	31.2	26.6	20.3	28.2	27.8	20.9
18	Copper	mg/kg	54.2	26.5	16.2	12.10	11.02	42.0
19	Zinc	mg/kg	23.0	31.0	24.62	29.42	33.36	42.52
20	Cadmium	mg/kg	ND	ND	ND	ND	ND	ND
21	Lead	mg/kg	ND	4.2	4.0	4.2	4.2	3.8
22	Mercury	mg/kg	ND	ND	ND	ND	ND	ND
23	Arsenic	mg/kg	ND	ND	ND	ND	ND	ND

\*Grab samples could not be collected due high current at KPT 2, Vadinar Jetty and Vadinar SBM

**REPORT**  
**ON**  
**ECOLOGICAL MONITORING**  
**OF MARINE ENVIRONMENT**  
**IN**  
**DPT HARBOUR AREA, NEAR BY CREEKS**  
**For**  
**DEENDAYAL PORT TRUST**

JUNE, 2021

**INTRODUCTION:**

The Forty Second Amendment to the Constitution in 1976 underscored the importance of 'green thinking'. Article 48A enjoins the state to protect and improve the environment and safeguard the forests and wildlife in the country. Further, Article 51A(g) states that the "fundamental duty of every citizen is to protect and improve the natural environment including forests, lakes, rivers and wildlife and to have compassion for living creatures".

Policy Statement for Abatement of Pollution (1992) has suggested developing relevant legislation and regulation, fiscal incentives, voluntary agreements and educational programs and information campaigns. It emphasizes the need for integration by incorporating environmental considerations into decision making at all levels by adopting frameworks namely, pollution prevention at source, application of best practicable solution, ensure polluter pays for control of pollution, focus on heavily polluted areas and river stretches and involve public in decision-making. The National Conservation Strategy and Policy Statement on Environment and Development, (1992) aimed at "integrating environmental concerns with developmental imperatives to meet the challenges by redirecting the thrust of our developmental process so that the basic needs of our people could be fulfilled by making judicious and sustainable use of natural resources." The priorities mentioned in this policy document include the sustainable use of land and water resources, prevention and control of pollution and preservation of biodiversity.

The National Water Policy, (2002) contains provisions for developing, conserving, sustainable utilizing and managing this important water resources and need to be governed by national perspectives.

**MARINE ENVIRONMENT:**

On national and state levels, we have several policies and regulation like Water (Prevention and Control of Pollution) Act, 1974, to regulate pollution discharges and restore water quality of our aquatic resources including the prescription of monitoring activities. One of the important provisions of the Water Act, 1974, is to maintain and restore the 'wholesomeness' of our aquatic resources. Water quality monitoring is one of the first steps required in the rational development and management of water resources. In the field of water quality management, there has been a steady evolution in procedures for designing system to obtain information on the changes of water quality. The monitoring comprises all activities to obtain 'information' with respect to the water system.



**Sampling Stations:**

The monitoring of marine environment for the study of biological and ecological Parameters was carried out on 10<sup>th</sup> June, 2021 in in harbour region of DPT, and on 11thJune, 2021 in creeks near by the port during spring tide .The monitoring of marine environment for the study of biological and ecological parameters was repeated again on 18<sup>th</sup> June, 2021 in harbour region of DPT and on19<sup>th</sup>June, 2021 in creeks near by the port during neap tidal condition .

Plankton samples from sub surface layer was collected both during high tide period and low tide period from 3 water quality monitoring stations of DPT harbour area andone stations in Nakti creek and one station in Khori creek. Sampling at second sampling station of Nakti creek was possible only during high tide period. Collected water samples were processed for estimation of Chlorophyll- a, Pheophytin- a, qualitative &quantitative evaluation of phytoplankton, qualitative &quantitative evaluation zooplanktons (density and their population).

**TABLE #1 SAMPLING LOCATIONS**

<b>monitoring requirement</b>	<b>Number of locations</b>
Kandla creek	3 in Kandla creek
Nakti creek	2 in Nakti creek
Khori Creek	1 in Khori creek
<b>Total Number of locations</b>	<b>6</b>

**Sampling methodology adopted:**

A marine sampling is an estimation of the body of information in the population. The theory of the sampling design is depending upon the underlying frequency distribution of the population of interest. The requirement for useful water sampling is to collect a representative sample of suitable volume from the specified depth and retain it free from contamination during retrieval. 50 liters of the water sample were collected from Sub surface by using bucket. From the collected water sample 1 liter of water sample were taken in an opaque plastic bottle for chlorophyll estimation, thereafter plankton samples were collected by using filtration assembly with nilyobolt cloth of 20 $\mu$ m mesh size.

**Samples Processing for chlorophyll estimation:**

Samples for the chlorophyll estimation were preserved in ice box on board in darkness to avoid degradation in opaque container covered with aluminium foil. Immediately after reaching the shore after sampling, 1 liter of collected water sample was filtered through GF/F filters (pore size 0.45  $\mu$ m) by using vacuum filtration assembly. After vacuum filtration the glass micro fiber filter paper was grunted in tissue grinder, macerating of glass fiber filter paper along with the filtrate was done in 90% aqueous Acetone in the glass tissue grinder with glass grinding tube. Glass fiber filter paper will assist breaking the cell during grinding and chlorophyll content was extracted with 10 ml of 90% Acetone, under cold dark conditions along with saturated magnesium carbonate solution in glass screw cap tubes. After an extraction period of 24 hours, the samples were transferred to calibrated centrifuge tubes and adjusted the volume to original volume with 90% aqueous acetone solution to make up the evaporation loss. The extract was clarified by using centrifuge in closed tubes. The clarified extracts were then decanted in clean cuvette and optical density was observed at wavelength 664, 665 nm. By using corrected optical density, Chlorophyll-a value was calculated as given in (APHA, 1998).

**PLANKTON:**

The entire area open water in the sea is the pelagic realm. Pelagic organisms live in the open sea. In contrast to the pelagic realm, the benthic realm comprises organisms and zone of the bottom of the sea. Vertically the pelagic realm can be dividing into two zones based on light penetration; upper photic or euphotic zone and lower dark water mass, aphotic zone below the photic zone. The term plankton is general term for organisms have such limited powers of locomotion that they are at the mercy of the prevailing water movement. Plankton is subdivided to phytoplankton and zooplankton. Phytoplankton is free floating organisms that are capable of photosynthesis and zooplankton is the various free floating animals.

Pelagic zone, represents the entire ocean water column from the surface to the deepest depths, is home to a diverse community of organisms. Differences in their locomotive ability categorize the organisms in the pelagic realm into two, **plankton** and **nekton** (Lalli and Parsons, 1997). **Plankton** consists of all organisms drifting in the water and is unable to swim against water currents, whereas **Nekton** includes organisms having strong locomotive power. Ecological studies on the plankton community, which form the base of the aquatic food chain, help in the better understanding of the dynamics and functioning of the marine ecosystem. The term 'Plankton' first coined by Victor Hensen (1887), Plankton, (Greek word: *planktos* meaning "passively drifting or wandering") is defined as drifting or free-floating organisms that inhabit the pelagic zone of water. Based on their mode of nutrition planktonic organisms are categorised into phytoplankton (organisms having an autotrophic mode of nutrition) and zooplankton (organisms having a heterotrophic mode of nutrition).

#### Phytoplankton in the marine environment:

Phytoplankton is free floating unicellular, filamentous and colonial eutrophic organisms that grow in aquatic environments whose movement is more or less dependent upon water currents. These micro flora acts as primary producers as well as the basis of food chain, source of protein, bio purifier and bio indicators of the aquatic ecosystems of which diverse array of the life depends. They are considered as an important component of aquatic flora, play a key role in maintaining equilibrium between abiotic and biotic components of aquatic ecosystem.

The phytoplankton includes a wide range of photosynthetic and phototrophic organisms. Marine phytoplankton is mostly microscopic and unicellular floating flora, which are the primary producers that support the pelagic food-chain. The two most prominent groups of phytoplankton are diatoms (Bacillariophyceae) and dinoflagellates (Dinophyceae). The phytoplankton those normally captured in the net from the Gulf of Kutch is normally dominated by these two major groups; diatoms and dinoflagellates. Phytoplankton also include numerous and diverse collection of extremely small, motile algae which are termed micro flagellates (naked flagellates) as well as and Cyanophytes (blue-green algae).

Algae are an ecologically important group in most aquatic ecosystems and have been an important component of biological monitoring programs. Algae are ideally suited for water quality assessment because they have rapid reproduction rates and very short life cycles, making them valuable indicators of short-term impacts.

Aquatic populations are impacted by anthropogenic stress, resulting in a variety of alterations in the biological integrity of aquatic systems. Algae can serve as an indicator of the degree of

deterioration of water quality, and many algal indicators have been used to assess environmental status.

### **Zooplankton in the marine environment:**

Zooplankton includes a taxonomically and morphologically diverse community of heterotrophic organisms that drift in the waters of the world's oceans. Qualitative and quantitative studies on zooplankton community is a prerequisite to delineate the ecological processes active in the marine ecosystem. Zooplankton community plays a pivotal role in the pelagic food web as the primary consumers of phytoplankton and act as the food source for organisms in the higher trophic levels, particularly the economically essential groups such as fish larvae and fishes. They also function in the cycling of elements in the marine ecosystem. The dynamics of the zooplankton community, their reproduction, growth and survival rate are all significant factors determining the recruitment and abundance of fish stocks as they form an essential food for larval, juvenile and adult fishes (Beaugrand et al., 2004). Zooplankton grazing in the marine environment controls the primary production and helps in determining the pelagic ecosystem (Banse, 1995). Through grazing in surface waters and following the production of sinking faecal matters and also by the active transportation of dissolved and particulate matter to deeper waters via vertical migration, they help in the transport of organic carbon to deep ocean layers and thus act as key drivers of biological pump' in the marine ecosystem. Zooplankton grazing and metabolism also, transform particulate organic matter into dissolved forms, promoting primary producer community, microbial demineralization, and particle export to the ocean's interior.

The categorisation of zooplankton into various ecological groups is based on several factors such as duration of planktonic life, size, food preferences and habitat. As they vary significantly in size from microscopic to metazoic forms, the classification of zooplankton based on size has paramount importance in the field of quantitative plankton research.

Based on the duration of planktonic life, zooplankton are categorised into Holoplankton (organisms which complete their entire lifecycle as plankton) and Meroplankton (organisms which are planktonic during the early part of their lives such as the larval stages of benthic and nektonic organisms). Tychoplankton are organisms which live a brief planktonic life, such as the benthic crustaceans (cumaceans, mysids, isopods) which ascend to the water column at night for feeding and certain ectoparasitic copepods, they leave the host and spend their life as plankton during their breeding cycle.


Zooplankton can be subdivided into holoplankton, i.e., permanent members of the plankton (e.g., Calanoid copepods), and meroplankton, i.e., temporary members in the plankton e.g., larvae of

fish, shrimp, and crab). The meroplankton group consists of larval and young stages of animals that will adopt a different lifestyle once they mature. In contrast to phytoplankton which consist of a relatively smaller variety of organisms, Zooplankton are extremely diverse, consist of a host of larval and adult forms representing many animal phylum.

Among the zooplankton one group always dominate than others; members of sub class copepods (Phylum Athropoda), and Tintinids (Phylum Protozoa) among the net planktons. These small animals are of vital importance in marine ecosystem as one of the primary herbivores animals in the sea, and it is they provide vital link between primary producer (autotrophs) and numerous small and large marine consumers.

As their community structure and function are highly susceptible to changes in the environmental conditions regular monitoring of their distribution as well as their interactions with various physicochemical parameters is inevitable for the sustainable management of the ecosystem (Kusum et al., 2014). Of all the marine zooplankton groups, copepods mainly calanoid copepods are the dominant groups in marine subtropical and tropical waters and exhibit considerable diversity in morphology and habitats they occupy (Madhupratap, 1991;)

It has been well established that potential of pelagic fishes viz. finfishes, crustaceans, molluscs and marine mammals either directly or indirectly depend on zooplankton. The herbivorous zooplankton is efficient grazers of the phytoplankton and is referred to as living machines transforming plant material into animal tissue. Hence they play an essential role as the intermediaries for nutrients/energy transfer between primary and tertiary trophic levels. Due to their large density, shorter lifespan, drifting nature, high group/species diversity and different tolerance to the stress, they used as the indicator organisms for the physical, chemical and biological processes in the aquatic ecosystem (Ghajbhiye, 2002).



### Spatial distribution of Plankton:

A characteristic of plankton population is that they tend to occur in patches, which are varying spatially on a scale of few meters to far as few kilometres in distance. They also vary in time scale, season as well as vertically in the water column. It is this patchiness and its constant changes in time and spot, that has made it so difficult for plankton biologist to learn about the ecology of plankton. The biological factors that causes this patchiness is due to the ability of zooplankton to migrate vertically and graze out the phytoplankton at a rapid rate that can create patchiness. Similarly the active swimming ability by certain zooplankton organisms can cause to aggregate in dense group.

At its most extreme, because the water in which plankton is suspended is constantly moving, each sample taken by the plankton biologists remain a different volume of water, so each sample is unique and replicate does not exist.

Plankton June also exhibit vertical patchiness. Physical factors contribute to this type of patchiness include light intensity, nutrients and density gradients in the water column.

Phytoplankton in particular tends to be unequally distributed vertically, which leads to the existence of different concentration of a chlorophyll value between photic zone and below the photic zone.

**Methodology adopted for Plankton sampling:**

Mixed plankton sample were obtained from the sub surface layer at each sampling locations by towing the net horizontally with the weight .After the tow of about 15-30minutes, plankton net was pulled up and washed down to the tail and collected the plankton adhered to plankton net in the collection bucket at the bottom by springing outer and inner surface of the net with sea water, while the net was hanging with the mouth upward. For quantitative evaluation 50 L water samples were collected from subsurface layer and filtered through 20µm mesh size net by using bucket and filtration assembly.

**Preservation and storage:**

Both filtered plankton and those collected from the plankton net were preserved with 5% buffered formalin and stored in 1L plastic container for further processing in the laboratory.

**Sample concentration:**

The collected plankton samples were concentrated by using centrifuge and made up to 50 ml with 5% formalin -Glycerine mixture.

**Taxonomic evaluation:**

Before processing, the sample was mixed carefully and a subsample was taken with a calibrated Stempel-pipette. 1 ml of the concentrated plankton samples were transferred on a glass slide with automatic pipette. The plankton sample on the glass slides were stained by using Lugol's iodine and added glycerine to avoid drying while observation. The plankton samples were identified by using Labex triangular Research microscope with photographic attachment. Microphotographs of the plankton samples were taken for record as well as for confirming the identification. The bigger sized zooplankton was observed through dissecting stereomicroscope with magnification of 20-30 x. Plankton organisms in the whole slide were identified to the lowest axon possible. A thorough literature search was conducted for the identification of the different groups of zooplankton that were encountered

**Cell counts by drop count method:**

The common glass slide mounted with a 1ml of concentrated phytoplankton/zooplankton sample in glycerol and covered with cover slip 22x 60mm was placed under the compound microscope provided with a mechanical stage. The plankton was then counted from the microscopic field of the left top corner of the slide. Then slide is moved horizontally along the right side and plankton

in each microscopic field was thus counted. When first microscopic field row was finished the next consecutive row was adjusted using the mechanical device of the stage. In this way all the plankton present in entire microscopic field are counted. From this total number in 1ml of the concentrated plankton, total number of plankton in the original volume of sample filtered was calculated as units/L.

#### **BENTHIC ORGANISMS:**

Benthos is those organisms that are associated with the sea bed or benthic habitats. Epi- benthic organisms live attached to a hard substratum or rooted to a shallow depth below the surface. In fauna organisms live below the sediment–water interface. Interstitial organisms live and move in pore water among sedimentary grains.

Because the benthic organisms are often collected and separated on sieves, a classification based on the overall size is used. Macro benthos include organisms whose shortest dimension is greater than or equal to 0.5 mm. Meio benthos are smaller than 0.5mm but larger than 42 $\mu$  in size.

The terms such as macro fauna and Meio fauna generally have little relevance with taxonomic classification. The terms Meio fauna and macro fauna depend on the size. Meio fauna were considered as good bioassay of community health and rather sensitive indicators of environmental changes

#### **SAMPLING METHODOLOGY ADOPTED FOR SUB TIDAL REGION:**

Van veen sampler (0.09m<sup>2</sup>) was used for sampling bottom sediments. Two sets of sediments were sampled from each location, one for macro fauna and other for Meio fauna. The macro fauna in the sediments were sieved on board to separate out the organisms. The fixation of Meio fauna is normally done by bulk fixation of the sediment sample. The bulk fixation is done by using 10% formalin (Buffered with borate). The organisms were preserved with seawater as diluting agent.

#### **Sample sieving:**

Sediments samples were sieved to extract the organisms. Sieving was performed carefully as possible to avoid any damage to the animals. The large portion of the sediment was split in to smaller portions and mixed with sea water in a bucket. The cohesive lumps were broken down by continuous stirring. The disaggregated sediments were then passed through the sieves.

#### **Sample staining:**

Sorting of the Meio fauna from the sieve is difficult task especially in the preserved material, because organisms are not easily detectable. To facilitate the animal detection the entire sample retained on the sieve after sieving operation were stained by immersing the sieve in a flat bottom



tub with 1% Rose Bengal stain; a protein stain. A staining period of 10-30 minutes is sufficient for sample detection.

#### **DIVERSITY INDICES:**

On the whole, diversity indices provide more information about community composition than simply species richness (number of species present); they also, take the relative abundances of different species into account. Based on this fact, diversity indices therefore depend not only on species richness but on the evenness, or equitability, with which individuals are distributed among the different species (Magurran, A. E. (1988)

A diversity index is a measure of species diversity within a community that consists of co-occurring populations of several (two or more) different species. It includes two components: richness and evenness. Richness is the measure of the number of different species within a sample showing that more the types of species in a community, the higher is the diversity or greater is the richness. Evenness is the measure of relative abundance of the different species with in a community.

The basic idea of diversity index is to obtain a quantitative estimate of biological variability that can be used to compare biological entities composed of discrete components in space and time ( Carol H.R. *etal.* 1998). Biodiversity is commonly expressed through indices based on species richness and species abundances (Whittaker 1972, Lande 1996, Purvis and Hector 2000). Biodiversity indices are a non-parametric tool used to describe the relationship between species number and abundance. The most widely used bio diversity indices are Shannon Weiner index and Simpson's index.

A diversity Index is a single statistic that incorporates in formation on richness and evenness. The diversity measures that incorporate the two concepts may be termed heterogeneity measures (Magurran, 2004).

Any study intended to interpret causes and effect of adverse impact on Biodiversity of communities require suitable measures to evaluate specie richness and Diversity. The former is number of species in community, while latter is a function of relative frequency of different species. Species richness is the iconic measure of biological diversity (Magurran, 2004). Several indices have been created to measure the diversity of species; however, the most widely used in the last decades are the Shannon (1948) and Simpson (1949) (Buzas and Hayek 1996; Gorelick 2006), with the components of diversity: richness ( $S$ ) and evenness ( $J$ )

### Simpson's diversity index

Simpson's index (**D**) is a measure of diversity, which takes into account both species richness, and evenness of abundance among the species present.. The Simpson index is one of the meaningful and robust biodiversity measures available.(Magurran ,2004).

The formula for calculating D is presented as:

$$D = \frac{\sum n_i(n_i - 1)}{N(N - 1)}$$

Where  $n_i$  = the total number of organisms of each individual species

$N$  = the total number of organisms of all species

The value of D ranges from 0 to 1. With this index, 0 represents infinite diversity and, 1, no diversity. When D increases diversity decreases. Simpson's index is therefore usually expressed as 1-D or 1/D. (Magurran, 2004)

Low species diversity suggests:

- relatively few successful species in the habitat
- the environment is quite stressful with relatively few ecological niches and only a few organisms are really well adapted to that environment
- food webs which are relatively simple
- change in the environment would probably have quite serious effects

High species diversity suggests:

- a greater number of successful species and a more stable ecosystem
- more ecological niches are available and the environment is less likely to be hostile complex food webs
- environmental change is less likely to be damaging to the ecosystem as a whole

### Species richness indices

The species richness(**S**) is simply the number of species present in an ecosystem. Species richness Indices of species richness are widely used to quantify or monitor the effects of anthropogenic disturbance. A decline in species richness may be concomitant with severe or chronic human-induced perturbation (Fair Fair weather 1990,) Species richness measures have traditionally been the mainstay in assessing the effects of environmental degradation on the biodiversity of natural assemblages of organisms (Clarke &Warwick, 2001)

Species richness is the iconic measure of biological diversity (Magurran, 2004). The species richness(**S**) is simply the number of species present in an ecosystem. This index makes no use of

relative abundances. The term species richness was coined by McIntosh (1967) and oldest and most intuitive measure of biological diversity (Magurran, 2004).

Margalef's diversity index is a species richness index. Margalef's Species richness index (d), or indices that describe the evenness of the distribution of the numbers of individuals among species, were derived.

The value of a diversity index increases both when the number of types increases and when evenness increases. For a given number of types, the value of diversity index is maximised when all types are equally abundant (Rosenzweig, M. L. (1995).

#### Shannon-Wiener's index:

An index of diversity commonly used in plankton community analyses is the Shannon-Wiener's index (**H**), which emphasizes not only the number of species (richness or variety), but also the apportionment of the numbers of individuals among the species (Odum 1971 and Reish 1984). Shannon-Wiener's index (**H**) reproduce community parameters to a single number by using an equation.

Shannon and Weiner index represents entropy. It is a diversity index taking into account the number of individuals as well as the number of taxa. It varies from 0 for communities with only single taxa to high values for community with many taxa each with few individuals. This index can also determine the pollution status of a water body. Normal values range from 0 to 4. This index is a combination of species present and the evenness of the species. Examining the diversity in the range of polluted and unpolluted ecosystems, Wilham and Dorris (1968) concluded that the values of the index greater than 3 indicate clean water, values in the range of 1 to 3 are characterized by moderate pollution and values less than 1 are characterized as heavily polluted

$$H' = - \sum_{j=1}^s \frac{n_j}{N} \ln \left( \frac{n_j}{N} \right)$$

#### RESULTS:

##### CHLOROPHYLL-a:

Water Samples for the chlorophyll estimation were collected from sub surface layer during high tide and low tide period of the tidal cycle for each sampling locations and analysed for Chlorophyll -a and after acidification for Pheophytin -a. Chlorophyll- a value was used as algal biomass indicator (APHA,1998) Algal biomass was estimated by converting Chlorophyll value.

In the sub surface water chlorophyll-a was varying from 0.314 -0.468mg/m<sup>3</sup>.in harbour region of DPT during sampling done in spring tide period of June, 2021. In the nearby creeks chlorophyll-a was varying from 0.329-0.739 mg/m<sup>3</sup>.Pheophytin –a level was below detectable limit- the all the sampling stations during spring in the harbour region of DPT.

In the sub surface water chlorophyll-a was varying from 0.527 -0.765 mg/m<sup>3</sup>.in harbour region of DPT during sampling done in neap tide period of June, 2021 . In the nearby creeks chlorophyll-a was varying from 0.425- 0.850 mg/m<sup>3</sup>.Pheophytin –a level was below detectable limit- the all the sampling stations during spring in the harbour region of DPT

**TABLE #2 VARIATIONS IN CHLOROPHYLL –a PHEOPHYTIN- a AND ALGAL BIOMASS FROM SAMPLING STATIONS IN DPT HARBOUR AREA ,NEAR BY CREEKS DURING SPRING TIDE IN JUNE,2021**

Sr.No.	Station	Tide	Chlorophyll-a (mg/m <sup>3</sup> )	Pheophytin- a (mg/m <sup>3</sup> )	Algal Biomass (Chlorophyll method) mg/m <sup>3</sup>
DPT HARBOUR AREA					
1	KPT1	High tide	0.381	BDL	25.53
		Low tide	0.440	BDL	29.48
2	KPT 2	High tide	0.314	BDL	21.04
		Low tide	0.417	BDL	27.94
3	KPT 3	High tide	0.468	BDL	31.36
		Low tide	0.424	BDL	28.41
CREEKS					
4	KPT-4 Khori-I	High tide	0.739	BDL	49.51
		Low tide	0.578	BDL	38.73
5	KPT-5 Nakti-I	High tide	0.637	BDL	42.68
		Low tide	0.409	BDL	27.40
6	KPT-5 Nakti-II	High tide	0.329	BDL	22.04

BDL: Below Detectable Limit.

**TABLE #3 VARIATIONS IN CHLOROPHYLL –aPHEOPHYTIN- a AND ALGAL BIOMASS FROM SAMPLING STATIONS IN DPT HARBOUR AREA ,NEAR BY CREEKS DURING NEAP TIDE IN JUNE,2021**

Sr.No.	Station	Tide	Chlorophyll-a (mg/m <sup>3</sup> )	Pheophytin- a (mg/m <sup>3</sup> )	Algal Biomass (Chlorophyll method) mg/m <sup>3</sup>
DPT HARBOUR AREA					
1	KPT1	High tide	0.631	BDL	42.28
		Low tide	0.765	BDL	51.25
2	KPT 2	High tide	0.731	BDL	48.98
		Low tide	0.614	BDL	41.14
3	KPT 3	High tide	0.527	BDL	35.31
		Low tide	0.615	BDL	41.21
CREEKS					
4	KPT-4 Khori-I	High tide	0.748	BDL	50.12
		Low tide	0.850	BDL	56.95
5	KPT-5 Nakti-I	High tide	0.715	BDL	47.90
		Low tide	0.715	BDL	47.90
6	KPT-5 Nakti-II	High tide	0.425	BDL	28.47

BDL: Below Detectable Limit.

### PHYTOPLANKTON POPULATION:

For the evaluation of the Phytoplankton population in DPT harbour area and within the immediate surroundings of the port, sampling was conducted from 5 sampling locations (3 in harbour area and two in Nakti creek) during high tide period and low tide period of spring tide. The phytoplankton community of the sub surface water in the harbour and nearby creeks was represented by, Diatoms and dinoflagellates during spring tide period. Diatoms were represented by 14 genera. Dinoflagellates were represented by one genera .during the sampling conducted in spring tide in June, 2021. Phytoplankton of the sampling stations at sub surface layer in the harbour area and nearby creeks was varying from 112-216 units/ L during high tide period and 147-172 units/ L during low tide of Spring Tide.

The phytoplankton community of the sub surface water in the harbour and nearby creeks was represented by Diatoms and dinoflagellates during spring tide period. Diatoms were represented by 15 genera and Dinoflagellates were represented one genera during the sampling conducted in Neap tide in June, 2021. Phytoplankton of the sampling stations at sub surface layer in the harbour area and nearby creeks was varying from 72-293 units/ L during high tide period and 202-375 units/ L during low tide of Neap Tide.

### Species Richness Indices and Diversity Indices :

#### Margalef's diversity index (Species Richness) S

At the organismal level, the most widely used biodiversity measures are those based on the number of species present, perhaps adjusted for the number of individuals sampled, Here Margalef's Species richness index (d), or indices that describe the evenness of the distribution of the numbers of individuals among species, are derived.

Margalef's diversity index (Species Richness) S of phytoplankton communities in the stations was varying from 1.907-2.660 with an average of 2.381 during the sampling conducted in High tide period of spring tide While .Margalef's diversity index (Species Richness) S of phytoplankton communities in the Kandla Harbour region and nearby creeks was varying from. 1.603-2.395 with an average of 2.140 during the consecutive in low tide period .

Margalef's diversity index (Species Richness) S of phytoplankton communities in the stations was varying from 1.775-2.614 with an average of 2.212 during the sampling conducted in High tide period of Neap tide While .Margalef's diversity index (Species Richness) S of phytoplankton communities in the Kandla Harbour region and nearby creeks was varying from. 1.695-2.193 with an average of 1.966 during the consecutive in low tide period .

#### Shannon-Wiener's index:

Shannon-Wiener's Index (H) of phytoplankton communities in the sampling stations was in the range of 0.794-0.908 ( $H'(\log_{10})$ ) between selected sampling stations with an average value of 0.844. during high tide period of spring tide .Shannon-Wiener's Index (H) of phytoplankton communities in the sampling stations was in the range of 0.788-0.845 ( $H'(\log_{10})$ ) between selected sampling stations with an average value of 0.813 during consecutive lowtide. Shannon-Wiener's Index (H) of phytoplankton communities in the sampling stations was in the range of 0.722-0.883 ( $H'(\log_{10})$ ) between selected sampling stations with an average value of 0.814. during high tide period of neap tide . Shannon-Wiener's Index (H) of phytoplankton communities in the sampling stations was in the range of 0.723-0.883 ( $H'(\log_{10})$ ) between selected sampling stations with an average value of 0.813 during consecutive low tide. Typical values are generally between 1.5 and 3.5 in most ecological studies, and the index is rarely greater than 4. The Shannon-Wiener's index increases as both the richness and the evenness of the community increase. This result indicates that diversity of phytoplankton of Kandla Harbour region and nearby creeks is less but with abundant population of few, with relatively few ecological niches and only very few opportunist organisms are really well adapted to this environment and thrive better than other species.

**Simpson's diversity index:**

Simpson's index (D) is a measure of diversity, which takes into account both species richness, and an evenness of abundance among the species present. The Simpson index is one of the meaningful and robust biodiversity measures available. (Magurran,2004).

Simpson diversity index (1-D) of phytoplankton communities was below 0.9 at all sampling stations in the Kandla Harbour region and nearby creeks, which was varying from 0.796- 0.840 between selected sampling stations with an average of 0.815 during high tide period of spring tide . Simpson diversity index (1-D) of phytoplankton communities was below 0.9 at all sampling stations in the Kandla Harbour region and nearby creeks, which was varying from 0.790- 0.821 between selected sampling stations with an average of 0.803 during consecutive low tide . Low species diversity suggests a relatively few successful species in this habitat. The environment is quite stressful with relatively few ecological niches and only a few organisms are really well adapted to that environment. Any change in the environment would probably have quite serious effects.

Simpson diversity index (1-D) of phytoplankton communities was below 0.9 at all sampling stations in Kandla Harbour region and nearby creeks, during high tide period and low tide period during neap tide also, which was varying from 0.741-0.831 with an average value of 0.800



**Environmental Monitoring Report of Deendayal Port Trust, JUNE-2021**

between selected sampling stations during high tide period and varying from 0.719-0.808 with an average value of 0.758 between selected sampling stations during consecutive low tide period. Low species diversity suggests a relatively few successful species in this habitat.

**Table # 4 PHYTOPLANKTON VARIATIONS IN ABUNDANCE AND DIVERSITY IN SUB SURFACE SAMPLING STATIONS IN DPT HARBOUR AREA ,NEAR BY CREEKS DURING SPRING TIDE IN JUNE,2021**

Tide	Sampling Station	Abundance In units/L	No of Species observed /total species	% of diversity	Margalef's diversity index (Species Richness S)	Shannon Weiner index H (log <sub>10</sub> )	Diversity Index (Simpson's Index) 1-D
HIGH TIDE	1	158	14/15	93.33	2.568	0.89	0.8401
	2	151	14/15	93.33	2.591	0.8397	0.8165
	3	170	12/15	80	2.142	0.7942	0.7962
	4	216	14/15	93.33	2.418	0.8223	0.8042
	5	193	15/15	100	2.66	0.9078	0.8326
	6	112	10/15	66.66	1.907	0.8103	0.8029
LOW TIDE	1	156	13/15	86.66	2.376	0.8446	0.8209
	2	147	9/15	60	1.603	0.7909	0.8148
	3	152	12/15	80	2.19	0.8051	0.791
	4	172	12/15	80	2.137	0.788	0.7904
	5	150	13/15	86.66	2.395	0.8371	0.7996

**Table # 5 PHYTOPLANKTON VARIATIONS IN ABUNDANCE AND DIVERSITY IN SUB SURFACE SAMPLING STATIONS IN DPT HARBOUR AREA ,NEAR BY CREEKS DURING NEAP TIDE IN JUNE,2021**

Tide	Sampling Station	Abundance In units/L	No of Species observed /total species	% of diversity	Margalef's diversity index (Species Richness S)	Shannon Weiner index H (log <sub>10</sub> )	Diversity Index (Simpson's Index) 1-D
HIGH TIDE	1	240	15/16	93.75	2.554	0.883	0.8308
	2	248	12/16	75	1.995	0.7916	0.7925
	3	212	15/16	93.75	2.614	0.8664	0.8285
	4	293	15/16	93.75	2.465	0.8666	0.8224
	5	280	11/16	68.75	1.775	0.7227	0.7413
	6	72	9/16	56.25	1.871	0.7522	0.7891
LOW TIDE	1	278	11/16	68.75	1.777	0.7379	0.7658
	2	206	12/16	75	2.065	0.7625	0.784
	3	202	10/16	62.50	1.695	0.7941	0.8008
	4	375	14/16	87.5	2.193	0.7182	0.7189
	5	303	13/16	81.25	2.1	0.7164	0.7232


**Table # 6 ABUNDANCE OF PHYTOPLANKTON SUBSURFACE SAMPLING STATIONS IN DPT HARBOUR AREA, NEAR BY CREEKS DURING SPRING TIDE IN JUNE,2021**

Tide	Surface	No of Sampling location	Group of phytoplankton	Phytoplankton Group range Units/L	Genera or species /total Phyto plankton	Taxon Diversity % (Group level)
HIGH TIDE	Sub surface	6	DIATOMS	112-214	14/15	93.33
			DINO FLAGELLATES	0-2	1/15	6.67
			TOTAL PHYTO PLANKTON	112-216	15	-
LOW TIDE	Sub surface	5	DIATOMS	147-171	14/15	93.33
			DINO FLAGELLATES	0-2	1/15	6.67
			TOTAL PHYTO PLANKTON	147-172	15	-


**Table # 7 ABUNDANCE OF PHYTOPLANKTON SUBSURFACE SAMPLING STATIONS IN DPT HARBOUR AREA, NEAR BY CREEKS DURING NEAP TIDE IN JUNE, 2021**

Tide	Surface	No of Sampling location	Group of phytoplankton	Phytoplankton Group range Units/L	Genera or species /total Phyto plankton	Taxon Diversity % (Group level)
HIGH TIDE	Sub surface	6	DIATOMS	72-291	15/16	93.75
			DINO FLAGELLATES	0-2	1/16	6.25
			TOTAL PHYTO PLANKTON	72-293	16	-
LOW TIDE	Sub surface	5	DIATOMS	202-374	15/16	93.75
			DINO FLAGELLATES	0-1	1/16	6.25
			TOTAL PHYTO PLANKTON	202-375	16	-

**Taxon Diversity % of Phytoplankton during High tide and Low tide period during spring tide**



**Taxon Diversity % of Phytoplankton during High tide and Low tide period during Neap tide**



**ZOOPLANKTON POPULATION:**

For the evaluation of the Zooplankton population in DPT harbour area and within the immediate surroundings of the port sampling was conducted from 6 sampling locations (3 in harbour area and two in Nakti creek and one in Khoricreek ) during high tide period and low tide period of spring tide and Neap tide in June 2021 . The Zooplankton community of the sub surface water in the harbour and nearby creeks during spring tide was represented by mainly four groups, Tintinids, Copepods, Foraminiferans and larval forms of Crustacea, Molluscans. The Zooplankton

community of the sub surface water in the harbour and nearby creeks during neap tide was represented by mainly five groups, Tintinids, Copepods, Arrow worms, Mysids and larval forms of Crustacea and Polychaetes.

Zooplankton of the sampling stations at sub surface layer in the DPT harbour area and nearby creek was varying from  $61-138 \times 10^3$  N/ m<sup>3</sup> during high tide and  $78-112 \times 10^3$  N/ m<sup>3</sup> during low tide of Spring Tide period. Zooplankton of the sampling stations at sub surface layer in the DPT harbour area and nearby creek was varying from  $47-176 \times 10^3$  N/ m<sup>3</sup> during high tide and  $80-157$  N/ L during low tide of Neap Tide period.

### Species Richness Indices and Diversity Indices:

#### Margalef's diversity index (Species Richness) S

At the organismal level, the most widely used biodiversity measures are those based on the number of species present, perhaps adjusted for the number of individuals sampled, Here Margalef's Species richness index (d), or indices that describe the evenness of the distribution of the numbers of individuals among species, are derived.

Margalef's diversity index (Species Richness) S of Zooplankton communities in the stations Kandla Harbour region and nearby creeks was varying from 2.563-3.067 with an average of 2.804 during the sampling conducted in High tide period. Margalef's diversity index (Species Richness) S of Zooplankton communities varying from 2.136-2.875 with an average of 2.485 during the sampling conducted in low tide period during Spring tide.

Margalef's diversity index (Species Richness) S of Zooplankton communities in the Kandla Harbour region and nearby creeks sampling stations was varying from 3.610-4.53 with an average of 4.016 during the sampling conducted in high tide and varying from 2.755-4.747 with an average of 3.779 during the sampling conducted in low tide during Neap tide period.

#### Shannon-Wiener's index:

Shannon-Wiener's Index (H) of Zooplankton communities in the sampling stations in Kandla Harbour region and nearby creeks was in the range of 0.912-1.017 ( $H'(\log_{10})$ ) between selected sampling stations with an average value of 0.947 ( $H'(\log_{10})$ ) during high tide period of spring tide.

Shannon-Wiener's Index (H) of Zooplankton communities in the sampling stations in Kandla Harbour region and nearby creeks was in the range of 0.872-0.939 ( $H'(\log_{10})$ ) between selected sampling stations with an average value of 0.908 ( $H'(\log_{10})$ ) during consecutive low tide period.

Shannon-Wiener's Index (H) of Zooplankton communities in the sampling stations in Kandla Harbour region and nearby creeks was in the range of 0.962-1.143 ( $H'(\log_{10})$ ) between selected

sampling stations with an average value of 1.071 ( $H'(\log_{10})$ ) during high tide period of Neap tide . Shannon-Wiener's Index (H) of Zooplankton communities in the sampling stations in Kandla Harbour region and nearby creeks was in the range of 0.952-1.168( $H'(\log_{10})$ ) between selected sampling stations with an average value of 1.051 ( $H'(\log_{10})$ ) during consecutive low tide period .Typical values are generally between 1.5 and 3.5 in most ecological studies, and the index is rarely greater than 4. The Shannon-Wiener's index increases as both the richness and the evenness of the community increase. This result indicates that diversity of Zooplankton of Kandla Harbour region and nearby creeks stations is slightly high with very minimum diverse population but very few opportunist organisms are really well adapted to this environment and thrive better than other species.

#### Simpson's diversity index:

Simpson's index (D) is a measure of diversity, which takes into account both species richness, and an evenness of abundance among the species present. The Simpson index is one of the meaningful and robust biodiversity measures available. (Magurran, 2004).

Simpson diversity index (1-D) of Zooplankton communities was below 0.9 at all sampling stations in the Kandla Harbour region and nearby creeks during high tide and low tide of spring tide period except few during high tide period, which was varying from 0.838-0.904 between selected sampling stations with an average of 0.862 during high tide period and was varying from 0.838-0.865 with an average value of 0.849 between selected sampling stations during low tide

Simpson diversity index (1-D) of Zooplankton communities was below 0.9 at all sampling stations in the Kandla Harbour region and nearby creeks during high tide and except one during high tide of Neap tide, which was varying from 0.853-0.905 between selected sampling stations with an average of 0.886 during high tide period and was varying from 0.840- 0.909 with an average value of 0.881 between selected sampling stations during consecutive low tide

This low species diversity suggests a relatively low number of successful species in this habitat. Environment is quite stressful with relatively few ecological niches and only few organisms are really well adapted to that environment. Any change in the environment would probably have quite serious effects.

**Table # 8 ZOOPLANKTON VARIATION IN ABUNDANCE AND DIVERSITY IN SUB SURFACE SAMPLING STATIONS IN DPT HARBOUR AREA ,NEAR BY CREEKS DURING SPRING TIDE IN JUNE,2021**

Tide	Sampling Station	Abundance In No / m <sup>3</sup>	No of Species/groups observed /total species/group	% of diversity	Margalef's diversity index (Species Richness S)	Shannon Weiner index H (log <sub>10</sub> )	Diversity Index (Simpson's Index) 1-D
HIGH TIDE	1	96 X10 <sup>3</sup>	15/17	88.24	3.067	1.01	0.8836
	2	77 X10 <sup>3</sup>	13/17	76.47	2.763	0.9118	0.8506
	3	92 X10 <sup>3</sup>	14/17	82.35	2.875	0.9144	0.8385
	4	138 X10 <sup>3</sup>	14/17	82.35	2.638	0.9177	0.8445
	5	108 X10 <sup>3</sup>	13/17	76.47	2.563	0.9144	0.852
	6	61 X10 <sup>3</sup>	13/17	76.47	2.919	1.017	0.9038
LOW TIDE	1	78 X10 <sup>3</sup>	11/17	64.70	2.295	0.8723	0.8382
	2	92 X10 <sup>3</sup>	14/17	82.35	2.875	0.9395	0.8538
	3	105 X10 <sup>3</sup>	12/17	70.58	2.364	0.8972	0.8443
	4	112 X10 <sup>3</sup>	14/17	82.35	2.755	0.9159	0.8468
	5	108 X10 <sup>3</sup>	11/17	64.70	2.136	0.9189	0.8654

**Table # 9 ZOOPLANKTON VARIATION IN ABUNDANCE AND DIVERSITY IN SUB SURFACE SAMPLING STATIONS IN DPT HARBOUR AREA ,NEAR BY CREEKS DURING NEAP TIDE IN JUNE,2021**

Tide	Sampling Station	Abundance In No / m <sup>3</sup>	No of Species/groups observed /total species/group	% of diversity	Margalef's diversity index (Species Richness S)	Shannon Weiner index H (log <sub>10</sub> )	Diversity Index (Simpson's Index) 1-D
HIGH TIDE	1	111 X10 <sup>3</sup>	18/26	69.23	3.61	0.9985	0.8526
	2	100 X10 <sup>3</sup>	19/26	73.07	3.909	1.068	0.8846
	3	103 X10 <sup>3</sup>	22/26	84.61	4.531	1.129	0.905
	4	176 X10 <sup>3</sup>	24/26	92.31	4.448	1.143	0.904
	5	155 X10 <sup>3</sup>	21/26	80.77	3.966	1.13	0.9041
	6	47 X10 <sup>3</sup>	15/26	57.69	3.636	0.9622	0.8668
LOW TIDE	1	80 X10 <sup>3</sup>	16/26	61.54	3.423	1	0.8684
	2	103 X10 <sup>3</sup>	17/26	65.38	3.452	0.9526	0.8401
	3	112 X10 <sup>3</sup>	14/26	53.85	2.755	1.005	0.8795
	4	157 X10 <sup>3</sup>	25/26	96.15	4.747	1.168	0.9082
	5	130 X10 <sup>3</sup>	23/26	88.46	4.52	1.131	0.9095


**Table # 10 ABUNDANCE OF ZOOPLANKTON IN SUBSURFACE SAMPLING STATIONS IN DPT HARBOUR AREA ,NEAR BY CREEKS DURING SPRING TIDE IN JUNE,2021**

Tide	Surface	No of Sampling locations	Group of Zooplankton	Abundance of Zooplankton $\times 10^3$ Group Range	Genera or species /total Zooplankton	Taxon Diversity % (Group level)
HIGH TIDE	Sub surface	6	Tintinids	10-16	4/17	23.53
			Copepods	31-72	9/17	52.95
			Foraminiferans	0-4	2/17	11.76
			Larval forms	15-50	2/17	11.76
			TOTAL ZOOPLANKTON NO/L	<b>61-138</b>	<b>17</b>	-
LOW TIDE	Sub surface	5	Tintinids	8-15	4/17	23.53
			Copepods	45-57	9/17	52.95
			Foraminiferans	0-2	2/17	11.76
			Larval forms	25-43	2/17	11.76
			TOTAL ZOOPLANKTON NO/L	<b>78-112</b>	<b>17</b>	-


**Table # 11 ABUNDANCE OF ZOOPLANKTON IN SUBSURFACE SAMPLING STATIONS IN DPT HARBOUR AREA , NEAR BY CREEKS DURING NEAP TIDE IN JUNE,2021**

Tide	Surface	No of Sampling locations	Group of Zooplankton	Abundance of Zooplankton $\times 10^3$ Group Range	Genera or species /total Zooplankton	Taxon Diversity % (Group level)
HIGH TIDE	Sub surface	6	Tintinids	4-15	6/26	23.07
			Copepods	25-98	10/26	38.46
			Mysids	1-2	1/26	3.85
			Arrow worms	1-2	1/26	3.85
			Foraminiferans	0-2	1/26	3.85
			Larval forms	17-59	7/26	26.92
			TOTAL ZOOPLANKTON NO/L	<b>47-176</b>	<b>26</b>	-
LOW TIDE	Sub surface	5	Tintinids	4-15	6/26	23.07
			Copepods	38-85	10/26	38.46
			Mysids	0-2	1/26	3.85
			Arrow worms	0-2	1/26	3.85
			Foraminiferans	0-1	1/26	3.85
			Larval forms	37-52	7/26	26.92
			TOTAL ZOOPLANKTON NO/L	<b>80-157</b>	<b>26</b>	-

Taxon Diversity % of Zooplankton during High tide and Low tide period of Spring tide



Taxon Diversity % of Zooplankton during High tide and Low tide period of Neap tide





**TABLE # 12 SYSTEMATIC ACCOUNT OF PHYTOPLANKTON IN THE SAMPLING LOCATIONS IN OF DPT HARBOUR AREA AND NEARBY CREEKS  
DURINGSPRING TIDE OF JUNE,2021**

GROUP	PHYLUM	CLASS	ORDER	FAMILY	GENUS/SPECIES	#	Relative Abundance
DIATOMS	Bacillariophyta	Coscinodiscophyceae	Thalassiosirales	Thalassiosiraceae	<i>Planktoniellasp</i>	D1	Rare
			Coscinodiscales	Coscinodiscaceae	<i>Coscinodiscus sp.</i>	D2	Abundant
			Triceratiales	Triceratiaceae	<i>Odontellasp</i>	D3	Rare
					<i>Triceratiumsp.</i>	D4	Occasional
			Biddulphiales	Biddulphiaceae	<i>Biddulphiasp</i>	D5	Frequent
			Rhizosoleniales	Rhizosoleniaceae	<i>Rhizosolenia sp.</i>	D6	Occasional
			Chaetocerotales	Chaetocerotaceae	<i>Chaetocerossp</i>	D7	Occasional
			Lithodesmiales	Lithodesmiaceae	<i>Ditylumsp</i>	D8	Dominant
		Thalassiosirales	Thalassiosiraceae	<i>Thalassiosirasp</i>	D9	Rare	
		Bacillariophyceae	Naviculales	Pleurosigmaaceae	<i>Pleurosigmasp</i>	D10	Rare
		Fragilariophyceae	Thalassionematales	Thalassionemataceae	<i>Thalassiothrix sp.</i>	D11	Frequent
					<i>Thalassionema sp.</i>	D12	Rare
			Fragilariales	Fragilariaceae	<i>Fragilariasp</i>	D13	Occasional
					<i>Synedrasp</i>	D14	Frequent
DINO FLAGELLATES	Dinoflagellata / Dinozoa	Dinophyceae	Gonyaulacales	Ceratiaceae	<i>Ceratiumfurca</i>	DF1	Rare

TABLE # 12 SYSTEMATIC ACCOUNT OF PHYTOPLANKTON IN THE SAMPLING LOCATIONS IN OF DPT HARBOUR AREA AND NEARBY CREEKS DURING AND NEAP TIDE OF JUNE,2021

GROUP	PHYLUM	CLASS	ORDER	FAMILY	GENUS/SPECIES	#	Relative Abundance
DIATOMS	Bacillariophyta	Coscinodiscophyceae	Thalassiosirales	Thalassiosiraceae	<i>Planktoniellasp</i>	D1	Occasional
			Coscinodiscales	Coscinodiscaceae	<i>Coscinodiscus sp.</i>	D2	Dominant
			Triceratiales	Triceratiaceae	<i>Triceratiumsp</i>	D3	Occasional
					<i>Odontellasp</i>	D4	Rare
			Biddulphiales	Biddulphiaceae	<i>Biddulphiasp</i>	D5	Abundant
			Rhizosoleniales	Rhizosoleniaceae	<i>Rhizosolenia sp.</i>	D6	Rare
			Chaetocerotales	Chaetocerotaceae	<i>Chaetocerossp</i>	D7	Rare
		Lithodesmiales	Lithodesmiaceae	<i>Ditylumsp</i>	D8	Abundant	
		Bacillariophyceae	Naviculales	Pleurosigmataceae	<i>Pleurosigmasp</i>	D9	Occasional
			Bacillariales	Bacillariaceae	<i>Bacillaria sp.</i>	D10	Rare
		Fragilariophyceae	Thalassionematales	Thalassionemataceae	<i>Thalassiothrix sp.</i>	D11	Frequent
					<i>Thalassionema sp.</i>	D12	Rare
			Fragilariales	Fragilariaceae	<i>Fragilariasp</i>	D13	Rare
					<i>Synedrasp</i>	D14	Frequent
					<i>Asterionellasp</i>	D15	Occasional
DINO FLAGELLATES	Dinoflagellata / Dinozoa	Dinophyceae	Peridinales	Protoperidiniaceae	<i>Protoperidinium sp.</i>	DF1	Rare

**TABLE #13 SYSTEMATIC ACCOUNT OF ZOOPLANKTON FROM THE SAMPLING LOCATIONS IN DPT HARBOUR AREA, AND NEAR BY CREEKS DURING SPRING TIDE OF JUNE,2021**

GROUP	PHYLUM	CLASS	ORDER	FAMILY	GENUS/SPECIES	#	RELATIVE ABUNDANCE
TINTINIDS	PROTOZOA	Spirotrichea	Tintinnida	Tintinnidiidae	<i>Leprotintinnussp.</i>	T1	Occasional
				Codonellidae	<i>Tintinnopsisfailakkaensis</i>	T2	Occasional
	<i>Tintinnopsisgracilis</i>				T3	Occasional	
	<i>Tintinnopsis radix</i>				T4	Rare	
COPEPODS	ARTHROPODA	Crustacea Sub class copepoda	Calanoida	Paracalanidae	<i>Acrocalanus sp.</i>	C1	Frequent
					<i>Bestiolina sp.</i>	C2	Rare
					<i>Parvocalanus sp.</i>	C3	Occasional
				Eucalanidae	<i>Pareucalanus sp.</i>	C4	Rare
				Clausocalanidae	<i>Clausocalanus sp.</i>	C5	Occasional
				Temoridae	<i>Temora sp.</i>	C6	Rare
			Cyclopoida	Oithonidae	<i>Oithona sp.</i>	C7	Abundant
			Harpacticoida	Ectinosomatidae	<i>Microsetellasp.</i>	C8	Frequent
			Poecilostomatatoida	Oncaeidae	<i>Oncaea sp.</i>	C9	Rare
CRUSTACEAN LARVAE	ARTHROPODA (CRUSTACEA)	Copepoda			Nauplius larvae of Copepods	L1	Dominant
BIVALVE LARVAE	MOLLUSCA	Pelecypoda			Veliger larvae of Bivalves	L2	Occasional
FORAMINIFERA	FORAMINIFERA	Globothalamea	Rotaliida	Globigerinidae	<i>Globigerina sp.</i>	F1	Rare
				Rotalliidae	<i>Rotalia sp.</i>	F2	Rare

**TABLE # 13 SYSTEMATIC ACCOUNT OF ZOOPLANKTON FROM THE SAMPLING LOCATIONS IN DPT HARBOUR AREA, AND NEAR BY CREEKS DURING NEAP TIDE OF JUNE,2021**

GROUP	PHYLUM	CLASS	ORDER	FAMILY	GENUS/SPECIES	#	RELATIVE ABUNDANCE
TINTINIDS	PROTOZOA CILIOPHORA	Spirotrichea	Tintinnida	Tintinnidiidae	<i>Leprotintinnus</i> sp.	T1	Rare
				Codonellidae	<i>Tintinnopsisaccuminata</i>	T2	Occasional
					<i>Tintinnopsisfailakkaensis</i>	T3	Occasional
					<i>Tintinnopsisgracilis</i>	T4	Rare
					<i>Tintinnopsisradix</i>	T5	Rare
				Codonellopsidae	<i>Codonellopsis</i> sp.	T6	Rare
COPEPODS	ATHROPODA	Crustacea Sub class copepoda	Calanoida	Paracalanidae	<i>Acrocalanus</i> sp.	C1	Abundant
					<i>Parvocalanus</i> sp.	C2	Rare
				Eucalanidae	<i>Pareucalanus</i> sp.	C3	Frequent
					<i>Subeucalanus</i> sp.	C4	Occasional
				Temoridae	<i>Temora</i> sp.	C5	Frequent
				Acartiidae	<i>Acartia</i> sp.	C6	Occasional
			Cyclopoida	Oithonidae	<i>Oithona</i> sp.	C7	Frequent
			Harpacticoida	Ectinosomatidae	<i>Microsetella</i> sp.	C8	Frequent
				Euterpinae	<i>Euterpina</i> sp.	C9	Frequent
			Poicilostomatatoida	Oncaeidae	<i>Oncaea</i> sp.	C10	Rare
ARROW WORMS	CHAETOGNATHA	Sagittoideae	Aphragmophora	Sagittidae	<i>Sagitta</i> sp.	A1	Rare
MYSIDS	ATHROPODA CRUSTACEA	Malacostraca	Mysida, Decapoda	Penaeidae	<i>Metapenaeus</i> sp.	M1	Rare

GROUP	PHYLUM	CLASS	ORDER	FAMILY	GENUS/SPECIES	#	RELATIVE ABUNDANCE
CRUSTACEAN LARVAE	ARTHROPODA (CRUSTACEA)	Copepoda			Nauplius larvae of Copepods	L1	Dominant
POLYCHAETE LARVA	ANNELIDA	Polychaeta			Trochophore larvae	L2	Occasional

## Environmental Monitoring Report Of Deendayal Port Trust, JUNE-2021

### BENTHIC ORGANISMS:

No Benthic organism was observed in the collected sediments by using the Van-veen grabs during the sampling conducted during spring tide period from DPT harbour region and nearby creek except few dead shells. Benthic organisms from the sample collected during Neap tide is represented by mainly Polychaetes, *Pontodrasp. Paronis sp.* and *Phalacophorus sp.* and few Amphipods. The benthic organisms at subtidal region of harbour region and creek varies from 30-100 N/m<sup>2</sup>

**Table # 14 BENTHIC FAUNA IN THE SAMPLING LOCATIONS IN DPT HARBOUR AREA CREEKS DURING NEAP TIDE IN JUNE, 2021**

Benthic fauna	ABUNDANCE IN NO/M <sup>2</sup> DIFFERENT SAMPLING STATIONS						
	REPRESENTATION BY GROUP						
	DPT HARBOUR			CREEKS			
POLYCHATES	KPT-1	KPT-2	KPT-3	KPT-4	KPT-5	KPT-6	
Family : Iospilidae <i>Pontodora sp.</i>	10	NS	0	20	30	NS	
Family : Syllidae <i>Syllis sp.</i>	20	NS	10	30	10	NS	
Family Glyceridae <i>Glycerasp.</i>	30	NS	0	0	0	NS	
<b>Total Polychaetes N/M<sup>2</sup></b>		NS				NS	
<b>Un identified Nematode worms</b>		NS		0		NS	
<b>Amhipods</b> Un identified	0	NS	0	50	0	NS	
<b>TOTAL Benthic Fauna NUMBER/ M<sup>2</sup></b>	60	NS	10	100	30	NS	

NS : No sample

## 7. Meteorological Data

Automatic Weather station have been installed in SevaSadan -3 at the Deendayal Port which records the data on Temperature (°C), Humidity (%), Wind (mph), Dew Point (°C), Wind Direction (°), Pressure, Solar radiation, heat Index and UVI.

### Temperature

The mean day time temperature for Deendayal Port was 29.77 °C. The day-time maximum temperature was 34.1 °C. The mean night time temperature was 32.53 °C. The minimum mean night time temperature recorded was 28.2 °C.

### Air Pressure

The mean absolute air pressure for the month of June was 1004.93 hpa, whereas the mean relative pressure was 1005.03 hpa. The maximum absolute air pressure recorded for the month of June was 1008.9 hpa.

### Heat Index

The mean day-time heat index for the month of June was 35.20 °C. The maximum heat index recorded was 44°C.

### Solar Radiation

The mean Solar Radiation in June was 208.28 w/m<sup>2</sup>. The maximum solar radiation recorded in the month of June was 654.8 w/m<sup>2</sup>.

### Humidity

The mean day-time humidity was 76.42 % for the month of June and mean night time humidity was 65.97%. Maximum humidity recorded during day-time was 84.0 % and maximum humidity recorded during night-time was 82.0%.

### Wind Velocity and Wind Direction

The mean wind velocity for the entire month of June was 9.72 km/hour (i.e. 2.7 mtr/sec). Maximum wind velocity recorded was 46.8 Km/hr (13 mtr/sec). The wind direction was mostly S to SW.

### Conclusive Summary and Remedial measures Suggested

- The AAQ monitoring at six locations of Deendayal Port indicates that the mean  $PM_{10}$  values at four locations viz. Coal storage area, Marine Bhavan and Oil Jetty area were found above the permissible standards ( $100 \mu\text{g}/\text{m}^3$ ) and  $PM_{2.5}$  was above permissible limits at Coal storage location (Limit  $60 \mu\text{g}/\text{m}^3$ ).
- Drinking water at all the twenty locations was found potable and was within permissible limits of BIS standards (IS 10500).
- Noise quality was also within the set permissible standards of an Industrial Area. The noise level observed during day time was  $>75 \text{ dB (A)}$  and at night time was  $>70 \text{ dB (A)}$  during the entire monitoring period.
- The sewage treated water of Deendayal Port Colony (Gopalpuri) was in line with the standards set by the Gujarat Pollution Control Board. The STP at Deendayal Port is not fully operational and STP at Vadinar Port was found non-operational.

### Reasons for higher Values of $PM_{10}$

- Large amount of coal is handled at Berth No. 6, 7, 8 and 9. The unloading of coal directly in the truck, using grabs cause coal to spread in air as well as coal dust to fall on ground. This settled coal dust again mixes with the air while trucks travel through it.
- Also, the coal laden trucks are not always covered with tarpaulin sheets, and these results in spillage of coal from trucks/dumpers during its transit from vessel to yard or storage site. This also increased PM values around marine Bhavan & Coal storage area.



### **Remedial Measures**

The values of PM<sub>10</sub> during the month of June, 2021 were observed beyond the permissible limit at four locations mentioned above. Given below are the remedial measures suggest to minimize the Air pollution at Deendayal Port.

- Guidelines for Coal Handling by GPCB should be strictly followed. (<http://gpcb.gov.in/pdf/coal-handling-guidelines.pdf>)
- Sewage Treatment Plan at Vadinar Port is not working. Hence, it is recommended to commission the sewage treatment plant at Vadinar immediately.
- Except for the higher values of PM<sub>10</sub> at Coal storage site, Oil Jetty, Tuna Port and Marine Bhavan locations, the monitoring results for the present month suggest that the overall Environment Quality of Deendayal Port is satisfactory.

**SOURCE OF LITERATURE AND ADDITIONAL REFERENCE FOR ECOLOGICAL STUDY**

- 1) ALBERT WEST PHAL (1976) Protozoa Blackwell , London
- 2) BANERJEE R.K. (1989) Heavy metals and Benthic foraminiferal distribution along Bombay coast India. Studies in benthic foraminifera. *Tokyo University Press* Tokyo pp 151-157
- 3) Banse K (1995) Zooplankton: Pivotal role in the control of ocean production: I. Biomass and production. *ICES J Mar Sci* 52: 265–277.
- 4) Beaugrand G, and Ibanez F (2004) Monitoring marine plankton ecosystems. II: long-term changes in North Sea calanoid copepods in relation to hydroclimatic variability. *Inter Res Mar Ecol Prog Ser* 284:35-47.
- 5) DAY F. (1889) The fauna of British India Ceylon and Burma- Fishes Vol-1- Vol-2 *Taylor and Francis* London
- 6) DESIKACHARY V. (1989) Atlas of diatoms, Madras Science Foundation
- 7) DESIKACHARY V. (1959) Cyanophyta ICAP Monographs on Algae *Indian Council of Agricultural research* New Delhi
- 8) FAIZAYOUSIF AL-YAMANI & MARIA A. SABUROVA (2010) illustrative guide on the flagellates of Intertidal soft sediment *Kuwait Institute for scientific Research* Kuwait
- 9) FAIZAYOUSIF AL-YAMANI, VALERIYSKRYABIN, ALEKSANDRA GUBANOVA, SERGEY KHVOROV AND IRINA PRUSOVA (2011), Marine zooplankton Practical guide from North western Arabian gulf Vol-1 and vol-2 *Kuwait Institute for scientific Research* Kuwait
- 10) FAUVEL P. (1953), The fauna of India Annelida - Polychaeta *Indian Press* Allahabad
- 11) Gajbhiye SN, Nair VR, and Desai BN (1984). Diurnal variation of zooplankton in Malad creek, Bombay. *Indian Journal of Marine Science*. 13:75-79.
- 12) HAYWARD P.J AND RYLAND J.S. (1995) Handbook of Marine fauna of north –West Europe *oxford University Press* London
- 13) HIGGINS R.P. HAJAMARTHIEL Eds. (1998) Introduction to the study of Meio Fauna
- 14) HORACE G. BARBER AND ELIZABETH Y. HAWORTH (1981) A guide to the Morphology of DIATOMS FRUSTULES.
- 15) INGRAM HENDEY (1964) An introductory account of smaller Algae of British coastal waters part-V. Bacillariophyceae
- 16) JOHN H. WICKSTEAD (1965) an Introduction to the study of Tropical Plankton . *Hutchinson Tropical Monographs*
- 17) JOYOTHIBABU, R. MADHU, N.V. MAHESHWARAN, P.A., NAIR K.K.C., VENUGOPL, P. BALASUBRAMANIAN T. (2005) Dominance of Dinoflagellates in micro zooplankton communities in the oceanic region Bay of Bengal and Andaman sea *Current science* vol.84. 10<sup>th</sup> May 2003
- 18) KASTURIRANGAN L.R. (1963) A key for the identification of the Common Planktonic Copepoda of Indian Coastal water
- 19) Kusum KK, Vineetha G, Raveendran TV, Nair VR, Muraleedharan KR, Achuthankutty CT and Joseph T (2014) Chaetognath community and their responses to varying environmental factors in the northern Indian ocean. *J Plankton Res* 36(4): 1146- 1152.
- 20) Lalli CM and Parsons TR (1997) Biological Oceanography: An Introduction. DOI <https://doi.org/10.1016/B978-0-7506-3384-0.X5056-7>.
- 21) Madhupratap M (1978) Studies on ecology of zooplankton of Cochin backwaters. *Mahasagar Bull Nat Inst Oceanogr* 11: 45-56.
- 22) Madhupratap M (1979) Distribution, community structure and species succession of copepods from Cochin Backwaters. *Indian J Mar Sci* 8: 1-8.
- 23) Madhupratap M (1987) Status and strategy of zooplankton of tropical Indian estuaries: A review. *Bull Plank Soc Jpn* 34: 65-81.

**Environmental Monitoring Report Of Deendayal Port Trust, JUNE-2021**

---

- 24) Madhupratap M (1999) Free living copepods of the Arabian Sea, Distribution and Research Perspectives. I J Mar Sci 146-149.
- 25) Madhupratap M and Haridas P (1986) Epipelagic calanoid copepods of the northern Indian Ocean. OceanologicaActa 9(2):105-117.
- 26) MANAL AL-KANDARI, FAIZA Y. AL-YAMANI , KHOLOOD AL-RIFAIE ( 2009) Marine phytoplankton Atlas of Kuwait's water *Kuwait Institute for scientific Research*
- 27) MPEDA (1998) Commercial Fishes and shell fishes of India
- 28) NEWEL G.E. & NEWELL R.C. (1963) Marine plankton a Practical Guide Hutchinson Educational
- 29) NIGAM R.C. AND CHATURVEDIS.K. (2000) Foraminiferal Study from KharoCreek , Kachchh ( Gujarat) North west coast of *India. Indian Journal of marine science* Vol.29 133-189
- 30) OLAV GIERE (1993) Meio benthology , Microscopic Fauna in Aquatic Sediments m Springer London
- 31) PERRAGALLO( 1965) Diatomees marines de franceA. *Asher & Co.* Amsterdam
- 32) Robert P.. Higgins (Eds.), (1985) An introduction to the study of Meuiu fauna Smithsons Institution press Washington DC
- 33) STERRER W. STERRERC.S Eds. Marine fauna and flora of Bermuda A systematic Guide to the identification of Marine Organisms. *John Wiely and Sons*New York
- 34) Suresh Gandhi . M. (2009) Distribution of certain ecological parameters and Foraminiferaldistribution in the depositional environment of Pak strait east coast of India .*Indian J. of Marine Science* Vol.33 pp 287-295
- 35) Venktaraman (1993 A systematic account of some south Indian diatoms . Proceeding of Indian Academy of Science Vol.X No.6 Sec.B.

\*\*\*\*\*

# ENVIRONMENTAL MONITORING REPORT FOR DEENDAYAL PORT TRUST



**REPORT NO.** : DCPL/DPT/20-21/15  
**Month** : July 2021  
**Issue No** : 01  
**Revision No** : 00  
**Prepared by** : DETOX CORPORATION PVT. LTD., SURAT

## TABLE OF CONTENTS

<b>Sr. No.</b>	<b>Particulars</b>	<b>Page No.</b>
<b>1</b>	<b>Ambient Air Quality Monitoring.....</b>	<b>1 - 17</b>
<b>2</b>	<b>Drinking Water Quality Monitoring.....</b>	<b>18 - 28</b>
<b>3</b>	<b>Noise Monitoring.....</b>	<b>29</b>
<b>4</b>	<b>Soil Monitoring.....</b>	<b>30 - 31</b>
<b>5</b>	<b>Sewage Treatment Plant Monitoring.....</b>	<b>32 - 38</b>
<b>6</b>	<b>Marine Water Monitoring.....</b>	<b>39 - 81</b>
<b>7</b>	<b>Meteorological Observations.....</b>	<b>82</b>
<b>8</b>	<b>Conclusive Summary &amp; Remedial Measures .....</b>	<b>83-84</b>
	<b>References.....</b>	<b>85- 86</b>

## **Introduction**

Monitoring of various environmental aspects of the Deendayal port by M/s Detox Corporation Pvt. Ltd. has been carried out through collection of samples, analysis of the same, comparing results with respect to the national standards and any other relevant standards by GBCB/CPCB/MoEF to identify non conformities in the Environment of the Deendayal Port. The results shall address the identified impacts and suggest measures to minimize the environmental impact due to various operations at Deendayal Port.

The environmental monitoring is carried out as per the Environment Management and Monitoring Plan submitted by Detox Corporation Pvt. Ltd.

## **1. Ambient Air Quality Monitoring**

As per the Environmental Monitoring Plan of Deendayal Port Trust, Air monitoring was carried out at six identified locations at Deendayal Port and two locations at Vadinar Port.

### **1.1 Air Quality Monitoring Methodology**

Air quality is measured in all the stations, for 24 hour for Total Suspended Particulate Matter (TSPM), PM<sub>10</sub>, PM<sub>2.5</sub>, SO<sub>2</sub>, NO<sub>x</sub>, NH<sub>3</sub> & Benzene, and Grab-sampling for CO & CO<sub>2</sub> measurements. The Air samplers are operated for a period of 24 hours and after a continuous operation of 8 hours of the sampler, the reagents were replaced to obtain 3 samples per day for each parameter namely, SO<sub>2</sub>, NO<sub>x</sub>. The EPM 2000 filter paper and PTFE Membrane bound filter paper are used for a period of 24 hours to obtain one sample each of TSPM, PM<sub>10</sub> & PM<sub>2.5</sub>.

The AAQ samples are collected twice a week from all the eight locations as per the EMP.

## 1.2 Results

The ambient air quality monitoring data for six stations, viz. Marine Bhavan, Oil Jetty, Port Colony, Gopalpuri Hospital, Tuna Port and Nr. Coal Storage Area for the month of July 2021 are given in Tables 1A to 6B. The ambient air quality monitoring data for two stations at Vadinar (Nr. Admin Building & Nr. Signal Building) are given in Tables 7A to 8B.

### Location 1: Marine Bhavan (AL1)

Table 1 : Results of Air Pollutant Concentration at Marine Bhavan										
Parameter	Date	TSPM [µg/m <sup>3</sup> ]	PM10 [µg/m <sup>3</sup> ]	PM2.5 [µg/m <sup>3</sup> ]	SO <sub>2</sub> [µg/m <sup>3</sup> ]		NO <sub>x</sub> [µg/m <sup>3</sup> ]		NH <sub>3</sub> [µg/m <sup>3</sup> ]	
Sampling Period	-	24hr	24hr	24hr	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
NAAQMS limit	-	NS	100 µg/m <sup>3</sup>	60 µg/m <sup>3</sup>		80 µg/m <sup>3</sup>		80 µg/m <sup>3</sup>		400 µg/m <sup>3</sup>
AL1 - 1	01.07.2021	417	302	96	3.08	2.78	59.07	48.70	13.53	15.32
					3.52		55.26		16.08	
					1.76		31.76		16.34	
AL1 - 2	05.07.2021	875	776	40	6.59	5.71	57.16	52.51	15.83	12.34
					5.71		47.64		10.21	
					4.84		52.72		10.98	
AL1 - 3	09.07.2021	769	693	11	8.79	7.62	33.03	34.30	13.79	13.87
					8.35		31.76		13.53	
					5.71		38.11		14.30	
AL1 - 4	14.07.2021	267	257	31	2.64	4.69	14.61	24.98	19.15	10.21
					5.28		28.58		5.11	
					6.15		31.76		6.38	
AL1 - 5	16.07.2021	234	143	8	10.55	10.11	13.34	20.11	9.19	12.34
					13.19		22.87		14.04	
					6.59		24.14		13.79	
AL1 - 6	21.07.2021	314	257	202	4.84	2.93	13.34	13.76	7.66	12.17
					1.32		15.24		12.25	
					2.64		12.70		16.59	
AL1 - 7	23.07.2021	387	256	163	1.76	3.08	38.11	28.16	16.34	61.10
					3.52		27.31		13.79	
					3.96		19.05		153.17	
AL1 - 8	27.07.2021	471	299	163	6.15	6.01	12.70	16.09	137.85	53.78
					6.59		18.42		10.98	
					5.28		17.15		12.51	
<b>Monthly Average</b>		467	373	89		5.37		29.83		23.89
<b>Standard Deviation</b>		234	230	78		2.58		14.43		20.85

NS: Not Specified

<b>Table 1B : Results of Air Pollutant Concentration at Marine Bhavan</b>					
<b>Parameter</b>	<b>Date</b>	<b>C<sub>6</sub>H<sub>6</sub> [µg/m<sup>3</sup>]</b>	<b>HC* ppm</b>	<b>CO [mg/m<sup>3</sup>]</b>	<b>CO<sub>2</sub> [ppm ]</b>
<b>Sampling Period</b>		<b>8 hr</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>
<b>NAAQMS limit</b>		<b>5.0 µg/m<sup>3</sup></b>	<b>NS</b>	<b>4.0 mg/m<sup>3</sup></b>	<b>NS</b>
AL1 – 1	01.07.2021	1.13	BDL	1.46	508
AL1 – 2	05.07.2021	1.1	BDL	1.52	489
AL1 – 3	09.07.2021	1.04	BDL	1.36	512
AL1 – 4	14.07.2021	1.14	BDL	1.48	562
AL1 – 5	16.07.2021	1.12	BDL	1.52	496
AL1 - 6	21.07.2021	1.05	BDL	1.48	485
AL1 – 7	23.07.2021	1.04	BDL	1.78	508
AL1 – 8	27.07.2021	1.1	BDL	1.69	495
<b>Monthly Average</b>		<b>1.09</b>	<b>-</b>	<b>1.54</b>	<b>507</b>
<b>Standard Deviation</b>		<b>0.04</b>	<b>-</b>	<b>0.13</b>	<b>24</b>

\* NMHC- Non- Methane Hydrocarbons

BDL- Below Detection Limit (Detection Limit – NMHC: 0.5ppm)

NS -Not Specified

At Marine Bhavan, the overall values of TSPM, PM<sub>10</sub>, PM<sub>2.5</sub>, SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> is attributed mainly by motor vehicle emission produced from various types of automobiles (both diesel and petrol driven). Moreover, the loading and unloading of Food Grains and Timber at Jetty no. 1 and 2 also contributes to the high levels of TSPM and PM<sub>10</sub>. The mean TSPM value at Marine Bhavan was 467 µg/m<sup>3</sup>, The mean PM<sub>10</sub> values were 373.0 µg/m<sup>3</sup>, which is above the permissible limit. PM<sub>2.5</sub> values were above the permissible limit (mean = 89 µg/m<sup>3</sup>). The average values of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were within the permissible limit. The average values of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were 5.37 µg/ m<sup>3</sup>, 29.83 µg/ m<sup>3</sup> & 23.89 µg/ m<sup>3</sup> respectively. These were within the permissible limit.

The levels of Benzene, Hydrocarbons (HC) and CO were within the permissible limit at Marine Bhavan. The mean Benzene concentration was 1.09 µg/m<sup>3</sup>, well below the permissible limit of 5.0 µg/m<sup>3</sup>. HC's were below the detectable limit and Carbon Monoxide concentration was 1.54 mg/m<sup>3</sup>, well below the permissible limit of 4.0 mg/m<sup>3</sup>.



**Location 2: Oil Jetty (AL2)**

Table 2 : Results of Air Pollutant Concentration at Oil Jetty										
Parameters	Date	TSPM [µg/m <sup>3</sup> ]	PM10 [µg/m <sup>3</sup> ]	PM2.5 [µg/m <sup>3</sup> ]	SO <sub>2</sub> [µg/m <sup>3</sup> ]		NO <sub>x</sub> [µg/m <sup>3</sup> ]		NH <sub>3</sub> [µg/m <sup>3</sup> ]	
					8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
Sampling Period	-	24hr	24hr	24hr	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
NAAQMS limit	-	NS	100 µg/m <sup>3</sup>	60 µg/m <sup>3</sup>	-	80 µg/m <sup>3</sup>	-	80 µg/m <sup>3</sup>	-	400 µg/m <sup>3</sup>
AL2 – 1	01.07.2021	265	392	127	2.20	1.47	55.26	45.94	7.66	9.96
					0.88		52.72		10.98	
					1.32		29.85		11.23	
AL2 – 2	05.07.2021	812	737	42	0.88	2.93	44.46	47.64	13.53	13.96
					2.64		47.64		13.53	
					5.28		50.81		14.81	
AL2 – 3	09.07.2021	807	707	35	5.28	8.35	17.15	24.56	7.91	10.98
					10.11		24.77		12.76	
					9.67		31.76		12.25	
AL2 – 4	14.07.2021	602	280	5	3.08	3.37	19.05	20.33	15.32	16.59
					2.64		17.15		16.08	
					4.40		24.77		18.38	
AL2 – 5	16.07.2021	578	539	6	4.40	4.10	16.51	17.15	6.13	6.98
					3.52		17.15		5.11	
					4.40		17.78		9.70	
AL2 – 6	21.07.2021	867	772	10	4.84	5.13	29.22	28.16	10.98	13.36
					4.40		32.39		12.76	
					6.15		22.87		16.34	
AL2 – 7	23.07.2021	244	194	76	2.20	2.20	23.50	26.25	13.79	15.40
					1.76		26.68		15.83	
					2.64		28.58		16.59	
AL2 – 8	27.07.2021	448	350	76	6.15	6.45	23.50	18.84	12.00	12.59
					7.03		14.61		13.02	
					6.15		18.42		12.76	
<b>Monthly Average</b>		578	496	47		4.25		28.61		12.48
<b>Standard Deviation</b>		244	224	43		2.30		11.83		3.10

NS: Not Specified

<b>Table 2B : Results of Air Pollutant Concentration at Oil Jetty</b>					
<b>Parameter</b>	<b>Date</b>	<b>C<sub>6</sub>H<sub>6</sub> [µg/m<sup>3</sup>]</b>	<b>HC* ppm</b>	<b>CO [mg/m<sup>3</sup>]</b>	<b>CO<sub>2</sub> [ppm ]</b>
<b>Sampling Period</b>		<b>8 hr</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>
<b>NAAQMS limit</b>		<b>5.0 µg/m<sup>3</sup></b>	<b>NS</b>	<b>4.0 mg/m<sup>3</sup></b>	<b>NS</b>
AL2 -1	01.07.2021	1.12	BDL	1.76	512
AL2 -2	05.07.2021	1.16	BDL	1.85	498
AL2 -3	09.07.2021	1.06	BDL	1.77	506
AL2 -4	14.07.2021	1.15	BDL	1.54	489
AL2 – 5	16.07.2021	1.14	BDL	1.78	490
AL2 – 6	21.07.2021	1.19	BDL	1.62	506
AL2 -7	23.07.2021	1.72	BDL	1.82	515
AL2 – 8	27.07.2021	1.58	BDL	1.78	510
<b>Monthly Average</b>		<b>1.27</b>	<b>-</b>	<b>1.74</b>	<b>503</b>
<b>Standard Deviation</b>		<b>0.24</b>	<b>-</b>	<b>0.11</b>	<b>10</b>

\* NMHC- Non- Methane Hydrocarbons

BDL- Below Detection Limit (Detection Limit – NMHC : 0.5ppm)

NS- Not Specified

The overall values of TSPM, PM<sub>10</sub>, PM<sub>2.5</sub>, SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> at Oil Jetty area was mainly by motor vehicle emission produced from various types of vehicles Oil Jetty Area. The mean TSPM values at Oil Jetty were 578 µg/m<sup>3</sup> The mean PM<sub>10</sub> values were 496 µg/m<sup>3</sup>, which is above the permissible limit. PM<sub>2.5</sub> values were below the permissible limit (mean = 47 µg/m<sup>3</sup>). The average values of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were within the permissible limit; The mean concentration of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were 4.25 µg/m<sup>3</sup>, 28.61 µg/m<sup>3</sup> and 12.48 µg/m<sup>3</sup> respectively.

The levels of Benzene, Hydrocarbons (HC) and CO were within the permissible limit at Oil Jetty. The mean Benzene concentration was 1.27 µg/m<sup>3</sup>. Well below the permissible limit of 5.0 µg/m<sup>3</sup>. , HC's were below the detectable limit and Carbon Monoxide concentration was 1.74 mg/m<sup>3</sup>, well below the permissible limit of 4.0 mg/m<sup>3</sup>.

**Location 3: Kandla Colony – Estate Office (AL-3)**

<b>Table 3 : Results of Air Pollutant Concentration at Estate Office</b>										
<b>Parameters</b>	<b>Date</b>	<b>TSPM [µg/m3]</b>	<b>PM10 [µg/m3]</b>	<b>PM2.5 [µg/m3]</b>	<b>SO2 [µg/m3]</b>		<b>NOx [µg/m3]</b>		<b>NH3 [µg/m3]</b>	
<b>Sampling Period</b>	-	<b>24hr</b>	<b>24hr</b>	<b>24hr</b>	<b>8 hr</b>	<b>24hr (Avg.)</b>	<b>8 hr</b>	<b>24hr (Avg.)</b>	<b>8 hr</b>	<b>24hr (Avg.)</b>
<b>NAAQMS limit</b>	-	<b>NS</b>	<b>100 µg/m3</b>	<b>60 µg/m3</b>	-	<b>80 µg/m3</b>	-	<b>80 µg/m3</b>	-	<b>400 µg/m3</b>
<b>AL3 – 1</b>	01.07.2021	168	153	55	3.52	3.81	20.96	21.38	14.30	10.21
					5.28		27.31		9.45	
					2.64		15.88		6.89	
<b>AL3 – 2</b>	05.07.2021	467	373	21	3.52	2.49	21.60	23.50	14.04	15.91
					1.32		18.42		15.83	
					2.64		30.49		17.87	
<b>AL3 – 3</b>	09.07.2021	297	139	37	3.08	4.98	23.50	24.77	9.19	7.66
					6.15		29.85		6.38	
					5.71		20.96		7.40	
<b>AL3 – 4</b>	14.07.2021	292	121	80	4.84	5.86	21.60	19.27	14.55	72.33
					5.71		18.42		186.35	
					7.03		17.78		16.08	
<b>AL3 – 5</b>	16.07.2021	629	566	96	17.58	10.11	17.15	14.82	13.53	12.00
					7.91		15.24		9.70	
					4.84		12.07		12.76	
<b>AL3 – 6</b>	21.07.2021	721	668	57	3.96	2.49	6.99	14.82	20.42	18.98
					1.32		15.88		21.44	
					2.20		21.60		15.06	
<b>AL3 – 7</b>	23.07.2021	490	406	51	2.64	2.49	22.87	23.29	11.23	11.91
					3.08		19.69		9.70	
					1.76		27.31		14.81	
<b>AL3 – 8</b>	27.07.2021	640	500	51	1.76	3.66	20.96	18.00	11.23	10.04
					4.40		17.15		8.17	
					4.84		15.88		10.72	
<b>Monthly Average</b>		463	366	56		4.49		19.98		19.88
<b>Standard Deviation</b>		196	210	23		2.59		3.89		21.49

NS: Not Specified

<b>Table 3B : Results of Air Pollutant Concentration at Kandla Port Colony</b>					
<b>Parameter</b>	<b>Date</b>	<b>C<sub>6</sub>H<sub>6</sub> [µg/m<sup>3</sup>]</b>	<b>HC*</b>	<b>CO [mg/m<sup>3</sup>]</b>	<b>CO<sub>2</sub> [ppm ]</b>
<b>Sampling Period</b>		<b>8 hr</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>
<b>NAAQMS limit</b>		<b>5.0 µg/m<sup>3</sup></b>	<b>NS</b>	<b>4.0 mg/m<sup>3</sup></b>	<b>NS</b>
AL3 -1	01.07.2021	1.12	BDL	1.78	510
AL3 -2	05.07.2021	1.22	BDL	1.84	526
AL3 -3	09.07.2021	1.16	BDL	1.96	520
AL3 -4	14.07.2021	1.26	BDL	1.88	542
AL3 -5	16.07.2021	1.18	BDL	1.78	533
AL3 -6	21.07.2021	1.26	BDL	1.6	525
AL3 -7	23.07.2021	1.21	BDL	1.58	542
AL3 -8	27.07.2021	1.11	BDL	1.78	502
<b>Monthly Average</b>		<b>1.19</b>	<b>-</b>	<b>1.78</b>	<b>525</b>
<b>Standard Deviation</b>		<b>0.06</b>	<b>-</b>	<b>0.13</b>	<b>14</b>

\* NMHC- Non- Methane Hydrocarbons

BDL- Below Detection Limit ( Detection Limit – NMHC : 0.5 ppm)

NS- Not Specified

The overall values of TSPM, PM<sub>10</sub>, PM<sub>2.5</sub>, SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> at Kandla Port Colony was attributed by vehicle emission produced from trucks and heavy duty vehicles that pass through the road outside Kandla Port Colony. The mean TSPM values at Oil Jetty were 463 µg/m<sup>3</sup>, The mean PM<sub>10</sub> values were 366 µg/m<sup>3</sup>, which is above the permissible limit. PM<sub>2.5</sub> values were slightly above the permissible limit (mean = 56 µg/m<sup>3</sup>). The average values of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were 4.49 µg/m<sup>3</sup>, 19.98 µg/m<sup>3</sup> and 19.88 µg/m<sup>3</sup> respectively and were all within the permissible limit.

The levels of Benzene, Hydrocarbons (HC) and CO were within the permissible limit at Kandla Port Colony. The mean Benzene concentration was 1.19 µg/m<sup>3</sup>, well below the permissible limit of 5.0 µg/m<sup>3</sup>. HC's were below the detectable limit and Carbon Monoxide concentration was 1.78 mg/m<sup>3</sup>, well below the permissible limit of 4.0 mg/m<sup>3</sup>.

**Location 4: Gopalpuri Hospital (AL-4)**

Table 4 : Results of Air Pollutant Concentration at Gopalpuri Hospital										
Parameter	Date	TSPM [µg/m <sup>3</sup> ]	PM10 [µg/m <sup>3</sup> ]	PM2.5 [µg/m <sup>3</sup> ]	SO <sub>2</sub> [µg/m <sup>3</sup> ]		NO <sub>x</sub> [µg/m <sup>3</sup> ]		NH <sub>3</sub> [µg/m <sup>3</sup> ]	
					8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
Sampling Period	-	24hr	24hr	24hr	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
NAAQMS limit	-	NS	100 µg/m <sup>3</sup>	60 µg/m <sup>3</sup>	-	80 µg/m <sup>3</sup>	-	80 µg/m <sup>3</sup>	-	400 µg/m <sup>3</sup>
AL4 -1	01.07.2021	148	138	21	1.32	2.20	12.70	13.34	3.57	5.36
					2.20		13.34		7.40	
					3.08		13.97		5.11	
AL4 -2	05.07.2021	313	277	115	3.52	1.76	24.14	19.48	5.36	6.72
					1.32		13.34		8.42	
					0.44		20.96		6.38	
AL4 -3	09.07.2021	287	152	40	1.32	2.49	12.70	30.06	5.87	5.79
					2.64		22.23		5.11	
					3.52		55.26		6.38	
AL4 -4	14.07.2021	143	77	8	1.32	0.88	13.34	11.64	11.74	9.19
					0.88		11.43		8.17	
					0.44		10.16		7.66	
AL4 -5	16.07.2021	196	119	83	1.32	2.93	20.33	15.24	5.62	7.49
					3.52		13.34		9.45	
					3.96		12.07		7.40	
AL4 -6	21.07.2021	228	128	100	2.64	1.90	22.87	17.57	7.15	6.89
					1.32		13.34		7.40	
					1.76		16.51		6.13	
AL4 -7	23.07.2021	338	200	109	0.88	1.32	19.05	26.25	7.15	9.36
					1.32		28.58		9.70	
					1.76		31.12		11.23	
AL4 -8	27.07.2021	806	746	27	1.76	2.49	19.05	16.94	6.89	6.47
					2.20		14.61		6.38	
					3.52		17.15		6.13	
<b>Monthly Average</b>		307	230	63		2.00		18.82		7.16
<b>Standard Deviation</b>		214	217	43		0.67		6.34		1.46

NS: Not Specified

<b>Table 4B : Results of Air Pollutant Concentration at Gopalpuri Hospital</b>					
<b>Parameter</b>	<b>Date</b>	<b>C<sub>6</sub>H<sub>6</sub> [µg/m<sup>3</sup>]</b>	<b>HC*</b>	<b>CO [mg/m<sup>3</sup>]</b>	<b>CO<sub>2</sub> [ppm ]</b>
<b>Sampling Period</b>		<b>8 hr</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>
<b>NAAQMS limit</b>		<b>5.0 µg/m<sup>3</sup></b>	<b>NS</b>	<b>4.0 mg/m<sup>3</sup></b>	<b>NS</b>
AL4 -1	01.07.2021	1.22	BDL	1.62	502
AL4 -2	05.07.2021	1.16	BDL	1.48	499
AL4 -3	09.07.2021	1.32	BDL	1.62	501
AL4 -4	14.07.2021	1.28	BDL	1.78	489
AL4 -5	16.07.2021	1.25	BDL	1.46	496
AL4 -6	21.07.2021	1.18	BDL	1.62	510
AL4 -7	23.07.2021	1.14	BDL	1.78	502
AL4 -8	27.07.2021	1.23	BDL	1.48	496
<b>Monthly Average</b>		<b>1.22</b>	<b>-</b>	<b>1.61</b>	<b>499</b>
<b>Standard Deviation</b>		<b>0.06</b>	<b>-</b>	<b>0.13</b>	<b>6</b>

\* NMHC- Non- Methane Hydrocarbons

BDL- Below Detection Limit (Detection Limit – NMHC: 0.5 ppm)

NS-Not Specified

The overall values of TSPM, PM<sub>10</sub>, PM<sub>2.5</sub>, SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> at Gopalpuri Hospital was attributed by vehicle emission produced from light motor vehicles of the colony residents. The mean TSPM values at Oil Jetty were 307 µg/m<sup>3</sup>, The mean PM<sub>10</sub> values were 230 µg/m<sup>3</sup>, which is above the permissible limit. PM<sub>2.5</sub> values were slight above the permissible limit (mean= 63 µg/m<sup>3</sup>). The average values of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were 2.00 µg/m<sup>3</sup>, 18.82 µg/m<sup>3</sup> and 7.16 µg/m<sup>3</sup> respectively and were all within the permissible limit.

The levels of Benzene, Hydrocarbons (HC) and CO were within the permissible limit at Gopalpuri Hospital. The mean Benzene concentration was 1.22 µg/m<sup>3</sup>, well below the permissible limit of 5.0 µg/m<sup>3</sup>. HC's were below the detectable limit and Carbon Monoxide concentration was 1.61 mg/m<sup>3</sup>, well below the permissible limit of 4.0 mg/m<sup>3</sup>.

**Location 5: Coal Storage Area (AL-5)**

Table 5 : Results of Air Pollutant Concentration at Coal Storage Area										
Parameters	Date	TSPM [µg/m <sup>3</sup> ]	PM10 [µg/m <sup>3</sup> ]	PM2.5 [µg/m <sup>3</sup> ]	SO <sub>2</sub> [µg/m <sup>3</sup> ]		NO <sub>x</sub> [µg/m <sup>3</sup> ]		NH <sub>3</sub> [µg/m <sup>3</sup> ]	
Sampling Period	-	24hr	24hr	24hr	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
NAAQMS limit	-	NS	100 µg/m <sup>3</sup>	60 µg/m <sup>3</sup>	-	80 µg/m <sup>3</sup>	-	80 µg/m <sup>3</sup>	-	400 µg/m <sup>3</sup>
AL5 – 1	01.07.2021	428	158	47	3.08	3.37	42.56	48.70	15.83	14.04
					4.84		50.81		12.76	
					2.20		52.72		13.53	
AL5 – 2	05.07.2021	496	150	44	9.67	6.01	57.16	54.84	10.21	14.98
					4.84		49.54		13.53	
					3.52		57.80		21.19	
AL5 – 3	09.07.2021	222	135	76	9.67	7.62	60.98	50.60	16.85	17.44
					3.52		57.16		18.89	
					9.67		33.66		16.59	
AL5 – 4	14.07.2021	349	309	21	17.58	9.23	22.87	31.97	9.45	15.32
					4.84		32.39		21.70	
					5.28		40.65		14.81	
AL5 – 5	16.07.2021	264	123	12	9.67	11.87	16.51	21.38	12.00	14.21
					13.19		22.23		14.04	
					12.75		25.41		16.59	
AL5 – 6	21.07.2021	358	303	33	4.40	5.28	22.87	19.69	16.85	18.47
					6.15		19.05		16.34	
					5.28		17.15		22.21	
AL5 – 7	23.07.2021	268	194	45	4.40	5.28	27.95	23.71	12.76	16.76
					5.28		20.96		16.59	
					6.15		22.23		20.93	
AL5 – 8	27.07.2021	446	273	45	6.15	6.89	14.61	17.15	10.21	13.19
					7.03		22.23		14.04	
					7.47		14.61		15.32	
<b>Monthly Average</b>		354	206	40		6.94		33.50		15.55
<b>Standard Deviation</b>		98	77	19		2.65		15.50		1.84

NS: Not Specified

<b>Table 5B : Results of Air Pollutant Concentration at Coal Storage Area</b>					
<b>Parameter</b>	<b>Date</b>	<b>C<sub>6</sub>H<sub>6</sub> [µg/m<sup>3</sup>]</b>	<b>HC*</b>	<b>CO [mg/m<sup>3</sup>]</b>	<b>CO<sub>2</sub> [ppm ]</b>
<b>Sampling Period</b>		<b>8 hr</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>
<b>NAAQMS limit</b>		<b>5.0 µg/m<sup>3</sup></b>	<b>NS</b>	<b>4.0 mg/m<sup>3</sup></b>	<b>NS</b>
AL5 – 1	01.07.2021	1.28	BDL	1.82	526
AL5 – 2	05.07.2021	1.11	BDL	1.78	522
AL5 – 3	09.07.2021	1.16	BDL	1.88	520
AL5 – 4	14.07.2021	1.32	BDL	1.78	530
AL5 – 5	16.07.2021	1.28	BDL	1.82	536
AL5 – 6	21.07.2021	1.22	BDL	1.77	522
AL5 – 7	23.07.2021	1.18	BDL	1.86	526
AL5 – 8	27.07.2021	1.26	BDL	1.9	530
<b>Monthly Average</b>		<b>1.23</b>	<b>-</b>	<b>1.83</b>	<b>527</b>
<b>Standard Deviation</b>		<b>0.07</b>	<b>-</b>	<b>0.05</b>	<b>5</b>

\* NMHC- Non- Methane Hydrocarbons

BDL- Below Detection Limit (Detection Limit – NMHC: 0.5 ppm)

NS-Not Specified

The overall values of TSPM, PM<sub>10</sub>, PM<sub>2.5</sub>, SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> at Coal Storage Area was comparatively highest among all the locations of Air Quality monitoring in Kandla Port. High values of TSPM, PM<sub>10</sub>, PM<sub>2.5</sub>, SO<sub>2</sub>, NO<sub>x</sub> at this location was due to lifting of coal with grab and other coal handling processes near Berth no. 6 & 7. Moreover, the traffic was also heavy around this place for transport of coal thus emissions produced from heavy vehicles. The mean TSPM values at Coal storage were 354 µg/m<sup>3</sup>. The mean PM<sub>10</sub> values were 206 µg/m<sup>3</sup>, which is well above the permissible limit. PM<sub>2.5</sub> values were below the permissible limit (mean = 40 µg/m<sup>3</sup>). The average values of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were 6.94 µg/m<sup>3</sup>, 33.50 µg/m<sup>3</sup> and 15.55 µg/m<sup>3</sup> respectively and were all within the permissible limit.

The levels of Benzene, Hydrocarbons (HC) and CO were within the permissible limit at Coal Storage Area. The mean Benzene concentration was 1.23 µg/m<sup>3</sup>, well below the permissible limit of 5.0 µg/m<sup>3</sup>. HC's were below the detectable limit and Carbon Monoxide concentration was 1.83 mg/m<sup>3</sup>, well below the permissible limit of 4.0 mg/m<sup>3</sup>.



**Location 6: Tuna Port (AL-6)**

**Table 6 : Results of Air Pollutant Concentration at Tuna Port**

Parameters	Date	TSPM [µg/m <sup>3</sup> ]	PM10 [µg/m <sup>3</sup> ]	PM2.5 [µg/m <sup>3</sup> ]	SO <sub>2</sub> [µg/m <sup>3</sup> ]		NO <sub>x</sub> [µg/m <sup>3</sup> ]		NH <sub>3</sub> [µg/m <sup>3</sup> ]	
					8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
Sampling Period	-	24hr	24hr	24hr	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
NAAQMS limit	-	NS	100 µg/m <sup>3</sup>	60 µg/m <sup>3</sup>	-	80 µg/m <sup>3</sup>	-	80 µg/m <sup>3</sup>	-	400 µg/m <sup>3</sup>
AL6 -1	01.07.2021	149	97	39	0.88	2.20	16.51	15.24	5.87	7.32
					2.20		17.15		7.91	
					3.52		12.07		8.17	
AL6 - 2	05.07.2021	270	169	97	2.20	2.20	13.97	17.36	12.76	12.00
					1.32		14.61		11.74	
					3.08		23.50		11.49	
AL6 - 3	09.07.2021	513	198	86	5.71	4.98	21.60	20.96	6.89	7.06
					6.15		17.15		6.64	
					3.08		24.14		7.66	
AL6 - 4	14.07.2021	230	97	98	2.20	3.08	8.26	9.53	7.40	8.76
					2.64		9.53		8.93	
					4.40		10.80		9.96	
AL6 - 5	16.07.2021	554	484	18	1.76	2.20	14.61	12.07	10.72	10.89
					3.52		12.07		10.98	
					1.32		9.53		10.98	
AL6 - 6	21.07.2021	405	302	98	2.20	2.64	6.35	9.95	16.34	14.89
					1.76		10.80		15.57	
					3.96		12.70		12.76	
AL6 - 7	23.07.2021	211	128	12	1.32	2.05	21.60	19.05	10.98	11.83
					2.20		13.34		13.27	
					2.64		22.23		11.23	
AL6 - 8	27.07.2021	645	524	12	0.88	2.05	14.61	18.84	10.21	9.36
					2.64		17.15		8.68	
					2.64		24.77		9.19	
<b>Monthly Average</b>		372	250	58		2.67		15.38		10.26
<b>Standard Deviation</b>		183	171	41		1.00		4.40		2.65

NS: Not Specified

<b>Table 6B : Results of Air Pollutant Concentration at Tuna Port</b>					
<b>Parameter</b>	<b>Date</b>	<b>C<sub>6</sub>H<sub>6</sub> [µg/m<sup>3</sup>]</b>	<b>HC*</b>	<b>CO [mg/m<sup>3</sup>]</b>	<b>CO<sub>2</sub> [ppm ]</b>
<b>Sampling Period</b>		<b>8 hr</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>
<b>NAAQMS limit</b>		<b>5.0 µg/m<sup>3</sup></b>	<b>NS</b>	<b>4.0 mg/m<sup>3</sup></b>	<b>NS</b>
AL6 -1	01.07.2021	1.2	BDL	1.79	510
AL6 – 2	05.07.2021	1.11	BDL	1.84	502
AL6 – 3	09.07.2021	1.19	BDL	1.72	511
AL6 – 4	14.07.2021	1.15	BDL	1.69	496
AL6 – 5	16.07.2021	1.06	BDL	1.88	499
AL6 – 6	21.07.2021	1.11	BDL	1.87	502
AL6 – 7	23.07.2021	1.06	BDL	1.74	506
AL6 – 8	27.07.2021	1.15	BDL	1.7	512
<b>Monthly Average</b>		<b>1.13</b>	<b>-</b>	<b>1.78</b>	<b>505</b>
<b>Standard Deviation</b>		<b>0.05</b>	<b>-</b>	<b>0.08</b>	<b>6</b>

\* NMHC- Non- Methane Hydrocarbons

BDL- Below Detection Limit (Detection Limit – NMHC : 0.5 ppm)

NS- Not Specified

The mean TSPM values at Tuna Port were 372 µg/m<sup>3</sup>, The mean PM<sub>10</sub> values were 250 µg/m<sup>3</sup>, which is above the permissible limit. PM<sub>2.5</sub> values were within the permissible limit (mean = 58 µg/m<sup>3</sup> µg/m<sup>3</sup>). The average values of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were 2.63 µg/m<sup>3</sup>, 15.38 µg/m<sup>3</sup> and 10.26 µg/m<sup>3</sup> respectively and were all within the permissible limit.

The levels of Benzene, Hydrocarbons (HC) and CO were within the permissible limit at Tuna Port. The mean Benzene concentration was 1.13 µg/m<sup>3</sup>, well below the permissible limit of 5.0 µg/m<sup>3</sup>. HC's were below the detectable limit and Carbon Monoxide concentration was 1.78 mg/m<sup>3</sup>, well below the permissible limit of 4.0 mg/m<sup>3</sup>.

**Location 7: Signal Building (Vadinar) (AL-7)**

Table 7 : Results of Air Pollutant Concentration at Signal Building										
Parameters	Date	TSPM [µg/m3]	PM10 [µg/m3]	PM2.5 [µg/m3]	SO2 [µg/m3]		NOx [µg/m3]		NH3 [µg/m3]	
					8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
Sampling Period	-	24hr	24hr	24hr	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
NAAQMS limit	-	NS	100 µg/m3	60 µg/m3	-	80 µg/m3	-	80 µg/m3	-	400 µg/m3
AL7 -1	01.07.2021	119	72	37	3.20	3.08	8.05	8.26	4.41	4.34
					2.97		8.49		4.42	
					3.06		8.24		4.19	
AL7 -2	05.07.2021	104	81	30	3.40	3.52	13.07	12.70	5.45	5.36
					3.18		12.38		5.29	
					3.99		12.65		5.35	
AL7 -3	09.07.2021	62	73	42	4.28	3.96	6.35	6.35	6.09	5.87
					3.60		6.50		5.82	
					4.01		6.20		5.69	
AL7 -4	14.07.2021	104	74	110	4.18	3.96	6.24	6.35	18.21	17.88
					3.87		6.47		17.45	
					3.84		6.34		17.97	
AL7 -5	16.07.2021	96	63	40	2.82	2.64	20.65	20.33	9.23	8.93
					2.65		19.80		8.86	
					2.45		20.54		8.69	
AL7 -6	21.07.2021	102	68	12	8.02	7.47	5.73	5.72	2.71	2.81
					7.18		6.03		2.59	
					7.22		5.40		3.13	
AL7 -7	23.07.2021	143	95	35	5.17	4.84	33.65	33.57	3.29	3.57
					5.10		33.82		3.8	
					4.26		33.23		3.62	
AL7 -8	27.07.2021	107	74	18	7.26	7.46	31.49	31.46	4.5	4.45
					7.49		31.78		4.75	
					7.62		31.12		4.1	
<b>Monthly Average</b>		105	75	40		5		16		7
<b>Standard Deviation</b>		23	10	30		2		12		5

NS: Not Specified

<b>Table 7B : Results of Air Pollutant Concentration at Signal Building</b>					
<b>Parameter</b>	<b>Date</b>	<b>C<sub>6</sub>H<sub>6</sub> [µg/m<sup>3</sup>]</b>	<b>HC*</b>	<b>CO [mg/m<sup>3</sup>]</b>	<b>CO<sub>2</sub> [ppm ]</b>
<b>Sampling Period</b>		<b>8 hr</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>
<b>NAAQMS limit</b>		<b>5.0 µg/m<sup>3</sup></b>	<b>NS</b>	<b>4.0 mg/m<sup>3</sup></b>	<b>NS</b>
AL7 -1	01.07.2021	1.1	BDL	1.56	489
AL7 – 2	05.07.2021	1.06	BDL	1.66	488
AL7 – 3	09.07.2021	1.02	BDL	1.72	479
AL7 – 4	14.07.2021	1.1	BDL	1.62	496
AL7 – 5	16.07.2021	1.11	BDL	1.68	488
AL7 – 6	21.07.2021	1.16	BDL	1.58	490
AL7 – 7	23.07.2021	1.12	BDL	1.66	481
AL7 – 8	27.07.2021	1.1	BDL	1.6	475
<b>Monthly Average</b>		<b>1.10</b>	<b>-</b>	<b>1.64</b>	<b>486</b>
<b>Standard Deviation</b>		<b>0.04</b>	<b>-</b>	<b>0.05</b>	<b>7</b>

\* NMHC- Non- Methane Hydrocarbons

BDL- Below Detection Limit (Detection Limit – NMHC : 0.5 ppm)

NS\_ Not Specified

The mean TSPM values at Vadinar Port were 105 µg/m<sup>3</sup>. The mean PM<sub>10</sub> values were 75 µg/m<sup>3</sup>, which is below the permissible limit. PM<sub>2.5</sub> values were also within the permissible limit (mean = 40 µg/m<sup>3</sup> µg/m<sup>3</sup>). The average values of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were 5.0 µg/m<sup>3</sup>, 16.0 µg/m<sup>3</sup> and 7.0 µg/m<sup>3</sup> respectively and were all within the permissible limit.

The levels of Benzene, Hydrocarbons (HC) and CO were within the permissible limit at Vadinar Port. The mean Benzene concentration was 1.10 µg/m<sup>3</sup>, well below the permissible limit of 5.0 µg/m<sup>3</sup>. HC's were below the detectable limit and Carbon Monoxide concentration was 1.64 mg/m<sup>3</sup>, well below the permissible limit of 4.0 mg/m<sup>3</sup>.

**Location 8: Admin Building (Vadinar) (AL-8)**

<b>Table 8 : Results of Air Pollutant Concentration at Admin Building</b>										
<b>Parameters</b>	<b>Date</b>	<b>TSPM [µg/m3]</b>	<b>PM10 [µg/m3]</b>	<b>PM2.5 [µg/m3]</b>	<b>SO2 [µg/m3]</b>		<b>NOx [µg/m3]</b>		<b>NH3 [µg/m3]</b>	
<b>Sampling Period</b>	-	<b>24hr</b>	<b>24hr</b>	<b>24hr</b>	<b>8 hr</b>	<b>24hr (Avg.)</b>	<b>8 hr</b>	<b>24hr (Avg.)</b>	<b>8 hr</b>	<b>24hr (Avg.)</b>
<b>NAAQMS limit</b>	-	<b>NS</b>	<b>100 µg/m3</b>	<b>60 µg/m3</b>	-	<b>80 µg/m3</b>	-	<b>80 µg/m3</b>	-	<b>400 µg/m3</b>
<b>AL8 -1</b>	01.07.2021	172	96	25	2.71	2.64	10.75	10.80	3.42	3.57
					2.64		10.55		3.71	
					2.58		11.09		3.58	
<b>AL8 -2</b>	05.07.2021	121	100	16	4.05	3.96	8.89	8.89	4.51	4.85
					3.95		8.81		5.18	
					3.88		8.96		4.86	
<b>AL8 -3</b>	09.07.2021	108	88	14	5.02	4.84	5.80	5.72	9.48	9.19
					4.79		5.70		8.94	
					4.72		5.67		9.15	
<b>AL8 -4</b>	14.07.2021	169	68	84	6.74	6.59	5.76	5.72	22.65	22.61
					6.16		5.52		23.06	
					6.88		5.89		22.12	
<b>AL8 -5</b>	16.07.2021	136	85	37	1.40	1.32	18.40	18.42	23.67	22.98
					1.23		18.53		22.46	
					1.32		18.33		22.81	
<b>AL8 -6</b>	21.07.2021	140	65	87	9.58	9.67	9.04	8.89	6.65	6.63
					9.80		8.86		6.72	
					9.62		8.76		6.52	
<b>AL8 -5</b>	23.07.2021	168	96	47	6.10	6.15	44.85	44.46	9.23	8.93
					6.24		44.21		8.46	
					6.10		44.32		9.1	
<b>AL8-6</b>	27.07.2021	153	53	40	3.46	3.52	45.00	44.46	3.95	4.08
					3.72		44.05		4.09	
					3.38		44.32		4.2	
<b>Monthly Average</b>		146	81	44		5		18		10
<b>Standard Deviation</b>		24	17	28		3		17		8

NS: Not Specified

<b>Table 8B : Results of Air Pollutant Concentration at Admin Building</b>					
<b>Parameter</b>	<b>Date</b>	<b>C<sub>6</sub>H<sub>6</sub> [µg/m<sup>3</sup>]</b>	<b>HC*</b>	<b>CO [mg/m<sup>3</sup>]</b>	<b>CO<sub>2</sub> [ppm ]</b>
<b>Sampling Period</b>		<b>8 hr</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>
<b>NAAQMS limit</b>		<b>5.0 µg/m<sup>3</sup></b>	<b>NS</b>	<b>4.0 mg/m<sup>3</sup></b>	<b>NS</b>
AL8 -1	01.07.2021	1.1	BDL	1.56	489
AL8-2	05.07.2021	1.06	BDL	1.66	488
AL8 -3	09.07.2021	1.02	BDL	1.72	479
AL8-4	14.07.2021	1.1	BDL	1.62	496
AL8 -5	16.07.2021	1.11	BDL	1.68	488
AL8-6	21.07.2021	1.16	BDL	1.58	490
AL8-7	23.07.2021	1.12	BDL	1.66	481
AL8-8	27.07.2021	1.1	BDL	1.6	475
Monthly Average		<b>1.10</b>	<b>-</b>	<b>1.64</b>	<b>486</b>
Standard Deviation		<b>0.04</b>	<b>-</b>	<b>0.05</b>	<b>7</b>

\* NMHC- Non- Methane Hydrocarbons

BDL- Below Detection Limit (Detection Limit – NMHC: 0.5 ppm)

NS-Not Specified

The overall values of TSPM, PM<sub>10</sub>, PM<sub>2.5</sub>, SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> at Admin Building Vadinar was comparatively low among all the locations of Air Quality monitoring in Kandla Port and Vadinar Port. The mean TSPM values at Vadinar Port were 146 µg/m<sup>3</sup>. The mean PM<sub>10</sub> values were 81 µg/m<sup>3</sup>, which is below the permissible limit. PM<sub>2.5</sub> values were also within the permissible limit (mean = 44.0 µg/m<sup>3</sup>). The average values of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were 5.0µg/m<sup>3</sup>, 18.0 µg/m<sup>3</sup> and 10.0 µg/m<sup>3</sup> respectively and were all within the permissible limit.

The levels of Benzene, Hydrocarbons (HC) and CO were within the permissible limit at Admin Building, Vadinar Port. The mean Benzene concentration was 1.10 µg/m<sup>3</sup>, well below the permissible limit of 5.0 µg/m<sup>3</sup>. HC's were below the detectable limit and Carbon Monoxide concentration was 1.64 mg/m<sup>3</sup>, well below the permissible limit of 4.0 mg/m<sup>3</sup>.

#### **1.4 Observations and Conclusion**

During the monitoring period, the overall Ambient Air Quality of the port area was found to be well within the desired levels for various pollutants. However, Near Coal storage area, Marine Bhavan and Oil Jetty area, PM<sub>10</sub> values was above the permissible standards. All other pollutants were recorded well below the prescribed limits.

## 2. Drinking Water Quality Monitoring

Drinking Water Quality Monitoring was carried out at twenty stations at Kandla, Vadinar & Township Area of Deendayal Port.

### 2.1 Drinking Water Monitoring Methodology

Drinking water samples were collected from 20 locations as prescribed in the tender document. Samples for physico-chemical analysis were collected in 1 liter carboys and samples for microbiological parameters were collected in sterilized bottles. These samples were then analyzed in laboratory for various drinking water parameters at Kandla Lab/Surat.

The Sampling and Analysis was done as per standard methods - IS 10500:2012. The water samples were analyzed for various parameters, viz. Color , Odor, Turbidity , Conductivity , pH , Chlorides , TDS, Total Hardness, Iron , Sulphate , Salinity , DO, BOD, Na, K, Ca, Mg, F, NO<sub>3</sub>, NO<sub>2</sub>, Mn, Cr-6, Cu, Cd, As, Hg, Pb, Zn, Bacterial Count (cfu) .

### 2.2 Results

The Drinking Water Quality monitoring data for 20 stations are given in below from table No. 9 to Table No. 15

**Table 9: Drinking Water Quality Monitoring Parameters for Nirman Building 1, P & C building & Main Gate (North) at Kandla**

Sr. No.	Parameter	Unit	Nirman Building 1	P & C building	Main Gate North	Acceptable Limits as per IS 10500 : 2012	Permissible Limits as per IS 10500 : 2012
1	pH	pH Unit	7.7	7.4	7.6	6.5 to 8.5	6.5 to 8.5
2	Total Dissolved Solids	mg/l	1290	1530	1180	500	2000
3	Turbidity	NTU	0	1	0	1.0	5.0
4	Odor	-	Odourless	Odourless	Odourless	Agreeable	Agreeable
5	Color	Hazen Units	Colourless	Colourless	Colourless	5.0	15.0
6	Conductivity	µs/cm	2500	3010	2200	NS*	NS*
7	Biochemical Oxygen Demand	mg/l	<2	<2	<2	NS*	NS*
8	Chloride as Cl	mg/l	416	436	451	250.0	1000.0
9	Ca as Ca	mg/l	72.14	52.10	64.13	75.0	200.0
10	Mg as Mg	mg/l	51.03	68.04	65.61	30.0	100.0
11	Total Hardness	mg/l	390	410	430	200.0	600.0
12	Iron as Fe	mg/l	<0.01	<0.01	<0.01	0.3	No Relaxation
13	Fluorides as F	mg/l	0.26	0.47	0.21	1.0	1.5
14	Sulphate as SO4	mg/l	140.52	166.8	156	200.0	400
15	Nitrite as NO2	mg/l	<0.01	<0.01	<0.01	NS*	NS*
16	Nitrate as NO3	mg/l	10.21	8.45	7.74	45.0	No Relaxation
17	Salinity	%	0.75	0.79	0.81	NS*	NS*
18	Sodium as Na	mg/l	170	168	148	NS*	NS*
19	Potassium as K	mg/l	2.2	2.3	2.8	NS*	NS*
20	Manganese	mg/l	<0.04	<0.04	<0.04	0.1	0.3
21	Hexavalent Chromium	mg/l	<0.03	<0.03	<0.03	NS*	NS*
22	Copper	mg/l	<0.05	<0.05	<0.05	0.05	1.5
23	Cadmium	mg/l	<0.002	<0.002	<0.002	0.003	0.003
24	Arsenic	mg/l	<0.01	<0.01	<0.01	0.01	0.05
25	Mercury	mg/l	<0.001	<0.001	<0.001	0.001	0.001
26	Lead	mg/l	<0.01	<0.01	<0.01	0.01	0.01
27	Zinc	mg/l	<0.1	<0.1	<0.1	5.0	15.0
28	Bacterial Count	CFU/100ml	Absent	Absent	Absent	Absent	Absent

\*NS: Not Specified



**Table 10: Drinking Water Quality Monitoring Parameters for Canteen, West Gate – I & Wharf Area at Kandla**

Sr. No.	Parameter	Unit	Canteen	West Gate - I	Wharf Area	Acceptable Limits as per IS 10500 : 2012	Permissible Limits as per IS 10500 : 2012
1	pH	pH Unit	7.5	7.3	7.6	6.5 to 8.5	6.5 to 8.5
2	Total Dissolved Solids	mg/l	1590	1190	1670	500	2000
3	Turbidity	NTU	0	0	1	1.0	5.0
4	Odor	-	Odourless	Odourless	Odourless	Agreeable	Agreeable
5	Color	Hazen Units	Colourless	Colourless	Colourless	5.0	15.0
6	Conductivity	µs/cm	3110	2330	3300	NS*	NS*
7	Biochemical Oxygen Demand	mg/l	<2	<2	<2	NS*	NS*
8	Chloride as Cl	mg/l	411	416	426	250.0	1000.0
9	Ca as Ca	mg/l	56.11	64.13	52.10	75.0	200.0
10	Mg as Mg	mg/l	60.75	48.60	63.18	30.0	100.0
11	Total Hardness	mg/l	390	360	390	200.0	600.0
12	Iron as Fe	mg/l	<0.01	<0.01	<0.01	0.3	No Relaxation
13	Fluorides as F	mg/l	0.18	0.53	1.05	1.0	1.5
14	Sulphate as SO <sub>4</sub>	mg/l	166.8	165.6	226.8	200.0	400
15	Nitrite as NO <sub>2</sub>	mg/l	<0.01	<0.01	<0.01	NS*	NS*
16	Nitrate as NO <sub>3</sub>	mg/l	10.56	11.97	7.53	45.0	No Relaxation
17	Salinity	%	0.74	0.75	0.77	NS*	NS*
18	Sodium as Na	mg/l	133	168	156	NS*	NS*
19	Potassium as K	mg/l	3	2.2	2.4	NS*	NS*
20	Manganese	mg/l	<0.04	<0.04	<0.04	0.1	0.3
21	Hexavalent Chromium	mg/l	<0.03	<0.03	<0.03	NS*	NS*
22	Copper	mg/l	<0.05	<0.05	<0.05	0.05	1.5
23	Cadmium	mg/l	<0.002	<0.002	<0.002	0.003	0.003
24	Arsenic	mg/l	<0.01	<0.01	<0.01	0.01	0.05
25	Mercury	mg/l	<0.001	<0.001	<0.001	0.001	0.001
26	Lead	mg/l	<0.01	<0.01	<0.01	0.01	0.01
27	Zinc	mg/l	<0.1	<0.1	<0.1	5.0	15.0
28	Bacterial Count	CFU/100ml	Absent	Absent	Absent	Absent	Absent

\*NS: Not Specified

**Table 11: Drinking Water Quality Monitoring Parameters for Sewa sadan – 3, Workshop I & Custom Building at Kandla**

Sr. No.	Parameter	Unit	SewaSadan – 3	Workshop	Custom Building	Acceptable Limits as per IS 10500 : 2012	Permissible Limits as per IS 10500 : 2012
1	pH	pH Unit	7.7	7.9	7.3	6.5 to 8.5	6.5 to 8.5
2	Total Dissolved Solids	mg/l	1490	1090	1330	500	2000
3	Turbidity	NTU	1	0	0	1.0	5.0
4	Odor	-	Odourless	Odourless	Odourless	Agreeable	Agreeable
5	Color	Hazen Units	Colourless	Colourless	Colourless	5.0	15.0
6	Conductivity	µs/cm	2990	2090	2680	NS*	NS*
7	Biochemical Oxygen Demand	mg/l	<2	<2	<2	NS*	NS*
8	Chloride	mg/l	451	456	461	250.0	1000.0
9	Ca as Ca	mg/l	60.12	56.11	64.13	75.0	200.0
10	Mg as Mg	mg/l	60.75	63.18	53.46	30.0	100.0
11	Total Hardness	mg/l	400	400	380	200.0	600.0
12	Iron as Fe	mg/l	<0.01	<0.01	<0.01	0.3	No Relaxation
13	Fluorides	mg/l	0.93	0.70	1.45	1.0	1.5
14	Sulphate	mg/l	156	171.6	195.6	200.0	400
15	Nitrite	mg/l	<0.01	<0.01	<0.01	NS*	NS*
16	Nitrate	mg/l	14.78	16.83	9.50	45.0	No Relaxation
17	Salinity	%	0.81	0.82	0.83	NS*	NS*
18	Sodium as Na	mg/l	162	152	162	NS*	NS*
19	Potassium as K	mg/l	2.3	2.4	2.8	NS*	NS*
20	Manganese	mg/l	<0.04	<0.04	<0.04	0.1	0.3
21	Hexavalent Chromium	mg/l	<0.03	<0.03	<0.03	NS*	NS*
22	Copper	mg/l	<0.05	<0.05	<0.05	0.05	1.5
23	Cadmium	mg/l	<0.002	<0.002	<0.002	0.003	0.003
24	Arsenic	mg/l	<0.01	<0.01	<0.01	0.01	0.05
25	Mercury	mg/l	<0.001	<0.001	<0.001	0.001	0.001
26	Lead	mg/l	<0.01	<0.01	<0.01	0.01	0.01
27	Zinc	mg/l	<0.1	<0.1	<0.1	5.0	15.0
28	Bacterial Count	CFU/100ml	Absent	Absent	Absent	Absent	Absent

\*NS: Not Specified

**Table 12: Drinking Water Quality Monitoring Parameters for Port Colony Kandla, Hospital Kandla & A.O. Building at Gandhidham**

Sr. No.	Parameter	Unit	Port Colony Kandla	Hospital Kandla	A.O. Building	Acceptable Limits as per IS 10500 : 2012	Permissible Limits as per IS 10500 : 2012
1	pH	pH Unit	7.8	7.7	7.0	6.5 to 8.5	6.5 to 8.5
2	Total Dissolved Solids	mg/l	1210	1450	1010	500	2000
3	Turbidity	NTU	1	2	2	1.0	5.0
4	Odor	-	Odourless	Odourless	Odourless	Agreeable	Agreeable
5	Color	Hazen Units	Colourless	Colourless	Colourless	5.0	15.0
6	Conductivity	µs/cm	2370	2880	2030	NS*	NS*
7	Biochemical Oxygen Demand	mg/l	<2	<2	<2	NS*	NS*
8	Chloride	mg/l	526	541	491	250.0	1000.0
9	Ca as Ca	mg/l	52.10	52.10	48.10	75.0	200.0
10	Mg as Mg	mg/l	58.32	68.04	75.33	30.0	100.0
11	Total Hardness	mg/l	370	410	430	200.0	600.0
12	Iron as Fe	mg/l	<0.01	<0.01	<0.01	0.3	No Relaxation
13	Fluorides	mg/l	1.05	1.16	0.93	1.0	1.5
14	Sulphate	mg/l	204	214.8	147.6	200.0	400
15	Nitrite	mg/l	<0.01	<0.01	<0.01	NS*	NS*
16	Nitrate	mg/l	9.01	9.72	9.15	45.0	No Relaxation
17	Salinity	%	0.95	0.98	0.89	NS*	NS*
18	Sodium as Na	mg/l	178	160	180	NS*	NS*
19	Potassium as K	mg/l	2.7	2.3	2.8	NS*	NS*
20	Manganese	mg/l	<0.04	<0.04	<0.04	0.1	0.3
21	Hexavalent Chromium	mg/l	<0.03	<0.03	<0.03	NS*	NS*
22	Copper	mg/l	<0.05	<0.05	<0.05	0.05	1.5
23	Cadmium	mg/l	<0.002	<0.002	<0.002	0.003	0.003
24	Arsenic	mg/l	<0.01	<0.01	<0.01	0.01	0.05
25	Mercury	mg/l	<0.001	<0.001	<0.001	0.001	0.001
26	Lead	mg/l	<0.01	<0.01	<0.01	0.01	0.01
27	Zinc	mg/l	<0.1	<0.1	<0.1	5.0	15.0
28	Bacterial Count	CFU/100ml	Absent	Absent	Absent	Absent	Absent

\*NS: Not Specified

**Table 13: Drinking Water Quality Monitoring Parameters for School Gopalpuri, Guest House & E - Type Quarter at Gopalpuri, Gandhidham**

Sr. No.	Parameter	Unit	School Gopalpuri	Guest House	E - Type Quarter	Acceptable Limits as per IS 10500 : 2012	Permissible Limits as per IS 10500 : 2012
1	pH	pH Unit	7.1	7.3	7.2	6.5 to 8.5	6.5 to 8.5
2	Total Dissolved Solids	mg/l	990	1410	1330	500	2000
3	Turbidity	NTU	1	0	1	1.0	5.0
4	Odor	-	Odourless	Odourless	Odourless	Agreeable	Agreeable
5	Color	Hazen Units	Colourless	Colourless	Colourless	5.0	15.0
6	Conductivity	µs/cm	1900	2900	2660	NS*	NS*
7	Biochemical Oxygen Demand	mg/l	<2	<2	<2	NS*	NS*
8	Chloride	mg/l	526	476	516	250.0	1000.0
9	Ca as Ca	mg/l	60.12	56.11	68.14	75.0	200.0
10	Mg as Mg	mg/l	55.89	53.46	53.46	30.0	100.0
11	Total Hardness	mg/l	380	360	390	200.0	600.0
12	Iron as Fe	mg/l	<0.01	<0.01	<0.01	0.3	No Relaxation
13	Fluorides	mg/l	1.08	0.82	1.14	1.0	1.5
14	Sulphate	mg/l	183.6	157.2	150	200.0	400
15	Nitrite	mg/l	<0.01	<0.01	<0.01	NS*	NS*
16	Nitrate	mg/l	10.35	11.48	10.35	45.0	No Relaxation
17	Salinity	%	0.95	0.86	0.93	NS*	NS*
18	Sodium as Na	mg/l	196	203	200	NS*	NS*
19	Potassium as K	mg/l	2.4	2.3	2.6	NS*	NS*
20	Manganese	mg/l	<0.04	<0.04	<0.04	0.1	0.3
21	Hexavalent Chromium	mg/l	<0.03	<0.03	<0.03	NS*	NS*
22	Copper	mg/l	<0.05	<0.05	<0.05	0.05	1.5
23	Cadmium	mg/l	<0.002	<0.002	<0.002	0.003	0.003
24	Arsenic	mg/l	<0.01	<0.01	<0.01	0.01	0.05
25	Mercury	mg/l	<0.001	<0.001	<0.001	0.001	0.001
26	Lead	mg/l	<0.01	<0.01	<0.01	0.01	0.01
27	Zinc	mg/l	<0.1	<0.1	<0.1	5.0	15.0
28	Bacterial Count	CFU/100ml	Absent	Absent	Absent	Absent	Absent

\*NS: Not Specified

**Table 14: Drinking Water Quality Monitoring Parameters for F - Type Quarter, Hospital Gopalpuri & Tuna Port**

Sr. No.	Parameter	Unit	F - Type Quarter	Hospital Gopalpuri	Tuna Port	Acceptable Limits as per IS 10500 : 2012	Permissible Limits as per IS 10500 : 2012
1	pH	pH Unit	7.5	7.2	7.3	6.5 to 8.5	6.5 to 8.5
2	Total Dissolved Solids	mg/l	1100	1020	1050	500	2000
3	Turbidity	NTU	1	1	0	1.0	5.0
4	Odor	-	Odourless	Odourless	Odorless	Agreeable	Agreeable
5	Color	Hazen Unit	Colourless	Colourless	Colorless	5.0	15.0
6	Conductivity	µs/cm	2200	2050	1940	NS*	NS*
7	Biochemical Oxygen Demand	mg/l	<2	<2	<2	NS*	NS*
8	Chloride	mg/l	506	546	592	250.0	1000.0
9	Ca as Ca	mg/l	64.13	72.14	72.14	75.0	200.0
10	Mg as Mg	mg/l	65.61	43.74	36.45	30.0	100.0
11	Total Hardness	mg/l	430	360	330	200.0	600.0
12	Iron as Fe+3	mg/l	<0.01	<0.01	<0.01	0.3	No Relaxation
13	Fluorides	mg/l	0.94	1.02	0.46	1.0	1.5
14	Sulphate	mg/l	165.6	159.6	120	200.0	400
15	Nitrite	mg/l	<0.01	<0.01	<0.01	NS*	NS*
16	Nitrate	mg/l	10.63	9.36	1.33	45.0	No Relaxation
17	Salinity	%	0.91	0.99	0.92	NS*	NS*
18	Sodium as Na	mg/l	180	180	188	NS*	NS*
19	Potassium as K	mg/l	2.5	2.4	2.6	NS*	NS*
20	Manganese	mg/l	<0.04	<0.04	<0.04	0.1	0.3
21	Hexavalent Chromium	mg/l	<0.03	<0.03	<0.03	NS*	NS*
22	Copper	mg/l	<0.05	<0.05	<0.05	0.05	1.5
23	Cadmium	mg/l	<0.002	<0.002	<0.002	0.003	0.003
24	Arsenic	mg/l	<0.01	<0.01	<0.01	0.01	0.05
25	Mercury	mg/l	<0.001	<0.001	<0.001	0.001	0.001
26	Lead	mg/l	<0.01	<0.01	<0.01	0.01	0.01
27	Zinc	mg/l	<0.1	<0.1	<0.1	5.0	15.0
28	Bacterial Count	CFU/100ml	Absent	Absent	Absent	Absent	Absent

\*NS: Not Specified

**Table 15: Drinking Water Quality Monitoring Parameters for Vadinar Jetty & Port Colony at Vadinar**

Sr. No.	Parameter	Unit	Vadinar Jetty	Port Colony Vadinar	Acceptable Limits as per IS 10500 : 2012	Permissible Limits as per IS 10500 : 2012
1	pH	pH Unit	7.9	7.6	6.5 to 8.5	6.5 to 8.5
2	Total Dissolved Solids	mg/l	950.0	620.0	500	2000
3	Turbidity	NTU	ND	ND	1.0	5.0
4	Odor	-	Odorless	Odorless	Agreeable	Agreeable
5	Color	Hazen Units	Colorless	Colorless	5.0	15.0
6	Conductivity	µs/cm	1580.0	1030.0	NS*	NS*
7	Biochemical Oxygen Demand	mg/l	<2	<2	NS*	NS*
8	Chloride	mg/l	445.99	466.04	250.0	1000.0
9	Ca as Ca	mg/l	60.12	52.10	75.0	200.0
10	Mg as Mg	mg/l	63.18	60.75	30.0	100.0
11	Total Hardness	mg/l	410	380	200.0	600.0
12	Iron as Fe+3	mg/l	<0.01	<0.01	0.3	No Relaxation
13	Fluorides	mg/l	0.99	0.94	1.0	1.5
14	Sulphate	mg/l	16.80	17.64	200.0	400
15	Nitrite	mg/l	<0.01	<0.01	NS*	NS*
16	Nitrate	mg/l	11.48	9.50	45.0	No Relaxation
17	Salinity	%	0.81	0.84	NS*	NS*
18	Sodium as Na	mg/l	142.0	156.0	NS*	NS*
19	Potassium as K	mg/l	2.3	2.4	NS*	NS*
20	Manganese	mg/l	<0.04	<0.04	0.1	0.3
21	Hexavalent Chromium	mg/l	<0.03	<0.03	NS*	NS*
22	Copper	mg/l	<0.05	<0.05	0.05	1.5
23	Cadmium	mg/l	<0.002	<0.002	0.003	0.003
24	Arsenic	mg/l	<0.01	<0.01	0.01	0.05
25	Mercury	mg/l	<0.001	<0.001	0.001	0.001
26	Lead	mg/l	<0.01	<0.01	0.01	0.01
27	Zinc	mg/l	<0.1	<0.1	5.0	15.0
28	Bacterial Count	CFU/100ml	Absent	Absent	Absent	Absent

\*NS: Not Specified

## 2.3 Results & Discussion

The colour of all drinking water samples was < 5 Hazen unit and odour of the samples was also agreeable. All parameters are found to be within the specified limit of the Drinking water Standard.

### pH

The limit of pH value for drinking water is specified as 6.5 to 8.5. pH value in the studied area varied from 7.0 to 7.9 pH unit. All the sampling points showed pH values within the prescribed limit by Indian Standards.

### Total Dissolved Solids (TDS)

TDS values in the studied area varied between 600 -1800 mg/l. None of the sampling points showed higher TDS values than the prescribed limit by Indian standards .

### Conductivity

Electrical Conductivity is the ability of a solution to transfer (conduct) electric current. Conductivity is used to measure the concentration of dissolved solids which have been ionized in a polar solution such as water. The conductivity in the samples collected during the month of July ranged from 1000-3300  $\mu\text{s}/\text{cm}$ . Electrical conductivity standards do not appear in BIS standards for drinking water.

### BOD

BOD value in the studied area was less than 2.0 mg/L. Indian standards does not show any standard values for BOD in drinking water.

### Chlorides

Excessive chloride concentration increase rates of corrosion of metals in the distribution system. This can lead to increased concentration of metals in the supply. Chloride value in the studied area varied between 400-600 mg/l and is found to be within the Permissible limit of the Drinking Water Standard.

### Calcium

Calcium value in the studied area varied between 45 - 80 mg/l and is found to be within the Permissible limit of the Drinking Water Standard. If calcium is present beyond the maximum acceptable limit, it causes incrustation of pipes.

### **Magnesium**

Magnesium value in the studied area varied between 30 – 80 mg/l. All the locations had Magnesium within the prescribed limits of 30-100 mg/L.

### **Total Hardness**

Hardness value in the studied area varied between 330-430 mg/l and is found to be within the Permissible limit of the Drinking Water Standard. The prescribed limit by Indian Standards is 200-600 mg/L.

### **Iron**

Iron value in the studied area was below 0.01mg/L and hence well below the permissible limit as per Indian Standards is 0.3 mg/L. The excess amount of iron causes slight toxicity; gives stringent taste to water.

### **Fluoride**

Fluoride value in the studied area varied between 0.1 – 1.4 mg/l and hence well below the permissible limit as per Indian Standards is 1.0-1.5 mg/L. Moderate amounts lead to dental effects, but long-term ingestion of large amounts can lead to potentially severe skeletal problems.

### **Sulphates**

Sulphate value in the studied area varied between 100 – 330 mg/l. All the sampling points showed sulphate values within the prescribed limits by Indian Standards (200-400 mg/L). Sulphate content in drinking water exceeding the 400 mg/L imparts bitter taste.

### **Nitrites (NO<sub>2</sub>) and Nitrates (NO<sub>3</sub>)**

Nitrite values in all the water samples were <0.1. There are no specified standard values for Nitrites in Drinking water. The mean Nitrate values in drinking water of KPT was 4.10 mg/l which is well within the permissible limit of the Drinking water Standard.

### **Salinity**

Salinity in drinking water in the present samples collected ranged from 0.6 to 0.9 %. There are no prescribed Indian standards for salinity in Drinking water.



### **Sodium and Potassium Salts**

Sodium values in the samples collected ranged from 100 - 2000 mg/l and Potassium salts ranged from 2.2 to 3.0 mg/l. There are no prescribed limits of Sodium and Potassium in Indian standards for Drinking water.

### **Heavy Metals in Drinking Water**

In the present study period drinking water samples were analyzed for Mn, Cr, Cu, Cd, As, Hg, Pb and Zn. All these heavy metals were well below the permissible limits prescribed by the Indian Standards.

### **Bacteriological Study**

Analysis of the bacteriological parameter at all location shows that Bacteria is not present and hence Bacterial count is in line with the permissible limit of drinking water. This shows that all the drinking water samples were safe from any bacteriological contamination.

## **2.4 Conclusions**

These results are compared with acceptable limits as prescribed in IS 10500:2012 – Drinking Water Specification. It is seen from the analysis data that during the study period the water was safe for human consumption at all drinking water monitoring stations.

### 3. Noise Level Monitoring

Noise sources in port operations include cargo handling, vehicular traffic, and loading / unloading containers and ships. Noise Monitoring was done at 13 stations at Kandla, Vadinar and Township area.

#### 3.1 Method of Monitoring

Sampling was done at all stations for 24 hour period. Data was recorded using automated sound level meter. The intensity of sound was measured in sound pressure level (SPL) and common unit of measurement is decibel (Db).

#### 3.2 Results

Table 16: Noise Monitoring data for ten locations of Deendayal Port and two locations of Vadinar Port

Sr. No.	Location	Day Time Average Noise Level (SPL) in dB(A)	Night Time Average Noise Level (SPL) in dB(A)
	<b>Sampling Time</b>	<b>6:00 am to 10:00 PM</b>	<b>10:00PM to 6:00 AM</b>
1	Marine Bhavan	60.71	55.49
2	Nirman Building 1	58.02	52.12
3	Tuna Port	53.16	46.37
4	Main Gate North	56.47	53.21
5	West Gate I	61.41	53.6
6	Canteen Area	56.78	48.45
7	Main Road	59.41	56.44
8	ATM Building	63.81	55.02
9	Wharf Area /Jetty Area	65.66	56.59
10	Port & Custom Office	53.59	49.22
	<b>Vadinar Port</b>		
11	Entrance Gate of Vadinar Port	56.32	54.2
12	Nr. Port Colony, Vadinar	55.5	54.8
13	Nr. Vadinar Jetty	58.76	55.4

**3.3 Conclusions-** Noise sources in port operations include cargo handling, vehicular traffic, and loading / unloading containers and ships. The Day Time Average Noise Level (SPL) in all ten locations at Deendayal Port ranged from 56.25 dB(A) to 69.51 dB(A) and it was within the permissible limits of 75 dB(A) for the industrial area for the daytime. The Night Time Average Noise Level (SPL) in all ten locations of Deendayal Port ranged from 48.28 dB to 62.33 dB(A) and it was within the permissible limits of 70 dB(A) for the industrial area for the night time.

#### 4. Soil Monitoring

Sampling and analysis of soil samples were undertaken at six locations within the study area (Deendayal Port and Vadinar Port) as a part of EMP. The soil sampling locations are initially decided based on the locations as provided in the tender document of the Deendayal Port.

##### 4.1 Methodology

The soil samples were collected in the month of July 2021. The samples collected from the all locations are homogeneous representative of each location. At random locations were identified at each location and soil was dug from 30 cm below the surface. It was uniformly mixed before homogenizing the soil samples. The samples were filled in polythene bags, labeled in the field with number and site name and sent to laboratory for analysis.

##### 4.2 Results

**Table-17: Chemical Characteristics of Soil in the Study Area**

Sr. No.	Parameter	Unit	Station Name					
			SL1	SL2	SL3	SL4	SL5	SL6
			Tuna Port	IFFCO Plant	Khori Creek	Nakti Creek	KPT Admin Site	KPT Colony
			Near main gate of Port	10 m away from main gate	Sand from creek at low tide		Vadinar	
1	Texture		Sandy Loam	Sandy Loam	Sandy Loam	Sandy Loam	Sandy Loam	Sandy Loam
2	pH	-	8.56	8.11	8.38	8.33	8.12	8.42
3	Electrical Conductivity	µs/cm	26,800.0	23,800.0	23,700.0	16,260.0	509.0	419.0
4	Moisture	%	23.66	22.09	24.41	23.65	9.44	7.59
5	Total Organic Carbon	%	0.16	0.24	0.32	0.10	0.20	0.12
6	Alkalinity	mg/kg	140.14	140.14	100.10	80.08	100.10	60.06
7	Total Nitrogen	%	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001
8	Chloride	mg/kg	3,908.6	4,309.5	6,114.0	3,959.0	39.3	68.7
9	Sulphate	mg/kg	203.0	177.9	113.8	93.8	13.4	15.5
10	Phosphorus	mg/kg	0.97	0.80	1.24	1.77	0.80	0.97
11	Potassium	mg/kg	779.4	644.4	1,135.8	766.8	129.6	180.0
12	Sodium	mg/kg	2,241.0	3,556.8	3,981.6	3,038.4	1,220.0	1,445.4
13	Calcium	mg/kg	144.29	128.22	168.30	224.40	104.20	56.11
14	Copper as Cu	mg/kg	42.6	61.2	38.2	22.6	16.2	23
15	Lead as Pb	mg/kg	4.2	3.2	3.6	3.8	ND	ND
16	Nickel as Ni	mg/kg	36.2	31.6	39.4	22.6	18.3	21.2
17	Zinc as Zn	mg/kg	58.60	39.25	52.4	46.60	46.80	38.20
18	Cadmium as Cd	mg/kg	ND	ND	ND	ND	ND	ND

### 4.3 Discussion

- The data shows that value of pH ranges from 8.11 at Nakti Creek to 8.56 at Tuna Creek indicating that all soil samples are neutral to slight basic. Tuna port samples showed maximum conductivity of 26,800 $\mu$ mhos/cm, while Nakti Creek location showed minimum conductivity of 16,260  $\mu$ mhos/cm. Conductivity at Vadinar Port was 509 and 419  $\mu$ mhos/cm at Admin site and Vadinar Port colony respectively.
- Total organic Carbon ranged from 0.1 % to 0.3 at Deendayal Port. At Vadinar Port, organic carbon content ranged from 0.1 % to 0.2 %.
- The concentration of Phosphorus and Potassium in the soil samples varies from 0.8 to 1.77 mg/kg and 600.0 to 1150 mg/kg respectively at Deendayal Port. The mean concentration of Phosphorous at Vadinar site was 0.89 mg/kg and mean concentration of Potassium at Vadinar site was 154.8 mg/kg.

These differences in NPK in soil at different locations are due to the dissimilar nature of soil at each of the locations. Samples SL3 & SL4 (Khori Creek & Nakti Creek) are of saline nature as they are coastal soil; where as other locations are inland locations and have different chemical properties.

### Heavy Metals in the Soil

Traces of Copper, Lead, Nickel and Zinc were observed in the soil samples collected from all the four locations of Deendayal Port and two locations of Vadinar Port. Cadmium metal was not detected in the Soil.

### 4.4 Conclusion

The soils of Deendayal Port and Vadinar Port appears to be neutral to basic with varying levels of Chloride, Sulphate, NPK and Calcium. As the nature of soil at different locations are different with respect to its proximity to the sea, the samples showed high degree of variations in their chemical properties.

## 5. Sewage Treatment Plant Monitoring

This involves safe collection of waste water (spent/used water) from wash areas, bathroom, industrial units, etc., waste from toilets of various buildings and its conveyance to the treatment plant and final disposal in conformity with the requirement and guide lines of State Pollution Control Board and other statutory bodies.

### 5.1 Methodology for STP Monitoring

To monitor the working efficiency of Sewage Treatment Plant (STP), STP Inlet and Outlet Samples were collected once a week. Locations selected are namely Gopalpuri Township, Deendayal Port and Vadinar. Samples were collected in 1 lit. Carboys and were analyzed in laboratory for various parameters.

### 5.2 Results

- **Kandla STP**

**Table 18: Sewage Water Monitoring at Kandla STP (1<sup>st</sup> Week)**

Date of Sampling		05.07.2021		
Sr. No.	Parameters	Unit	Results	
			KPT STP I/L	KPT STP O/L
1	pH	pH unit	7.5	7.8
2	Total Suspended Solids	mg/l	125.4	64.3
3	Residual Chlorine	mg/l	<1.0	<0.5
4	COD	mg/l	333.3	102
5	BOD @ 27 °C	mg/l	110.0	26.0
<b>Aeration Tank</b>				
6	MLSS	mg/l	18.0	
7	MLVSS	%	88.0	

**Table 19: Sewage Water Monitoring at Kandla STP (2<sup>nd</sup> Week)**

Date of Sampling		15.07.2021		
Sr. No.	Parameters	Unit	Results	
			KPT STP I/L	KPT STP O/L
1	pH	pH unit	7.6	7.68
2	Total Suspended Solids	mg/l	350	46
3	Residual Chlorine	mg/l	<1.0	<0.5
4	COD	mg/l	585	98
5	BOD @ 27 °C	mg/l	196.0	26.0
<b>Aeration Tank</b>				
6	MLSS	mg/l	24.0	
7	MLVSS	%	82.0	

**Table 20: Sewage Water Monitoring at Kandla STP (3<sup>rd</sup> Week)**

Date of Sampling		20.07.2021		
Sr. No.	Parameters	Unit	Results	
			KPT STP I/L	KPT STP O/L
1	pH	pH unit	7.73	7.61
2	Total Suspended Solids	mg/l	192.6	62
3	Residual Chlorine	mg/l	<1.0	<0.5
4	COD	mg/l	222	20
5	BOD @ 27 °C	mg/l	68.0	8.0
<b>Aeration Tank</b>				
6	MLSS	mg/l	16.0	
7	MLVSS	%	86.0	

**Table 21: Sewage Water Monitoring at Kandla STP (4<sup>th</sup> Week)**

Date of Sampling		26.07.2021		
Sr. No.	Parameters	Unit	Results	
			KPT STP I/L	KPT STP O/L
1	pH	pH unit	Plant was not working	
2	Total Suspended Solids	mg/l		
3	Residual Chlorine	mg/l		
4	COD	mg/l		
5	BOD @ 27 °C	mg/l		
<b>Aeration Tank</b>				
6	MLSS	mg/l	-	
7	MLVSS	%	-	

- Gopalpuri Colony STP**

**Table 22: Sewage Water Monitoring at Gopalpuri STP (1<sup>st</sup> Week)**

Date of Sampling		05.07.2021		
Sr. No.	Parameters	Unit	Results	
			Gopalpuri STP I/L	Gopalpuri STP O/L
1	pH	pH unit	7.7	7.63
2	Total Suspended Solids	mg/l	408.3	38.3
3	Residual Chlorine	mg/l	<1.0	<0.5
4	COD	mg/l	262.6	102
5	BOD @ 27 °C	mg/l	82.0	28.0
<b>Aeration Tank</b>				
6	MLSS	mg/l	12.0	
7	MLVSS	%	92.0	

**Table 23: Sewage Water Monitoring at Gopalpuri STP (2<sup>nd</sup> Week)**

Date of Sampling		15.07.2021		
Sr. No.	Parameters	Unit	Results	
			Gopalpuri STP I/L	Gopalpuri STP O/L
1	pH	pH unit	7.3	7.43
2	Total Suspended Solids	mg/l	333	69
3	Residual Chlorine	mg/l	<1.0	<0.5
4	COD	mg/l	444.4	103
5	BOD @ 27 °C	mg/l	142.0	28.0
<b>Aeration Tank</b>				
6	MLSS	mg/l	16.0	
7	MLVSS	%	89.0	

**Table 24: Sewage Water Monitoring at Gopalpuri STP (3<sup>rd</sup> Week)**

Date of Sampling		20.07.2021		
Sr. No.	Parameters	Unit	Results	
			Gopalpuri STP I/L	Gopalpuri STP O/L
1	pH	pH unit	7.39	7.43
2	Total Suspended Solids	mg/l	166.6	36.7
3	Residual Chlorine	mg/l	<1.0	<0.5
4	COD	mg/l	230	58
5	BOD @ 27 °C	mg/l	70.0	19.0
<b>Aeration Tank</b>				
6	MLSS	mg/l	12.0	
7	MLVSS	%	92.0	



**Table 25: Sewage Water Monitoring at Gopalpuri STP (4<sup>th</sup> Week)**

Date of Sampling		26.07.2021		
Sr. No.	Parameters	Unit	Results	
			Gopalpuri STP I/L	Gopalpuri STP O/L
1	pH	pH unit	7.28	7.4
2	Total Suspended Solids	mg/l	160	38
3	Residual Chlorine	mg/l	<1.0	<0.5
4	COD	mg/l	210	62
5	BOD @ 27 °C	mg/l	62.0	19.0
<b>Aeration Tank</b>				
6	MLSS	mg/l	11.0	
7	MLVSS	%	96.0	

- **Vadinar STP**

**Table 26: Sewage Water Monitoring at Vadinar STP (1<sup>st</sup> Week)**

Date of Sampling		05.07.2021		
Sr. No.	Parameters	Unit	Results	
			Vadinar STP I/L	Vadinar O/L
1	pH	pH unit	7.23	NOT WORKING
2	Total Suspended Solids	mg/l	8	
3	Residual Chlorine	mg/l	70.0	
4	COD	mg/l	86.0	
5	BOD @ 27 °C	mg/l	27.0	

**Table 27: Sewage Water Monitoring at Vadinar STP (2<sup>nd</sup> Week)**

<b>Date of Sampling</b>	<b>15.07.2021</b>
-------------------------	-------------------

Sr. No.	Parameters	Unit	Results	
			Vadinar STP I/L	Vadinar O/L
1	pH	pH unit	7.22	NOT WORKING
2	Total Suspended Solids	mg/l	62	
3	Residual Chlorine	mg/l	<1.0	
4	COD	mg/l	82.0	
5	BOD @ 27 °C	mg/l	27.0	

**Table 28: Sewage Water Monitoring at Vadinar STP (3<sup>rd</sup> Week)**

<b>Date of Sampling</b>	<b>20.07.2021</b>
-------------------------	-------------------

Sr. No.	Parameters	Unit	Results	
			Vadinar STP I/L	Vadinar O/L
1	pH	pH unit	7.22	NOT WORKING
2	Total Suspended Solids	mg/l	62	
3	Residual Chlorine	mg/l	<1.0	
4	COD	mg/l	82.0	
5	BOD @ 27 °C	mg/l	27.0	

**Table 29: Sewage Water Monitoring at Vadinar STP (4<sup>th</sup> Week)**

<b>Date of Sampling</b>	<b>26.07.2021</b>
-------------------------	-------------------

Sr. No.	Parameters	Unit	Results	
			Vadinar STP I/L	Vadinar O/L
1	pH	pH unit	7.18	NOT WORKING
2	Total Suspended Solids	mg/l	72	
3	Residual Chlorine	mg/l	<1.0	
4	COD	mg/l	80.0	
5	BOD @ 27 °C	mg/l	26.0	

### 5.3 Conclusions:

The GPCB standards of BOD, TSS and Residual Chlorine for STP outlet are 20 mg/lit, 30 mg/lit & 0.5 mg/lit respectively. It is suggested to do treatment on regular basis to avoid flow of contaminated/polluted water into the sea. Also, the STP at Vadinar is also non-functional and thus, steps should be taken to commission the STP at Vadinar Port. Hence, currently only inlet samples are collected and analysed. And the sample of kandla stp was not collected in the last week of July 2021 as plant was not working.

## 6. Marine Water Monitoring

The Forty Second Amendment to the Constitution in 1976 underscored the importance of 'green thinking'. Article 48A enjoins the state to protect and improve the environment and safeguard the forests and wildlife in the country. Further, Article 51A(g) states that the "fundamental duty of every citizen is to protect and improve the natural environment including forests, lakes, rivers and wildlife and to have compassion for living creatures".

Policy Statement for Abatement of Pollution (1992) has suggested developing relevant legislation and regulation, fiscal incentives, voluntary agreements and educational programs and information campaigns. It emphasizes the need for integration by incorporating environmental considerations into decision making at all levels by adopting frameworks namely, pollution prevention at source, application of best practicable solution, ensure polluter pays for control of pollution, focus on heavily polluted areas and river stretches and involve public in decision-making. The National Conservation Strategy and Policy Statement on Environment and Development, (1992) aimed at "integrating environmental concerns with developmental imperatives to meet the challenges by redirecting the thrust of our developmental process so that the basic needs of our people could be fulfilled by making judicious and sustainable use of natural resources." The priorities mentioned in this policy document include the sustainable use of land and water resources, prevention and control of pollution and preservation of biodiversity.

The National Water Policy, (2002) contains provisions for developing, conserving, sustainable utilizing and managing this important water resources and need to be governed by national perspectives.

### Marine Environment

On national and state levels, we have several policies and regulation like Water (Prevention and Control of Pollution) Act, 1974, to regulate pollution discharges and restore water quality of our aquatic resources including the prescription of monitoring activities. One of the important provisions of the Water Act, 1974, is to maintain and restore the 'wholesomeness' of our aquatic resources. Water quality monitoring is one of the first steps required in the rational development and management of water resources. In the field of water quality management, there has been a steady evolution in procedures for designing system to obtain information on the changes of water quality. The monitoring comprises all activities to obtain 'information' with respect to the water system.

## Sampling Stations

The monitoring of marine environment for the study of biological and ecological parameters was carried out on 10<sup>th</sup>& 12<sup>th</sup> July -2021 in harbor regions of KPT and on 10<sup>th</sup> July-2021 at Vadinar during spring tide period of New moon phase of Lunar Cycle. The monitoring of marine environment for the study of biological and ecological parameters was repeated again on 17<sup>th</sup>& 19<sup>th</sup> July 2021 in harbor regions of KPT. 17<sup>th</sup> July -2021 in Vadinar during Neap tide period first quarter of Lunar Cycle..

Plankton samples from sub surface layer was collected both during high tide period and low tide period from 3 water quality monitoring stations of KPT harbour area and two stations in Nakti creek and one station in Khori creek. The same sampling schedule was repeated during consecutive spring tide and neap tide in same month. Plankton samples from sub surface layer was collected both during high tide period and low tide period from 1 water quality monitoring stations near Vadinar jetty area during spring tide and neap tide in this month .Collected water samples were processed for estimation of Chlorophyll- a, Pheophytin- a, qualitative & quantitative evaluation of phytoplankton, qualitative & quantitative evaluation zooplanktons (density and their population).

### Sampling Locations

Offshore monitoring requirement	Number of locations
Offshore Installations	3 in Kandla creek 2 in Nakti creek 1 in Khori creek 1 near Vadinar Jetty 1 near 1 <sup>st</sup> SBM
<b>Total Number of locations</b>	<b>8</b>

#### 6.1 Marine Water Quality

Marine water quality of marine waters of Deendayal Port Harbor waters, Khori and Nakti Creeks and two locations of Vadinar are monitored for various physico-chemical parameters during spring and neap tide of each month.

The results of marine water quality and Marine sediments are as below;

**Table 30: Marine Water Quality Monitoring Parameters for location near KPT colony**

Sr. No.	Parameters	Unit	Kandla Creek Near KPT colony (1)			
			23°0'58"N 70°13'22."E			
			Spring Tide		Neap Tide	
			Tide →	High Tide	Low Tide	High Tide
1	pH	pH unit	7.29	7.25	7.13	7.15
2	Color	-	Colorless	Colorless	Colorless	Colorless
3	Odor	-	Odorless	Odorless	Odorless	Odorless
4	Salinity	ppt	32.0	31.8	32.1	32.6
5	Turbidity	NTU	30	28	35	27
6	Total Dissolved Solids	mg/l	37802.0	23743	43720.0	43881.0
7	Total Suspended Solids	mg/l	624	412	409	261
8	Total Solids	mg/l	38426.2	24155.4	44129.0	44142.0
9	DO	mg/l	4.5	5	4.9	5.3
10	COD	mg/l	72.0	68.0	74.0	76.0
11	BOD	mg/l	<2	<2	<2	<2
12	Silica	mg/l	0.23	0.38	0.30	0.48
13	Phosphate	mg/l	0.31	0.28	0.19	0.35
14	Sulphate	mg/l	2856	2556	2076	2160
15	Nitrate	mg/l	2.10	2.04	2.40	2.04
16	Nitrite	mg/l	<0.05	<0.05	<0.05	<0.05
17	Calcium	mg/l	521.04	561.12	521.04	440.88
18	Magnesium	mg/l	1798.2	1798.2	1773.9	1871.1
19	Sodium	mg/l	14122.0	14820.0	10110.0	10872.0
20	Potassium	mg/l	325.0	289.0	321.0	289.0
21	Iron	mg/l	1.12	1.42	1.52	1.45
22	Chromium	mg/l	0.12	0.13	0.12	0.11
23	Copper	mg/l	0.12	0.19	0.06	0.08
24	Arsenic	mg/l	<0.01	<0.01	<0.01	<0.01
25	Cadmium	mg/l	0.07	0.06	0.05	0.06
26	Mercury	mg/l	<0.001	<0.001	<0.001	<0.001
27	Lead	mg/l	0.20	0.19	0.11	0.12
28	Zinc	mg/l	0.05	0.06	0.06	0.07

**Table 31: Marine Water Quality Monitoring Parameters for location near passenger Jetty One at Kandla**

Sr. No.	Parameters	Unit	Near passenger Jetty One (2)			
			23° 0'18 "N 70°13'31"E			
			Spring Tide		Neap Tide	
			Tide →	High Tide	Low Tide	High Tide
1	pH	pH unit	5.3	4.7	7.3	7.27
2	Color	-	80.0	76.0	Colorless	Colorless
3	Odor	-	<2	<2	Odorless	Odorless
4	Salinity	ppt	0.32	0.68	33.0	32.6
5	Turbidity	NTU	0.32	0.26	22	28
6	Total Dissolved Solids	mg/l	2976	2748	46102.0	47052.0
7	Total Suspended Solids	mg/l	2.25	2.03	211	312
8	Total Solids	mg/l	<0.05	<0.05	46313.0	47364.0
9	DO	mg/l	521.04	521.04	4.6	4.9
10	COD	mg/l	1846.8	1773.9	86.0	79.0
11	BOD	mg/l	11052.0	13425.0	<2	<2
12	Silica	mg/l	325.0	306.0	0.39	0.72
13	Phosphate	mg/l	1.55	1.62	0.34	0.30
14	Sulphate	mg/l	0.12	0.14	1956	2520
15	Nitrate	mg/l	0.18	0.16	1.74	2.52
16	Nitrite	mg/l	<0.01	<0.01	<0.05	<0.05
17	Calcium	mg/l	0.07	0.05	480.96	480.96
18	Magnesium	mg/l	<0.001	<0.001	1822.5	1822.5
19	Sodium	mg/l	0.28	0.16	11011.0	10452.0
20	Potassium	mg/l	0.05	0.06	333.0	315.0
21	Iron	mg/l	5.3	4.7	1.56	1.89
22	Chromium	mg/l	80.0	76.0	0.16	0.14
23	Copper	mg/l	<2	<2	0.09	0.08
24	Arsenic	mg/l	0.32	0.68	<0.01	<0.01
25	Cadmium	mg/l	0.32	0.26	0.06	0.07
26	Mercury	mg/l	2976	2748	<0.001	<0.001
27	Lead	mg/l	2.25	2.03	0.16	0.19
28	Zinc	mg/l	<0.05	<0.05	0.06	0.08

**Table 32: Marine Water Quality Monitoring Parameters for location Near Coal Berth**

Sr. No.	Parameters	Unit	Near Coal Berth			
			22°59'12"N 70°13'40"E			
			Spring Tide		Neap Tide	
			Tide →	High Tide	Low Tide	High Tide
1	pH	pH unit	7.51	7.30	7.29	7.5
2	Color	-	Colorless	Colorless	Colorless	Colorless
3	Odor	-	Odorless	Odorless	Odorless	Odorless
4	Salinity	ppt	32.6	32.8	32.4	33.1
5	Turbidity	NTU	35	47	35	47
6	Total Dissolved Solids	mg/l	40788	35363	41086.0	42830.0
7	Total Suspended Solids	mg/l	563	601	215	161
8	Total Solids	mg/l	41351.3	35964.2	41301.0	42991.0
9	DO	mg/l	4.8	5	4.8	5
10	COD	mg/l	88.0	70.0	90.0	79.0
11	BOD	mg/l	<2	<2	<2	<2
12	Silica	mg/l	0.30	0.34	0.42	0.36
13	Phosphate	mg/l	0.28	0.32	0.35	0.38
14	Sulphate	mg/l	2580	3444	3156	3240
15	Nitrate	mg/l	1.93	2.10	2.56	2.46
16	Nitrite	mg/l	<0.05	<0.05	<0.05	<0.05
17	Calcium	mg/l	561.12	480.96	561.12	601.2
18	Magnesium	mg/l	1725.3	1798.2	1725.3	1725.3
19	Sodium	mg/l	15555.0	13252.0	11052.0	11412.0
20	Potassium	mg/l	389.0	296.0	315.0	296.0
21	Iron	mg/l	1.47	2.02	2.10	2.02
22	Chromium	mg/l	0.19	0.15	0.12	0.20
23	Copper	mg/l	0.14	0.12	0.06	0.10
24	Arsenic	mg/l	<0.01	<0.01	<0.01	<0.01
25	Cadmium	mg/l	0.04	0.06	0.08	0.07
26	Mercury	mg/l	<0.001	<0.001	<0.001	<0.001
27	Lead	mg/l	0.20	0.18	0.10	0.12
28	Zinc	mg/l	0.08	0.06	0.07	0.06



**Table 33: Marine Water Quality Monitoring Parameters for location Khori creek at Kandla**

Sr. No.	Parameters	Unit	KPT 4			
			Near 15/16 Berth			
			Spring Tide		Neap Tide	
			Tide →	High Tide	Low Tide	High Tide
1	pH	pH unit	7.25	7.20	7.39	7.45
2	Color	-	Colorless	Colorless	Colorless	Colorless
3	Odor	-	Odorless	Odorless	Odorless	Odorless
4	Salinity	ppt	32.4	32.8	32.4	32.2
5	Turbidity	NTU	50	29	51	55
6	Total Dissolved Solids	mg/l	35588	33113	43563.0	44059.0
7	Total Suspended Solids	mg/l	407	420	213	265
8	Total Solids	mg/l	35995.3	33533.4	43776.0	44324.0
9	DO	mg/l	5.2	4.8	5.3	4.7
10	COD	mg/l	68.0	79.0	76.0	86.0
11	BOD	mg/l	<2	<2	<2	<2
12	Silica	mg/l	0.81	0.29	0.79	0.37
13	Phosphate	mg/l	0.26	0.34	0.43	0.42
14	Sulphate	mg/l	2388	2652	2280	2376
15	Nitrate	mg/l	1.74	1.96	2.10	2.57
16	Nitrite	mg/l	<0.05	<0.05	<0.05	<0.05
17	Calcium	mg/l	601.20	561.12	601.2	561.12
18	Magnesium	mg/l	1749.6	1822.5	1725.3	1798.2
19	Sodium	mg/l	10026.0	11252.0	10512.0	9899.0
20	Potassium	mg/l	302.0	378.0	266.0	275.0
21	Iron	mg/l	1.66	1.48	1.45	1.60
22	Chromium	mg/l	0.16	0.16	0.18	0.16
23	Copper	mg/l	0.15	0.10	0.12	0.10
24	Arsenic	mg/l	<0.01	<0.01	<0.01	<0.01
25	Cadmium	mg/l	0.07	0.06	0.06	0.05
26	Mercury	mg/l	<0.001	<0.001	<0.001	<0.001
27	Lead	mg/l	0.16	0.2	0.08	0.10
28	Zinc	mg/l	0.07	0.08	0.05	0.05

**Table 34: Marine Water Quality Monitoring Parameters for location Nakti Creek near Tuna Port**

Sr. No.	Parameters	Unit	Nakti Creek Near Tuna Port			
			22°57'49."N 70° 7'0.67"E			
			Spring Tide		Neap Tide	
			Tide →	High Tide	Low Tide	High Tide
1	pH	pH unit	7.29	7.35	7.2	7.28
2	Color	-	Colorless	Colorless	Colorless	Colorless
3	Odor	-	Odorless	Odorless	Odorless	Odorless
4	Salinity	ppt	32.6	33.2	33.6	33.0
5	Turbidity	NTU	35	29	29	29
6	Total Dissolved Solids	mg/l	38200	18212	46852.0	47695.0
7	Total Suspended Solids	mg/l	324	214	200	196
8	Total Solids	mg/l	38524.4	18426.2	47052.0	47891.0
9	DO	mg/l	5.1	5.1	4.9	5
10	COD	mg/l	80.0	68.0	89.0	78.0
11	BOD	mg/l	<2	<2	<2	<2
12	Silica	mg/l	0.26	0.29	0.62	0.30
13	Phosphate	mg/l	0.28	0.26	0.31	0.38
14	Sulphate	mg/l	2964	3408	3240	3156
15	Nitrate	mg/l	1.95	2.18	2.56	2.49
16	Nitrite	mg/l	<0.05	<0.05	<0.05	<0.05
17	Calcium	mg/l	561.12	521.04	601.2	440.88
18	Magnesium	mg/l	1822.5	1749.6	1798.2	1822.5
19	Sodium	mg/l	11256.0	12625.0	11021.0	11425.0
20	Potassium	mg/l	302.0	366.0	396.0	378.0
21	Iron	mg/l	1.83	1.76	2.02	2.11
22	Chromium	mg/l	0.15	0.18	0.20	0.18
23	Copper	mg/l	0.12	0.11	0.16	0.10
24	Arsenic	mg/l	<0.01	<0.01	<0.01	<0.01
25	Cadmium	mg/l	0.04	0.06	0.07	0.08
26	Mercury	mg/l	<0.001	<0.001	<0.001	<0.001
27	Lead	mg/l	0.18	0.19	0.12	0.16
28	Zinc	mg/l	0.06	0.05	0.06	0.07

**Table 35: Marine Water Quality Monitoring Parameters for location Nakti Creek Near NH-8A at Kandla**

Sr. No.	Parameters	Unit	Nakti Creek Near NH-8A			
			23° 02'01"N 70° 09'31"E			
			Spring Tide		Neap Tide	
			Tide →		High Tide	Low Tide
1	pH	pH unit	7.37	Sampling not possible during Low Tide	7.37	Sampling not possible during Low Tide
2	Color	-	Colorless		Colorless	
3	Odor	-	Odorless		Odorless	
4	Salinity	ppt	33.4		32.4	
5	Turbidity	NTU	27		33	
6	Total Dissolved Solids	mg/l	35166		42125.0	
7	Total Suspended Solids	mg/l	180		164.3	
8	Total Solids	mg/l	35346.3		42289.3	
9	DO	mg/l	5		5.5	
10	COD	mg/l	72.0		79.0	
11	BOD	mg/l	<2		<2	
12	Silica	mg/l	0.61		0.62	
13	Phosphate	mg/l	0.30		0.39	
14	Sulphate	mg/l	2988		3036	
15	Nitrate	mg/l	2.43		2.72	
16	Nitrite	mg/l	<0.05		<0.05	
17	Calcium	mg/l	601.20		521.04	
18	Magnesium	mg/l	1749.6		1749.6	
19	Sodium	mg/l	14485.0		11528.0	
20	Potassium	mg/l	396.0		311.0	
21	Iron	mg/l	1.52		2.06	
22	Chromium	mg/l	0.16		0.19	
23	Copper	mg/l	0.16		0.11	
24	Arsenic	mg/l	<0.01		<0.01	
25	Cadmium	mg/l	0.04		0.06	
26	Mercury	mg/l	<0.001		<0.001	
27	Lead	mg/l	0.17		0.10	
28	Zinc	mg/l	0.06		0.07	

**Table 36: Marine Water Quality Monitoring Parameters for locations Nr. Vadinar Jetty**

Sr. No.	Parameters	Unit	Nr.Vadinar Jetty			
			22°26'25.26"N 69°40'20.41"E			
			Spring Tide		Neap Tide	
			Tide →	High Tide	Low Tide	High Tide
1	pH	pH unit	7.60	7.45	7.5	7.8
2	Color	-	Colorless	Colorless	Colorless	Colorless
3	Odor	-	Odorless	Odorless	Odorless	Odorless
4	Salinity	ppt	33.2	32.6	33.0	33.1
5	Turbidity	NTU	32	28	35	25
6	Total Dissolved Solids	mg/l	37530	35780	43940.0	46623.0
7	Total Suspended Solids	mg/l	327	417	405.5	399.5
8	Total Solids	mg/l	37856.5	36197.4	44345.5	47022.5
9	DO	mg/l	5.2	5.1	5.2	5.1
10	COD	mg/l	68.0	72.0	78.0	79.0
11	BOD	mg/l	<2	<2	<2	<2
12	Silica	mg/l	0.32	0.31	0.42	0.62
13	Phosphate	mg/l	0.26	0.26	0.35	0.33
14	Sulphate	mg/l	2136	2352	2220	2304
15	Nitrate	mg/l	2.72	2.80	2.09	2.44
16	Nitrite	mg/l	<0.05	<0.05	<0.05	<0.05
17	Calcium	mg/l	480.96	561.12	561.12	521.04
18	Magnesium	mg/l	1749.6	1749.6	1579.5	1555.2
19	Sodium	mg/l	16458.0	15555.0	11425.0	12021.0
20	Potassium	mg/l	345.0	388.0	316.0	296.0
21	Iron	mg/l	2.06	2.10	2.45	2.3
22	Chromium	mg/l	0.16	0.20	0.15	0.16
23	Copper	mg/l	0.17	0.18	0.09	0.08
24	Arsenic	mg/l	<0.01	<0.01	<0.01	<0.01
25	Cadmium	mg/l	0.05	0.06	0.05	0.06
26	Mercury	mg/l	<0.001	<0.001	<0.001	<0.001
27	Lead	mg/l	0.14	0.13	0.10	0.10
28	Zinc	mg/l	0.08	0.09	0.05	0.06

### **6.1.1 Marine Sediments**

Sediment samples were collected with Van Veen Grab from the six locations in Kandla Port Waters and two locations in Vadinar Port. Samples were collected and preserved in silver foil in ice box to prevent the contamination/decaying of the samples.

## 6.2 Results

The Sediment Quality results are given in below from table no. 34 A & B.

**Table 34A: Results of Analysis of Sediment of Kandla & Vadinar Port (Spring Tide)**

Sr. No.	Parameters	Unit	KPT - 1	KPT - 2	Khori - 1	Nakti - 1 (Near NH-8A )	Jetty
1	Texture	-	Sandy Loam	Sandy Loam	Sandy Loam	Sandy Loam	Sandy Loam
2	Organic Matter	mg/kg	0.64	1.12	0.39	1.03	1.06
3	Organic Carbon	mg/kg	0.37	0.65	0.22	0.60	0.52
4	Inorganic Phosphate	mg/kg	126.0	125.0	136.0	146.0	152.0
5	Moisture	%	11.70	18.10	6.60	26.1	23.50
6	Aluminium	mg/kg	ND	ND	ND	ND	ND
7	Silica	mg/kg	26.0	24.0	42.0	49.0	41.2
8	Phosphate	mg/kg	9.88	7.82	8.80	9.70	18.00
9	Sulphate	mg/kg	170.0	192.0	259.0	259.0	362.0
10	Nitrite	mg/kg	0.12	0.13	0.11	0.11	0.11
11	Nitrate	mg/kg	9.23	7.82	9.25	9.25	7.52
12	Calcium	mg/kg	144.3	148.0	132.0	124.0	169.0
13	Magnesium	mg/kg	165.2	214.0	122.0	136.0	162.0
14	Sodium	mg/kg	2221.0	1686.0	1882.0	1775.0	3785.0
15	Potassium	mg/kg	641.0	542.0	738.0	562.0	658.0
16	Chromium	mg/kg	123	145	126	130	162
17	Nickel	mg/kg	24.8	22.5	18.9	26.02	38
18	Copper	mg/kg	48	42	20.6	27.5	23.6
19	Zinc	mg/kg	32.60	36.00	30.40	36.00	32.00
20	Cadmium	mg/kg	ND	ND	ND	ND	ND
21	Lead	mg/kg	2.8	1.8	1.2	4.5	5.8
22	Mercury	mg/kg	ND	ND	ND	ND	ND
23	Arsenic	mg/kg	ND	ND	ND	ND	ND

\*Grab samples could not be collected due high current at KPT 3, Natki Creek Near Tuna port & Vadinar SBM

**Table 34B: Results of Analysis of Sediment of Kandla & Vadinar Port (Neap Tide)**

Sr. No.	Parameters	Unit	KPT - 1	KPT - 2	Jetty
1	Texture	-	Sandy loam	Sandy loam	Sandy loam
2	Organic Matter	mg/kg	0.74	1.10	1.10
3	Organic Carbon	mg/kg	0.52	0.62	0.63
4	Inorganic Phosphate	mg/kg	162.0	142.0	162.0
5	Moisture	%	15.62	14.20	21.52
6	Aluminium	mg/kg	ND	ND	ND
7	Silica	mg/kg	16.60	20.30	39.2
8	Phosphate	mg/kg	9.8	7.26	16.66
9	Sulphate	mg/kg	342.0	280.0	289.0
10	Nitrite	mg/kg	0.10	0.11	0.1
11	Nitrate	mg/kg	10.6	9.8	8.02
12	Calcium	mg/kg	141.0	152.0	178.0
13	Magnesium	mg/kg	156.0	214.0	206.0
14	Sodium	mg/kg	2210.0	1786.0	3682.0
15	Potassium	mg/kg	590.0	562.0	666.0
16	Chromium	mg/kg	136	149	158
17	Nickel	mg/kg	26.2	23.5	32
18	Copper	mg/kg	52	46	18.2
19	Zinc	mg/kg	33.20	34.00	22.00
20	Cadmium	mg/kg	ND	ND	ND
21	Lead	mg/kg	2.4	2.2	4.6
22	Mercury	mg/kg	ND	ND	ND
23	Arsenic	mg/kg	ND	ND	ND

\*Grab samples could not be collected due high current at KPT 3,Khori, Natki Creek Near Tuna Port, Vadinar Jetty and Vadinar SBM

**REPORT**  
**ON**  
**ECOLOGICAL MONITORING**  
**OF MARINE ENVIRONMENT**  
**IN**  
**DPT HARBOUR AREA, NEAR BY CREEKS**  
**For**  
**DEENDAYAL PORT TRUST**

JULY, 2021



**INTRODUCTION:**

The Forty Second Amendment to the Constitution in 1976 underscored the importance of 'green thinking'. Article 48A enjoins the state to protect and improve the environment and safeguard the forests and wildlife in the country. Further, Article 51A(g) states that the "fundamental duty of every citizen is to protect and improve the natural environment including forests, lakes, rivers and wildlife and to have compassion for living creatures".

Policy Statement for Abatement of Pollution (1992) has suggested developing relevant legislation and regulation, fiscal incentives, voluntary agreements and educational programs and information campaigns. It emphasizes the need for integration by incorporating environmental considerations into decision making at all levels by adopting frameworks namely, pollution prevention at source, application of best practicable solution, ensure polluter pays for control of pollution, focus on heavily polluted areas and river stretches and involve public in decision-making. The National Conservation Strategy and Policy Statement on Environment and Development, (1992) aimed at "integrating environmental concerns with developmental imperatives to meet the challenges by redirecting the thrust of our developmental process so that the basic needs of our people could be fulfilled by making judicious and sustainable use of natural resources." The priorities mentioned in this policy document include the sustainable use of land and water resources, prevention and control of pollution and preservation of biodiversity.

The National Water Policy, (2002) contains provisions for developing, conserving, sustainable utilizing and managing this important water resources and need to be governed by national perspectives.

**MARINE ENVIRONMENT:**

On national and state levels, we have several policies and regulation like Water (Prevention and Control of Pollution) Act, 1974, to regulate pollution discharges and restore water quality of our aquatic resources including the prescription of monitoring activities. One of the important provisions of the Water Act, 1974, is to maintain and restore the 'wholesomeness' of our aquatic resources. Water quality monitoring is one of the first steps required in the rational development and management of water resources. In the field of water quality management, there has been a steady evolution in procedures for designing system to obtain information on the changes of water quality. The monitoring comprises all activities to obtain 'information' with respect to the water system.

**Sampling Stations:**

The monitoring of marine environment for the study of biological and ecological Parameters was carried out on 10th July, 2021 in in harbour region of DPT, and on 12<sup>th</sup>July, 2021 in creeks near by the port during spring tide .The monitoring of marine environment for the study of biological and ecological parameters was repeated again on 17<sup>th</sup> July, 2021 in harbour region of DPT and on 19<sup>th</sup>July, 2021 in creeks near by the port during neap tidal condition.

Plankton samples from sub surface layer was collected both during high tide period and low tide period from 3 water quality monitoring stations of DPT harbour area and one stations in Nakti creek and one station in Khori creek. Sampling at second sampling station of Nakti creek was possible only during high tide period. Collected water samples were processed for estimation of Chlorophyll- a, Pheophytin- a, qualitative & quantitative evaluation of phytoplankton, qualitative & quantitative evaluation zooplanktons(density and their population).

**TABLE #1 SAMPLING LOCATIONS**

<b>monitoring requirement</b>	<b>Number of locations</b>
Kandla creek	3 in Kandla creek
Nakti creek	2 in Nakti creek
Khori Creek	1 in Khori creek
<b>Total Number of locations</b>	<b>6</b>

### Sampling methodology adopted:

A marine sampling is an estimation of the body of information in the population. The theory of the sampling design is depending upon the underlying frequency distribution of the population of interest. The requirement for useful water sampling is to collect a representative sample of suitable volume from the specified depth and retain it free from contamination during retrieval.

50 liters of the water sample were collected from Sub surface by using bucket. From the collected water sample 1 liter of water sample were taken in an opaque plastic bottle for chlorophyll estimation, thereafter plankton samples were collected by using filtration assembly with nilyobolt cloth of 20 $\mu$ m mesh size.

### Samples Processing for chlorophyll estimation:

Samples for the chlorophyll estimation were preserved in ice box on board in darkness to avoid degradation in opaque container covered with aluminium foil. Immediately after reaching the shore after sampling, 1 liter of collected water sample was filtered through GF/F filters (pore size 0.45  $\mu$ m) by using vacuum filtration assembly. After vacuum filtration the glass micro fiber filter paper was grunted in tissue grinder, macerating of glass fiber filter paper along with the filtrate was done in 90% aqueous Acetone in the glass tissue grinder with glass grinding tube. Glass fiber filter paper will assist breaking the cell during grinding and chlorophyll content was extracted with 10 ml of 90% Acetone, under cold dark conditions along with saturated magnesium carbonate solution in glass screw cap tubes. After an extraction period of 24 hours, the samples were transferred to calibrated centrifuge tubes and adjusted the volume to original volume with 90% aqueous acetone solution to make up the evaporation loss. The extract was clarified by using centrifuge in closed tubes. The clarified extracts were then decanted in clean cuvette and optical density was observed at wavelength 664, 665 nm. By using corrected optical density, Chlorophyll-a value was calculated as given in (APHA, 1998).

### PLANKTON:

The entire area open water in the sea is the pelagic realm. Pelagic organisms live in the open sea. In contrast to the pelagic realm, the benthic realm comprises organisms and zone of the bottom of the sea. Vertically the pelagic realm can be dividing into two zones based on light penetration; upper photic or euphotic zone and lower dark water mass, aphotic zone below the photic zone.

The term plankton is general term for organisms have such limited powers of locomotion that they are at the mercy of the prevailing water movement. Plankton is subdivided to phytoplankton and

zooplankton. Phytoplankton is free floating organisms that are capable of photosynthesis and zooplankton is the various free floating animals.

Pelagic zone, represents the entire ocean water column from the surface to the deepest depths, is home to a diverse community of organisms. Differences in their locomotive ability categorize the organisms in the pelagic realm into two, **plankton** and **nekton** (Lalli and Parsons, 1997). **Plankton** consists of all organisms drifting in the water and is unable to swim against water currents, whereas **Nekton** includes organisms having strong locomotive power. Ecological studies on the plankton community, which form the base of the aquatic food chain, help in the better understanding of the dynamics and functioning of the marine ecosystem. The term 'Plankton' first coined by Victor Hensen (1887), Plankton, (Greek word: *planktos* meaning "passively drifting or wandering") is defined as drifting or free-floating organisms that inhabit the pelagic zone of water. Based on their mode of nutrition planktonic organisms are categorised into phytoplankton (organisms having an autotrophic mode of nutrition) and zooplankton (organisms having a heterotrophic mode of nutrition).

#### Phytoplankton in the marine environment:

Phytoplankton is free floating unicellular, filamentous and colonial eutrophic organisms that grow in aquatic environments whose movement is more or less dependent upon water currents. These micro flora acts as primary producers as well as the basis of food chain, source of protein, bio purifier and bio indicators of the aquatic ecosystems of which diverse array of the life depends .They are considered as an important component of aquatic flora, play a key role in maintaining equilibrium between abiotic and biotic components of aquatic ecosystem.

The phytoplankton includes a wide range of photosynthetic and phototrophic organisms. Marine phytoplankton is mostly microscopic and unicellular floating flora, which are the primary producers that support the pelagic food-chain. The two most prominent groups of phytoplankton are diatoms (Bacillariophyceae) and dinoflagellates (Dinophyceae).The phytoplankton those normally captured in the net from the Gulf of Kutch is normally dominated by these two major groups; diatoms and dinoflagellates. Phytoplankton also include numerous and diverse collection of extremely small, motile algae which are termed micro flagellates (naked flagellates) as well as and Cyanophytes (blue-green algae).

Algae are an ecologically important group in most aquatic ecosystems and have been an important component of biological monitoring programs. Algae are ideally suited for water quality assessment because they have rapid reproduction rates and very short life cycles, making them valuable indicators of short-term impacts.

Aquatic populations are impacted by anthropogenic stress, resulting in a variety of alterations in the biological integrity of aquatic systems. Algae can serve as an indicator of the degree of deterioration of water quality, and many algal indicators have been used to assess environmental status.

### **Zooplankton in the marine environment:**

Zooplankton includes a taxonomically and morphologically diverse community of heterotrophic organisms that drift in the waters of the world's oceans. Qualitative and quantitative studies on zooplankton community is a prerequisite to delineate the ecological processes active in the marine ecosystem. Zooplankton community plays a pivotal role in the pelagic food web as the primary consumers of phytoplankton and act as the food source for organisms in the higher trophic levels, particularly the economically essential groups such as fish larvae and fishes. They also function in the cycling of elements in the marine ecosystem. The dynamics of the zooplankton community, their reproduction, growth and survival rate are all significant factors determining the recruitment and abundance of fish stocks as they form an essential food for larval, juvenile and adult fishes (Beaugrand et al., 2004). Zooplankton grazing in the marine environment controls the primary production and helps in determining the pelagic ecosystem (Banse, 1995). Through grazing in surface waters and following the production of sinking faecal matters and also by the active transportation of dissolved and particulate matter to deeper waters via vertical migration, they help in the transport of organic carbon to deep ocean layers and thus act as key drivers of biological pump' in the marine ecosystem. Zooplankton grazing and metabolism also, transform particulate organic matter into dissolved forms, promoting primary producer community, microbial demineralization, and particle export to the ocean's interior.

The categorisation of zooplankton into various ecological groups is based on several factors such as duration of planktonic life, size, food preferences and habitat. As they vary significantly in size from microscopic to metazoic forms, the classification of zooplankton based on size has paramount importance in the field of quantitative plankton research.


Based on the duration of planktonic life, zooplankton are categorised into Holoplankton (organisms which complete their entire lifecycle as plankton) and Meroplankton (organisms which are planktonic during the early part of their lives such as the larval stages of benthic and nektonic organisms). Tychoplankton are organisms which live a brief planktonic life, such as the benthic crustaceans (cumaceans, mysids, isopods) which ascend to the water column at night for feeding and certain ectoparasitic copepods, they leave the host and spend their life as plankton during their breeding cycle.

Zooplankton can be subdivided into holoplankton, i.e., permanent members of the plankton (e.g., Calanoid copepods), and meroplankton, i.e., temporary members in the plankton e.g., larvae of fish, shrimp, and crab). The meroplankton group consists of larval and young stages of animals that will adopt a different lifestyle once they mature. In contrast to phytoplankton which consist of a relatively smaller variety of organisms, Zooplankton are extremely divers, consist of a host of larval and adult forms representing many animal phylum.

Among the zooplankton one group always dominate than others; members of sub class copepods (Phylum Athropoda), and Tintinids (Phylum Protozoa) among the net planktons. These small animals are of vital importance in marine ecosystem as one of the primary herbivores animals in the sea, and it is they provide vital link between primary producer (autotrophs) and numerous small and large marine consumers.

As their community structure and function are highly susceptible to changes in the environmental conditions regular monitoring of their distribution as well as their interactions with various physicochemical parameters is inevitable for the sustainable management of the ecosystem (Kusum et al., 2014). Of all the marine zooplankton groups, copepods mainly calanoid copepods are the dominant groups in marine subtropical and tropical waters and exhibit considerable diversity in morphology and habitats they occupy (Madhupratap, 1991;)

It has been well established that potential of pelagic fishes viz. finfishes, crustaceans, molluscs and marine mammals either directly or indirectly depend on zooplankton. The herbivorous zooplankton is efficient grazers of the phytoplankton and is referred to as living machines transforming plant material into animal tissue. Hence they play an essential role as the intermediaries for nutrients/energy transfer between primary and tertiary trophic levels. Due to their large density, shorter lifespan, drifting nature, high group/species diversity and different tolerance to the stress, they used as the indicator organisms for the physical, chemical and biological processes in the aquatic ecosystem (Ghajibhiye, 2002).



**Spatial distribution of Plankton:**

A characteristic of plankton population is that they tend to occur in patches, which are varying spatially on a scale of few meters to far as few kilometres in distance. They also vary in time scale, season as well as vertically in the water column. It is this patchiness and its constant changes in time and spot, that has made it so difficult for plankton biologist to learn about the ecology of plankton. The biological factors that causes this patchiness is due to the ability of zooplankton to migrate vertically and graze out the phytoplankton at a rapid rate that can create patchiness. Similarly the active swimming ability by certain zooplankton organisms can cause to aggregate in dense group.

At its most extreme, because the water in which plankton is suspended is constantly moving, each sample taken by the plankton biologists remain a different volume of water, so each sample is unique and replicate does not exist.

Plankton may also exhibit vertical patchiness. Physical factors contribute to this type of patchiness include light intensity, nutrients and density gradients in the water column.

Phytoplankton in particular tends to be unequally distributed vertically, which leads to the existence of different concentration of a chlorophyll value between photic zone and below the photic zone.

**Methodology adopted for Plankton sampling:**

Mixed plankton sample were obtained from the sub surface layer at each sampling locations by towing the net horizontally with the weight .After the tow of about 15-30minutes, plankton net was pulled up and washed down to the tail and collected the plankton adhered to plankton net in the collection bucket at the bottom by springing outer and inner surface of the net with sea water, while the net was hanging with the mouth upward. For quantitative evaluation 50 L water samples were collected from subsurface layer and filtered through 20µm mesh size net by using bucket and filtration assembly.

**Preservation and storage:**

Both filtered plankton and those collected from the plankton net were preserved with 5% buffered formalin and stored in 1L plastic container for further processing in the laboratory.

**Sample concentration:**

The collected plankton samples were concentrated by using centrifuge and made up to 50 ml with 5% formalin -Glycerine mixture.

**Taxonomic evaluation:**

Before processing, the sample was mixed carefully and a subsample was taken with a calibrated Stempel-pipette. 1 ml of the concentrated plankton samples were transferred on a glass slide with automatic pipette. The plankton sample on the glass slides were stained by using Lugol's iodine and added glycerine to avoid drying while observation. The plankton samples were identified by using Labex triangular Research microscope with photographic attachment. Microphotographs of the plankton samples were taken for record as well as for confirming the identification. The bigger sized zooplankton was observed through dissecting stereomicroscope with magnification of 20-30 x. Plankton organisms in the whole slide were identified to the lowest axon possible. A thorough literature search was conducted for the identification of the different groups of zooplankton that were encountered

**Cell counts by drop count method:**

The common glass slide mounted with a 1ml of concentrated phytoplankton/zooplankton sample in glycerol and covered with cover slip 22x 60mm was placed under the compound microscope provided with a mechanical stage. The plankton was then counted from the microscopic field of the left top corner of the slide. Then slide is moved horizontally along the right side and plankton in each microscopic field was thus counted. When first microscopic field row was finished the next



consecutive row was adjusted using the mechanical device of the stage. In this way all the plankton present in entire microscopic field are counted. From this total number in 1ml of the concentrated plankton, total number of plankton in the original volume of sample filtered was calculated as units/L.

#### **BENTHIC ORGANISMS:**

Benthos is those organisms that are associated with the sea bed or benthic habitats. Epi- benthic organisms live attached to a hard substratum or rooted to a shallow depth below the surface. In fauna organisms live below the sediment–water interface. Interstitial organisms live and move in pore water among sedimentary grains.

Because the benthic organisms are often collected and separated on sieves, a classification based on the overall size is used. Macro benthos include organisms whose shortest dimension is greater than or equal to 0.5 mm. Meio benthos are smaller than 0.5mm but larger than 42 $\mu$  in size.

The terms such as macro fauna and Meio fauna generally have little relevance with taxonomic classification. The terms Meio fauna and macro fauna depend on the size. Meio fauna were considered as good bioassay of community health and rather sensitive indicators of environmental changes

#### **SAMPLING METHODOLOGY ADOPTED FOR SUB TIDAL REGION:**

Van veen sampler (0.09m<sup>2</sup>) was used for sampling bottom sediments. Two sets of sediments were sampled from each location, one for macro fauna and other for Meio fauna. The macro fauna in the sediments were sieved on board to separate out the organisms. The fixation of Meio fauna is normally done by bulk fixation of the sediment sample. The bulk fixation is done by using 10% formalin (Buffered with borate). The organisms were preserved with seawater as diluting agent.

#### **Sample sieving:**

Sediments samples were sieved to extract the organisms. Sieving was performed carefully as possible to avoid any damage to the animals. The large portion of the sediment was split in to smaller portions and mixed with sea water in a bucket. The cohesive lumps were broken down by continuous stirring. The disaggregated sediments were then passed through the sieves.

#### **Sample staining:**

Sorting of the Meio fauna from the sieve is difficult task especially in the preserved material, because organisms are not easily detectable. To facilitate the animal detection the entire sample retained on the sieve after sieving operation were stained by immersing the sieve in a flat bottom

tub with 1% Rose Bengal stain; a protein stain. A staining period of 10-30 minutes is sufficient for sample detection.

#### **DIVERSITY INDICES:**

On the whole, diversity indices provide more information about community composition than simply species richness (number of species present); they also, take the relative abundances of different species into account. Based on this fact, diversity indices therefore depend not only on species richness but on the evenness, or equitability, with which individuals are distributed among the different species (Magurram, A. E. (1988)

A diversity index is a measure of species diversity within a community that consists of co-occurring populations of several (two or more) different species. It includes two components: richness and evenness. Richness is the measure of the number of different species within a sample showing that more the types of species in a community, the higher is the diversity or greater is the richness. Evenness is the measure of relative abundance of the different species with in a community.

The basic idea of diversity index is to obtain a quantitative estimate of biological variability that can be used to compare biological entities composed of discrete components in space and time ( Carol H.R. *etal.* 1998). Biodiversity is commonly expressed through indices based on species richness and species abundances (Whittaker 1972, Lande 1996, Purvis and Hector 2000). Biodiversity indices are a non-parametric tool used to describe the relationship between species number and abundance. The most widely used bio diversity indices are Shannon Weiner index and Simpson's index.

A diversity Index is a single statistic that incorporates in formation on richness and evenness. The diversity measures that incorporate the two concepts may be termed heterogeneity measures (Magurran, 2004).

Any study intended to interpret causes and effect of adverse impact on Biodiversity of communities require suitable measures to evaluate specie richness and Diversity. The former is number of species in community, while latter is a function of relative frequency of different species. Species richness is the iconic measure of biological diversity (Magurran, 2004). Several indices have been created to measure the diversity of species; however, the most widely used in the last decades are the Shannon (1948) and Simpson (1949) (Buzas and Hayek 1996; Gorelick 2006), with the components of diversity: richness ( $S$ ) and evenness ( $J$ )

### Simpson's diversity index

Simpson's index (**D**) is a measure of diversity, which takes into account both species richness, and evenness of abundance among the species present.. The Simpson index is one of the meaningful and robust biodiversity measures available.(Magurran ,2004).

The formula for calculating D is presented as:

$$D = \frac{\sum n_i(n_i - 1)}{N(N - 1)}$$

Where  $n_i$  = the total number of organisms of each individual species

$N$  = the total number of organisms of all species

The value of D ranges from 0 to 1. With this index, 0 represents infinite diversity and, 1, no diversity. When D increases diversity decreases. Simpson's index is therefore usually expressed as 1-D or 1/D. (Magurran, 2004)

Low species diversity suggests:

- relatively few successful species in the habitat
- the environment is quite stressful with relatively few ecological niches and only a few organisms are really well adapted to that environment
- food webs which are relatively simple
- change in the environment would probably have quite serious effects

High species diversity suggests:

- a greater number of successful species and a more stable ecosystem
- more ecological niches are available and the environment is less likely to be hostile complex food webs
- environmental change is less likely to be damaging to the ecosystem as a whole

### Species richness indices

The species richness(**S**) is simply the number of species present in an ecosystem. Species richness Indices of species richness are widely used to quantify or monitor the effects of anthropogenic disturbance. A decline in species richness may be concomitant with severe or chronic human-induced perturbation (Fair Fair weather 1990,) Species richness measures have traditionally been the mainstay in assessing the effects of environmental degradation on the biodiversity of natural assemblages of organisms (Clarke &Warwick, 2001)

Species richness is the iconic measure of biological diversity (Magurran, 2004). The species richness(**S**) is simply the number of species present in an ecosystem. This index makes no use of

relative abundances. The term species richness was coined by McIntosh (1967) and oldest and most intuitive measure of biological diversity (Magurran, 2004).

Margalef's diversity index is a species richness index. Margalef's Species richness index (d), or indices that describe the evenness of the distribution of the numbers of individuals among species, were derived.

The value of a diversity index increases both when the number of types increases and when evenness increases. For a given number of types, the value of diversity index is maximised when all types are equally abundant (Rosenzweig, M. L. (1995).

### Shannon-Wiener's index:

An index of diversity commonly used in plankton community analyses is the Shannon-Wiener's index (**H**), which emphasizes not only the number of species (richness or variety), but also the apportionment of the numbers of individuals among the species (Odum 1971 and Reish 1984). Shannon-Wiener's index (**H**) reproduce community parameters to a single number by using an equation.

Shannon and Weiner index represents entropy. It is a diversity index taking into account the number of individuals as well as the number of taxa. It varies from 0 for communities with only single taxa to high values for community with many taxa each with few individuals. This index can also determine the pollution status of a water body. Normal values range from 0 to 4. This index is a combination of species present and the evenness of the species. Examining the diversity in the range of polluted and unpolluted ecosystems, Wilham and Dorris (1968) concluded that the values of the index greater than 3 indicate clean water, values in the range of 1 to 3 are characterized by moderate pollution and values less than 1 are characterized as heavily polluted

$$H' = - \sum_{j=1}^s \frac{n_j}{N} \ln \left( \frac{n_j}{N} \right)$$

### RESULTS:

#### CHLOROPHYLL-a:

Water Samples for the chlorophyll estimation were collected from sub surface layer during high tide and low tide period of the tidal cycle for each sampling locations and analysed for Chlorophyll -a and after acidification for Pheophytin -a. Chlorophyll- a value was used as algal biomass indicator (APHA,1998) Algal biomass was estimated by converting Chlorophyll value.

In the sub surface water chlorophyll-a was varying from 0.511 -0.921mg/m<sup>3</sup>.in harbour region of DPT during sampling done in spring tide period of July, 2021. In the nearby creeks chlorophyll-a was

**DCPL/DPT/20-21/15 -JULY - 2021**

varying from 0.173-0.980 mg/m<sup>3</sup>.Pheophytin –a level was below detectable limit- the all the sampling stations during spring in the harbour region of DPT.

In the sub surface water chlorophyll-a was varying from 0.391 -0.835mg/m<sup>3</sup>.in harbour region of DPT during sampling done in neap tide period of July, 2021 . In the nearby creeks chlorophyll-a was varying from 0.308-0.991 mg/m<sup>3</sup>.Pheophytin –a level was below detectable limit- the all the sampling stations during spring in the harbour region ofDPT

**TABLE #2 VARIATIONS IN CHLOROPHYLL –a PHEOPHYTIN- a AND ALGAL BIOMASS FROM SAMPLING STATIONS IN DPT HARBOUR AREA ,NEAR BY CREEKS DURING SPRING TIDE IN JULY,2021**

Sr.No.	Station	Tide	Chlorophyll-a (mg/m <sup>3</sup> )	Pheophytin- a (mg/m <sup>3</sup> )	Algal Biomass (Chlorophyll method) mg/m <sup>3</sup>
DPT HARBOUR AREA					
1	KPT1	High tide	0.629	BDL	42.14
		Low tide	0.921	BDL	61.71
2	KPT 2	High tide	0.745	BDL	49.92
		Low tide	0.558	BDL	37.39
3	KPT 3	High tide	0.511	BDL	34.24
		Low tide	0.598	BDL	40.06
CREEKS					
4	KPT-4 Khori-I	High tide	0.425	BDL	28.48
		Low tide	0.473	BDL	31.69
5	KPT-5 Nakti-I	High tide	0.714	BDL	47.84
		Low tide	0.980	BDL	65.66
6	KPT-5 Nakti-II	High tide	0.173	BDL	11.59

BDL: Below Detectable Limit.

**TABLE #3 VARIATIONS IN CHLOROPHYLL –aPHEOPHYTIN- a AND ALGAL BIOMASS FROM SAMPLING STATIONS IN DPT HARBOUR AREA ,NEAR BY CREEKS DURING NEAP TIDE IN JULY,2021**

Sr.No.	Station	Tide	Chlorophyll-a (mg/m <sup>3</sup> )	Pheophytin- a (mg/m <sup>3</sup> )	Algal Biomass (Chlorophyll method) mg/m <sup>3</sup>
DPT HARBOUR AREA					
1	KPT1	High tide	0.730	BDL	48.91
		Low tide	0.835	BDL	55.94
2	KPT 2	High tide	0.391	BDL	26.20
		Low tide	0.484	BDL	32.43
3	KPT 3	High tide	0.612	BDL	41.00
		Low tide	0.513	BDL	34.37
CREEKS					
4	KPT-4 Khori-I	High tide	0.385	BDL	25.80
		Low tide	0.497	BDL	33.30
5	KPT-5 Nakti-I	High tide	0.991	BDL	66.39
		Low tide	0.692	BDL	46.36
6	KPT-5 Nakti-II	High tide	0.308	BDL	20.64

BDL: Below Detectable Limit.

**PHYTOPLANKTON POPULATION:**

For the evaluation of the Phytoplankton population in DPT harbour area and within the immediate surroundings of the port, sampling was conducted from 5 sampling locations (3 in harbour area and two in Nakti creek) during high tide period and low tide period of spring tide. The phytoplankton community of the sub surface water in the harbour and nearby creeks was represented by, Diatoms and blue green algae during spring tide period. Diatoms were represented by 14 genera. Blue green were represented by one genera. During the sampling conducted in spring tide in July, 2021. Phytoplankton of the sampling stations at sub surface layer in the harbour area and nearby creeks was varying from 68 -196 units/ L during high tide period and 171-212 units/ L during low tide of Spring Tide.

The phytoplankton community of the sub surface water in the harbour and nearby creeks was represented by Diatoms and Blue green algae during spring tide period. Diatoms were represented by 14 genera and Blue green algae were represented two genera during the sampling conducted in Neap tide in July, 2021. Phytoplankton of the sampling stations at sub surface layer in the harbour area and nearby creeks was varying from 86-224 units/ L during high tide period and 222-254 units/ L during low tide of Neap Tide.

#### **Species Richness Indices and Diversity Indices :**

##### **Margalef's diversity index (Species Richness) S**

At the organismal level, the most widely used biodiversity measures are those based on the number of species present, perhaps adjusted for the number of individuals sampled, Here Margalef's Species richness index (d), or indices that describe the evenness of the distribution of the numbers of individuals among species, are derived.

Margalef's diversity index (Species Richness) S of phytoplankton communities in the stations was varying from 1.896 -2.495 with an average of 2.315 during the sampling conducted in High tide period of spring tide. While Margalef's diversity index (Species Richness) S of phytoplankton communities in the Kandla Harbour region and nearby creeks was varying from 2.054-2.334 with an average of 2.170 during the consecutive in low tide period .

Margalef's diversity index (Species Richness) S of phytoplankton communities in the stations was varying from 2.245-2.630 with an average of 2.495 during the sampling conducted in High tide period of Neap tide While Margalef's diversity index (Species Richness) S of phytoplankton communities in the Kandla Harbour region and nearby creeks was varying from 2.003-2.709 with an average of 2.232 during the consecutive in low tide period .

##### **Shannon-Wiener's index:**

Shannon-Wiener's Index (H) of phytoplankton communities in the sampling stations was in the range of 0.878-0.959 ( $H'(\log_{10})$ ) between selected sampling stations with an average value of 0.905 during high tide period of spring tide . Shannon-Wiener's Index (H) of phytoplankton communities in the sampling stations was in the range of 0.863-0.904 ( $H'(\log_{10})$ ) between selected sampling stations with an average value of 0.892 during consecutive low tide. Shannon-Wiener's Index (H) of phytoplankton communities in the sampling stations was in the range of 0.960-1.025 ( $H'(\log_{10})$ ) between selected sampling stations with an average value of 0.990. during high tide period of neap tide . Shannon-Wiener's Index (H) of phytoplankton communities in the sampling stations was in the range of 0.969-1.008 ( $H'(\log_{10})$ ) between selected sampling stations with an average value of 0.990 during consecutive low tide. Typical values are generally between 1.5 and 3.5 in most ecological

studies, and the index is rarely greater than 4. The Shannon-Wiener's index increases as both the richness and the evenness of the community increase. This result indicates that diversity of phytoplankton of Kandla Harbour region and nearby creeks is less but with abundant population of few, with relatively few ecological niches and only very few opportunist organisms are really well adapted to this environment and thrive better than other species.

**Simpson's diversity index:**

Simpson's index (D) is a measure of diversity, which takes into account both species richness, and an evenness of abundance among the species present. The Simpson index is one of the meaningful and robust biodiversity measures available. (Magurran,2004).

Simpson diversity index (1-D) of phytoplankton communities was below 0.9 at all sampling stations in the Kandla Harbour region and nearby creeks, which was varying from 0.837- 0.878 between selected sampling stations with an average of 0.855 during high tide period of spring tide . Simpson diversity index (1-D) of phytoplankton communities was below 0.9 at all sampling stations in the Kandla Harbour region and nearby creeks, which was varying from 0.840- 0.856 between selected sampling stations with an average of 0.849 during consecutive low tide . Low species diversity suggests a relatively few successful species in this habitat. The environment is quite stressful with relatively few ecological niches and only a few organisms are really well adapted to that environment. Any change in the environment would probably have quite serious effects.

Simpson diversity index (1-D) of phytoplankton communities was below 0.9 at all sampling stations in Kandla Harbour region and nearby creeks, during high tide period and low tideperiod during neap tide also, which was varying from 0.872-0.891 with an average value of 0.881 between selected sampling stations during high tide period and varying from 0.882-0.889 with an average value of 0.885 between selected sampling stations during consecutive low tide period Low species diversity suggests a relatively few successful species in this habitat.



**Table # 4 PHYTOPLANKTON VARIATIONS IN ABUNDANCE AND DIVERSITY IN SUB SURFACE SAMPLING STATIONS IN DPT HARBOUR AREA ,NEAR BY CREEKS DURING SPRING TIDE IN JULY,2021**

Tide	Sampling Station	Abundance In units/L	No of Species observed /total species	% of diversity	Margalef's diversity index (Species Richness S)	Shannon Weiner index H (log <sub>10</sub> )	Diversity Index (Simpson's Index) 1-D
HIGH TIDE	1	183	14/15	93.33	2.495	0.906	0.8502
	2	154	13/15	86.66	2.382	0.8957	0.8537
	3	159	13/15	86.66	2.367	0.8948	0.8424
	4	188	13/15	86.66	2.292	0.8783	0.8372
	5	196	14/15	93.33	2.463	0.9587	0.8667
	6	68	9/15	60	1.896	0.899	0.8784
LOW TIDE	1	171	13/15	86.66	2.334	0.9041	0.8535
	2	212	12/15	80	2.054	0.8992	0.8565
	3	197	13/15	86.66	2.271	0.89	0.8406
	4	203	12/15	80	2.07	0.8634	0.8401
	5	179	12/15	80	2.121	0.9037	0.856

**Table # 5 PHYTOPLANKTON VARIATIONS IN ABUNDANCE AND DIVERSITY IN SUB SURFACE SAMPLING STATIONS IN DPT HARBOUR AREA ,NEAR BY CREEKS DURING NEAP TIDE IN JULY2021**

Tide	Sampling Station	Abundance In units/L	No of Species observed /total species	% of diversity	Margalef's diversity index (Species Richness S)	Shannon Weiner index H (log <sub>10</sub> )	Diversity Index (Simpson's Index) 1-D
HIGH TIDE	1	205	15/16	93.75	2.63	1.002	0.8735
	2	184	14/16	87.5	2.493	0.9603	0.872
	3	221	14/16	87.5	2.408	0.9762	0.8773
	4	213	15/16	93.75	2.611	1.025	0.8905
	5	224	15/16	93.75	2.587	1.011	0.8859
	6	86	11/16	68.75	2.245	0.9685	0.8914
LOW TIDE	1	243	12/16	75	2.003	0.9696	0.8823
	2	222	12/16	75	2.036	0.9893	0.8893
	3	222	13/16	81.25	2.221	1.001	0.8872
	4	254	16/16	100	2.709	1.008	0.883
	5	239	13/16	81.25	2.191	0.985	0.8864


**Table # 6 ABUNDANCE OF PHYTOPLANKTON SUBSURFACE SAMPLING STATIONS IN DPT HARBOUR AREA, NEAR BY CREEKS DURING SPRING TIDE IN JULY,2021**

Tide	Surface	No of Sampling location	Group of phytoplankton	Phytoplankton Group range Units/L	Genera or species /total Phyto plankton	Taxon Diversity % (Group level)
HIGH TIDE	Sub surface	6	DIATOMS	68-194	14/15	93.33
			BLUE GREEN	0-2	1/15	6.67
			TOTAL PHYTO PLANKTON	68-196	15	-
LOW TIDE	Sub surface	5	DIATOMS	170-211	14/15	93.33
			BLUE GREEN	0-1	1/15	6.67
			TOTAL PHYTO PLANKTON	171-212	15	-


**Table # 7 ABUNDANCE OF PHYTOPLANKTON SUBSURFACE SAMPLING STATIONS IN DPT HARBOUR AREA, NEAR BY CREEKS DURING NEAP TIDE IN JULY,2021**

Tide	Surface	No of Sampling location	Group of phytoplankton	Phytoplankton Group range Units/L	Genera or species /total Phyto plankton	Taxon Diversity % (Group level)
HIGH TIDE	Sub surface	6	DIATOMS	74-202	14/16	87.5
			BLUE GREEN	12-26	2/16	12.5
			TOTAL PHYTO PLANKTON	86-224	16	-
LOW TIDE	Sub surface	5	DIATOMS	201-236	14/16	87.5
			BLUE GREEN	16-21	2/16	12.5
			TOTAL PHYTO PLANKTON	222-254	16	-

**Taxon Diversity % of Phytoplankton during High tide and Low tide period during spring tide**



**Taxon Diversity % of Phytoplankton during High tide and Low tide period during Neap tide**



**ZOOPLANKTON POPULATION:**

For the evaluation of the Zooplankton population in DPT harbour area and within the immediate surroundings of the port sampling was conducted from 6 sampling locations (3 in harbour area and two in Nakti creek and one in Khoricreek ) during high tide period and low tide period of spring tide and Neap tide in July 2021 . The Zooplankton community of the sub surface water in the harbour

and nearby creeks during spring tide was represented by mainly four groups, Tintinids, Copepods, Foraminiferans and larval forms of Crustacea, Molluscs and Polychaetes. The Zooplankton community of the sub surface water in the harbour and nearby creeks during neap tide was represented by mainly five groups, Tintinids, Copepods, Arrow worms, Mysids and larval forms of Crustaceans, Molluscs and Polychaetes,.

Zooplankton of the sampling stations at sub surface layer in the DPT harbour area and nearby creek was varying from  $59-142 \times 10^3$  N/ m<sup>3</sup> during high tide and  $123-147 \times 10^3$  N/ m<sup>3</sup> during low tide of Spring Tide period. Zooplankton of the sampling stations at sub surface layer in the DPT harbour area and nearby creek was varying from  $59-147 \times 10^3$  N/ m<sup>3</sup> during high tide and 141-164 N/ L during low tide of Neap Tide period.

### **Species Richness Indices and Diversity Indices:**

#### **Margalef's diversity index (Species Richness) S**

At the organismal level, the most widely used biodiversity measures are those based on the number of species present, perhaps adjusted for the number of individuals sampled, Here Margalef's Species richness index (d), or indices that describe the evenness of the distribution of the numbers of individuals among species, are derived.

Margalef's diversity index (Species Richness) S of Zooplankton communities in the stations Kandla Harbour region and nearby creeks was varying from 2.850 -3.366 with an average of 3.040 during the sampling conducted in High tide period. Margalef's diversity index (Species Richness) S of Zooplankton communities varying from 2.263-2.701 with an average of 2.562 during the sampling conducted in low tide period during Spring tide.

Margalef's diversity index (Species Richness) S of Zooplankton communities in the Kandla Harbour region and nearby creeks sampling stations was varying from 3.188-4.133 with an average of 3.754 during the sampling conducted in high tide and varying from 2.802 -4.314 with an average of 3.548 during the sampling conducted in low tide during Neap tide period.

**Shannon-Wiener's index:**  
Shannon-Wiener's Index (H) of Zooplankton communities in the sampling stations in Kandla Harbour region and nearby creeks was in the range of 1.011-1.080 (H'(log10)) between selected sampling stations with an average value of 1.050 (H'(log10)) during high tide period of spring tide. Shannon-Wiener's Index (H) of Zooplankton communities in the sampling stations in Kandla Harbour region and nearby creeks was in the range of 0.953 -1.011 (H'(log10)) between selected sampling stations with an average value of 0.988 (H'(log10)) during consecutive low tide period.

Shannon-Wiener's Index (H) of Zooplankton communities in the sampling stations in Kandla Harbour region and nearby creeks was in the range of 0.884-1.145 (H'(log10)) between selected sampling

stations with an average value of 1.075 ( $H'(\log_{10})$ ) during high tide period of Neap tide. Shannon-Wiener's Index (H) of Zooplankton communities in the sampling stations in Kandla Harbour region and nearby creeks was in the range of 1.004- 1.177 ( $H'(\log_{10})$ ) between selected sampling stations with an average value of 1.056 ( $H'(\log_{10})$ ) during consecutive low tide period. Typical values are generally between 1.5 and 3.5 in most ecological studies, and the index is rarely greater than 4. The Shannon-Wiener's index increases as both the richness and the evenness of the community increase. This result indicates that diversity of Zooplankton of Kandla Harbour region and nearby creeks stations is slightly high with very minimum diverse population but very few opportunist organisms are really well adapted to this environment and thrive better than other species.

**Simpson's diversity index:**

Simpson's index (D) is a measure of diversity, which takes into account both species richness, and an evenness of abundance among the species present. The Simpson index is one of the meaningful and robust biodiversity measures available. (Magurran, 2004).

Simpson diversity index (1-D) of Zooplankton communities was below 0.9 most of sampling stations except few in the Kandla Harbour region and nearby creeks during high tide and low tide of spring tide period, which was varying from 0.882-0.911 between selected sampling stations with an average of 0.899 during high tide period and was varying from 0.875- 0.888 with an average value of 0.882 between selected sampling stations during low tide

Simpson diversity index (1-D) of Zooplankton communities was below 0.9 at all sampling stations in the Kandla Harbour region and nearby creeks during high tide and low tide period except few, which was varying from 0.829-0.907 between selected sampling stations with an average of 0.887 during high tide period and was varying from 0.872- 0.913 with an average value of 0.886 between selected sampling stations during consecutive low tide

This low species diversity suggests a relatively low number of successful species in this habitat. Environment is quite stressful with relatively few ecological niches and only few organisms are really well adapted to that environment. Any change in the environment would probably have quite serious effects.

**Table # 8 ZOOPLANKTON VARIATION IN ABUNDANCE AND DIVERSITY IN SUB SURFACE SAMPLING STATIONS IN DPT HARBOUR AREA ,NEAR BY CREEKS DURING SPRING TIDE IN JULY,2021**

Tide	Sampling Station	Abundance In No / m <sup>3</sup>	No of Species/groups observed /total species/group	% of diversity	Margalef's diversity index (Species Richness S)	Shannon Weiner index H (log <sub>10</sub> )	Diversity Index (Simpson's Index) 1-D
HIGH TIDE	1	125 X10 <sup>3</sup>	15/19	78.95	2.9	1.02	0.8906
	2	116 X10 <sup>3</sup>	16/19	84.21	3.156	1.08	0.9076
	3	116 X10 <sup>3</sup>	17/19	89.47	3.366	1.076	0.8961
	4	142 X10 <sup>3</sup>	16/19	84.21	3.027	1.011	0.8821
	5	136 X10 <sup>3</sup>	15/19	78.95	2.85	1.077	0.9077
	6	59 X10 <sup>3</sup>	13/19	68.42	2.943	1.037	0.9112
LOW TIDE	1	129 X10 <sup>3</sup>	12/19	63.16	2.263	0.9534	0.8751
	2	123 X10 <sup>3</sup>	14/19	73.68	2.701	0.9887	0.8835
	3	145 X10 <sup>3</sup>	14/19	73.68	2.612	1.011	0.8879
	4	147 X10 <sup>3</sup>	14/19	73.68	2.605	0.9919	0.8823
	5	140 X10 <sup>3</sup>	14/19	73.68	2.631	0.9951	0.8808

**Table # 9 ZOOPLANKTON VARIATION IN ABUNDANCE AND DIVERSITY IN SUB SURFACE SAMPLING STATIONS IN DPT HARBOUR AREA ,NEAR BY CREEKS DURING NEAP TIDE IN JULY,2021**

Tide	Sampling Station	Abundance In No / m <sup>3</sup>	No of Species/groups observed /total species/group	% of diversity	Margalef's diversity index (Species Richness S)	Shannon Weiner index H (log <sub>10</sub> )	Diversity Index (Simpson's Index) 1-D
HIGH TIDE	1	162 X10 <sup>3</sup>	19/23	82.61	3.538	1.094	0.895
	2	152 X10	20/23	86.96	3.782	1.088	0.8906
	3	146 X10	20/23	86.96	3.812	1.106	0.9011
	4	174 X10	22/23	95.65	4.071	1.134	0.9015
	5	161 X10	22/23	95.65	4.133	1.145	0.9069
	6	59 X10	14/23	60.86	3.188	0.8842	0.8299
LOW TIDE	1	141 X10	17/23	73.91	3.233	1.004	0.8719
	2	142 X10	18/23	78.26	3.43	1.022	0.8797
	3	148 X10	15/23	65.22	2.802	1.034	0.8911
	4	164 X10	23/23	100	4.314	1.177	0.9134
	5	156 X10	21/23	91.30	3.961	1.046	0.8781


**Table # 10 ABUNDANCE OF ZOOPLANKTON IN SUBSURFACE SAMPLING STATIONS IN DPT  
HARBOUR AREA ,NEAR BY CREEKS DURING SPRING TIDE IN JULY2021**

Tide	Surface	No of Sampling locations	Group of Zooplankton	Abundance of Zooplankton $\times 10^3$ Group Range	Genera or species /total Zooplankton	Taxon Diversity % (Group level)
HIGH TIDE	Sub surface	6	Tintinids	3-10	4/19	21.05
			Copepods	34-77	7/19	36.84
			Foraminiferans	2-6	2/19	10.53
			Larval forms	20-57	6/19	31.58
			TOTAL ZOOPLANKTON NO/L	<b>59-142</b>	<b>19</b>	-
LOW TIDE	Sub surface	5	Tintinids	3-8	4/19	21.05
			Copepods	76-80	7/19	36.84
			Foraminiferans	0-2	2/19	10.53
			Larval forms	40-63	6/19	31.58
			TOTAL ZOOPLANKTON NO/L	<b>123-147</b>	<b>19</b>	-


**Table # 11 ABUNDANCE OF ZOOPLANKTON IN SUBSURFACE SAMPLING STATIONS IN DPT  
HARBOUR AREA , NEAR BY CREEKS DURING NEAPTIDE IN JULY,2021**

Tide	Surface	No of Sampling locations	Group of Zooplankton	Abundance of Zooplankton $\times 10^3$ Group Range	Genera or species /total Zooplankton	Taxon Diversity % (Group level)
HIGH TIDE	Sub surface	6	Tintinids	4-13	4/23	17.39
			Copepods	37-83	8/23	34.78
			Mysids	0-2	1/23	4.35
			Arrow worms	1-2	1/23	4.35
			Foraminiferans	0-4	1/23	4.35
			Larval forms	17-74	8/23	34.78
			TOTAL ZOOPLANKTON NO/L	<b>59-173</b>	<b>23</b>	-
LOW TIDE	Sub surface	5	Tintinids	3-13	4/23	17.39
			Copepods	70-84	8/23	34.78
			Mysids	0-2	1/23	4.35
			Arrow worms	0-2	1/23	4.35
			Foraminiferans	0-2	1/23	4.35
			Larval forms	60-70	8/23	34.78
			TOTAL ZOOPLANKTON NO/L	<b>140-164</b>	<b>23</b>	-

Taxon Diversity % of Zooplankton during High tide and Low tide period of Spring tide



Taxon Diversity % of Zooplankton during High tide and Low tide period of Neap tide





**TABLE # 12 SYSTEMATIC ACCOUNT OF PHYTOPLANKTON IN THE SAMPLING LOCATIONS IN OF DPT HARBOUR AREA AND NEARBY CREEKS DURING SPRING TIDE OF JULY, 2021**

GROUP	PHYLUM	CLASS	ORDER	FAMILY	GENUS/SPECIES	#	Relative Abundance
BLUE GREEN ALAGE	Cynophyta	Cynophyceae	Stigonematales	Stigonemataceae	<i>Stigonemasp</i>	B1	Rare
DIATOMS	Bacillariophyta	Coscinodiscophyceae	Thalassiosirales	Thalassiosiraceae	<i>Planktoniellasp</i>	D1	Rare
			Coscinodiscales	Coscinodiscaceae	<i>Coscinodiscus sp.</i>	D2	Abundant
					<i>Palmeriasp</i>	D3	Occasional
			Triceratiales	Triceratiaceae	<i>Odontellasp</i>	D4	Frequent
					<i>Triceratiumsp.</i>	D5	Frequent
			Biddulphiales	Biddulphiaceae	<i>Biddulphiasp</i>	D6	Abundant
			Hemiaulales	Bellerocheaceae	<i>Bellerocheasp</i>	D7	Occasional
				Hemiaulaceae	<i>Eucampiasp</i>	D8	Rare
			Lithodesmiales	Lithodesmiaceae	<i>Ditylumsp</i>	D9	Frequent
			Thalassiosirales	Thalassiosiraceae	<i>Thalassiosirasp</i>	D10	Rare
		Bacillariophyceae	Naviculales	Pleurosigmaaceae	<i>Pleurosigmasp</i>	D11	Rare
			Bacillariales	Bacillariaceae	<i>Nitzschiasp</i>	D12	Rare
		Fragilariophyceae	Thalassionematales	Thalassionemataceae	<i>Thalassiothrix sp.</i>	D13	Dominant
			Fragilariales	Fragilariaceae	<i>Synedrasp</i>	D14	Occasional

**TABLE # 12 SYSTEMATIC ACCOUNT OF PHYTOPLANKTON IN THE SAMPLING LOCATIONS IN OF DPT HARBOUR AREA AND NEARBY CREEKS DURING AND NEAP TIDE OF JULY,2021**

GROUP	PHYLUM	CLASS	ORDER	FAMILY	GENUS/SPECIES	#	Relative Abundance
BLUE GREEN ALAGE	Cynophyta	Cynophyceae	Chlorococcales	Chroococcaceae	<i>Microcystis sp.</i>	B1	Occasional
			Stigonematales	Stigonemataceae	<i>Stigonemasp</i>	B2	Frequent
DIATOMS	Bacillariophyta	Coccinodiscophyceae	Thalassiosirales	Thalassiosiraceae	<i>Planktoniellasp</i>	D1	Rare
			Coscinodiscales	Coscinodiscaceae	<i>Coscinodiscus sp.</i>	D2	Abundant
					<i>Palmeriasp</i>	D3	Occasional
			Triceratiales	Triceratiaceae	<i>Odontellasp</i>	D4	Frequent
					<i>Triceratiumsp.</i>	D5	Abundant
			Biddulphiales	Biddulphiaceae	<i>Biddulphiasp</i>	D6	Dominant
			Hemiaulales	Belleracheaceae	<i>Belleracheasp</i>	D7	Occasional
				Hemiaulaceae	<i>Eucampiasp</i>	D8	Rare
			Lithodesmiales	Lithodesmiaceae	<i>Ditylumsp</i>	D9	Abundant
			Thalassiosirales	Thalassiosiraceae	<i>Thalassiosirasp</i>	D10	Frequent
		Bacillariophyceae	Naviculales	Pleurosigmataceae	<i>Pleurosigmasp</i>	D11	Rare
			Bacillariales	Bacillariaceae	<i>Nitzschiasp</i>	D12	Rare
		Fragilariophyceae	Thalassionematales	Thalassionemataceae	<i>Thalassiothrix sp.</i>	D13	Frequent
			Fragilariales	Fragilariaceae	<i>Synedrasp</i>	D14	Frequent

**TABLE #13 SYSTEMATIC ACCOUNT OF ZOOPLANKTON FROM THE SAMPLING LOCATIONS IN DPT HARBOUR AREA, AND NEAR BY CREEKS DURING SPRING TIDE OF JULY,2021**

GROUP	PHYLUM	CLASS	ORDER	FAMILY	GENUS/SPECIES	#	RELATIVE ABUNDANCE		
TINTINIDS	PROTOZOA CILIOPHORA	Spirotrichea	Tintinnida	Tintinnidiidae	<i>Leprotintinnussp.</i>	T1	Rare		
				Codonellidae	<i>Tintinnopsisfailakkaensis</i>	T2	Rare		
					<i>Tintinnopsisgracilis</i>	T3	Rare		
					<i>Tintinnopsis radix</i>	T4	Rare		
COPEPODS	ATHROPODA	Crustacea Sub class copepoda	Calanoida	Paracalanidae	<i>Acrocalanus sp.</i>	C1	Abundant		
					<i>Bestiolina sp.</i>	C2	Rare		
					<i>Parvocalanus sp.</i>	C3	Occasional		
						Temoridae	<i>Temora sp.</i>	C4	Frequent
					Cyclopoida	Oithonidae	<i>Oithona sp.</i>	C5	Frequent
					Harpacticoida	Ectinosomatidae	<i>Microsetellasp.</i>	C6	Abundant
						Euterpinae	<i>Euterpina</i>	C7	Occasional
CRUSTACEAN LARVAE	ARTHROPODA (CRUSTACEA)	Copepoda			Nauplius larvae of Copepods	L1	Dominant		
(Brachyuraian LARVAE	ARTHROPODA CRUSTACEA	DECAPODA (BRACHYURA)			Zoea larvae	L2	Rare		
CYPHONAUTES LARVAE	BRYOZOA				Cyphonautes larvae	L3	Occasional		
BIVALVE LARVAE	MOLLUSCA	Pelecypoda			Veliger larvae of Bivalves	L4	Rare		
MOLLUSCAN LARVAE	MOLLUSCA	Gastropoda Streptoneura			Opisthobranchia larvae	L5	Rare		
POLYCHAETE LARVAE	ANNELIDA				Trochophore larvae	L6	Frequent		
FORAMINIFERA	FORAMINIFERA	Globothalamea	Rotaliida	Globigerinidae	<i>Globigerina sp.</i>	F1	Rare		
				Rotaliidae	<i>Rotalia sp.</i>	F2	Rare		

**TABLE # 13 SYSTEMATIC ACCOUNT OF ZOOPLANKTON FROM THE SAMPLING LOCATIONS IN DPT HARBOUR AREA, AND NEAR BY CREEKS DURING NEAP TIDE OF JULY,2021**

GROUP	PHYLUM	CLASS	ORDER	FAMILY	GENUS/SPECIES	#	RELATIVE ABUNDANCE
TINTINIDS	PROTOZOA CILIOPHORA	Spirotrichea	Tintinnida	Tintinnidiidae	<i>Leprotintinnus</i> sp.	T1	Rare
				Codonellidae	<i>Tintinnopsis failakkaensis</i>	T2	Occasional
					<i>Tintinnopsis gracilis</i>	T3	Occasional
					<i>Tintinnopsis radix</i>	T4	Rare
COPEPODS	ARTHROPODA	Crustacea Sub class copepoda	Calanoida	Paracalanidae	<i>Acrocalanus</i> sp.	C1	Abundant
					<i>Parvocalanus</i> sp.	C2	Rare
				Eucalanidae	<i>Subeucalanus</i> sp.	C3	Frequent
				Temoridae	<i>Temora</i> sp.	C5	Frequent
				Acartiidae	<i>Acartia</i> sp.	C6	Occasional
			Cyclopoida	Oithonidae	<i>Oithona</i> sp.	C7	Frequent
			Harpacticoida	Ectinosomatidae	<i>Microsetella</i> sp.	C8	Abundant
				Euterpinae	<i>Euterpina</i> sp.	C9	Frequent
ARROW WORMS	CHAETOGNATHA	Sagittoideae	Aphragmophora	Sagittidae	<i>Sagitta</i> sp.	A1	Rare
MYSIDS	ARTHROPODA CRUSTACEA	Malacostraca	Mysida, Decapoda	Penaeidae	<i>Metapenaeus</i> sp.	M1	Rare
CRUSTACEAN LARVAE	ARTHROPODA (CRUSTACEA)	Copepoda			Nauplius larvae of Copepods	L1	Dominant
POLYCHAETE LARVA	ANNELIDA	Polychaeta			Trochophore larvae	L2	Occasional
BARNACLE LARVAE	ARTHROPODA CRUSTACEA	Maxillopoda Thecostraca			Cirripede larvae	L3	Rare
BIVALVE LARVAE	MOLLUSCA	Pelecypoda			Veliger larvae of Bivalves	L4	Occasional

**Environmental Monitoring Report of Deendayal Port Trust, JULY-2021**

GROUP	PHYLUM	CLASS	ORDER	FAMILY	GENUS/SPECIES	#	RELATIVE ABUNDANCE
MOLLUSCAN LARVAE	MOLLUSCA	Gastropoda Streptoneura			Opisthobranchia larvae	L5	Rare
BRACHYURAIAN LARVAE	ARTHROPODA CRUSTACEA	DECAPODA (BRACHYURA)			Zoea larvae	L6	Occasional
CYPHONAUTES LARVAE	BRYOZOA				Cyphonautes larvae	L7	Occasional
ECHINODERMATA larvae	ECHINODERMATA	Ophiuroidea			Ophiopluetus larvae	L8	Occasional
FORAMINIFERA	FORAMINIFERA	Globothalamea	Rotalliida	Rotalliidae	<i>Rotalia</i> sp.	F1	Rare

**BENTHIC ORGANISMS:**

No Benthic organisms were observed in the collected sediments by using the Van-veen grabs during the sampling conducted IN spring tide period as well as Neap tide period from DPT harbour region and nearby creek except few dead shells.

## 7. Meteorological Data

Automatic Weather station have been installed in SevaSadan -3 at the Deendayal Port which records the data on Temperature (°C), Humidity (%), Wind (mph), Dew Point (°C), Wind Direction (°), Pressure, Solar radiation, heat Index and UVI.

### Temperature

The mean day time temperature for Deendayal Port was 28.5 °C. The day-time maximum temperature was 32.1 °C. The mean night time temperature was 30.3 °C. The minimum mean night time temperature recorded was 27.8 °C.

### Air Pressure

The mean absolute air pressure for the month of July was 1002.4 hpa, whereas the mean relative pressure was 1000.2 hpa. The maximum absolute air pressure recorded for the month of July was 1004.1 hpa.

### Heat Index

The mean day-time heat index for the month of July was 36.1 °C. The maximum heat index recorded was 43°C.

### Solar Radiation

The mean Solar Radiation in July was 158.4 w/m<sup>2</sup>. The maximum solar radiation recorded in the month of July was 751.7 w/m<sup>2</sup>.

### Humidity

The mean day-time humidity was 80.3 % for the month of July and mean night time humidity was 71.2%. Maximum humidity recorded during day-time was 89.0 % and maximum humidity recorded during night-time was 85.0%.

### Wind Velocity and Wind Direction

The mean wind velocity for the entire month of July was 11.72 km/hour (i.e. 2.7 mtr/sec). Maximum wind velocity recorded was 47.2 Km/hr (13 mtr/sec). The wind direction was mostly S to SW.

### Rainfall

The mean Rainfall in July was 58.1 mm. The maximum Rainfall recorded in the month of July was 132.7 mm.

### **Conclusive Summary and Remedial measures Suggested**

- The AAQ monitoring at six locations of Deendayal Port indicates that the mean PM<sub>10</sub> values at four locations viz. Coal storage area, Marine Bhavan and Oil Jetty area were found above the permissible standards (100 µg/m<sup>3</sup>) and PM<sub>2.5</sub> was above permissible limits at Coal storage location (Limit 60 µg/m<sup>3</sup>).
- Drinking water at all the twenty locations was found potable and was within permissible limits of BIS standards (IS 10500).
- Noise quality was also within the set permissible standards of an Industrial Area. The noise level observed during day time was >75 dB (A) and at night time was >70 dB (A) during the entire monitoring period.
- The sewage treated water of Deendayal Port Colony (Gopalpuri) was in line with the standards set by the Gujarat Pollution Control Board. The STP at Deendayal Port is not fully operational and STP at Vadinar Port was found non-operational.

### **Reasons for higher Values of PM<sub>10</sub>**

- Large amount of coal is handled at Berth No. 6, 7, 8 and 9. The unloading of coal directly in the truck, using grabs cause coal to spread in air as well as coal dust to fall on ground. This settled coal dust again mixes with the air while trucks travel through it.
- Also, the coal laden trucks are not always covered with tarpaulin sheets and these results in spillage of coal from trucks/dumpers during its transit from vessel to yard or storage site. This also increased PM values around marine Bhavan & Coal storage area.



### **Remedial Measures**

The values of PM<sub>10</sub> during the month of July, 2021 were observed beyond the permissible limit at four locations mentioned above. Given below are the remedial measures suggest to minimize the Air pollution at Deendayal Port.

- Guidelines for Coal Handling by GPCB should be strictly followed. (<http://gpcb.gov.in/pdf/coal-handling-guidelines.pdf>)
- Sewage Treatment Plan at Vadinar Port is not working. Hence, it is recommended to commission the sewage treatment plant at Vadinar immediately.
- Except for the higher values of PM<sub>10</sub> at Coal storage site, Oil Jetty, Tuna Port and Marine Bhavan locations, the monitoring results for the present month suggest that the overall Environment Quality of Deendayal Port is satisfactory.

**SOURCE OF LITERATURE AND ADDITIONAL REFERENCE FOR ECOLOGICAL STUDY**

- 1) ALBERT WEST PHAL (1976) Protozoa Blackwell , London
- 2) BANERJEE R.K. (1989) Heavy metals and Benthic foraminiferal distribution along Bombay coast India. Studies in benthic foraminifera. *Tokyo University Press* Tokyo pp 151-157
- 3) Banse K (1995) Zooplankton: Pivotal role in the control of ocean production: I. Biomass and production. *ICES J Mar Sci* 52: 265–277.
- 4) Beaugrand G, and Ibanez F (2004) Monitoring marine plankton ecosystems. II: long-term changes in North Sea calanoid copepods in relation to hydroclimatic variability. *Inter Res Mar Ecol Prog Ser* 284:35-47.
- 5) DAY F. (1889) The fauna of British India Ceylon and Burma- Fishes Vol-1- Vol-2 *Taylor and Francis* London
- 6) DESIKACHARY T.V. (1989) Atlas of diatoms, Madras Science Foundation
- 7) DESIKACHARY T.V. (1959) Cyanophyta ICAP Monographs on Algae *Indian Council of Agricultural research* New Delhi
- 8) FAIZAYOUSIF AL-YAMANI & MARIA A. SABUROVA (2010) illustrative guide on the flagellates of Intertidal soft sediment *Kuwait Institute for scientific Research* Kuwait
- 9) FAIZAYOUSIF AL-YAMANI, VALERIYSKRYABIN, ALEKSANDRA GUBANOVA, SERGEY KHVOROV AND IRINA PRUSOVA (2011), Marine zooplankton Practical guide from North western Arabian gulf Vol-1 and vol-2 *Kuwait Institute for scientific Research* Kuwait
- 10) FAUVEL P. (1953), The fauna of India Annelida - Polychaeta *Indian Press* Allahabad
- 11) Gajbhiye SN, Nair VR, and Desai BN (1984). Diurnal variation of zooplankton in Malad creek, Bombay. *Indian Journal of Marine Science*. 13:75-79.
- 12) HAYWARD P.J AND RYLAND J.S. (1995) Handbook of Marine fauna of north –West Europe *oxford University Press* London
- 13) HIGGINS R.P. HAJAMARTHIEL Eds. (1998) Introduction to the study of Meio Fauna
- 14) HORACE G. BARBER AND ELIZABETH Y. HAWORTH (1981) A guide to the Morphology of DIATOMS FRUSTULES.
- 15) INGRAM HENDEY (1964) An introductory account of smaller Algae of British coastal waters part-V. Bacillariophyceae
- 16) JOHN H. WICKSTEAD (1965) an Introduction to the study of Tropical Plankton .Hutchinson Tropical Monographs
- 17) JOYOTHIBABU, R. MADHU, N.V. MAHESHWARAN, P.A., NAIR K.K.C., VENUGOPL, P. BALASUBRAMANIAN T. (2005) Dominance of Dinoflagellates in micro zooplankton communities in the oceanic region Bay of Bengal and Andaman sea *Current science* vol.84. 10<sup>th</sup> May 2003
- 18) KASTURIRANGAN L.R. (1963) A key for the identification of the Common Planktonic Copepoda of Indian Coastal water
- 19) Kusum KK, Vineetha G, Raveendran TV, Nair VR, Muraleedharan KR, Achuthankutty CT and Joseph T (2014) Chaetognath community and their responses to varying environmental factors in the northern Indian ocean. *J Plankton Res* 36(4): 1146- 1152.
- 20) Lalli CM and Parsons TR (1997) Biological Oceanography: An Introduction. DOI <https://doi.org/10.1016/B978-0-7506-3384-0.X5056-7>.
- 21) Madhupratap M (1978) Studies on ecology of zooplankton of Cochin backwaters. *Mahasagar Bull Nat Inst Oceanogr* 11: 45-56.
- 22) Madhupratap M (1979) Distribution, community structure and species succession of copepods from Cochin Backwaters. *Indian J Mar Sci* 8: 1-8.
- 23) Madhupratap M (1987) Status and strategy of zooplankton of tropical Indian estuaries: A review. *Bull Plank Soc Jpn* 34: 65-81.

## Environmental Monitoring Report Of Deendayal Port Trust, JULY-2021

---

- 24) Madhupratap M (1999) Free living copepods of the Arabian Sea, Distribution and Research Perspectives. I J Mar Sci 146-149.
- 25) Madhupratap M and Haridas P (1986) Epipelagic calanoid copepods of the northern Indian Ocean. OceanologicaActa 9(2):105-117.
- 26) MANAL AL-KANDARI, FAIZA Y. AL-YAMANI , KHOLOOD AL-RIFAIE ( 2009) Marine phytoplankton Atlas of Kuwait's water *Kuwait Institute for scientific Research*
- 27) MPEDA (1998) Commercial Fishes and shell fishes of India
- 28) NEWEL G.E. & NEWELL R.C. (1963) Marine plankton a Practical Guide Hutchinson Educational
- 29) NIGAM R.C. AND CHATURVEDIS.K. (2000) Foraminiferal Study from KharoCreek , Kachchh ( Gujarat) North west coast of India. *Indian Journal of marine science* Vol.29 133-189
- 30) OLAV GIERE (1993) Meio benthology , Microscopic Fauna in Aquatic Sediments m Springer London
- 31) PERRAGALLO( 1965) Diatomees marines de franceA. *Asher & Co.* Amsterdam
- 32) Robert P.. Higgins (Eds.), (1985) An introduction to the study of Meuo fauna Smithsons Institution press Washington DC
- 33) STERRER W. STERRERC.S Eds. Marine fauna and flora of Bermuda A systematic Guide to the identification of Marine Organisms. *John Wiely and Sons*New York
- 34) Suresh Gandhi. M. (2009) Distribution of certain ecological parameters and Foraminiferaldistribution in the depositional environment of Pak strait east coast of India .*Indian J. of Marine Science* Vol.33 pp 287-295
- 35) Venktaraman (1993 A systematic account of some south Indian diatoms . Proceeding of Indian Academy of Science Vol.X No.6 Sec.B.

\*\*\*\*\*

# ENVIRONMENTAL MONITORING REPORT FOR DEENDAYAL PORT TRUST



REPORT NO. : DCPL/DPT/20-21/16  
Month : Aug 2021  
Issue No : 01  
Revision No : 00  
Prepared by : DETOX CORPORATION PVT. LTD., SURAT

## TABLE OF CONTENTS

<b>Sr. No.</b>	<b>Particulars</b>	<b>Page No.</b>
<b>1</b>	<b>Ambient Air Quality Monitoring.....</b>	<b>1 - 17</b>
<b>2</b>	<b>Drinking Water Quality Monitoring.....</b>	<b>18 - 28</b>
<b>3</b>	<b>Noise Monitoring.....</b>	<b>29</b>
<b>4</b>	<b>Soil Monitoring.....</b>	<b>30 - 31</b>
<b>5</b>	<b>Sewage Treatment Plant Monitoring.....</b>	<b>32 - 38</b>
<b>6</b>	<b>Marine Water Monitoring.....</b>	<b>39 - 79</b>
<b>7</b>	<b>Meteorological Observations.....</b>	<b>80</b>
<b>8</b>	<b>Conclusive Summary &amp; Remedial Measures .....</b>	<b>81-82</b>
	<b>References.....</b>	<b>83- 84</b>

## **Introduction**

Monitoring of various environmental aspects of the Deendayal port by M/s Detox Corporation Pvt. Ltd. has been carried out through collection of samples, analysis of the same, comparing results with respect to the national standards and any other relevant standards by GBCB/CPCB/MoEF to identify non conformity in the Environment of the Deendayal Port. The results shall address the identified impacts and suggest measures to minimize the environmental impact due to various operations at Deendayal Port.

The environmental monitoring is carried out as per the Environment Management and Monitoring Plan submitted by Detox Corporation Pvt. Ltd.

## **1. Ambient Air Quality Monitoring**

As per the Environmental Monitoring Plan of Deendayal Port Trust, Air monitoring was carried out at six identified locations at Deendayal Port and two locations at Vadinar Port.

### **1.1 Air Quality Monitoring Methodology**

Air quality is measured in all the stations, for 24 hour for Total Suspended Particulate Matter (TSPM), PM<sub>10</sub>, PM<sub>2.5</sub>, SO<sub>2</sub>, NO<sub>x</sub>, NH<sub>3</sub> & Benzene, and Grab-sampling for CO & CO<sub>2</sub> measurements. The Air samplers are operated for a period of 24 hours and after a continuous operation of 8 hours of the sampler, the reagents were replaced to obtain 3 samples per day for each parameter namely, SO<sub>2</sub>, NO<sub>x</sub>. The EPM 2000 filter paper and PTFE Membrane bound filter paper are used for a period of 24 hours to obtain one sample each of TSPM, PM<sub>10</sub> & PM<sub>2.5</sub>.

The AAQ samples are collected twice a week from all the eight locations as per the EMP.

## 1.2 Results

The ambient air quality monitoring data for six stations, viz. Marine Bhavan, Oil Jetty, Port Colony, Gopalpuri Hospital, Tuna Port and Nr. Coal Storage Area for the month of August 2021 are given in Tables 1A to 6B. The ambient air quality monitoring data for two stations at Vadinar (Nr. Admin Building & Nr. Signal Building) are given in Tables 7A to 8B.

### Location 1: Marine Bhavan (AL1)

Table 1 : Results of Air Pollutant Concentration at Marine Bhavan										
Parameter	Date	TSPM [µg/m <sup>3</sup> ]	PM10 [µg/m <sup>3</sup> ]	PM2.5 [µg/m <sup>3</sup> ]	SO <sub>2</sub> [µg/m <sup>3</sup> ]		NO <sub>x</sub> [µg/m <sup>3</sup> ]		NH <sub>3</sub> [µg/m <sup>3</sup> ]	
Sampling Period	-	24hr	24hr	24hr	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
NAAQMS limit	-	NS	100 µg/m <sup>3</sup>	60 µg/m <sup>3</sup>		80 µg/m <sup>3</sup>		80 µg/m <sup>3</sup>		400 µg/m <sup>3</sup>
AL1 – 1	04.08.2021	328	179	68	2.64	2.40	20.33	21.17	9.45	9.87
					0.62		19.05		9.70	
					3.96		24.14		10.47	
AL1 – 2	06.08.2021	659	211	75	7.03	6.15	14.61	20.11	13.79	13.70
					5.71		15.88		13.53	
					5.71		29.85		13.79	
AL1 – 3	11.08.2021	813	247	70	8.35	7.03	29.85	27.10	12.00	11.49
					7.91		31.76		13.02	
					4.84		19.69		9.45	
AL1 – 4	13.08.2021	549	272	89	2.20	2.05	18.42	17.15	14.55	15.68
					1.76		15.88		17.69	
					2.20		17.15		14.81	
AL1 – 5	18.08.2021	442	300	45	3.96	3.66	19.69	21.38	5.36	9.62
					4.40		20.33		12.00	
					2.64		24.14		11.49	
AL1 - 6	20.08.2021	360	299	88	3.08	3.22	17.78	16.51	10.47	6.13
					4.40		21.60		5.36	
					2.20		10.16		2.55	
AL1 - 7	25.08.2021	340	290	72	2.64	2.64	13.34	18.00	14.81	11.57
					3.52		22.23		10.47	
					1.76		18.42		9.45	
AL1 – 8	27.08.2021	471	299	63	3.08	2.93	27.31	26.25	10.98	7.83
					1.76		30.49		5.62	
					3.96		20.96		6.89	
<b>Monthly Average</b>		495	262	71		3.76		20.96		10.74
<b>Standard Deviation</b>		171	46	14		1.83		3.96		3.08

NS: Not Specified

<b>Table 1B : Results of Air Pollutant Concentration at Marine Bhavan</b>					
<b>Parameter</b>	<b>Date</b>	<b>C<sub>6</sub>H<sub>6</sub> [µg/m<sup>3</sup>]</b>	<b>HC* ppm</b>	<b>CO [mg/m<sup>3</sup>]</b>	<b>CO<sub>2</sub> [ppm ]</b>
<b>Sampling Period</b>		<b>8 hr</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>
<b>NAAQMS limit</b>		<b>5.0 µg/m<sup>3</sup></b>	<b>NS</b>	<b>4.0 mg/m<sup>3</sup></b>	<b>NS</b>
AL1 – 1	04.08.2021	1.06	BDL	1.86	488
AL1 – 2	06.08.2021	1.22	BDL	1.74	496
AL1 – 3	11.08.2021	1.28	BDL	1.7	499
AL1 – 4	13.08.2021	1.2	BDL	1.68	501
AL1 – 5	18.08.2021	1.21	BDL	1.72	490
AL1 - 6	20.08.2021	1.06	BDL	1.62	497
AL1 – 7	25.08.2021	1.12	BDL	1.52	488
AL1 – 8	27.08.2021	1.06	BDL	1.72	496
Monthly Average		<b>1.15</b>	-	<b>1.70</b>	<b>494</b>
Standard Deviation		<b>0.09</b>	-	<b>0.10</b>	<b>5</b>

\* NMHC- Non- Methane Hydrocarbons

BDL- Below Detection Limit (Detection Limit – NMHC: 0.5ppm)

NS -Not Specified

At Marine Bhavan, the overall values of TSPM, PM<sub>10</sub>, PM<sub>2.5</sub>, SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> is attributed mainly by motor vehicle emission produced from various types of automobiles (both diesel and petrol driven). Moreover, the loading and unloading of Food Grains and Timber at Jetty no. 1 and 2 also contributes to the high levels of TSPM and PM<sub>10</sub>. The mean TSPM value at Marine Bhavan was 495 µg/m<sup>3</sup>, The mean PM<sub>10</sub> values were 262.0 µg/m<sup>3</sup>, which is above the permissible limit. PM<sub>2.5</sub> values were slightly above the permissible limit (mean = 71 µg/m<sup>3</sup>). The average values of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were within the permissible limit. The average values of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were 3.76 µg/ m<sup>3</sup>, 20.96 µg/ m<sup>3</sup> & 10.74 µg/ m<sup>3</sup> respectively. These were within the permissible limit.

The levels of Benzene, Hydrocarbons (HC) and CO were within the permissible limit at Marine Bhavan. The mean Benzene concentration was 1.15 µg/m<sup>3</sup>, well below the permissible limit of 5.0 µg/m<sup>3</sup>. HC's were below the detectable limit and Carbon Monoxide concentration was 1.70 mg/m<sup>3</sup>, well below the permissible limit of 4.0 mg/m<sup>3</sup>.



**Location 2: Oil Jetty (AL2)**

**Table 2 : Results of Air Pollutant Concentration at Oil Jetty**

Parameters	Date	TSPM [µg/m <sup>3</sup> ]	PM10 [µg/m <sup>3</sup> ]	PM2.5 [µg/m <sup>3</sup> ]	SO <sub>2</sub> [µg/m <sup>3</sup> ]		NO <sub>x</sub> [µg/m <sup>3</sup> ]		NH <sub>3</sub> [µg/m <sup>3</sup> ]	
					8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
Sampling Period	-	24hr	24hr	24hr	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
NAAQMS limit	-	NS	100 µg/m <sup>3</sup>	60 µg/m <sup>3</sup>	-	80 µg/m <sup>3</sup>	-	80 µg/m <sup>3</sup>	-	400 µg/m <sup>3</sup>
AL2 – 1	04.08.2021	299	222	55	3.52	3.81	18.42	16.30	13.53	14.81
					4.84		12.70		14.81	
					3.08		17.78		16.08	
AL2 – 2	06.08.2021	837	394	89	1.32	3.08	12.07	12.28	7.40	10.47
					3.08		10.80		11.74	
					4.84		13.97		12.25	
AL2 – 3	11.08.2021	403	350	49	8.35	9.38	33.66	25.62	4.08	6.89
					8.79		19.05		6.89	
					10.99		24.14		9.70	
AL2 – 4	13.08.2021	511	327	82	3.08	2.05	17.78	16.30	7.15	9.10
					1.76		15.88		10.72	
					1.32		15.24		9.45	
AL2 – 5	18.08.2021	567	281	75	3.08	2.20	17.15	18.63	9.70	7.83
					1.32		26.04		5.36	
					2.20		12.70		8.42	
AL2 – 6	20.08.2021	728	490	90	6.15	5.86	22.87	15.88	5.36	8.00
					7.91		8.89		8.42	
					3.52		15.88		10.21	
AL2 – 7	25.08.2021	344	237	67	0.88	1.17	24.14	20.75	9.96	10.38
					0.88		15.88		12.76	
					1.76		22.23		8.42	
AL2 – 8	27.08.2021	475	278	76	1.32	2.20	15.88	17.78	5.87	8.51
					1.76		24.14		9.19	
					3.52		13.34		10.47	
<b>Monthly Average</b>		520	322	73		3.72		17.94		9.50
<b>Standard Deviation</b>		186	88	15		2.70		3.95		2.47

NS: Not Specified

<b>Table 2B : Results of Air Pollutant Concentration at Oil Jetty</b>					
<b>Parameter</b>	<b>Date</b>	<b>C<sub>6</sub>H<sub>6</sub> [µg/m<sup>3</sup>]</b>	<b>HC* ppm</b>	<b>CO [mg/m<sup>3</sup>]</b>	<b>CO<sub>2</sub> [ppm ]</b>
<b>Sampling Period</b>		<b>8 hr</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>
<b>NAAQMS limit</b>		<b>5.0 µg/m<sup>3</sup></b>	<b>NS</b>	<b>4.0 mg/m<sup>3</sup></b>	<b>NS</b>
AL2 -1	04.08.2021	1.22	BDL	1.86	492
AL2 -2	06.08.2021	1.06	BDL	1.72	496
AL2 -3	11.08.2021	1.26	BDL	1.76	489
AL2 -4	13.08.2021	1.23	BDL	1.66	500
AL2 – 5	18.08.2021	1.2	BDL	1.84	496
AL2 – 6	20.08.2021	1.16	BDL	1.74	489
AL2 -7	25.08.2021	1.18	BDL	1.76	476
AL2 – 8	27.08.2021	1.23	BDL	1.7	490
<b>Monthly Average</b>		<b>1.19</b>	<b>-</b>	<b>1.76</b>	<b>491</b>
<b>Standard Deviation</b>		<b>0.06</b>	<b>-</b>	<b>0.07</b>	<b>7</b>

\* NMHC- Non- Methane Hydrocarbons

BDL- Below Detection Limit (Detection Limit – NMHC : 0.5ppm)

NS- Not Specified

The overall values of TSPM, PM<sub>10</sub>, PM<sub>2.5</sub>, SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> at Oil Jetty area was mainly by motor vehicle emission produced from various types of vehicles Oil Jetty Area. The mean TSPM values at Oil Jetty were 520 µg/m<sup>3</sup>. The mean PM<sub>10</sub> values were 322 µg/m<sup>3</sup>, which is above the permissible limit. PM<sub>2.5</sub> values were slightly above the permissible limit (mean = 73 µg/m<sup>3</sup>). The average values of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were within the permissible limit; The mean concentration of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were 3.72 µg/m<sup>3</sup>, 17.94 µg/m<sup>3</sup> and 9.50 µg/m<sup>3</sup> respectively.

The levels of Benzene, Hydrocarbons (HC) and CO were within the permissible limit at Oil Jetty. The mean Benzene concentration was 1.19 µg/m<sup>3</sup>. Well below the permissible limit of 5.0 µg/m<sup>3</sup>. , HC's were below the detectable limit and Carbon Monoxide concentration was 1.76 mg/m<sup>3</sup>, well below the permissible limit of 4.0 mg/m<sup>3</sup>.

**Location 3: Kandla Colony – Estate Office (AL-3)**

Table 3 : Results of Air Pollutant Concentration at Estate Office										
Parameters	Date	TSPM [µg/m <sup>3</sup> ]	PM10 [µg/m <sup>3</sup> ]	PM2.5 [µg/m <sup>3</sup> ]	SO <sub>2</sub> [µg/m <sup>3</sup> ]		NO <sub>x</sub> [µg/m <sup>3</sup> ]		NH <sub>3</sub> [µg/m <sup>3</sup> ]	
Sampling Period	-	24hr	24hr	24hr	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
NAAQMS limit	-	NS	100 µg/m <sup>3</sup>	60 µg/m <sup>3</sup>	-	80 µg/m <sup>3</sup>	-	80 µg/m <sup>3</sup>	-	400 µg/m <sup>3</sup>
AL3 – 1	04.08.2021	159	97	35	1.32	2.05	25.41	23.92	14.04	16.68
					2.20		33.66		17.36	
					2.64		12.70		18.64	
AL3 – 2	06.08.2021	473	176	60	3.96	9.96	12.70	13.13	8.68	7.40
					21.98		10.80		7.15	
					3.96		15.88		6.38	
AL3 – 3	11.08.2021	379	253	74	3.96	4.25	15.24	20.75	8.42	6.47
					5.28		20.96		7.15	
					3.52		26.04		3.83	
AL3 – 4	13.08.2021	652	331	67	4.84	2.49	9.53	9.32	12.76	9.62
					1.76		9.53		9.70	
					0.88		8.89		6.38	
AL3 – 5	18.08.2021	643	457	92	4.84	3.52	24.14	24.77	9.70	38.21
					3.52		34.30		10.47	
					2.20		15.88		94.45	
AL3 – 6	20.08.2021	721	389	75	4.84	4.25	20.96	20.96	10.21	8.25
					2.20		15.88		9.45	
					5.71		26.04		5.11	
AL3 – 7	25.08.2021	298	208	68	4.40	3.22	22.23	19.05	12.00	11.66
					3.52		17.78		12.00	
					1.76		17.15		10.98	
AL3 – 8	27.08.2021	574	300	96	2.64	2.93	16.51	17.57	11.49	8.76
					4.40		17.15		9.45	
					1.76		19.05		5.36	
<b>Monthly Average</b>		488	276	71		4.08		18.68		13.38
<b>Standard Deviation</b>		196	117	19		2.50		5.26		10.53

NS: Not Specified

<b>Table 3B : Results of Air Pollutant Concentration at Kandla Port Colony</b>					
<b>Parameter</b>	<b>Date</b>	<b>C<sub>6</sub>H<sub>6</sub> [µg/m<sup>3</sup>]</b>	<b>HC*</b>	<b>CO [mg/m<sup>3</sup>]</b>	<b>CO<sub>2</sub> [ppm ]</b>
<b>Sampling Period</b>		<b>8 hr</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>
<b>NAAQMS limit</b>		<b>5.0 µg/m<sup>3</sup></b>	<b>NS</b>	<b>4.0 mg/m<sup>3</sup></b>	<b>NS</b>
AL3 -1	04.08.2021	1.12	BDL	1.84	480
AL3 -2	06.08.2021	1.16	BDL	1.76	488
AL3 -3	11.08.2021	1.22	BDL	1.8	496
AL3 -4	13.08.2021	1.26	BDL	1.74	490
AL3 – 5	18.08.2021	1.2	BDL	1.79	496
AL3 – 6	20.08.2021	1.06	BDL	1.82	499
AL3 – 7	25.08.2021	1.11	BDL	1.8	500
AL3 – 8	27.08.2021	1.07	BDL	1.76	490
<b>Monthly Average</b>		<b>1.15</b>	<b>-</b>	<b>1.79</b>	<b>492</b>
<b>Standard Deviation</b>		<b>0.07</b>	<b>-</b>	<b>0.03</b>	<b>7</b>

\* NMHC- Non- Methane Hydrocarbons

BDL- Below Detection Limit ( Detection Limit – NMHC : 0.5 ppm)

NS- Not Specified

The overall values of TSPM, PM<sub>10</sub>, PM<sub>2.5</sub>, SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> at Kandla Port Colony was attributed by vehicle emission produced from trucks and heavy duty vehicles that pass through the road outside Kandla Port Colony. The mean TSPM values at Oil Jetty were 488 µg/m<sup>3</sup>, The mean PM<sub>10</sub> values were 276 µg/m<sup>3</sup>, which is above the permissible limit. PM<sub>2.5</sub> values were slightly above the permissible limit (mean = 71 µg/m<sup>3</sup>). The average values of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were 4.08 µg/m<sup>3</sup>, 18.68 µg/m<sup>3</sup> and 13.38 µg/m<sup>3</sup> respectively and were all within the permissible limit.

The levels of Benzene, Hydrocarbons (HC) and CO were within the permissible limit at Kandla Port Colony. The mean Benzene concentration was 1.15 µg/m<sup>3</sup>, well below the permissible limit of 5.0 µg/m<sup>3</sup>. HC's were below the detectable limit and Carbon Monoxide concentration was 1.79 mg/m<sup>3</sup>, well below the permissible limit of 4.0 mg/m<sup>3</sup>.

**Location 4: Gopalpuri Hospital (AL-4)**

<b>Table 4 : Results of Air Pollutant Concentration at Gopalpuri Hospital</b>										
<b>Parameter</b>	<b>Date</b>	<b>TSPM [µg/m3]</b>	<b>PM10 [µg/m3]</b>	<b>PM2.5 [µg/m3]</b>	<b>SO2 [µg/m3]</b>		<b>NOx [µg/m3]</b>		<b>NH3 [µg/m3]</b>	
<b>Sampling Period</b>	-	<b>24hr</b>	<b>24hr</b>	<b>24hr</b>	<b>8 hr</b>	<b>24hr (Avg.)</b>	<b>8 hr</b>	<b>24hr (Avg.)</b>	<b>8 hr</b>	<b>24hr (Avg.)</b>
<b>NAAQMS limit</b>	-	<b>NS</b>	<b>100 µg/m3</b>	<b>60 µg/m3</b>	-	<b>80 µg/m3</b>	-	<b>80 µg/m3</b>	-	<b>400 µg/m3</b>
<b>AL4 -1</b>	04.08.2021	128	77	28	3.08	3.08	7.62	12.70	9.70	8.76
					2.64		17.15		10.21	
					3.52		13.34		6.38	
<b>AL4 -2</b>	06.08.2021	180	115	38	3.52	2.05	6.35	10.16	5.36	5.02
					0.88		11.43		5.11	
					1.76		12.70		4.60	
<b>AL4 -3</b>	11.08.2021	228	110	48	1.76	3.08	11.43	9.95	4.34	4.85
					3.96		6.35		5.36	
					3.52		12.07		4.85	
<b>AL4 -4</b>	13.08.2021	327	260	60	3.08	3.08	11.43	9.74	7.15	5.62
					3.52		10.80		4.08	
					2.64		6.99		5.62	
<b>AL4 -5</b>	18.08.2021	269	156	70	2.20	3.08	15.88	14.61	5.36	6.89
					3.96		8.89		8.93	
					3.08		19.05		6.38	
<b>AL4 -6</b>	20.08.2021	228	113	86	3.52	2.34	13.97	12.91	5.36	9.02
					2.20		8.89		9.70	
					1.32		15.88		12.00	
<b>AL4 -7</b>	25.08.2021	222	116	49	2.20	2.78	19.05	17.15	6.89	6.72
					3.52		14.61		8.42	
					2.64		17.78		4.85	
<b>AL4 -8</b>	27.08.2021	249	119	30	2.64	3.08	12.07	12.07	7.91	8.93
					3.08		12.70		9.19	
					3.52		11.43		9.70	
<b>Monthly Average</b>		229	133	51		2.82		12.41		6.98
<b>Standard Deviation</b>		59	56	20		0.41		2.56		1.75

NS: Not Specified

<b>Table 4B : Results of Air Pollutant Concentration at Gopalpuri Hospital</b>					
<b>Parameter</b>	<b>Date</b>	<b>C<sub>6</sub>H<sub>6</sub> [µg/m<sup>3</sup>]</b>	<b>HC*</b>	<b>CO [mg/m<sup>3</sup>]</b>	<b>CO<sub>2</sub> [ppm ]</b>
<b>Sampling Period</b>		<b>8 hr</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>
<b>NAAQMS limit</b>		<b>5.0 µg/m<sup>3</sup></b>	<b>NS</b>	<b>4.0 mg/m<sup>3</sup></b>	<b>NS</b>
AL4 -1	04.08.2021	1.12	BDL	1.88	490
AL4 -2	06.08.2021	1.18	BDL	1.76	488
AL4 -3	11.08.2021	1.26	BDL	1.72	496
AL4 -4	13.08.2021	1.21	BDL	1.8	500
AL4 -5	18.08.2021	1.28	BDL	1.79	482
AL4 -6	20.08.2021	1.2	BDL	1.84	493
AL4 -7	25.08.2021	1.18	BDL	1.86	498
AL4 -8	27.08.2021	1.16	BDL	1.8	490
<b>Monthly Average</b>		<b>1.20</b>	<b>-</b>	<b>1.81</b>	<b>492</b>
<b>Standard Deviation</b>		<b>0.05</b>	<b>-</b>	<b>0.05</b>	<b>6</b>

\* NMHC- Non- Methane Hydrocarbons

BDL- Below Detection Limit (Detection Limit – NMHC: 0.5 ppm)

NS-Not Specified

The overall values of TSPM, PM<sub>10</sub>, PM<sub>2.5</sub>, SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> at Gopalpuri Hospital was attributed by vehicle emission produced from light motor vehicles of the colony residents. The mean TSPM values at Oil Jetty were 229 µg/m<sup>3</sup>, The mean PM<sub>10</sub> values were 133 µg/m<sup>3</sup>, which is above the permissible limit. PM<sub>2.5</sub> values were slightly above the permissible limit (mean= 51 µg/m<sup>3</sup>). The average values of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were 2.82 µg/m<sup>3</sup>, 12.41 µg/m<sup>3</sup> and 6.98 µg/m<sup>3</sup> respectively and were all within the permissible limit.

The levels of Benzene, Hydrocarbons (HC) and CO were within the permissible limit at Gopalpuri Hospital. The mean Benzene concentration was 1.20 µg/m<sup>3</sup>, well below the permissible limit of 5.0 µg/m<sup>3</sup>. HC's were below the detectable limit and Carbon Monoxide concentration was 1.81 mg/m<sup>3</sup>, well below the permissible limit of 4.0 mg/m<sup>3</sup>.

**Location 5: Coal Storage Area (AL-5)**

Table 5 : Results of Air Pollutant Concentration at Coal Storage Area										
Parameters	Date	TSPM [µg/m <sup>3</sup> ]	PM10 [µg/m <sup>3</sup> ]	PM2.5 [µg/m <sup>3</sup> ]	SO <sub>2</sub> [µg/m <sup>3</sup> ]		NO <sub>x</sub> [µg/m <sup>3</sup> ]		NH <sub>3</sub> [µg/m <sup>3</sup> ]	
Sampling Period	-	24hr	24hr	24hr	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
NAAQMS limit	-	NS	100 µg/m <sup>3</sup>	60 µg/m <sup>3</sup>	-	80 µg/m <sup>3</sup>	-	80 µg/m <sup>3</sup>	-	400 µg/m <sup>3</sup>
AL5 – 1	04.08.2021	312	167	69	3.08	3.66	22.23	22.23	9.45	9.53
					3.52		23.50		7.15	
					4.40		20.96		12.00	
AL5 – 2	06.08.2021	530	333	77	9.23	6.15	20.96	20.11	16.59	17.02
					5.71		24.77		17.87	
					3.52		14.61		16.59	
AL5 – 3	11.08.2021	759	394	92	10.99	9.23	24.14	27.74	8.42	7.74
					7.47		25.41		7.15	
					9.23		33.66		7.66	
AL5 – 4	13.08.2021	813	435	94	1.76	1.61	17.78	18.84	13.02	10.30
					1.32		19.05		8.93	
					1.76		19.69		8.93	
AL5 – 5	18.08.2021	700	471	79	4.40	3.96	21.60	22.02	12.00	11.66
					4.40		19.05		10.47	
					3.08		25.41		12.51	
AL5 – 6	20.08.2021	566	427	80	3.08	3.96	16.51	18.00	16.85	15.66
					3.52		15.24		16.34	
					5.28		22.23		13.79	
AL5 – 7	25.08.2021	456	224	76	3.96	4.10	13.97	17.15	10.47	7.04
					4.40		19.69		9.70	
					3.96		17.78		0.94	
AL5 – 8	27.08.2021	249	164	70	4.40	3.66	23.50	27.52	11.49	13.44
					3.52		28.58		14.04	
					3.08		30.49		14.81	
<b>Monthly Average</b>		548	327	80		4.54		21.70		11.55
<b>Standard Deviation</b>		204	125	9		2.26		4.07		3.60

NS: Not Specified

<b>Table 5B : Results of Air Pollutant Concentration at Coal Storage Area</b>					
<b>Parameter</b>	<b>Date</b>	<b>C<sub>6</sub>H<sub>6</sub> [µg/m<sup>3</sup>]</b>	<b>HC*</b>	<b>CO [mg/m<sup>3</sup>]</b>	<b>CO<sub>2</sub> [ppm ]</b>
<b>Sampling Period</b>		<b>8 hr</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>
<b>NAAQMS limit</b>		<b>5.0 µg/m<sup>3</sup></b>	<b>NS</b>	<b>4.0 mg/m<sup>3</sup></b>	<b>NS</b>
AL5 – 1	04.08.2021	1.06	BDL	1.96	460
AL5 – 2	06.08.2021	1.11	BDL	1.88	458
AL5 – 3	11.08.2021	1.26	BDL	1.9	456
AL5 – 4	13.08.2021	1.3	BDL	1.82	460
AL5 – 5	18.08.2021	1.26	BDL	1.96	456
AL5 – 6	20.08.2021	1.22	BDL	1.93	474
AL5 – 7	25.08.2021	1.38	BDL	1.89	470
AL5 – 8	27.08.2021	1.30	BDL	1.9	468
<b>Monthly Average</b>		<b>1.24</b>	<b>-</b>	<b>1.91</b>	<b>463</b>
<b>Standard Deviation</b>		<b>0.11</b>	<b>-</b>	<b>0.05</b>	<b>7</b>

\* NMHC- Non- Methane Hydrocarbons

BDL- Below Detection Limit (Detection Limit – NMHC: 0.5 ppm)

NS-Not Specified

The overall values of TSPM, PM<sub>10</sub>, PM<sub>2.5</sub>, SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> at Coal Storage Area was comparatively highest among all the locations of Air Quality monitoring in Kandla Port. High values of TSPM, PM<sub>10</sub>, PM<sub>2.5</sub>, SO<sub>2</sub>, NO<sub>x</sub> at this location was due to lifting of coal with grab and other coal handling processes near Berth no. 6 & 7. Moreover, the traffic was also heavy around this place for transport of coal thus emissions produced from heavy vehicles. The mean TSPM values at Coal storage were 548 µg/m<sup>3</sup>. The mean PM<sub>10</sub> values were 327 µg/m<sup>3</sup>, which is well above the permissible limit. PM<sub>2.5</sub> values were above the permissible limit (mean = 80 µg/m<sup>3</sup>). The average values of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were 4.54 µg/m<sup>3</sup>, 21.70 µg/m<sup>3</sup> and 11.55 µg/m<sup>3</sup> respectively and were all within the permissible limit.

The levels of Benzene, Hydrocarbons (HC) and CO were within the permissible limit at Coal Storage Area. The mean Benzene concentration was 1.24 µg/m<sup>3</sup>, well below the permissible limit of 5.0 µg/m<sup>3</sup>. HC's were below the detectable limit and Carbon Monoxide concentration was 1.91 mg/m<sup>3</sup>, well below the permissible limit of 4.0 mg/m<sup>3</sup>.



**Location 6: Tuna Port (AL-6)**

Table 6 : Results of Air Pollutant Concentration at Tuna Port										
Parameters	Date	TSPM [µg/m3]	PM10 [µg/m3]	PM2.5 [µg/m3]	SO2 [µg/m3]		NOx [µg/m3]		NH3 [µg/m3]	
Sampling Period	-	24hr	24hr	24hr	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
NAAQMS limit	-	NS	100 µg/m3	60 µg/m3	-	80 µg/m3	-	80 µg/m3	-	400 µg/m3
AL6 -1	04.08.2021	133	75	26	3.96	2.93	22.23	15.46	15.06	12.17
					2.64		13.34		12.25	
					2.20		10.80		9.19	
AL6 - 2	06.08.2021	203	149	67	2.20	2.05	8.26	9.10	5.87	6.81
					2.20		10.16		6.38	
					1.76		8.89		8.17	
AL6 - 3	11.08.2021	316	166	44	5.28	4.40	17.78	18.63	6.89	6.47
					4.84		22.23		4.60	
					3.08		15.88		7.91	
AL6 - 4	13.08.2021	530	342	83	3.08	2.05	5.72	7.83	5.36	6.55
					1.32		9.53		7.91	
					1.76		8.26		6.38	
AL6 - 5	18.08.2021	468	291	84	0.88	2.05	20.96	17.15	12.76	12.08
					1.76		12.70		12.25	
					3.52		17.78		11.23	
AL6 - 6	20.08.2021	319	181	63	4.40	3.08	33.03	28.58	10.47	12.93
					1.32		22.87		15.57	
					3.52		29.85		12.76	
AL6 - 7	25.08.2021	256	156	58	3.08	3.22	15.88	17.15	9.96	10.21
					2.64		17.78		9.45	
					3.96		17.78		11.23	
AL6 - 8	27.08.2021	554	375	80	2.64	3.08	17.15	16.30	10.47	9.96
					3.08		12.07		8.42	
					3.52		19.69		10.98	
<b>Monthly Average</b>		347	217	63		2.86		16.28		9.65
<b>Standard Deviation</b>		155	106	20		0.81		6.35		2.71

NS: Not Specified

<b>Table 6B : Results of Air Pollutant Concentration at Tuna Port</b>					
<b>Parameter</b>	<b>Date</b>	<b>C<sub>6</sub>H<sub>6</sub> [µg/m<sup>3</sup>]</b>	<b>HC*</b>	<b>CO [mg/m<sup>3</sup>]</b>	<b>CO<sub>2</sub> [ppm ]</b>
<b>Sampling Period</b>		<b>8 hr</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>
<b>NAAQMS limit</b>		<b>5.0 µg/m<sup>3</sup></b>	<b>NS</b>	<b>4.0 mg/m<sup>3</sup></b>	<b>NS</b>
AL6 -1	04.08.2021	1.11	BDL	1.74	460
AL6 – 2	06.08.2021	1.26	BDL	1.89	470
AL6 – 3	11.08.2021	1.2	BDL	1.88	472
AL6 – 4	13.08.2021	1.16	BDL	1.9	466
AL6 – 5	18.08.2021	1.07	BDL	1.97	460
AL6 – 6	20.08.2021	1.11	BDL	1.89	451
AL6 – 7	25.08.2021	1.2	BDL	1.8	460
AL6 – 8	27.08.2021	1.21	BDL	1.82	470
<b>Monthly Average</b>		<b>1.17</b>	<b>-</b>	<b>1.86</b>	<b>464</b>
<b>Standard Deviation</b>		<b>0.06</b>	<b>-</b>	<b>0.07</b>	<b>7</b>

\* NMHC- Non- Methane Hydrocarbons

BDL- Below Detection Limit (Detection Limit – NMHC : 0.5 ppm)

NS- Not Specified

The mean TSPM values at Tuna Port were 347 µg/m<sup>3</sup>, The mean PM<sub>10</sub> values were 217 µg/m<sup>3</sup>, which is above the permissible limit. PM<sub>2.5</sub> values were slightly the permissible limit (mean = 63 µg/m<sup>3</sup>). The average values of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were 2.86 µg/m<sup>3</sup>, 16.28 µg/m<sup>3</sup> and 9.65 µg/m<sup>3</sup> respectively and were all within the permissible limit.

The levels of Benzene, Hydrocarbons (HC) and CO were within the permissible limit at Tuna Port. The mean Benzene concentration was 1.17 µg/m<sup>3</sup>, well below the permissible limit of 5.0 µg/m<sup>3</sup>. HC's were below the detectable limit and Carbon Monoxide concentration was 1.86 mg/m<sup>3</sup>, well below the permissible limit of 4.0 mg/m<sup>3</sup>.

**Environmental Monitoring Report of Deendayal Port Trust, AUGUST-2021**

**Location 7: Signal Building (Vadinar) (AL-7)**

**Table 7 : Results of Air Pollutant Concentration at Signal Building**

Parameters	Date	TSPM [µg/m <sup>3</sup> ]	PM10 [µg/m <sup>3</sup> ]	PM2.5 [µg/m <sup>3</sup> ]	SO <sub>2</sub> [µg/m <sup>3</sup> ]		NO <sub>x</sub> [µg/m <sup>3</sup> ]		NH <sub>3</sub> [µg/m <sup>3</sup> ]	
					8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
Sampling Period	-	24hr	24hr	24hr	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
NAAQMS limit	-	NS	100 µg/m <sup>3</sup>	60 µg/m <sup>3</sup>	-	80 µg/m <sup>3</sup>	-	80 µg/m <sup>3</sup>	-	400 µg/m <sup>3</sup>
AL7 -1	04.08.2021	144	97	30	3.08	3.08	7.62	11.86	7.66	5.53
					2.64		14.61		5.36	
					3.52		13.34		3.57	
AL7 -2	06.08.2021	180	120	38	3.96	4.98	28.58	17.78	4.60	6.55
					4.84		14.61		10.47	
					6.15		10.16		4.60	
AL7 -3	11.08.2021	148	85	29	2.20	2.34	8.89	16.51	5.36	8.68
					3.08		26.04		11.49	
					1.76		14.61		9.19	
AL7 -4	13.08.2021	165	115	32	2.64	2.07	20.33	14.19	9.19	6.47
					0.48		13.34		3.57	
					3.08		8.89		6.64	
AL7 -5	18.08.2021	151	99	35	4.84	3.96	14.61	16.30	8.93	7.49
					3.08		21.60		6.38	
					3.96		12.70		7.15	
AL7 -6	20.08.2021	173	104	64	3.96	3.81	9.53	11.22	4.85	4.68
					3.08		8.89		3.57	
					4.40		15.24		5.62	
AL7 -7	25.08.2021	168	114	44	3.96	1.67	13.34	13.55	16.85	10.89
					0.44		6.99		12.00	
					0.62		20.33		3.83	
AL7 -8	27.08.2021	113	54	37	6.15	2.78	7.62	12.70	9.70	8.76
					1.76		17.15		9.45	
					0.44		13.34		7.15	
<b>Monthly Average</b>		155	98	39		3.1		14.3		7.4
<b>Standard Deviation</b>		21	21	11		1.1		2.4		2.0

NS: Not Specified

<b>Table 7B : Results of Air Pollutant Concentration at Signal Building</b>					
<b>Parameter</b>	<b>Date</b>	<b>C<sub>6</sub>H<sub>6</sub> [µg/m<sup>3</sup>]</b>	<b>HC*</b>	<b>CO [mg/m<sup>3</sup>]</b>	<b>CO<sub>2</sub> [ppm ]</b>
<b>Sampling Period</b>		<b>8 hr</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>
<b>NAAQMS limit</b>		<b>5.0 µg/m<sup>3</sup></b>	<b>NS</b>	<b>4.0 mg/m<sup>3</sup></b>	<b>NS</b>
AL7 -1	04.08.2021	1.11	BDL	1.9	460
AL7 – 2	06.08.2021	1.2	BDL	1.86	472
AL7 – 3	11.08.2021	1.18	BDL	1.79	460
AL7 – 4	13.08.2021	1.08	BDL	1.86	461
AL7 – 5	18.08.2021	1.12	BDL	1.96	456
AL7 – 6	20.08.2021	1.2	BDL	1.9	460
AL7 – 7	25.08.2021	1.18	BDL	1.88	470
AL7 – 8	27.08.2021	1.1	BDL	1.82	465
<b>Monthly Average</b>		<b>1.15</b>	<b>-</b>	<b>1.87</b>	<b>463</b>
<b>Standard Deviation</b>		<b>0.05</b>	<b>-</b>	<b>0.05</b>	<b>6</b>

\* NMHC- Non- Methane Hydrocarbons

BDL- Below Detection Limit (Detection Limit – NMHC : 0.5 ppm)

NS\_ Not Specified

The mean TSPM values at Vadinar Port were 155 µg/m<sup>3</sup>. The mean PM<sub>10</sub> values were 98 µg/m<sup>3</sup>, which is below the permissible limit. PM<sub>2.5</sub> values were also within the permissible limit (mean = 39 µg/m<sup>3</sup>). The average values of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were 3.1 µg/m<sup>3</sup>, 14.3 µg/m<sup>3</sup> and 7.4 µg/m<sup>3</sup> respectively and were all within the permissible limit.

The levels of Benzene, Hydrocarbons (HC) and CO were within the permissible limit at Vadinar Port. The mean Benzene concentration was 1.15 µg/m<sup>3</sup>, well below the permissible limit of 5.0 µg/m<sup>3</sup>. HC's were below the detectable limit and Carbon Monoxide concentration was 1.87 mg/m<sup>3</sup>, well below the permissible limit of 4.0 mg/m<sup>3</sup>.

**Location 8: Admin Building (Vadinar) (AL-8)**

Table 8 : Results of Air Pollutant Concentration at Admin Building										
Parameters	Date	TSPM [µg/m3]	PM10 [µg/m3]	PM2.5 [µg/m3]	SO2 [µg/m3]		NOx [µg/m3]		NH3 [µg/m3]	
Sampling Period	-	24hr	24hr	24hr	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
NAAQMS limit	-	NS	100 µg/m3	60 µg/m3	-	80 µg/m3	-	80 µg/m3	-	400 µg/m3
AL8 -1	04.08.2021	119	55	28	4.84	3.52	19.05	18.42	7.15	6.30
					2.64		22.87		6.64	
					3.08		13.34		5.11	
AL8 -2	06.08.2021	111	56	47	17.58	6.30	17.78	16.73	7.91	8.00
					0.44		19.05		5.62	
					0.88		13.34		10.47	
AL8 -3	11.08.2021	180	100	56	1.76	2.78	15.24	15.24	4.34	5.19
					3.52		22.87		4.85	
					3.08		7.62		6.38	
AL8 -4	13.08.2021	130	77	42	3.96	4.54	13.97	11.86	8.17	7.15
					6.15		10.16		10.47	
					3.52		11.43		2.81	
AL8 -5	18.08.2021	100	68	29	3.96	2.64	7.62	8.89	7.40	7.40
					0.88		8.89		9.45	
					3.08		10.16		5.36	
AL8 -6	20.08.2021	160	97	58	3.52	4.98	12.70	12.70	8.93	8.42
					5.28		10.80		9.19	
					6.15		14.61		7.15	
AL8 -5	25.08.2021	143	65	49	3.52	3.96	6.99	12.49	12.00	8.42
					3.96		17.15		4.34	
					4.40		13.34		8.93	
AL8-6	27.08.2021	160	100	53	2.20	2.93	7.62	11.43	8.17	6.30
					3.08		19.05		4.60	
					3.52		7.62		6.13	
<b>Monthly Average</b>		138	77	45		4.0		13.5		7.1
<b>Standard Deviation</b>		28	19	12		1.3		3.1		1.2

NS: Not Specified

<b>Table 8B : Results of Air Pollutant Concentration at Admin Building</b>					
<b>Parameter</b>	<b>Date</b>	<b>C<sub>6</sub>H<sub>6</sub> [µg/m<sup>3</sup>]</b>	<b>HC*</b>	<b>CO [mg/m<sup>3</sup>]</b>	<b>CO<sub>2</sub> [ppm ]</b>
<b>Sampling Period</b>		<b>8 hr</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>
<b>NAAQMS limit</b>		<b>5.0 µg/m<sup>3</sup></b>	<b>NS</b>	<b>4.0 mg/m<sup>3</sup></b>	<b>NS</b>
AL8 -1	04.08.2021	1.12	BDL	1.96	460
AL8-2	06.08.2021	1.06	BDL	1.86	456
AL8 -3	11.08.2021	1.11	BDL	1.88	466
AL8-4	13.08.2021	1.18	BDL	1.9	470
AL8 -5	18.08.2021	1.26	BDL	1.92	466
AL8-6	20.08.2021	1.16	BDL	1.96	460
AL8-7	25.08.2021	1.2	BDL	1.86	456
AL8-8	27.08.2021	1.26	BDL	1.8	462
Monthly Average		<b>1.17</b>	-	<b>1.89</b>	<b>462</b>
Standard Deviation		<b>0.07</b>	-	<b>0.05</b>	<b>5</b>

\* NMHC- Non- Methane Hydrocarbons

BDL- Below Detection Limit (Detection Limit – NMHC: 0.5 ppm)

NS-Not Specified

The overall values of TSPM, PM<sub>10</sub>, PM<sub>2.5</sub>, SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> at Admin Building Vadinar was comparatively low among all the locations of Air Quality monitoring in Kandla Port and Vadinar Port. The mean TSPM values at Vadinar Port were 138 µg/m<sup>3</sup>. The mean PM<sub>10</sub> values were 77 µg/m<sup>3</sup>, which is below the permissible limit. PM<sub>2.5</sub> values were also within the permissible limit (mean = 45.0 µg/m<sup>3</sup>). The average values of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were 4.0µg/m<sup>3</sup>, 13.5 µg/m<sup>3</sup> and 7.1 µg/m<sup>3</sup> respectively and were all within the permissible limit.

The levels of Benzene, Hydrocarbons (HC) and CO were within the permissible limit at Admin Building, Vadinar Port. The mean Benzene concentration was 1.17 µg/m<sup>3</sup>, well below the permissible limit of 5.0 µg/m<sup>3</sup>. HC's were below the detectable limit and Carbon Monoxide concentration was 1.89 mg/m<sup>3</sup>, well below the permissible limit of 4.0 mg/m<sup>3</sup>.

#### **1.4 Observations and Conclusion**

During the monitoring period, the overall Ambient Air Quality of the port area was found to be well within the desired levels for various gaseous pollutants. However, Particulate matter as PM<sub>10</sub> and PM<sub>2.5</sub> was found to exceed the limits at locations like Near Coal storage area, Marine Bhavan, Estate Office , Tuna Port and Oil Jetty area.

## 2. Drinking Water Quality Monitoring

Drinking Water Quality Monitoring was carried out at twenty stations at Kandla, Vadinar & Township Area of Deendayal Port.

### 2.1 Drinking Water Monitoring Methodology

Drinking water samples were collected from 20 locations as prescribed in the tender document. Samples for physico-chemical analysis were collected in 1 liter carboys and samples for microbiological parameters were collected in sterilized bottles. These samples were then analyzed in laboratory for various drinking water parameters at Kandla Lab/Surat.

The Sampling and Analysis was done as per standard methods - IS 10500:2012. The water samples were analyzed for various parameters, viz. Color , Odor, Turbidity , Conductivity , pH , Chlorides , TDS, Total Hardness, Iron , Sulphate , Salinity , DO, BOD, Na, K, Ca, Mg, F, NO<sub>3</sub>, NO<sub>2</sub>, Mn, Cr-6, Cu, Cd, As, Hg, Pb, Zn, Bacterial Count (cfu) .

### 2.2 Results

The Drinking Water Quality monitoring data for 20 stations are given in below from table No. 9 to Table No. 15

**Table 9: Drinking Water Quality Monitoring Parameters for Nirman Building 1, P & C building & Main Gate (North) at Kandla**

Sr. No.	Parameter	Unit	Nirman Building 1	P & C building	Main Gate North	Acceptable Limits as per IS 10500 : 2012	Permissible Limits in the absence of Alternate Source as per IS 10500 : 2012
1	pH	pH Unit	7.4	7.3	7.2	6.5 to 8.5	6.5 to 8.5
2	Total Dissolved Solids	mg/l	930	1250	890	500	2000
3	Turbidity	NTU	0	1	0	1.0	5.0
4	Odor	-	Odorless	Odorless	Odorless	Agreeable	Agreeable
5	Color	Hazen Units	Colorless	Colorless	Colorless	5.0	15.0
6	Conductivity	µs/cm	1850	2460	1700	NS*	NS*
7	Biochemical Oxygen Demand	mg/l	<2	<2	<2	NS*	NS*
8	Chloride as Cl	mg/l	626	656	541	250.0	1000.0
9	Ca as Ca	mg/l	68.14	60.12	76.15	75.0	200.0
10	Mg as Mg	mg/l	58.32	72.90	68.04	30.0	100.0
11	Total Hardness	mg/l	390	410	430	200.0	600.0
12	Iron as Fe	mg/l	<0.01	<0.01	<0.01	0.3	No Relaxation
13	Fluorides as F	mg/l	0.34	0.41	0.13	1.0	1.5
14	Sulphate as SO4	mg/l	290.4	175.2	200.4	200.0	400
15	Nitrite as NO2	mg/l	<0.01	<0.01	<0.01	NS*	NS*
16	Nitrate as NO3	mg/l	6.27	8.10	13.38	45.0	No Relaxation
17	Salinity	%	1.13	1.19	0.98	NS*	NS*
18	Sodium as Na	mg/l	160	178	150	NS*	NS*
19	Potassium as K	mg/l	2.2	2.3	2.8	NS*	NS*
20	Manganese	mg/l	<0.04	<0.04	<0.04	0.1	0.3
21	Hexavalent Chromium	mg/l	<0.03	<0.03	<0.03	NS*	NS*
22	Copper	mg/l	<0.05	<0.05	<0.05	0.05	1.5
23	Cadmium	mg/l	<0.002	<0.002	<0.002	0.003	0.003
24	Arsenic	mg/l	<0.01	<0.01	<0.01	0.01	0.05
25	Mercury	mg/l	<0.001	<0.001	<0.001	0.001	0.001
26	Lead	mg/l	<0.01	<0.01	<0.01	0.01	0.01
27	Zinc	mg/l	<0.1	<0.1	<0.1	5.0	15.0
28	Bacterial Count	CFU/100ml	Absent	Absent	Absent	Absent	Absent

\*NS: Not Specified



**Table 10: Drinking Water Quality Monitoring Parameters for Canteen, West Gate – I & Wharf Area at Kandla**

Sr. No.	Parameter	Unit	Canteen	West Gate – I	Wharf Area	Acceptable Limits as per IS 10500 : 2012	Permissible Limits in the absence of Alternate Source as per IS 10500 : 2012
1	pH	pH Unit	7.6	7.3	7.2	6.5 to 8.5	6.5 to 8.5
2	Total Dissolved Solids	mg/l	1320	990	1030	500	2000
3	Turbidity	NTU	0	0	1	1.0	5.0
4	Odor	-	Odorless	Odorless	Odorless	Agreeable	Agreeable
5	Color	Hazen Units	Colorless	Colorless	Colorless	5.0	15.0
6	Conductivity	µs/cm	2590	1890	2010	NS*	NS*
7	Biochemical Oxygen Demand	mg/l	<2	<2	<2	NS*	NS*
8	Chloride as Cl	mg/l	717	596	616	250.0	1000.0
9	Ca as Ca	mg/l	64.13	60.12	56.11	75.0	200.0
10	Mg as Mg	mg/l	72.90	70.47	68.04	30.0	100.0
11	Total Hardness	mg/l	390	360	390	200.0	600.0
12	Iron as Fe	mg/l	<0.01	<0.01	<0.01	0.3	No Relaxation
13	Fluorides as F	mg/l	0.44	0.49	0.51	1.0	1.5
14	Sulphate as SO <sub>4</sub>	mg/l	190.8	198	289.2	200.0	400
15	Nitrite as NO <sub>2</sub>	mg/l	<0.01	<0.01	<0.01	NS*	NS*
16	Nitrate as NO <sub>3</sub>	mg/l	8.80	10.42	9.50	45.0	No Relaxation
17	Salinity	%	1.29	1.08	1.11	NS*	NS*
18	Sodium as Na	mg/l	130	168	158	NS*	NS*
19	Potassium as K	mg/l	3	2.2	2.4	NS*	NS*
20	Manganese	mg/l	<0.04	<0.04	<0.04	0.1	0.3
21	Hexavalent Chromium	mg/l	<0.03	<0.03	<0.03	NS*	NS*
22	Copper	mg/l	<0.05	<0.05	<0.05	0.05	1.5
23	Cadmium	mg/l	<0.002	<0.002	<0.002	0.003	0.003
24	Arsenic	mg/l	<0.01	<0.01	<0.01	0.01	0.05
25	Mercury	mg/l	<0.001	<0.001	<0.001	0.001	0.001
26	Lead	mg/l	<0.01	<0.01	<0.01	0.01	0.01
27	Zinc	mg/l	<0.1	<0.1	<0.1	5.0	15.0
28	Bacterial Count	CFU/100ml	Absent	Absent	Absent	Absent	Absent

\*NS: Not Specified

**Table 11: Drinking Water Quality Monitoring Parameters for Sewa sadan – 3, Workshop I & Custom Building at Kandla**

Sr. No.	Parameter	Unit	SewaSadan – 3	Workshop	Custom Building	Acceptable Limits as per IS 10500 : 2012	Permissible Limits in the absence of Alternate Source as per IS 10500 : 2012
1	pH	pH Unit	7.3	7.8	7.4	6.5 to 8.5	6.5 to 8.5
2	Total Dissolved Solids	mg/l	950	1050	1100	500	2000
3	Turbidity	NTU	1	1	0	1.0	5.0
4	Odor	-	Odorless	Odorless	Odorless	Agreeable	Agreeable
5	Color	Hazen Units	Colorless	Colorless	Colorless	5.0	15.0
6	Conductivity	µs/cm	1890	2080	2150	NS*	NS*
7	Biochemical Oxygen Demand	mg/l	<2	<2	<2	NS*	NS*
8	Chloride	mg/l	742	692	576	250.0	1000.0
9	Ca as Ca	mg/l	76.15	60.12	52.10	75.0	200.0
10	Mg as Mg	mg/l	58.32	68.04	68.04	30.0	100.0
11	Total Hardness	mg/l	400	400	380	200.0	600.0
12	Iron as Fe	mg/l	<0.01	<0.01	<0.01	0.3	No Relaxation
13	Fluorides	mg/l	0.88	0.59	0.59	1.0	1.5
14	Sulphate	mg/l	219.6	207.6	174	200.0	400
15	Nitrite	mg/l	<0.01	<0.01	<0.01	NS*	NS*
16	Nitrate	mg/l	13.73	10.21	12.88	45.0	No Relaxation
17	Salinity	%	1.34	1.25	1.04	NS*	NS*
18	Sodium as Na	mg/l	148	150	166	NS*	NS*
19	Potassium as K	mg/l	2.3	2.4	2.8	NS*	NS*
20	Manganese	mg/l	<0.04	<0.04	<0.04	0.1	0.3
21	Hexavalent Chromium	mg/l	<0.03	<0.03	<0.03	NS*	NS*
22	Copper	mg/l	<0.05	<0.05	<0.05	0.05	1.5
23	Cadmium	mg/l	<0.002	<0.002	<0.002	0.003	0.003
24	Arsenic	mg/l	<0.01	<0.01	<0.01	0.01	0.05
25	Mercury	mg/l	<0.001	<0.001	<0.001	0.001	0.001
26	Lead	mg/l	<0.01	<0.01	<0.01	0.01	0.01
27	Zinc	mg/l	<0.1	<0.1	<0.1	5.0	15.0
28	Bacterial Count	CFU/100ml	Absent	Absent	Absent	Absent	Absent

\*NS: Not Specified

**Table 12: Drinking Water Quality Monitoring Parameters for Port Colony Kandla, Hospital Kandla & A.O. Building at Gandhidham**

Sr. No.	Parameter	Unit	Port Colony Kandla	Hospital Kandla	A.O. Building	Acceptable Limits as per IS 10500 : 2012	Permissible Limits in the absence of Alternate Source as per IS 10500 : 2012
1	pH	pH Unit	7.5	7.4	7.1	6.5 to 8.5	6.5 to 8.5
2	Total Dissolved Solids	mg/l	1080	1350	950	500	2000
3	Turbidity	NTU	0	0	1	1.0	5.0
4	Odor	-	Odorless	Odorless	Odorless	Agreeable	Agreeable
5	Color	Hazen Units	Colorless	Colorless	Colorless	5.0	15.0
6	Conductivity	µs/cm	2100	2670	1890	NS*	NS*
7	Biochemical Oxygen Demand	mg/l	<2	<2	<2	NS*	NS*
8	Chloride	mg/l	767	712	641	250.0	1000.0
9	Ca as Ca	mg/l	60.12	64.13	56.11	75.0	200.0
10	Mg as Mg	mg/l	70.47	72.90	82.62	30.0	100.0
11	Total Hardness	mg/l	370	410	430	200.0	600.0
12	Iron as Fe	mg/l	<0.01	<0.01	<0.01	0.3	No Relaxation
13	Fluorides	mg/l	0.85	0.65	0.93	1.0	1.5
14	Sulphate	mg/l	178.8	202.8	207.6	200.0	400
15	Nitrite	mg/l	<0.01	<0.01	<0.01	NS*	NS*
16	Nitrate	mg/l	9.50	10.21	11.62	45.0	No Relaxation
17	Salinity	%	1.39	1.29	1.16	NS*	NS*
18	Sodium as Na	mg/l	170	164	178	NS*	NS*
19	Potassium as K	mg/l	2.7	2.3	2.8	NS*	NS*
20	Manganese	mg/l	<0.04	<0.04	<0.04	0.1	0.3
21	Hexavalent Chromium	mg/l	<0.03	<0.03	<0.03	NS*	NS*
22	Copper	mg/l	<0.05	<0.05	<0.05	0.05	1.5
23	Cadmium	mg/l	<0.002	<0.002	<0.002	0.003	0.003
24	Arsenic	mg/l	<0.01	<0.01	<0.01	0.01	0.05
25	Mercury	mg/l	<0.001	<0.001	<0.001	0.001	0.001
26	Lead	mg/l	<0.01	<0.01	<0.01	0.01	0.01
27	Zinc	mg/l	<0.1	<0.1	<0.1	5.0	15.0
28	Bacterial Count	CFU/100ml	Absent	Absent	Absent	Absent	Absent

\*NS: Not Specified

**Table 13: Drinking Water Quality Monitoring Parameters for School Gopalpuri, Guest House & E - Type Quarter at Gopalpuri, Gandhidham**

Sr. No.	Parameter	Unit	School Gopalpuri	Guest House	E - Type Quarter	Acceptable Limits as per IS 10500 : 2012	Permissible Limits in the absence of Alternate Source as per IS 10500 : 2012
1	pH	pH Unit	7.5	7.5	7.4	6.5 to 8.5	6.5 to 8.5
2	Total Dissolved Solids	mg/l	1010	1350	1080	500	2000
3	Turbidity	NTU	1	0	1	1.0	5.0
4	Odor	-	Odorless	Odorless	Odorless	Agreeable	Agreeable
5	Color	Hazen Units	Colorless	Colorless	Colorless	5.0	15.0
6	Conductivity	µs/cm	1990	2670	2120	NS*	NS*
7	Biochemical Oxygen Demand	mg/l	<2	<2	<2	NS*	NS*
8	Chloride	mg/l	697	496	586	250.0	1000.0
9	Ca as Ca	mg/l	56.11	64.13	72.14	75.0	200.0
10	Mg as Mg	mg/l	70.47	53.46	58.32	30.0	100.0
11	Total Hardness	mg/l	380	360	390	200.0	600.0
12	Iron as Fe	mg/l	<0.01	<0.01	<0.01	0.3	No Relaxation
13	Fluorides	mg/l	0.58	0.80	1.05	1.0	1.5
14	Sulphate	mg/l	175.2	170.4	165.6	200.0	400
15	Nitrite	mg/l	<0.01	<0.01	<0.01	NS*	NS*
16	Nitrate	mg/l	9.50	10.70	9.50	45.0	No Relaxation
17	Salinity	%	1.26	0.90	1.06	NS*	NS*
18	Sodium as Na	mg/l	190	186	189	NS*	NS*
19	Potassium as K	mg/l	2.4	2.3	2.6	NS*	NS*
20	Manganese	mg/l	<0.04	<0.04	<0.04	0.1	0.3
21	Hexavalent Chromium	mg/l	<0.03	<0.03	<0.03	NS*	NS*
22	Copper	mg/l	<0.05	<0.05	<0.05	0.05	1.5
23	Cadmium	mg/l	<0.002	<0.002	<0.002	0.003	0.003
24	Arsenic	mg/l	<0.01	<0.01	<0.01	0.01	0.05
25	Mercury	mg/l	<0.001	<0.001	<0.001	0.001	0.001
26	Lead	mg/l	<0.01	<0.01	<0.01	0.01	0.01
27	Zinc	mg/l	<0.1	<0.1	<0.1	5.0	15.0
28	Bacterial Count	CFU/100ml	Absent	Absent	Absent	Absent	Absent

\*NS: Not Specified

**Table 14: Drinking Water Quality Monitoring Parameters for F - Type Quarter, Hospital Gopalpuri & Tuna Port**

Sr. No.	Parameter	Unit	F - Type Quarter	Hospital Gopalpuri	Tuna Port	Acceptable Limits as per IS 10500 : 2012	Permissible Limits in the absence of Alternate Source as per IS 10500 : 2012
1	pH	pH Unit	7.7	7.5	7.32	6.5 to 8.5	6.5 to 8.5
2	Total Dissolved Solids	mg/l	1050	1080	1020	500	2000
3	Turbidity	NTU	0	1	0	1.0	5.0
4	Odor	-	Odorless	Odorless	Odorless	Agreeable	Agreeable
5	Color	Hazen Unit	Colorless	Colorless	Colorless	5.0	15.0
6	Conductivity	µs/cm	1990	2150	2000	NS*	NS*
7	Biochemical Oxygen Demand	mg/l	<2	<2	<2	NS*	NS*
8	Chloride	mg/l	626	641	604	250.0	1000.0
9	Ca as Ca	mg/l	76.15	80.16	80.16	75.0	200.0
10	Mg as Mg	mg/l	51.03	60.75	60.75	30.0	100.0
11	Total Hardness	mg/l	430	360	330	200.0	600.0
12	Iron as Fe+3	mg/l	<0.01	<0.01	<0.01	0.3	No Relaxation
13	Fluorides	mg/l	1.05	0.92	0.46	1.0	1.5
14	Sulphate	mg/l	138	190.8	180	200.0	400
15	Nitrite	mg/l	<0.01	<0.01	<0.01	NS*	NS*
16	Nitrate	mg/l	10.00	7.96	8.2	45.0	No Relaxation
17	Salinity	%	1.13	1.16	1.09	NS*	NS*
18	Sodium as Na	mg/l	190	186	188	NS*	NS*
19	Potassium as K	mg/l	2.5	2.4	2.6	NS*	NS*
20	Manganese	mg/l	<0.04	<0.04	<0.04	0.1	0.3
21	Hexavalent Chromium	mg/l	<0.03	<0.03	<0.03	NS*	NS*
22	Copper	mg/l	<0.05	<0.05	<0.05	0.05	1.5
23	Cadmium	mg/l	<0.002	<0.002	<0.002	0.003	0.003
24	Arsenic	mg/l	<0.01	<0.01	<0.01	0.01	0.05
25	Mercury	mg/l	<0.001	<0.001	<0.001	0.001	0.001
26	Lead	mg/l	<0.01	<0.01	<0.01	0.01	0.01
27	Zinc	mg/l	<0.1	<0.1	<0.1	5.0	15.0
28	Bacterial Count	CFU/100ml	Absent	Absent	Absent	Absent	Absent

\*NS: Not Specified

**Table 15: Drinking Water Quality Monitoring Parameters for Vadinar Jetty & Port Colony at Vadinar**

Sr. No.	Parameter	Unit	Vadinar Jetty	Port Colony Vadinar	Acceptable Limits as per IS 10500 : 2012	Permissible Limits in the absence of Alternate Source as per IS 10500 : 2012
1	pH	pH Unit	7.6	7.4	6.5 to 8.5	6.5 to 8.5
2	Total Dissolved Solids	mg/l	1010	990	500	2000
3	Turbidity	NTU	ND	ND	1.0	5.0
4	Odor	-	Odorless	Odorless	Agreeable	Agreeable
5	Color	Hazen Units	Colorless	Colorless	5.0	15.0
6	Conductivity	µs/cm	210.0	990.0	NS*	NS*
7	Biochemical Oxygen Demand	mg/l	<2	<2	NS*	NS*
8	Chloride	mg/l	476	491	250.0	1000.0
9	Ca as Ca	mg/l	64.13	56.11	75.0	200.0
10	Mg as Mg	mg/l	75.33	68.04	30.0	100.0
11	Total Hardness	mg/l	470	420	200.0	600.0
12	Iron as Fe+3	mg/l	<0.01	<0.01	0.3	No Relaxation
13	Fluorides	mg/l	0.99	0.94	1.0	1.5
14	Sulphate	mg/l	16.80	17.64	200.0	400
15	Nitrite	mg/l	<0.01	<0.01	NS*	NS*
16	Nitrate	mg/l	11.48	9.50	45.0	No Relaxation
17	Salinity	%	0.86	0.89	NS*	NS*
18	Sodium as Na	mg/l	140.0	146.0	NS*	NS*
19	Potassium as K	mg/l	2.2	2.3	NS*	NS*
20	Manganese	mg/l	<0.04	<0.04	0.1	0.3
21	Hexavalent Chromium	mg/l	<0.03	<0.03	NS*	NS*
22	Copper	mg/l	<0.05	<0.05	0.05	1.5
23	Cadmium	mg/l	<0.002	<0.002	0.003	0.003
24	Arsenic	mg/l	<0.01	<0.01	0.01	0.05
25	Mercury	mg/l	<0.001	<0.001	0.001	0.001
26	Lead	mg/l	<0.01	<0.01	0.01	0.01
27	Zinc	mg/l	<0.1	<0.1	5.0	15.0
28	Bacterial Count	CFU/100ml	Absent	Absent	Absent	Absent

\*NS: Not Specified

### 2.3 Results & Discussion

The colour of all drinking water samples was < 5 Hazen unit and odour of the samples was also agreeable. All parameters are found to be within the specified limit of the Drinking water Standard.

#### pH

The limit of pH value for drinking water is specified as 6.5 to 8.5. pH value in the studied area varied from 7.0 to 8.0 pH unit. All the sampling points showed pH values within the prescribed limit by Indian Standards.

#### Total Dissolved Solids (TDS)

TDS values in the studied area varied between 600 -1800 mg/l. None of the sampling points showed higher TDS values than the prescribed limit by Indian standards .

#### Conductivity

Electrical Conductivity is the ability of a solution to transfer (conduct) electric current. Conductivity is used to measure the concentration of dissolved solids which have been ionized in a polar solution such as water. The conductivity in the samples collected during the month of August ranged from 1000-3300 µs/cm. Electrical conductivity standards do not appear in BIS standards for drinking water.

#### BOD

BOD value in the studied area was less than 2.0 mg/L. Indian standards does not show any standard values for BOD in drinking water.

#### Chlorides

Excessive chloride concentration increase rates of corrosion of metals in the distribution system. This can lead to increased concentration of metals in the supply. Chloride value in the studied area varied between 400-800 mg/l and is found to be within the Permissible limit of the Drinking Water Standard.

#### Calcium

Calcium value in the studied area varied between 45 - 80 mg/l and is found to be within the Permissible limit of the Drinking Water Standard. If calcium is present beyond the maximum acceptable limit, it causes incrustation of pipes.

### **Magnesium**

Magnesium value in the studied area varied between 30 – 85 mg/l. All the locations had Magnesium within the prescribed limits of 30-100 mg/L.

### **Total Hardness**

Hardness value in the studied area varied between 330-470 mg/l and is found to be within the Permissible limit of the Drinking Water Standard. The prescribed limit by Indian Standards is 200-600 mg/L.

### **Iron**

Iron value in the studied area was below 0.01mg/L and hence well below the permissible limit as per Indian Standards is 0.3 mg/L. The excess amount of iron causes slight toxicity; gives stringent taste to water.

### **Fluoride**

Fluoride value in the studied area varied between 0.1 – 1.0 mg/l and hence well below the permissible limit as per Indian Standards is 1.0-1.5 mg/L. Moderate amounts lead to dental effects, but long-term ingestion of large amounts can lead to potentially severe skeletal problems.

### **Sulphates**

Sulphate value in the studied area varied between 10 – 300 mg/l. All the sampling points showed sulphate values within the prescribed limits by Indian Standards (200-400 mg/L). Sulphate content in drinking water exceeding the 400 mg/L imparts bitter taste.

### **Nitrites (NO<sub>2</sub>) and Nitrates (NO<sub>3</sub>)**

Nitrite values in all the water samples were <0.1. There are no specified standard values for Nitrites in Drinking water. The mean Nitrate values in drinking water of KPT was 6.27 mg/l which is well within the permissible limit of the Drinking water Standard.

### **Salinity**

Salinity in drinking water in the present samples collected ranged from 0.8 to 1.3 %. There are no prescribed Indian standards for salinity in Drinking water.



### **Sodium and Potassium Salts**

Sodium values in the samples collected ranged from 100 - 2000 mg/l and Potassium salts ranged from 2.2 to 3.0 mg/l. There are no prescribed limits of Sodium and Potassium in Indian standards for Drinking water.

### **Heavy Metals in Drinking Water**

In the present study period drinking water samples were analyzed for Mn, Cr, Cu, Cd, As, Hg, Pb and Zn. All these heavy metals were well below the permissible limits prescribed by the Indian Standards.

### **Bacteriological Study**

Analysis of the bacteriological parameter at all location shows that Bacteria is not present and hence Bacterial count is in line with the permissible limit of drinking water. This shows that all the drinking water samples were safe from any bacteriological contamination.

## **2.4 Conclusions**

These results are compared with acceptable limits as prescribed in IS 10500:2012 – Drinking Water Specification. It is seen from the analysis data that during the study period the water was safe for human consumption at all drinking water monitoring stations.

### 3. Noise Level Monitoring

Noise sources in port operations include cargo handling, vehicular traffic, and loading / unloading containers and ships. Noise Monitoring was done at 13 stations at Kandla, Vadinar and Township area.

#### 3.1 Method of Monitoring

Sampling was done at all stations for 24 hour period. Data was recorded using automated sound level meter. The intensity of sound was measured in sound pressure level (SPL) and common unit of measurement is decibel (Db).

#### 3.2 Results

Table 16: Noise Monitoring data for ten locations of Deendayal Port and two locations of Vadinar Port

Sr. No.	Location	Day Time Average Noise Level (SPL) in dB(A)	Night Time Average Noise Level (SPL) in dB(A)
	<b>Sampling Time</b>	<b>6:00 am to 10:00 PM</b>	<b>10:00PM to 6:00 AM</b>
1	Marine Bhavan	52.0	48.9
2	Nirman Building 1	52.7	46.9
3	Tuna Port	56.2	49.8
4	Main Gate North	66.8	60.7
5	West Gate I	70.4	63.0
6	Canteen Area	54.8	44.7
7	Main Road	65.9	51.1
8	ATM Building	66.4	56.6
9	Wharf Area /Jetty Area	72.2	67.7
10	Port & Custom Office	51.5	46.3
	<b>Vadinar Port</b>		
11	Entrance Gate of Vadinar Port	66.8	53.7
12	Nr. Port Colony, Vadinar	60.4	52.8
13	Nr. Vadinar Jetty	72.5	63.7

**3.3 Conclusions-** Noise sources in port operations include cargo handling, vehicular traffic, and loading / unloading containers and ships. The Day Time Average Noise Level (SPL) in all ten locations at Deendayal Port ranged from 52.0 dB(A) to 72.2 dB(A) and it was within the permissible limits of 75 dB(A) for the industrial area for the daytime. The Night Time Average Noise Level (SPL) in all ten locations of Deendayal Port ranged from 44.7 dB to 67.7 dB(A) and it was within the permissible limits of 70 dB(A) for the industrial area for the night time.

#### 4. Soil Monitoring

Sampling and analysis of soil samples were undertaken at six locations within the study area (Deendayal Port and Vadinar Port) as a part of EMP. The soil sampling locations are initially decided based on the locations as provided in the tender document of the Deendayal Port.

##### 4.1 Methodology

The soil samples were collected in the month of August 2021. The samples collected from the all locations are homogeneous representative of each location. At random locations were identified at each location and soil was dug from 30 cm below the surface. It was uniformly mixed before homogenizing the soil samples. The samples were filled in polythene bags, labeled in the field with number and site name and sent to laboratory for analysis.

##### 4.2 Results

**Table-17: Chemical Characteristics of Soil in the Study Area**

Sr. No.	Parameter	Unit	Station Name					
			SL1	SL2	SL3	SL4	SL5	SL6
			Tuna Port	IFFCO Plant	Khori Creek	Nakti Creek	KPT Admin Site	KPT Colony
			Near main gate of Port	10 m away from main gate	Sand from creek at low tide		Vadinar	
1	Texture		Sandy Loam	Sandy Loam	Sandy Loam	Sandy Loam	Sandy Loam	Sandy Loam
2	pH	-	8.60	8.10	8.42	8.30	8.09	8.32
3	Electrical Conductivity	µs/cm	23,400.0	20,420.0	23,700.0	17,200.0	510.0	400.0
4	Moisture	%	20.42	21.16	23.22	20.12	9.04	8.22
5	Total Organic Carbon	%	0.18	0.18	0.25	0.11	0.21	0.16
6	Alkalinity	mg/kg	60.06	140.04	140.04	60.06	100.10	80.04
7	Total Nitrogen	%	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001
8	Chloride	mg/kg	4,010.0	4,324.0	5,982.0	4,001.0	42.2	67.8
9	Sulphate	mg/kg	188.0	179.2	110.0	100.0	14.0	16.2
10	Phosphorus	mg/kg	0.90	0.86	1.04	1.62	0.78	0.88
11	Potassium	mg/kg	786.0	656.0	1,162.0	780.0	130.0	182.0
12	Sodium	mg/kg	2,341.0	3,618.0	4,220.0	3,122.0	1,224.0	1,400.0
13	Calcium	mg/kg	160.00	130.00	170.00	220.00	110.00	68.00
14	Copper as Cu	mg/kg	32.2	58.2	42.2	23.4	17.4	23
15	Lead as Pb	mg/kg	3.8	3.8	3.6	4.1	BQL	BQL
16	Nickel as Ni	mg/kg	37.2	32.4	41.2	24.5	19.3	20.4
17	Zinc as Zn	mg/kg	59.36	38.32	53.4	48.50	49.20	40.40
18	Cadmium as Cd	mg/kg	ND	ND	ND	ND	ND	ND

### 4.3 Discussion

- The data shows that value of pH ranges from 8.42 at Nakti Creek to 8.60 at Tuna Creek indicating that all soil samples are neutral to slight basic. Tuna port samples showed maximum conductivity of 23,400µmhos/cm, while Nakti Creek location showed minimum conductivity of 17,200 µmhos/cm. Conductivity at Vadinar Port was 510 and 400 µmhos/cm at Admin site and Vadinar Port colony respectively.
- Total organic Carbon ranged from 0.1 % to 0.3 at Deendayal Port. At Vadinar Port, organic carbon content ranged from 0.1 % to 0.2 %.
- The concentration of Phosphorus and Potassium in the soil samples varies from 0.8 to 1.62 mg/kg and 600.0 to 1170 mg/kg respectively at Deendayal Port. The mean concentration of Phosphorous at Vadinar site was 0.80 mg/kg and mean concentration of Potassium at Vadinar site was 156 mg/kg.

These differences in NPK in soil at different locations are due to the dissimilar nature of soil at each of the locations. Samples SL3 & SL4 (Khorī Creek & Nakti Creek) are of saline nature as they are coastal soil; where as other locations are inland locations and have different chemical properties.

### Heavy Metals in the Soil

Traces of Copper, Lead, Nickel and Zinc were observed in the soil samples collected from all the four locations of Deendayal Port and two locations of Vadinar Port. Cadmium metal was not detected in the Soil.

### 4.4 Conclusion

The soils of Deendayal Port and Vadinar Port appears to be neutral to basic with varying levels of Chloride, Sulphate, NPK and Calcium. As the nature of soil at different locations are different with respect to its proximity to the sea, the samples showed high degree of variations in their chemical properties.

## 5. Sewage Treatment Plant Monitoring

This involves safe collection of waste water (spent/used water) from wash areas, bathroom, industrial units, etc., waste from toilets of various buildings and its conveyance to the treatment plant and final disposal in conformity with the requirement and guide lines of State Pollution Control Board and other statutory bodies.

### 5.1 Methodology for STP Monitoring

To monitor the working efficiency of Sewage Treatment Plant (STP), STP Inlet and Outlet Samples were collected once a week. Locations selected are namely Gopalpuri Township, Deendayal Port and Vadinar. Samples were collected in 1 lit. Carboys and were analyzed in laboratory for various parameters.

### 5.2 Results

- **Kandla STP**

**Table 18: Sewage Water Monitoring at Kandla STP (1<sup>st</sup> Week)**

Date of Sampling		05.08.2021		
Sr. No.	Parameters	Unit	Results	
			KPT STP I/L	KPT STP O/L
1	pH	pH unit	7.9	7.5
2	Total Suspended Solids	mg/l	107	101
3	Residual Chlorine	mg/l	<1.0	<0.5
4	COD	mg/l	334	108
5	BOD @ 27 °C	mg/l	118.0	27.0
<b>Aeration Tank</b>				
6	MLSS	mg/l	12.0	
7	MLVSS	%	88.0	

**Table 19: Sewage Water Monitoring at Kandla STP (2<sup>nd</sup> Week)**

Date of Sampling		12.08.2021		
Sr. No.	Parameters	Unit	Results	
			KPT STP I/L	KPT STP O/L
1	pH	pH unit	7.6	7.68
2	Total Suspended Solids	mg/l	193	101
3	Residual Chlorine	mg/l	<1.0	<0.5
4	COD	mg/l	414	104
5	BOD @ 27 °C	mg/l	136.0	27.0
<b>Aeration Tank</b>				
6	MLSS	mg/l	9.0	
7	MLVSS	%	97.0	

**Table 20: Sewage Water Monitoring at Kandla STP (3<sup>rd</sup> Week)**

Date of Sampling		19.08.2021		
Sr. No.	Parameters	Unit	Results	
			KPT STP I/L	KPT STP O/L
1	pH	pH unit	7.86	7.48
2	Total Suspended Solids	mg/l	204	104
3	Residual Chlorine	mg/l	<1.0	<0.5
4	COD	mg/l	261	70
5	BOD @ 27 °C	mg/l	87.0	20.0
<b>Aeration Tank</b>				
6	MLSS	mg/l	10.0	
7	MLVSS	%	90.0	

**Table 21: Sewage Water Monitoring at Kandla STP (4<sup>th</sup> Week)**

Date of Sampling		23.08.2021		
Sr. No.	Parameters	Unit	Results	
			KPT STP I/L	KPT STP O/L
1	pH	pH unit	7.43	7.16
2	Total Suspended Solids	mg/l	403.3	150.4
3	Residual Chlorine	mg/l	<1.0	<1.0
4	COD	mg/l	313.1	151.5
5	BOD @ 27 °C	mg/l	106.0	52.0
6.	Fecal Coliform	MPN Index / 100 ml	-	>1600
<b>Aeration Tank</b>				
7.	MLSS	mg/l	33.0	
8	MLVSS	%	81.0	

- **Gopalpuri Colony STP**

**Table 22: Sewage Water Monitoring at Gopalpuri STP (1<sup>st</sup> Week)**

Date of Sampling		05.08.2021		
Sr. No.	Parameters	Unit	Results	
			Gopalpuri STP I/L	Gopalpuri STP O/L
1	pH	pH unit	7.76	7.34
2	Total Suspended Solids	mg/l	98.1	62.4
3	Residual Chlorine	mg/l	<1.0	<0.5
4	COD	mg/l	324	102
5	BOD @ 27 °C	mg/l	110.0	28.0
<b>Aeration Tank</b>				
6	MLSS	mg/l	12.0	
7	MLVSS	%	92.0	

**Table 23: Sewage Water Monitoring at Gopalpuri STP (2<sup>nd</sup> Week)**

Date of Sampling		12.08.2021		
Sr. No.	Parameters	Unit	Results	
			Gopalpuri STP I/L	Gopalpuri STP O/L
1	pH	pH unit	7.72	7.37
2	Total Suspended Solids	mg/l	406	107
3	Residual Chlorine	mg/l	<1.0	<0.5
4	COD	mg/l	320	103
5	BOD @ 27 °C	mg/l	110.0	26.0
<b>Aeration Tank</b>				
6	MLSS	mg/l	14.0	
7	MLVSS	%	90.0	

**Table 24: Sewage Water Monitoring at Gopalpuri STP (3<sup>rd</sup> Week)**

Date of Sampling		19.08.2021		
Sr. No.	Parameters	Unit	Results	
			Gopalpuri STP I/L	Gopalpuri STP O/L
1	pH	pH unit	7.71	7.34
2	Total Suspended Solids	mg/l	404	107
3	Residual Chlorine	mg/l	<1.0	<0.5
4	COD	mg/l	351	101
5	BOD @ 27 °C	mg/l	115.0	23.0
<b>Aeration Tank</b>				
6	MLSS	mg/l	16.0	
7	MLVSS	%	88.0	



**Table 25: Sewage Water Monitoring at Gopalpuri STP (4<sup>th</sup> Week)**

Date of Sampling		23.08.2021		
Sr. No.	Parameters	Unit	Results	
			Gopalpuri STP I/L	Gopalpuri STP O/L
1	pH	pH unit	7.7	7.35
2	Total Suspended Solids	mg/l	405	107
3	Residual Chlorine	mg/l	<1.0	<0.5
4	COD	mg/l	242	101
5	BOD @ 27 °C	mg/l	80.0	23.0
6.	Fecal Coliform	MPN Index / 100 ml	-	>1600
<b>Aeration Tank</b>				
7.	MLSS	mg/l	18.0	
8.	MLVSS	%	88.0	

- **Vadinar STP**

**Table 26: Sewage Water Monitoring at Vadinar STP (1<sup>st</sup> Week)**

Date of Sampling		05.08.2021		
Sr. No.	Parameters	Unit	Results	
			Vadinar STP I/L	Vadinar O/L
1	pH	pH unit	7.23	NOT WORKING
2	Total Suspended Solids	mg/l	18	
3	Residual Chlorine	mg/l	<1.0	
4	COD	mg/l	89.0	
5	BOD @ 27 °C	mg/l	28.0	

**Table 27: Sewage Water Monitoring at Vadinar STP (2<sup>nd</sup> Week)**

Date of Sampling		12.08.2021		
Sr. No.	Parameters	Unit	Results	
			Vadinar STP I/L	Vadinar O/L
1	pH	pH unit	7.28	NOT WORKING
2	Total Suspended Solids	mg/l	60	
3	Residual Chlorine	mg/l	<1.0	
4	COD	mg/l	78.0	
5	BOD @ 27 °C	mg/l	28.0	

**Table 28: Sewage Water Monitoring at Vadinar STP (3<sup>rd</sup> Week)**

Date of Sampling		19.08.2021		
Sr. No.	Parameters	Unit	Results	
			Vadinar STP I/L	Vadinar O/L
1	pH	pH unit	7.32	NOT WORKING
2	Total Suspended Solids	mg/l	60	
3	Residual Chlorine	mg/l	<1.0	
4	COD	mg/l	80.0	
5	BOD @ 27 °C	mg/l	26.0	

**Table 29: Sewage Water Monitoring at Vadinar STP (4<sup>th</sup> Week)**

<b>Date of Sampling</b>	<b>23.08.2021</b>
-------------------------	-------------------

Sr. No.	Parameters	Unit	Results	
			Vadinar STP I/L	Vadinar O/L
1	pH	pH unit	7.18	NOT WORKING
2	Total Suspended Solids	mg/l	72	
3	Residual Chlorine	mg/l	<1.0	
4	COD	mg/l	80.0	
5	BOD @ 27 °C	mg/l	26.0	

**5.3 Conclusions:**

The GPCB standards of BOD, TSS and Residual Chlorine for STP outlet are 20 mg/lit, 30 mg/lit & 0.5 mg/lit respectively. It is suggested to do treatment on regular basis to avoid flow of contaminated/polluted water into the sea. Also, the STP at Vadinar is also non-functional and thus, steps should be taken to commission the STP at Vadinar Port. Hence, currently only inlet samples are collected and analysed.

## **6. Marine Water Monitoring**

The Forty Second Amendment to the Constitution in 1976 underscored the importance of 'green thinking'. Article 48A enjoins the state to protect and improve the environment and safeguard the forests and wildlife in the country. Further, Article 51A(g) states that the "fundamental duty of every citizen is to protect and improve the natural environment including forests, lakes, rivers and wildlife and to have compassion for living creatures".

Policy Statement for Abatement of Pollution (1992) has suggested developing relevant legislation and regulation, fiscal incentives, voluntary agreements and educational programs and information campaigns. It emphasizes the need for integration by incorporating environmental considerations into decision making at all levels by adopting frameworks namely, pollution prevention at source, application of best practicable solution, ensure polluter pays for control of pollution, focus on heavily polluted areas and river stretches and involve public in decision-making. The National Conservation Strategy and Policy Statement on Environment and Development, (1992) aimed at "integrating environmental concerns with developmental imperatives to meet the challenges by redirecting the thrust of our developmental process so that the basic needs of our people could be fulfilled by making judicious and sustainable use of natural resources." The priorities mentioned in this policy document include the sustainable use of land and water resources, prevention and control of pollution and preservation of biodiversity.

The National Water Policy, (2002) contains provisions for developing, conserving, sustainable utilizing and managing this important water resources and need to be governed by national perspectives.

### **Marine Environment**

On national and state levels, we have several policies and regulation like Water (Prevention and Control of Pollution) Act, 1974, to regulate pollution discharges and restore water quality of our aquatic resources including the prescription of monitoring activities. One of the important provisions of the Water Act, 1974, is to maintain and restore the 'wholesomeness' of our aquatic resources. Water quality monitoring is one of the first steps required in the rational development and management of water resources. In the field of water quality management, there has been a steady evolution in procedures for designing system to obtain information on the changes of water quality. The monitoring comprises all activities to obtain 'information' with respect to the water system.

## Sampling Stations

The monitoring of marine environment for the study of biological and ecological parameters was carried out on 9<sup>th</sup>& 10<sup>th</sup> August-2021 in harbor regions of KPT and on 9<sup>th</sup> August-2021 at Vadinar during spring tide period of New moon phase of Lunar Cycle. The monitoring of marine environment for the study of biological and ecological parameters was repeated again on 16<sup>th</sup>& 17<sup>th</sup> August 2021 in harbor regions of KPT. 16<sup>th</sup> August -2021 in Vadinar during Neap tide period first quarter of Lunar Cycle..

Plankton samples from sub surface layer was collected both during high tide period and low tide period from 3 water quality monitoring stations of KPT harbour area and two stations in Nakti creek and one station in Khori creek. The same sampling schedule was repeated during consecutive spring tide and neap tide in same month. Plankton samples from sub surface layer was collected both during high tide period and low tide period from 1 water quality monitoring stations near Vadinar jetty area during spring tide and neap tide in this month .Collected water samples were processed for estimation of Chlorophyll- a, Pheophytin- a, qualitative & quantitative evaluation of phytoplankton, qualitative & quantitative evaluation zooplanktons (density and their population).

### Sampling Locations

Offshore monitoring requirement	Number of locations
Offshore Installations	3 in Kandla creek 2 in Nakti creek 1 in Khori creek 1 near Vadinar Jetty 1 near 1 <sup>st</sup> SBM
<b>Total Number of locations</b>	<b>8</b>

#### 6.1 Marine Water Quality

Marine water quality of marine waters of Deendayal Port Harbor waters, Khori and Nakti Creeks and two locations of Vadinar are monitored for various physico-chemical parameters during spring and neap tide of each month.

The results of marine water quality and Marine sediments are as below;

**Table 30: Marine Water Quality Monitoring Parameters for location near KPT colony**

Sr. No.	Parameters	Unit	Kandla Creek Near KPT colony (1)			
			23°0'58"N 70°13'22."E			
			Spring Tide		Neap Tide	
Tide →	High Tide	Low Tide	High Tide	Low Tide		
1	pH	pH unit	7.16	7.14	7.3	7.26
2	Color	-	Colorless	Colorless	Colorless	Colorless
3	Odor	-	Odorless	Odorless	Odorless	Odorless
4	Salinity	ppt	32.2	32.0	32.0	31.8
5	Turbidity	NTU	39	28	32	29
6	Total Dissolved Solids	mg/l	42660	41056	37802.0	43665.0
7	Total Suspended Solids	mg/l	675	979	614.2	372.4
8	Total Solids	mg/l	46346	44350	46346.0	44369.4
9	DO	mg/l	4.5	3.9	4.6	5.1
10	COD	mg/l	80.0	78.0	78.0	80.0
11	BOD	mg/l	<2.0	<2.0	<2.0	<2.0
12	Silica	mg/l	7.82	6.09	0.49	0.73
13	Phosphate	mg/l	0.57	0.14	0.16	0.17
14	Sulphate	mg/l	2628	1656	2352	2076
15	Nitrate	mg/l	2.22	2.03	2.53	3.77
16	Nitrite	mg/l	<0.05	<0.05	<0.05	<0.05
17	Calcium	mg/l	1322.64	1242.48	601.2	480.96
18	Magnesium	mg/l	1239.3	1336.5	1749.6	1773.9
19	Sodium	mg/l	11012.0	10828.0	11022.0	10202.0
20	Potassium	mg/l	340.0	300.0	320.0	302.0
21	Iron	mg/l	1.32	1.40	1.20	1.30
22	Chromium	mg/l	0.16	0.14	0.12	0.11
23	Copper	mg/l	0.06	0.07	0.14	0.18
24	Arsenic	mg/l	<0.01	<0.01	<0.01	<0.01
25	Cadmium	mg/l	0.05	0.06	0.07	0.06
26	Mercury	mg/l	<0.001	<0.001	<0.001	<0.001
27	Lead	mg/l	0.11	0.12	0.18	0.19
28	Zinc	mg/l	0.08	0.06	0.07	0.06

**Table 31: Marine Water Quality Monitoring Parameters for location near passenger Jetty One at Kandla**

Sr. No.	Parameters	Unit	Near passenger Jetty One (2)			
			23° 0'18 "N 70°13'31"E			
			Spring Tide		Neap Tide	
			Tide →	High Tide	Low Tide	High Tide
1	pH	pH unit	7.28	7.25	7.39	7.42
2	Color	-	Colorless	Colorless	Colorless	Colorless
3	Odor	-	Odorless	Odorless	Odorless	Odorless
4	Salinity	ppt	31.7	32.6	32.0	32.2
5	Turbidity	NTU	37	27	42	47
6	Total Dissolved Solids	mg/l	41612	45181	41735.0	36900.0
7	Total Suspended Solids	mg/l	717	808	414	432.9
8	Total Solids	mg/l	47224	44028	37224.0	44028.0
9	DO	mg/l	4.4	4.1	5.4	4.8
10	COD	mg/l	90.0	86.0	86.0	82.0
11	BOD	mg/l	<2.0	<2.0	<2.0	<2.0
12	Silica	mg/l	9.44	7.84	0.53	0.64
13	Phosphate	mg/l	0.06	0.11	0.18	0.19
14	Sulphate	mg/l	2760	1572	2652	2616
15	Nitrate	mg/l	2.36	2.25	3.45	4.29
16	Nitrite	mg/l	<0.05	<0.05	<0.05	<0.05
17	Calcium	mg/l	1202.40	1122.24	561.12	480.96
18	Magnesium	mg/l	1336.5	1385.1	1798.2	1749.6
19	Sodium	mg/l	11752.0	10652.0	11120.0	12120.0
20	Potassium	mg/l	306.0	290.0	289.0	322.0
21	Iron	mg/l	1.56	1.66	1.50	1.40
22	Chromium	mg/l	0.13	0.12	0.10	0.12
23	Copper	mg/l	0.08	0.09	0.15	0.16
24	Arsenic	mg/l	<0.01	<0.01	<0.01	<0.01
25	Cadmium	mg/l	0.06	0.07	0.07	0.08
26	Mercury	mg/l	<0.001	<0.001	<0.001	<0.001
27	Lead	mg/l	0.10	0.09	0.18	0.17
28	Zinc	mg/l	0.07	0.06	0.08	0.06

**Table 32: Marine Water Quality Monitoring Parameters for location Near Coal Berth**

Sr. No.	Parameters	Unit	Near Coal Berth			
			22°59'12"N 70°13'40"E			
			Spring Tide		Neap Tide	
			Tide →	High Tide	Low Tide	High Tide
1	pH	pH unit	7.30	7.51	7.53	7.32
2	Color	-	Colorless	Colorless	Colorless	Colorless
3	Odor	-	Odorless	Odorless	Odorless	Odorless
4	Salinity	ppt	32.0	32.5	32.0	31.8
5	Turbidity	NTU	33	25	37	45
6	Total Dissolved Solids	mg/l	48590	39430	45812.0	35363.0
7	Total Suspended Solids	mg/l	555	809	587.3	591.2
8	Total Solids	mg/l	45108	41100	41720.0	40200.0
9	DO	mg/l	3.8	4	4.9	5.1
10	COD	mg/l	88.0	90.0	90.0	82.0
11	BOD	mg/l	<2.0	<2.0	<2.0	<2.0
12	Silica	mg/l	5.98	3.98	0.42	0.85
13	Phosphate	mg/l	0.10	0.08	0.15	0.19
14	Sulphate	mg/l	2856	2988	2736	2208
15	Nitrate	mg/l	2.73	2.33	4.75	3.79
16	Nitrite	mg/l	<0.05	<0.05	<0.05	<0.05
17	Calcium	mg/l	1362.72	1322.64	480.96	601.2
18	Magnesium	mg/l	1190.7	1239.3	1822.5	1846.8
19	Sodium	mg/l	11452.0	10890.0	11125.0	10890.0
20	Potassium	mg/l	311.0	269.0	345.0	400.0
21	Iron	mg/l	1.80	1.92	1.30	2.01
22	Chromium	mg/l	0.11	0.12	0.18	0.19
23	Copper	mg/l	0.07	0.06	0.18	0.16
24	Arsenic	mg/l	<0.01	<0.01	<0.01	<0.01
25	Cadmium	mg/l	0.05	0.06	0.06	0.07
26	Mercury	mg/l	<0.001	<0.001	<0.001	<0.001
27	Lead	mg/l	0.10	0.11	0.20	0.19
28	Zinc	mg/l	0.08	0.06	0.07	0.06



**Table 33: Marine Water Quality Monitoring Parameters for location Khori creek at Kandla**

Sr. No.	Parameters	Unit	KPT 4			
			Near 15/16 Berth			
			Spring Tide		Neap Tide	
			Tide →	High Tide	Low Tide	High Tide
1	pH	pH unit	7.37	7.42	7.26	7.22
2	Color	-	Colorless	Colorless	Colorless	Colorless
3	Odor	-	Odorless	Odorless	Odorless	Odorless
4	Salinity	ppt	32.6	31.6	31.8	31.6
5	Turbidity	NTU	45	37	52	28
6	Total Dissolved Solids	mg/l	42420	38440	33550.0	33133.0
7	Total Suspended Solids	mg/l	654	624	701.5	490.4
8	Total Solids	mg/l	44940	40080	44940.0	40080.0
9	DO	mg/l	4.4	4.3	5.3	5.9
10	COD	mg/l	92.0	88.0	88.0	92.0
11	BOD	mg/l	<2.0	<2.0	<2.0	<2.0
12	Silica	mg/l	2.45	7.45	0.69	0.51
13	Phosphate	mg/l	0.10	0.02	0.24	0.16
14	Sulphate	mg/l	1668	2268	2616	2580
15	Nitrate	mg/l	1.96	1.53	3.34	4.86
16	Nitrite	mg/l	<0.05	<0.05	<0.05	<0.05
17	Calcium	mg/l	1242.48	1282.56	521.04	480.96
18	Magnesium	mg/l	1287.9	1336.5	1725.3	1725.3
19	Sodium	mg/l	12152.0	13020.0	12162.0	11782.0
20	Potassium	mg/l	288.0	316.0	389.0	380.0
21	Iron	mg/l	1.60	1.55	1.48	1.38
22	Chromium	mg/l	0.15	0.16	0.20	0.18
23	Copper	mg/l	0.08	0.10	0.15	0.11
24	Arsenic	mg/l	<0.01	<0.01	<0.01	<0.01
25	Cadmium	mg/l	0.08	0.06	0.08	0.07
26	Mercury	mg/l	<0.001	<0.001	<0.001	<0.001
27	Lead	mg/l	0.09	0.11	0.18	0.17
28	Zinc	mg/l	0.07	0.05	0.08	0.06

**Table 34: Marine Water Quality Monitoring Parameters for location Nakti Creek near Tuna Port**

Sr. No.	Parameters	Unit	Nakti Creek Near Tuna Port			
			22°57'49."N 70° 7'0.67"E			
			Spring Tide		Neap Tide	
			Tide →	High Tide	Low Tide	High Tide
1	pH	pH unit	7.18	7.30	7.3	7.37
2	Color	-	Colorless	Colorless	Colorless	Colorless
3	Odor	-	Odorless	Odorless	Odorless	Odorless
4	Salinity	ppt	31.6	33.0	31.8	31.6
5	Turbidity	NTU	36	48	36	28
6	Total Dissolved Solids	mg/l	47540	37880	38200.0	37205.0
7	Total Suspended Solids	mg/l	885	852	332.5	474
8	Total Solids	mg/l	46280	38780	38280.0	49040.0
9	DO	mg/l	4.2	4.3	5.3	5.2
10	COD	mg/l	76.0	78.0	90.0	92.0
11	BOD	mg/l	<2.0	<2.0	<2.0	<2.0
12	Silica	mg/l	4.33	2.45	0.91	0.73
13	Phosphate	mg/l	0.08	0.10	0.18	0.18
14	Sulphate	mg/l	2052	4500	2628	2268
15	Nitrate	mg/l	2.17	2.47	5.14	5.70
16	Nitrite	mg/l	<0.05	<0.05	<0.05	<0.05
17	Calcium	mg/l	1362.72	1282.56	561.12	561.12
18	Magnesium	mg/l	1215	1239.3	1773.9	1773.9
19	Sodium	mg/l	11582.0	11262.0	10589.0	10110.0
20	Potassium	mg/l	326.0	366.0	347.0	311.0
21	Iron	mg/l	2.02	2.00	1.60	1.58
22	Chromium	mg/l	0.20	0.19	0.16	0.15
23	Copper	mg/l	0.10	0.08	0.12	0.10
24	Arsenic	mg/l	<0.01	<0.01	<0.01	<0.01
25	Cadmium	mg/l	0.06	0.07	0.08	0.06
26	Mercury	mg/l	<0.001	<0.001	<0.001	<0.001
27	Lead	mg/l	0.11	0.12	0.32	0.62
28	Zinc	mg/l	0.06	0.07	0.07	0.06

**Table 35: Marine Water Quality Monitoring Parameters for location Nakti Creek Near NH-8A at Kandla**

Sr. No.	Parameters	Unit	Nakti Creek Near NH-8A			
			23° 02'01"N 70° 09'31"E			
			Spring Tide		Neap Tide	
			Tide →	High Tide	Low Tide	High Tide
1	pH	pH unit	7.39	Sampling not possible during Low Tide	7.39	Sampling not possible during Low Tide
2	Color	-	Colorless		Colorless	
3	Odor	-	Odorless		Odorless	
4	Salinity	ppt	32.9		31.8	
5	Turbidity	NTU	36		35	
6	Total Dissolved Solids	mg/l	36020		35465.0	
7	Total Suspended Solids	mg/l	666		380.3	
8	Total Solids	mg/l	44660		46002.0	
9	DO	mg/l	4.7		5.5	
10	COD	mg/l	80.0		88.0	
11	BOD	mg/l	<2.0		<2.0	
12	Silica	mg/l	7.73		0.45	
13	Phosphate	mg/l	0.08		0.17	
14	Sulphate	mg/l	3660		2280	
15	Nitrate	mg/l	2.74		4.15	
16	Nitrite	mg/l	<0.05		<0.05	
17	Calcium	mg/l	1402.80		561.12	
18	Magnesium	mg/l	1190.7		1773.9	
19	Sodium	mg/l	13030.0		11120.0	
20	Potassium	mg/l	348.0		320.0	
21	Iron	mg/l	1.89		1.50	
22	Chromium	mg/l	0.17		0.17	
23	Copper	mg/l	0.09		0.11	
24	Arsenic	mg/l	<0.01		<0.01	
25	Cadmium	mg/l	0.08		0.07	
26	Mercury	mg/l	<0.001		<0.001	
27	Lead	mg/l	0.09		0.2	
28	Zinc	mg/l	0.08		0.08	

**Table 36: Marine Water Quality Monitoring Parameters for locations Nr. Vadinar Jetty**

Sr. No.	Parameters	Unit	Nr.Vadinar Jetty			
			22°26'25.26"N 69°40'20.41"E			
			Spring Tide		Neap Tide	
			Tide →		High Tide	Low Tide
1	pH	pH unit	7.60	7.90	7.38	7.25
2	Color	-	Colorless	Colorless	Colorless	Colorless
3	Odor	-	Odorless	Odorless	Odorless	Odorless
4	Salinity	ppt	33.0	32.5	32.0	32.0
5	Turbidity	NTU	48	37	47	40
6	Total Dissolved Solids	mg/l	38810	36220	37902.0	35080.0
7	Total Suspended Solids	mg/l	405	380	456.9	395.5
8	Total Solids	mg/l	42180	42020	38990.0	38620.0
9	DO	mg/l	4.3	4.7	4.5	4.9
10	COD	mg/l	90.0	88.0	82.0	78.0
11	BOD	mg/l	<2.0	<2.0	<2.0	<2.0
12	Silica	mg/l	6.00	7.65	0.76	0.93
13	Phosphate	mg/l	0.56	0.68	0.20	0.17
14	Sulphate	mg/l	2628	2268	2520	2376
15	Nitrate	mg/l	2.05	2.15	3.03	3.04
16	Nitrite	mg/l	<0.05	<0.05	<0.05	<0.05
17	Calcium	mg/l	1242.48	1362.72	641.28	521.04
18	Magnesium	mg/l	1239.3	1239.3	1798.2	1798.2
19	Sodium	mg/l	14025.0	13879.0	11012.0	11212.0
20	Potassium	mg/l	326.0	300.0	342.0	333.0
21	Iron	mg/l	1.88	1.79	1.60	1.30
22	Chromium	mg/l	0.18	0.18	0.18	0.12
23	Copper	mg/l	0.08	0.08	0.18	0.20
24	Arsenic	mg/l	<0.01	<0.01	<0.01	<0.01
25	Cadmium	mg/l	0.06	0.07	0.06	0.07
26	Mercury	mg/l	<0.001	<0.001	<0.001	<0.001
27	Lead	mg/l	0.10	0.09	0.16	0.2
28	Zinc	mg/l	0.06	0.06	0.06	0.07

### 6.1.1 Marine Sediments

Sediment samples were collected with Van Veen Grab from the six locations in Kandla Port Waters and two locations in Vadinar Port. Samples were collected and preserved in silver foil in ice box to prevent the contamination/decaying of the samples.

## 6.2 Results

The Sediment Quality results are given in below from table no. 34 A & B.

**Table 34A: Results of Analysis of Sediment of Kandla & Vadinar Port (Spring Tide)**

Sr. No.	Parameters	Unit	KPT - 1	KPT - 2	KPT - 3	KPT - 5	Jetty
1	Texture	-	Sandy Loam	Sandy Loam	Sandy Loam	Sandy Loam	Sandy Loam
2	Organic Matter	mg/kg	1.20	1.08	1.20	1.86	1.46
3	Organic Carbon	mg/kg	0.70	0.96	0.87	0.65	0.68
4	Inorganic Phosphate	mg/kg	120.0	132.0	142.0	162.0	160.0
5	Moisture	%	20.20	23.10	21.88	21.2	23.80
6	Aluminium	mg/kg	ND	ND	ND	ND	ND
7	Silica	mg/kg	28.0	21.0	24.0	36.0	23.0
8	Phosphate	mg/kg	10.50	11.20	9.80	9.60	10.20
9	Sulphate	mg/kg	210.0	242.0	160.0	170.0	140.0
10	Nitrite	mg/kg	0.11	0.12	0.1	0.11	0.12
11	Nitrate	mg/kg	9.80	7.44	10.80	9.20	8.40
12	Calcium	mg/kg	342.0	270.0	325.0	309.0	322.0
13	Magnesium	mg/kg	186.0	145.0	178.0	152.0	202.0
14	Sodium	mg/kg	8824.0	7242.0	9452.0	7122.0	8777.0
15	Potassium	mg/kg	396.0	388.0	460.0	680.0	780.0
16	Chromium	mg/kg	88	60	72.2	68.8	70.2
17	Nickel	mg/kg	20.4	30.4	19.5	21.3	30
18	Copper	mg/kg	60	34	21.5	18.2	23.4
19	Zinc	mg/kg	30.20	32.50	33.20	40.00	28.00
20	Cadmium	mg/kg	ND	ND	ND	ND	ND
21	Lead	mg/kg	2.8	2.4	3.9	5.2	3.8
22	Mercury	mg/kg	ND	ND	ND	ND	ND
23	Arsenic	mg/kg	ND	ND	ND	ND	ND

\*Grab samples could not be collected due high current at KPT 3, Natki Creek Near Tuna port & Vadinar SBM

**Table 34B: Results of Analysis of Sediment of Kandla & Vadinar Port (Neap Tide)**

Sr. No.	Parameters	Unit	KPT - 1	KPT - 2	KPT - 3	KPT - 4	KPT - 5	Jetty
1	Texture	-	Sandy loam	Sandy loam	Sandy loam	Sandy loam	Sandy loam	Sandy loam
2	Organic Matter	mg/kg	1.20	1.12	1.20	1.80	1.62	1.10
3	Organic Carbon	mg/kg	0.69	0.65	0.69	1.04	0.94	0.64
4	Inorganic Phosphate	mg/kg	120.0	142.0	116.0	136.0	142.0	152.0
5	Moisture	%	20.08	21.52	23.05	24.55	28.88	22.02
6	Aluminium	mg/kg	ND	ND	ND	ND	ND	ND
7	Silica	mg/kg	21.20	18.88	21.0	18.8	16.20	13.60
8	Phosphate	mg/kg	8.8	8.9	7.70	8.20	8.40	6.2
9	Sulphate	mg/kg	180.0	196.0	142.0	166.0	120.0	210.0
10	Nitrite	mg/kg	0.1	0.11	0.1	0.12	0.11	0.13
11	Nitrate	mg/kg	9.80	6.89	8.99	8.80	7.93	10
12	Calcium	mg/kg	322.0	266.0	320.0	296.0	300.0	288.0
13	Magnesium	mg/kg	180.0	145.0	180.0	142.0	212.0	196.0
14	Sodium	mg/kg	8242.0	7002.0	8942.0	6641.0	8041.0	9424.0
15	Potassium	mg/kg	380.0	396.0	422.0	644.0	621.0	386.0
16	Chromium	mg/kg	79	54	74.2	64.7	58.4	66
17	Nickel	mg/kg	18.2	28.2	20.6	19.4	28.4	18.8
18	Copper	mg/kg	54	20	22.5	16.8	18.6	74.2
19	Zinc	mg/kg	28.20	18.80	28.40	34.50	18.60	75.00
20	Cadmium	mg/kg	ND	ND	ND	ND	ND	ND
21	Lead	mg/kg	2	2.1	2.8	3.8	2.4	ND
22	Mercury	mg/kg	ND	ND	ND	ND	ND	ND
23	Arsenic	mg/kg	ND	ND	ND	ND	ND	ND

**REPORT**  
**ON**  
**ECOLOGICAL MONITORING**  
**OF MARINE ENVIRONMENT**  
**IN**  
**DPT HARBOUR AREA, NEAR BY CREEKS**  
**For**  
**DEENDAYAL PORT TRUST**

AUGUST, 2021



**Sampling Stations:**

The monitoring of marine environment for the study of biological and ecological Parameters was carried out on 9<sup>th</sup> August, 2021 in harbour region of DPT, and on 10<sup>th</sup> August, 2021 in creeks near by the port during spring tide. The monitoring of marine environment for the study of biological and ecological parameters was repeated again on 15<sup>th</sup> August, 2021 in harbour region of DPT and on 16<sup>th</sup> August, 2021 in creeks near by the port during neap tidal condition.

Plankton samples from sub surface layer was collected both during high tide period and low tide period from 3 water quality monitoring stations of DPT harbour area and one stations in Nakti creek and one station in Khori creek. Sampling at second sampling station of Nakti creek was possible only during high tide period. Collected water samples were processed for estimation of Chlorophyll- a, Pheophytin- a, qualitative & quantitative evaluation of phytoplankton, qualitative & quantitative evaluation zooplanktons (density and their population).

**TABLE #1 SAMPLING LOCATIONS**

<b>monitoring requirement</b>	<b>Number of locations</b>
Kandla creek	3 in Kandla creek
Nakti creek	2 in Nakti creek
Khori Creek	1 in Khori creek
<b>Total Number of locations</b>	<b>6</b>

### Sampling methodology adopted:

A marine sampling is an estimation of the body of information in the population. The theory of the sampling design is depending upon the underlying frequency distribution of the population of interest. The requirement for useful water sampling is to collect a representative sample of suitable volume from the specified depth and retain it free from contamination during retrieval.

50 litres of the water sample were collected from Sub surface by using bucket. From the collected water sample 1 litres of water sample were taken in an opaque plastic bottle for chlorophyll estimation, thereafter plankton samples were collected by using filtration assembly with nilyobolt cloth of 20 $\mu$ m mesh size.

### Samples Processing for chlorophyll estimation:

Samples for the chlorophyll estimation were preserved in ice box on board in darkness to avoid degradation in opaque container covered with aluminium foil. Immediately after reaching the shore after sampling, 1 litres of collected water sample was filtered through GF/F filters (pore size 0.45  $\mu$ m) by using vacuum filtration assembly. After vacuum filtration the glass micro fiber filter paper was grunted in tissue grinder, macerating of glass fiber filter paper along with the filtrate was done in 90% aqueous Acetone in the glass tissue grinder with glass grinding tube. Glass fiber filter paper will assist breaking the cell during grinding and chlorophyll content was extracted with 10 ml of 90% Acetone, under cold dark conditions along with saturated magnesium carbonate solution in glass screw cap tubes. After an extraction period of 24 hours, the samples were transferred to calibrated centrifuge tubes and adjusted the volume to original volume with 90% aqueous acetone solution to make up the evaporation loss. The extract was clarified by using centrifuge in closed tubes. The clarified extracts were then decanted in clean cuvette and optical density was observed at wavelength 664, 665 nm. By using corrected optical density, Chlorophyll-a value was calculated as given in (APHA, 1998).

### PLANKTON:

The entire area open water in the sea is the pelagic realm. Pelagic organisms live in the open sea. In contrast to the pelagic realm, the benthic realm comprises organisms and zone of the bottom of the sea. Vertically the pelagic realm can be dividing into two zones based on light penetration; upper photic or euphotic zone and lower dark water mass, aphotic zone below the photic zone.

The term plankton is general term for organisms have such limited powers of locomotion that they are at the mercy of the prevailing water movement. Plankton is subdivided to phytoplankton and zooplankton. Phytoplankton is free floating organisms that are capable of photosynthesis and zooplankton is the various free floating animals.

Pelagic zone, represents the entire ocean water column from the surface to the deepest depths, is home to a diverse community of organisms. Differences in their locomotive ability categorize the organisms in the pelagic realm into two, **plankton** and **nekton** (Lalli and Parsons, 1997). **Plankton** consists of all organisms drifting in the water and is unable to swim against water currents, whereas **Nekton** includes organisms having strong locomotive power. Ecological studies on the plankton community, which form the base of the aquatic food chain, help in the better understanding of the dynamics and functioning of the marine ecosystem. The term 'Plankton' first coined by Victor Hensen (1887), Plankton, (Greek word: *planktos* meaning "passively drifting or wandering") is defined as drifting or free-floating organisms that inhabit the pelagic zone of water. Based on their mode of nutrition planktonic organisms are categorised into phytoplankton (organisms having an autotrophic mode of nutrition) and zooplankton (organisms having a heterotrophic mode of nutrition).

#### Phytoplankton in the marine environment:

Phytoplankton is free floating unicellular, filamentous and colonial eutrophic organisms that grow in aquatic environments whose movement is more or less dependent upon water currents. These micro flora acts as primary producers as well as the basis of food chain, source of protein, bio purifier and bio indicators of the aquatic ecosystems of which diverse array of the life depends .They are considered as an important component of aquatic flora, play a key role in maintaining equilibrium between abiotic and biotic components of aquatic ecosystem.

The phytoplankton includes a wide range of photosynthetic and phototrophic organisms. Marine phytoplankton is mostly microscopic and unicellular floating flora, which are the primary producers that support the pelagic food-chain. The two most prominent groups of phytoplankton are diatoms (Bacillariophyceae) and dinoflagellates (Dinophyceae).The phytoplankton those normally captured in the net from the Gulf of Kutch is normally dominated by these two major groups; diatoms and dinoflagellates. Phytoplankton also include numerous and diverse collection of extremely small, motile algae which are termed micro flagellates (naked flagellates) as well as and Cyanophytes (blue-green algae).

Algae are an ecologically important group in most aquatic ecosystems and have been an important component of biological monitoring programs. Algae are ideally suited for water quality assessment because they have rapid reproduction rates and very short life cycles, making them valuable indicators of short-term impacts.

Aquatic populations are impacted by anthropogenic stress, resulting in a variety of alterations in the biological integrity of aquatic systems. Algae can serve as an indicator of the degree of deterioration of water quality, and many algal indicators have been used to assess environmental status.

### **Zooplankton in the marine environment:**

Zooplankton includes a taxonomically and morphologically diverse community of heterotrophic organisms that drift in the waters of the world's oceans. Qualitative and quantitative studies on zooplankton community are a prerequisite to delineate the ecological processes active in the marine ecosystem. Zooplankton community plays a pivotal role in the pelagic food web as the primary consumers of phytoplankton and act as the food source for organisms in the higher trophic levels, particularly the economically essential groups such as fish larvae and fishes. They also function in the cycling of elements in the marine ecosystem. The dynamics of the zooplankton community, their reproduction, growth and survival rate are all significant factors determining the recruitment and abundance of fish stocks as they form an essential food for larval, juvenile and adult fishes (Beaugrand et al., 2004). Zooplankton grazing in the marine environment controls the primary production and helps in determining the pelagic ecosystem (Banse, 1995). Through grazing in surface waters and following the production of sinking faecal matters and also by the active transportation of dissolved and particulate matter to deeper waters via vertical migration, they help in the transport of organic carbon to deep ocean layers and thus act as key drivers of biological pump' in the marine ecosystem. Zooplankton grazing and metabolism also, transform particulate organic matter into dissolved forms, promoting primary producer community, microbial demineralization, and particle export to the ocean's interior.

The categorisation of zooplankton into various ecological groups is based on several factors such as duration of planktonic life, size, food preferences and habitat. As they vary significantly in size from microscopic to metazoic forms, the classification of zooplankton based on size has paramount importance in the field of quantitative plankton research.

Based on the duration of planktonic life, zooplankton are categorised into Holoplankton (organisms which complete their entire lifecycle as plankton) and Meroplankton (organisms which are planktonic during the early part of their lives such as the larval stages of benthic and nektonic organisms). Tychoplankton are organisms which live a brief planktonic life, such as the benthic crustaceans (cumaceans, mysids, isopods) which ascend to the water column at night for feeding and certain ectoparasitic copepods, they leave the host and spend their life as plankton during their breeding cycle.


Zooplankton can be subdivided into holoplankton, i.e., permanent members of the plankton (e.g., Calanoid copepods), and meroplankton, i.e., temporary members in the plankton e.g., larvae of fish,

shrimp, and crab). The meroplankton group consists of larval and young stages of animals that will adopt a different lifestyle once they mature. In contrast to phytoplankton which consist of a relatively smaller variety of organisms, Zooplankton are extremely divers, consist of a host of larval and adult forms representing many animal phylum.

Among the zooplankton one group always dominate than others; members of sub class copepods (Phylum Athropoda), and Tintinids (Phylum Protozoa) among the net planktons. These small animals are of vital importance in marine ecosystem as one of the primary herbivores animals in the sea, and it is they provide vital link between primary producer (autotrophs) and numerous small and large marine consumers.

As their community structure and function are highly susceptible to changes in the environmental conditions regular monitoring of their distribution as well as their interactions with various physicochemical parameters is inevitable for the sustainable management of the ecosystem (Kusum et al., 2014). Of all the marine zooplankton groups, copepods mainly calanoid copepods are the dominant groups in marine subtropical and tropical waters and exhibit considerable diversity in morphology and habitats they occupy (Madhupratap, 1991;)

It has been well established that potential of pelagic fishes viz. finfishes, crustaceans, molluscs and marine mammals either directly or indirectly depend on zooplankton. The herbivorous zooplankton is efficient grazers of the phytoplankton and is referred to as living machines transforming plant material into animal tissue. Hence they play an essential role as the intermediaries for nutrients/energy transfer between primary and tertiary trophic levels. Due to their large density, shorter lifespan, drifting nature, high group/species diversity and different tolerance to the stress, they used as the indicator organisms for the physical, chemical and biological processes in the aquatic ecosystem (Ghajbhiye, 2002).



**Spatial distribution of Plankton:**

A characteristic of plankton population is that they tend to occur in patches, which are varying spatially on a scale of few meters to far as few kilometres in distance. They also vary in time scale, season as well as vertically in the water column. It is this patchiness and its constant changes in time and spot, that has made it so difficult for plankton biologist to learn about the ecology of plankton. The biological factors that causes this patchiness is due to the ability of zooplankton to migrate vertically and graze out the phytoplankton at a rapid rate that can create patchiness. Similarly the active swimming ability by certain zooplankton organisms can cause to aggregate in dense group.

At its most extreme, because the water in which plankton is suspended is constantly moving, each sample taken by the plankton biologists remain a different volume of water, so each sample is unique and replicate does not exist.

Plankton may also exhibit vertical patchiness. Physical factors contribute to this type of patchiness include light intensity, nutrients and density gradients in the water column.

Phytoplankton in particular tends to be unequally distributed vertically, which leads to the existence of different concentration of a chlorophyll value between photic zone and below the photic zone.

**Methodology adopted for Plankton sampling:**

Mixed plankton sample were obtained from the sub surface layer at each sampling locations by towing the net horizontally with the weight .After the tow of about 15-30minutes, plankton net was pulled up and washed down to the tail and collected the plankton adhered to plankton net in the collection bucket at the bottom by springing outer and inner surface of the net with sea water, while the net was hanging with the mouth upward. For quantitative evaluation 50 L water samples were collected from subsurface layer and filtered through 20µm mesh size net by using bucket and filtration assembly.

**Preservation and storage:**

Both filtered plankton and those collected from the plankton net were preserved with 5% buffered formalin and stored in 1L plastic container for further processing in the laboratory.

**Sample concentration:**

The collected plankton samples were concentrated by using centrifuge and made up to 50 ml with 5% formalin -Glycerine mixture.

**Taxonomic evaluation:**

Before processing, the sample was mixed carefully and a subsample was taken with a calibrated Stempel-pipette. 1 ml of the concentrated plankton samples were transferred on a glass slide with automatic pipette. The plankton sample on the glass slides were stained by using Lugol's iodine and added glycerine to avoid drying while observation. The plankton samples were identified by using Labex triangular Research microscope with photographic attachment. Microphotographs of the plankton samples were taken for record as well as for confirming the identification. The bigger sized zooplankton was observed through dissecting stereomicroscope with magnification of 20-30 x. Plankton organisms in the whole slide were identified to the lowest axon possible. A thorough literature search was conducted for the identification of the different groups of zooplankton that were encountered

**Cell counts by drop count method:**

The common glass slide mounted with a 1ml of concentrated phytoplankton/zooplankton sample in glycerol and covered with cover slip 22x 60mm was placed under the compound microscope provided with a mechanical stage. The plankton was then counted from the microscopic field of the left top corner of the slide. Then slide is moved horizontally along the right side and plankton in each microscopic field was thus counted. When first microscopic field row was finished the next consecutive row was adjusted using the mechanical device of the stage. In this way all the plankton present in entire microscopic field are counted. From this total number in 1ml of the concentrated

plankton, total number of plankton in the original volume of sample filtered was calculated as units/L.

#### **BENTHIC ORGANISMS:**

Benthos is those organisms that are associated with the sea bed or benthic habitats. Epi- benthic organisms live attached to a hard substratum or rooted to a shallow depth below the surface. In fauna organisms live below the sediment–water interface. Interstitial organisms live and move in pore water among sedimentary grains.

Because the benthic organisms are often collected and separated on sieves, a classification based on the overall size is used. Macro benthos include organisms whose shortest dimension is greater than or equal to 0.5 mm. Meio benthos are smaller than 0.5mm but larger than 42 $\mu$  in size.

The terms such as macro fauna and Meio fauna generally have little relevance with taxonomic classification. The terms Meio fauna and macro fauna depend on the size. Meio fauna were considered as good bioassay of community health and rather sensitive indicators of environmental changes

#### **SAMPLING METHODOLOGY ADOPTED FOR SUB TIDAL REGION:**

Van veen sampler (0.09m<sup>2</sup>) was used for sampling bottom sediments. Two sets of sediments were sampled from each location, one for macro fauna and other for Meio fauna. The macro fauna in the sediments were sieved on board to separate out the organisms. The fixation of Meio fauna is normally done by bulk fixation of the sediment sample. The bulk fixation is done by using 10% formalin (Buffered with borate). The organisms were preserved with seawater as diluting agent.

#### **Sample sieving:**

Sediments samples were sieved to extract the organisms. Sieving was performed carefully as possible to avoid any damage to the animals. The large portion of the sediment was split in to smaller portions and mixed with sea water in a bucket. The cohesive lumps were broken down by continuous stirring. The disaggregated sediments were then passed through the sieves.

#### **Sample staining:**

Sorting of the Meio fauna from the sieve is difficult task especially in the preserved material, because organisms are not easily detectable. To facilitate the animal detection the entire sample retained on the sieve after sieving operation were stained by immersing the sieve in a flat bottom tub with 1% Rose Bengal stain; a protein stain. A staining period of 10-30 minutes is sufficient for sample detection.



### DIVERSITY INDICES:

On the whole, diversity indices provide more information about community composition than simply species richness (number of species present); they also, take the relative abundances of different species into account. Based on this fact, diversity indices therefore depend not only on species richness but on the evenness, or equitability, with which individuals are distributed among the different species (Magurram, A. E. (1988)

A diversity index is a measure of species diversity within a community that consists of co-occurring populations of several (two or more) different species. It includes two components: richness and evenness. Richness is the measure of the number of different species within a sample showing that more the types of species in a community, the higher is the diversity or greater is the richness. Evenness is the measure of relative abundance of the different species with in a community.

The basic idea of diversity index is to obtain a quantitative estimate of biological variability that can be used to compare biological entities composed of discrete components in space and time ( Carol H.R. *etal.* 1998). Biodiversity is commonly expressed through indices based on species richness and species abundances (Whittaker 1972, Lande 1996, Purvis and Hector 2000). Biodiversity indices are a non-parametric tool used to describe the relationship between species number and abundance. The most widely used bio diversity indices are Shannon Weiner index and Simpson's index.

A diversity Index is a single statistic that incorporates in formation on richness and evenness. The diversity measures that incorporate the two concepts may be termed heterogeneity measures (Magurran, 2004).

Any study intended to interpret causes and effect of adverse impact on Biodiversity of communities require suitable measures to evaluate specie richness and Diversity. The former is number of species in community, while latter is a function of relative frequency of different species. Species richness is the iconic measure of biological diversity (Magurran, 2004). Several indices have been created to measure the diversity of species; however, the most widely used in the last decades are the Shannon (1948) and Simpson (1949) (Buzas and Hayek 1996; Gorelick 2006), with the components of diversity: richness ( $S$ ) and evenness ( $J$ )

### Simpson's diversity index

Simpson's index ( $D$ ) is a measure of diversity, which takes into account both species richness, and evenness of abundance among the species present.. The Simson index is one of the meaningful and robust biodiversity measures available.(Magurran ,2004).

The formula for calculating  $D$  is presented as:

$$D = \frac{\sum n_i(n_i - 1)}{N(N - 1)}$$

Where  $n_i$  = the total number of organisms of each individual species

$N$  = the total number of organisms of all species

The value of  $D$  ranges from 0 to 1. With this index, 0 represents infinite diversity and, 1, no diversity. When  $D$  increases diversity decreases. Simpson's index is therefore usually expressed as  $1-D$  or  $1/D$ . (Magurran, 2004)

Low species diversity suggests:

- relatively few successful species in the habitat
- the environment is quite stressful with relatively few ecological niches and only a few organisms are really well adapted to that environment
- food webs which are relatively simple
- change in the environment would probably have quite serious effects

High species diversity suggests:

- a greater number of successful species and a more stable ecosystem
- more ecological niches are available and the environment is less likely to be hostile complex food webs
- environmental change is less likely to be damaging to the ecosystem as a whole

### Species richness indices

The species richness ( $S$ ) is simply the number of species present in an ecosystem. Species richness Indices of species richness are widely used to quantify or monitor the effects of anthropogenic disturbance. A decline in species richness may be concomitant with severe or chronic human-induced perturbation (Fair Fair weather 1990,) Species richness measures have traditionally been the mainstay in assessing the effects of environmental degradation on the biodiversity of natural assemblages of organisms (Clarke & Warwick, 2001)

Species richness is the iconic measure of biological diversity (Magurran, 2004). The species richness ( $S$ ) is simply the number of species present in an ecosystem. This index makes no use of relative abundances. The term species richness was coined by McIntosh (1967) and oldest and most intuitive measure of biological diversity (Magurran, 2004).

Margalef's diversity index is a species richness index. Margalef's Species richness index ( $d$ ), or indices that describe the evenness of the distribution of the numbers of individuals among species, were derived.

The value of a diversity index increases both when the number of types increases and when evenness increases. For a given number of types, the value of diversity index is maximised when all types are equally abundant (Rosenzweig, M. L. (1995).

#### **Shannon-Wiener's index:**

An index of diversity commonly used in plankton community analyses is the Shannon-Wiener's index (**H**), which emphasizes not only the number of species (richness or variety), but also the apportionment of the numbers of individuals among the species (Odum 1971 and Reish 1984). Shannon-Wiener's index (**H**) reproduce community parameters to a single number by using an equation.

Shannon and Weiner index represents entropy. It is a diversity index taking into account the number of individuals as well as the number of taxa. It varies from 0 for communities with only single taxa to high values for community with many taxa each with few individuals. This index can also determine the pollution status of a water body. Normal values range from 0 to 4. This index is a combination of species present and the evenness of the species. Examining the diversity in the range of polluted and unpolluted ecosystems, Wilham and Dorris (1968) concluded that the values of the index greater than 3 indicate clean water, values in the range of 1 to 3 are characterized by moderate pollution and values less than 1 are characterized as heavily polluted

$$H' = - \sum_{j=1}^s \frac{n_j}{N} \ln \left( \frac{n_j}{N} \right)$$

#### **RESULTS:**

##### **CHLOROPHYLL-a:**

Water Samples for the chlorophyll estimation were collected from sub surface layer during high tide and low tide period of the tidal cycle for each sampling locations and analysed for Chlorophyll -a and after acidification for Pheophytin -a. Chlorophyll- a value was used as algal biomass indicator (APHA,1998) Algal biomass was estimated by converting Chlorophyll value.

In the sub surface water chlorophyll-a was varying from 0.305 -0.543mg/m<sup>3</sup>.in harbour region of DPT during sampling done in spring tide period of August, 2021. In the nearby creeks chlorophyll-a was varying from 0.290-0.732 mg/m<sup>3</sup>.Pheophytin -a level was below detectable limit- the all the sampling stations during springtide in the harbour region of DPT.

In the sub surface water chlorophyll-a was varying from 0.220 -0.748mg/m<sup>3</sup>.in harbour region of DPT during sampling done in neap tide period of August, 2021 . In the nearby creeks chlorophyll-a was varying from BDL-0.862 mg/m<sup>3</sup>.Pheophytin -a level was below detectable limit- the all the sampling stations during neap tide in the harbour region of DPT.

**DCPL/DPT/20-21/16 -AUGUST - 2021**

---

**TABLE #2 VARIATIONS IN CHLOROPHYLL –a PHEOPHYTIN- a AND ALGAL BIOMASS FROM SAMPLING STATIONS IN DPT HARBOUR AREA ,NEAR BY CREEKS DURING SPRING TIDE IN AUGUST,2021**

Sr. No.	Station	Tide	Chlorophyll-a (mg/m <sup>3</sup> )	Pheophytin- a (mg/m <sup>3</sup> )	Algal Biomass (Chlorophyll method) mg/m <sup>3</sup>
DPTHARBOUR AREA					
1	KPT1	High tide	0.425	BDL	28.48
		Low tide	0.307	BDL	20.57
2	KPT 2	High tide	0.305	BDL	20.43
		Low tide	0.543	BDL	36.38
3	KPT 3	High tide	0.527	BDL	35.31
		Low tide	0.425	BDL	28.47
CREEKS					
4	KPT-4 Khori-I	High tide	0.543	BDL	36.38
		Low tide	0.527	BDL	35.31
5	KPT-5 Nakti-I	High tide	0.409	BDL	27.40
		Low tide	0.732	BDL	49.04
6	KPT-5 Nakti-II	High tide	0.290	BDL	19.43

BDL: Below Detectable Limit.

**TABLE #3 VARIATIONS IN CHLOROPHYLL –a PHEOPHYTIN- a AND ALGAL BIOMASS FROM SAMPLING STATIONS IN DPT HARBOUR AREA ,NEAR BY CREEKS DURING NEAP TIDE IN AUGUST,2021**

Sr. No.	Station	Tide	Chlorophyll-a (mg/m <sup>3</sup> )	Pheophytin- a (mg/m <sup>3</sup> )	Algal Biomass (Chlorophyll method) mg/m <sup>3</sup>
DPTHARBOUR AREA					
1	KPT1	High tide	0.220	BDL	14.74
		Low tide	0.308	BDL	20.64
2	KPT 2	High tide	0.748	BDL	50.11
		Low tide	0.731	BDL	48.98
3	KPT 3	High tide	0.307	BDL	20.56
		Low tide	0.221	BDL	14.81
CREEKS					
4	KPT-4 Khori-I	High tide	0.543	BDL	36.38
		Low tide	0.221	BDL	14.81
5	KPT-5 Nakti-I	High tide	0.862	BDL	57.75
		Low tide	0.216	BDL	14.47
6	KPT-5 Nakti-II	High tide	BDL	BDL	-

BDL: Below Detectable Limit.

### PHYTOPLANKTON POPULATION:

For the evaluation of the Phytoplankton population in DPT harbour area and within the immediate surroundings of the port, sampling was conducted from 5 sampling locations (3 in harbour area and two in Nakti creek) during high tide period and low tide period of spring tide and neap tide.

The phytoplankton community of the sub surface water in the harbour and nearby creeks was represented by, Diatoms blue green algae and dinoflagellates during spring tide period. Diatoms were represented by 14 genera. Blue green were represented by three genera and two genera of Dinoflagellates during the sampling conducted in spring tide in August, 2021. Phytoplankton of the sampling stations at sub surface layer in the harbour area and nearby creeks was varying from 98-226 units/ L during high tide period and 191-259 units/ L during low tide of Spring Tide.

The phytoplankton community of the sub surface water in the harbour and nearby creeks was represented by Diatoms, Blue green algae and Dinoflagellates during Neap tide period. Diatoms were represented by 15 genera and Blue green algae were represented two genera and Dinoflagellates were represented by three genera during the sampling conducted in Neap tide in August, 2021. Phytoplankton of the sampling stations at sub surface layer in the harbour area and nearby creeks was varying from 65-307 units/ L during high tide period and 238-281 units/ L during low tide of Neap Tide.

### Species Richness Indices and Diversity Indices:

#### Margalef's diversity index (Species Richness) S

At the organismal level, the most widely used biodiversity measures are those based on the number of species present, perhaps adjusted for the number of individuals sampled, Here Margalef's Species richness index (d), or indices that describe the evenness of the distribution of the numbers of individuals among species, are derived.

Margalef's diversity index (Species Richness) S of phytoplankton communities in the stations was varying from 1.527-3.091 with an average of 2.420 during the sampling conducted in High tide period of spring tide. While Margalef's diversity index (Species Richness) S of phytoplankton communities in the Kandla Harbour region and nearby creeks was varying from 1.679-2.621 with an average of 2.225 during the consecutive low tide period.

Margalef's diversity index (Species Richness) S of phytoplankton communities in the stations was varying from 2.344 -3.188 with an average of 2.887 during the sampling conducted in High tide period of Neap tide. While Margalef's diversity index (Species Richness) S of phytoplankton communities in the Kandla Harbour region and nearby creeks was varying from 2.526-3.246 with an average of 2.887 during the consecutive low tide period.

**Shannon-Wiener's index:**

Shannon-Wiener's Index (H) of phytoplankton communities in the sampling stations was in the range of 0.728 -0.860 ( $H'(\log_{10})$ ) between selected sampling stations with an average value of 0.807 during high tide period of spring tide .Shannon-Wiener's Index (H) of phytoplankton communities in the sampling stations was in the range of 0.726-0.836 ( $H'(\log_{10})$ ) between selected sampling stations with an average value of 0.773 during consecutive lowtide. Shannon-Wiener's Index (H) of phytoplankton communities in the sampling stations was in the range of 0.880-0.959 ( $H'(\log_{10})$ ) between selected sampling stations with an average value of 0.909 during high tide period of neap tide. Shannon-Wiener's Index (H) of phytoplankton communities in the sampling stations was in the range of 0.819-0.911 ( $H'(\log_{10})$ ) between selected sampling stations with an average value of 0.887 during consecutive low tide. Typical values are generally between 1.5 and 3.5 in most ecological studies, and the index is rarely greater than 4. The Shannon-Wiener's index increases as both the richness and the evenness of the community increase. This result indicates that diversity of phytoplankton of Kandla Harbour region and nearby creeks is less but with abundant population of few, with relatively few ecological niches and only very few opportunist organisms are really well adapted to this environment and thrive better than other species.

**Simpson's diversity index:**

Simpson's index (D) is a measure of diversity, which takes into account both species richness, and an evenness of abundance among the species present. The Simpson index is one of the meaningful and robust biodiversity measures available. (Magurran,2004).

Simpson diversity index (1-D) of phytoplankton communities was below 0.9 at all sampling stations in the Kandla Harbour region and nearby creeks, which was varying from 0.785- 0.823 between selected sampling stations with an average of 0.801 during high tide period of spring tide . Simpson diversity index (1-D) of phytoplankton communities was below 0.9 at all sampling stations in the Kandla Harbour region and nearby creeks, which was varying from 0.770- 0.820 between selected sampling stations with an average of 0.787 during consecutive low tide . Low species diversity suggests a relatively few successful species in this habitat. The environment is quite stressful with relatively few ecological niches and only a few organisms are really well adapted to that environment. Any change in the environment would probably have quite serious effects.

Simpson diversity index (1-D) of phytoplankton communities was below 0.9 at all sampling stations in Kandla Harbour region and nearby creeks, during high tide period and low tide period during neap tide also, which was varying from 0.811-0.886 with an average value of 0.836 between selected sampling stations during high tide period and varying from 0.774-0.826 with an average

value of 0.813 between selected sampling stations during consecutive low tide period Low species diversity suggests a relatively few successful species in this habitat.

**Table # 4 PHYTOPLANKTON VARIATIONS IN ABUNDANCE AND DIVERSITY IN SUB SURFACE SAMPLING STATIONS IN DPT HARBOUR AREA ,NEAR BY CREEKS DURING SPRING TIDE IN AUGUST,2021**

Tide	Sampling Station	Abundance In units/L	No of Species observed /total species	% of diversity	Margalef's diversity index (Species Richness S)	Shannon Weiner index H (log <sub>10</sub> )	Diversity Index (Simpson's Index) 1-D
HIGH TIDE	1	210	14/19	73.68	2.431	0.7923	0.7938
	2	177	17/19	89.47	3.091	0.8603	0.8182
	3	226	12/19	63.16	2.029	0.7883	0.7853
	4	221	17/19	89.47	2.964	0.8243	0.7993
	5	190	14/19	73.68	2.478	0.8531	0.8227
	6	98	8/19	42.11	1.527	0.7279	0.7886
LOW TIDE	1	191	13/19	68.42	2.285	0.7812	0.7901
	2	233	13/19	68.42	2.201	0.7658	0.7795
	3	209	15/19	78.94	2.621	0.8367	0.8208
	4	213	10/19	52.63	1.679	0.7264	0.7732
	5	259	14/19	73.68	2.339	0.7547	0.7702

**Table # 5 PHYTOPLANKTON VARIATIONS IN ABUNDANCE AND DIVERSITY IN SUB SURFACE SAMPLING STATIONS IN DPT HARBOUR AREA ,NEAR BY CREEKS DURING NEAP TIDE IN AUGUST,2021**

Tide	Sampling Station	Abundance In units/L	No of Species observed /total species	% of diversity	Margalef's diversity index (Species Richness S)	Shannon Weiner index H (log <sub>10</sub> )	Diversity Index (Simpson's Index) 1-D
HIGH TIDE	1	221	18/20	90	3.149	0.9462	0.8522
	2	283	19/20	95	3.188	0.8844	0.811
	3	268	17/20	85	2.862	0.899	0.8255
	4	256	14/20	70	2.344	0.8803	0.8328
	5	307	19/20	95	3.143	0.8857	0.8113
	6	65	12/20	60	2.635	0.9594	0.8861
LOW TIDE	1	238	15/20	75	2.558	0.8192	0.7738
	2	281	19/20	95	3.192	0.9106	0.8188
	3	256	19/20	95	3.246	0.9023	0.8241
	4	242	17/20	85	2.915	0.9102	0.8263
	5	255	15/20	75	2.526	0.8939	0.824

**Table # 6 ABUNDANCE OF PHYTOPLANKTON SUBSURFACE SAMPLING STATIONS IN DPT HARBOUR AREA, NEAR BY CREEKS DURING SPRING TIDE IN AUGUST, 2021**


Tide	Surface	No of Sampling location	Group of phytoplankton	Phytoplankton Group range Units/L	Genera or species /total Phyto plankton	Taxon Diversity % (Group level)
HIGH TIDE	Sub surface	6	DIATOMS	94-215	14/19	73.68
			BLUE GREEN	4-13	3/19	15.79
			DINOFLAGELLATES	0-1	2/19	10.53
			TOTAL PHYTO PLANKTON	98-226	19	-
LOW TIDE	Sub surface	5	DIATOMS	182-250	14/19	73.68
			BLUE GREEN	8-12	3/19	15.79
			DINOFLAGELLATES	0-1	2/19	10.53
			TOTAL PHYTO PLANKTON	191-259	19	-

**Table # 7 ABUNDANCE OF PHYTOPLANKTON SUBSURFACE SAMPLING STATIONS IN DPT HARBOUR AREA, NEAR BY CREEKS DURING NEAP TIDE IN AUGUST, 2021**


Tide	Surface	No of Sampling location	Group of phytoplankton	Phytoplankton Group range Units/L	Genera or species /total Phyto plankton	Taxon Diversity % (Group level)
HIGH TIDE	Sub surface	6	DIATOMS	64-298	15/20	75
			BLUE GREEN	0-6	2/20	10
			DINOFLAGELLATES	0-5	3/20	15
			TOTAL PHYTO PLANKTON	65-307	20	-
LOW TIDE	Sub surface	5	DIATOMS	236-274	15/20	75
			BLUE GREEN	1-5	2/20	10
			DINOFLAGELLATES	0-4	3/20	15
			TOTAL PHYTO PLANKTON	238-281	20	-



**Taxon Diversity % of Phytoplankton during High tide and Low tide period during spring tide**



**Taxon Diversity % of Phytoplankton during High tide and Low tide period during Neap tide**



**ZOOPLANKTON POPULATION:**

For the evaluation of the Zooplankton population in DPT harbour area and within the immediate surroundings of the port sampling was conducted from 6 sampling locations (3 in harbour area and two in Nakti creek and one in Khoricreek) during high tide period and low tide period of spring tide and Neap tide in August,2021. The Zooplankton community of the sub surface water in the harbour and nearby creeks during spring tide was represented by mainly five groups, Tintinids, Copepods,

Ciliates Foraminiferans and larval forms of Crustacea, Molluscans and Polychaetes. The Zooplankton community of the sub surface water in the harbour and nearby creeks during neap tide was represented by mainly four groups, Tintinids, Copepods, Mysids and larval forms of Crustaceans, Molluscans and Polychaetes.

Zooplankton of the sampling stations at sub surface layer in the DPT harbour area and nearby creek was varying from 34-109x10<sup>3</sup> N/ m<sup>3</sup> during high tide and 109-123 x10<sup>3</sup> N/ m<sup>3</sup> during low tide of Spring Tide period. Zooplankton of the sampling stations at sub surface layer in the DPT harbour area and nearby creek was varying from 23-109 x10<sup>3</sup> N/ m<sup>3</sup> during high tide and 86-103x10<sup>3</sup> N/ m<sup>3</sup> during low tide of Neap Tide period.

### **Species Richness Indices and Diversity Indices:**

#### **Margalef's diversity index (Species Richness) S**

At the organismal level, the most widely used biodiversity measures are those based on the number of species present, perhaps adjusted for the number of individuals sampled, Here Margalef's Species richness index (d), or indices that describe the evenness of the distribution of the numbers of individuals among species, are derived.

Margalef's diversity index (Species Richness) S of Zooplankton communities in the stations Kandla Harbour region and nearby creeks was varying from 2.269-3.505 with an average of 3.009 during the sampling conducted in High tide period. Margalef's diversity index (Species Richness) S of Zooplankton communities varying from 2.701-3.354 with an average of 3.033 during the sampling conducted in low tide period during Spring tide.

Margalef's diversity index (Species Richness) S of Zooplankton communities in the Kandla Harbour region and nearby creeks sampling stations was varying from 1.914- 3.962 with an average of 2.754 during the sampling conducted in high tide and varying from 1.972-3.236 with an average of 2.640 during the sampling conducted in low tide during Neap tide period.

#### **Shannon-Wiener's index:**

Shannon-Wiener's Index (H) of Zooplankton communities in the sampling stations in Kandla Harbour region and nearby creeks was in the range of 0.813-1.052 (H'(log10)) between selected sampling stations with an average value of 0.995 (H'(log10)) during high tide period of spring tide. Shannon-Wiener's Index (H) of Zooplankton communities in the sampling stations in Kandla Harbour region and nearby creeks was in the range of 0.991-1.067(H'(log10)) between selected sampling stations with an average value of 1.035 (H'(log10)) during consecutive low tide period.

Shannon-Wiener's Index (H) of Zooplankton communities in the sampling stations in Kandla Harbour region and nearby creeks was in the range of 0.696-1.007 (H'(log10)) between selected sampling stations with an average value of 0.897 (H'(log10)) during high tide period of Neap tide.

Wiener's Index (H) of Zooplankton communities in the sampling stations in Kandla Harbour region and nearby creeks was in the range of 0.785-0.983 ( $H'(\log_{10})$ ) between selected sampling stations with an average value of 0.883 ( $H'(\log_{10})$ ) during consecutive low tide period. Typical values are generally between 1.5 and 3.5 in most ecological studies, and the index is rarely greater than 4. The Shannon-Wiener's index increases as both the richness and the evenness of the community increase. This result indicates that diversity of Zooplankton of Kandla Harbour region and nearby creeks stations is slightly high with very minimum diverse population but very few opportunist organisms are really well adapted to this environment and thrive better than other species.

**Simpson's diversity index:**

Simpson's index (D) is a measure of diversity, which takes into account both species richness, and an evenness of abundance among the species present. The Simpson index is one of the meaningful and robust biodiversity measures available. (Magurran, 2004).

Simpson diversity index (1-D) of Zooplankton communities was below 0.9 most of sampling stations in the Kandla Harbour region and nearby creeks during high tide and low tide of spring tide period, which was varying from 0.839-0.899 between selected sampling stations with an average of 0.884 during high tide period and was varying from 0.887- 0.908 with an average value of 0.897 between selected sampling stations during low tide

Simpson diversity index (1-D) of Zooplankton communities was below 0.9 at all sampling stations in the Kandla Harbour region and nearby creeks during high tide and low tide period except few, which was varying from 0.771-0.869 between selected sampling stations with an average of 0.833 during high tide period and was varying from 0.787- 0.863 with an average value of 0.826 between selected sampling stations during consecutive low tide

This low species diversity suggests a relatively low number of successful species in this habitat. Environment is quite stressful with relatively few ecological niches and only few organisms are really well adapted to that environment. Any change in the environment would probably have quite serious effects.

**Table # 8 ZOOPLANKTON VARIATION IN ABUNDANCE AND DIVERSITY IN SUB SURFACE SAMPLING STATIONS IN DPT HARBOUR AREA ,NEAR BY CREEKS DURING SPRING TIDE IN AUGUST,2021**

Tide	Sampling Station	Abundance In No / m <sup>3</sup>	No of Species/groups observed /total species/group	% of diversity	Margalef's diversity index (Species Richness S)	Shannon Weiner index H (log <sub>10</sub> )	Diversity Index (Simpson's Index) 1-D
HIGH TIDE	1	93 X10 <sup>3</sup>	16/19	84.21	3.309	1.052	0.8955
	2	96 X10 <sup>3</sup>	17/19	89.47	3.505	1.032	0.8899
	3	105 X10 <sup>3</sup>	15/19	78.95	3.008	1.031	0.8958
	4	109 X10 <sup>3</sup>	15/19	78.95	2.984	1.037	0.8991
	5	109 X10 <sup>3</sup>	15/19	78.95	2.984	1.008	0.8865
	6	34 X10 <sup>3</sup>	9/19	47.37	2.269	0.8131	0.8396
LOW TIDE	1	110 X10 <sup>3</sup>	15/19	78.95	2.978	1.001	0.8881
	2	118 X10 <sup>3</sup>	17/19	89.47	3.354	1.067	0.8984
	3	123 X10 <sup>3</sup>	14/19	73.68	2.701	0.9911	0.887
	4	117 X10 <sup>3</sup>	16/19	84.21	3.15	1.065	0.9088
	5	109 X10 <sup>3</sup>	15/19	78.95	2.984	1.051	0.904

**Table # 9 ZOOPLANKTON VARIATION IN ABUNDANCE AND DIVERSITY IN SUB SURFACE SAMPLING STATIONS IN DPT HARBOUR AREA ,NEAR BY CREEKS DURING NEAP TIDE IN AUGUST,2021**

Tide	Sampling Station	Abundance In No / m <sup>3</sup>	No of Species/groups observed /total species/group	% of diversity	Margalef's diversity index (Species Richness S)	Shannon Weiner index H (log <sub>10</sub> )	Diversity Index (Simpson's Index) 1-D
HIGH TIDE	1	99 X10 <sup>3</sup>	14/20	70	2.829	0.9755	0.8695
	2	94 X10 <sup>3</sup>	19/20	95	3.962	1.007	0.8646
	3	91 X10 <sup>3</sup>	13/20	65	2.66	0.9544	0.8698
	4	101 X10 <sup>3</sup>	14/20	70	2.817	0.8993	0.8176
	5	109 X10 <sup>3</sup>	12/20	60	2.345	0.8501	0.8089
	6	23 X10 <sup>3</sup>	7/20	35	1.914	0.6965	0.7708
LOW TIDE	1	89 X10 <sup>3</sup>	11/20	55	2.228	0.8172	0.7878
	2	103 X10 <sup>3</sup>	16/20	80	3.236	0.9831	0.8633
	3	96 X10 <sup>3</sup>	14/20	70	2.848	0.92	0.8412
	4	86 X10 <sup>3</sup>	14/20	70	2.918	0.9071	0.8375
	5	96 X10 <sup>3</sup>	10/20	50	1.972	0.7875	0.7987


**Table # 10 ABUNDANCE OF ZOOPLANKTON IN SUBSURFACE SAMPLING STATIONS IN DPT HARBOUR AREA ,NEAR BY CREEKS DURING SPRING TIDE IN AUGUST,2021**

Tide	Surface	No of Sampling locations	Group of Zooplankton	Abundance of Zooplankton $\times 10^3$ Group Range	Genera or species /total Zooplankton	Taxon Diversity % (Group level)
HIGH TIDE	Sub surface	6	Tintinids	1-28	5/19	26.32
			Copepods	17-49	6/19	31.58
			Ciliates	1-6	1/19	5.26
			Larval forms	9-46	6/19	31.58
			Foraminiferans	0-2	1/19	5.26
			TOTAL ZOOPLANKTON NO/L	<b>34-109</b>	<b>19</b>	
LOW TIDE	Sub surface	5	Tintinids	20-27	5/19	26.32
			Copepods	39-55	6/19	31.58
			Ciliates	1-7	1/19	5.26
			Larval forms	40-46	6/19	31.58
			Foraminiferans	0-1	1/19	5.26
			TOTAL ZOOPLANKTON NO/L	<b>109-123</b>	<b>19</b>	


**Table # 11 ABUNDANCE OF ZOOPLANKTON IN SUBSURFACE SAMPLING STATIONS IN DPT HARBOUR AREA , NEAR BY CREEKS DURING NEAPTIDE IN AUGUST,2021**

Tide	Surface	No of Sampling locations	Group of Zooplankton	Abundance of Zooplankton $\times 10^3$ Group Range	Genera or species /total Zooplankton	Taxon Diversity % (Group level)
HIGH TIDE	Sub surface	6	Tintinids	1-11	5/20	25
			Copepods	7-26	6/20	30
			Mysids	0-6	2/20	10
			Larval forms	15-84	7/20	35
			TOTAL ZOOPLANKTON NO/L	<b>23-109</b>	<b>20</b>	-
LOW TIDE	Sub surface	5	Tintinids	6-12	5/20	25
			Copepods	5-23	6/20	30
			Mysids	1-4	2/20	10
			Larval forms	57-74	7/20	35
			TOTAL ZOOPLANKTON NO/L	<b>86-103</b>	<b>20</b>	-

**Taxon Diversity % of Zooplankton during High tide and Low tide period of Spring tide**



**Taxon Diversity % of Zooplankton during High tide and Low tide period of Neap tide**



**TABLE # 12 SYSTEMATIC ACCOUNT OF PHYTOPLANKTON IN THE SAMPLING LOCATIONS IN OF DPT HARBOUR AREA AND NEARBY CREEKS DURING SPRING TIDE OF AUGUST, 2021**

GROUP	PHYLUM	CLASS	ORDER	FAMILY	GENUS/SPECIES	#	Relative Abundance
BLUE GREEN ALGAE	Cyanophyta	Cyanophyceae	Nostocales	Oscillatoriaceae	<i>Arthospirasp.</i>	B1	Rare
					<i>Lyngbya sp.</i>	B2	Rare
			Stigonematales	Stigonemataceae	<i>Stigonema sp.</i>	B3	Occasional
DIATOMS	Bacillariophyta	Coscinodiscophyceae	Thalassiosirales	Thalassiosiraceae	<i>Planktoniellasp</i>	D1	Rare
			Coscinodiscales	Coscinodiscaceae	<i>Coscinodiscus sp.</i>	D2	Abundant
			Triceratiales	Triceratiaceae	<i>Odontellasp</i>	D3	Occasional
					<i>Triceratiumsp.</i>	D4	Rare
			Biddulphiales	Biddulphiaceae	<i>Biddulphiasp</i>	D5	Dominant
			Hemiaulales	Belleracheaceae	<i>Belleracheasp</i>	D6	Rare
			Chaetocerotales	Chaetocerotaceae	<i>Chaetocerossp</i>	D7	Occasional
			Lithodesmiales	Lithodesmiaceae	<i>Ditylumsp</i>	D8	Abundant
		Bacillariophyceae	Naviculales	Pleurosigmaaceae	<i>Pleurosigmasp</i>	D9	Rare
		Fragilariophyceae	Thalassionematales	Thalassionemataceae	<i>Thalassiothrix sp.</i>	D10	Abundant
					<i>Thalassionema sp.</i>	D11	Rare
			Fragilariales	Fragilariaceae	<i>Asterionelopsis sp..</i>	D12	Rare
					<i>Fragilariasp</i>	D13	Occasional
					<i>Synedrasp</i>	D14	Rare
DINO FLAGELLATES	Dinoflagellata / Dinozoa	Dinophyceae	Peridinales	Proto-peridiniaceae	<i>Proto-peridinium sp.</i>	DF1	Rare
			Gonyaulacales	Ceratiaceae	<i>Ceratiumfurca</i>	DF2	Rare

**TABLE # 12 SYSTEMATIC ACCOUNT OF PHYTOPLANKTON IN THE SAMPLING LOCATIONS IN OF DPT HARBOUR AREA AND NEARBY CREEKS DURING AND NEAP TIDE OF AUGUST,2021**

GROUP	PHYLUM	CLASS	ORDER	FAMILY	GENUS/SPECIES	#	Relative Abundance
BLUE GREEN ALGAE	Cyanophyta	Cyanophyceae	Nostocales	Oscillatoriaceae	<i>Oscillatoria sp.</i>	B1	Rare
			Stigonematales	Stigonemataceae	<i>Stigonema sp.</i>	B2	Rare
DIATOMS	Bacillariophyta	Coccinodiscophyceae	Coccinodiscales	Coccinodiscaceae	<i>Coccinodiscus sp.</i>	D1	Dominant
			Triceratiales	Triceratiaceae	<i>Odontellasp</i>	D2	Occasional
					<i>Triceratiumsp.</i>	D3	Rare
					<i>Biddulphi</i> asp	D4	Abundant
			Biddulphiales	Biddulphiaceae	<i>Biddulphi</i> asp	D4	Abundant
			Hemiaulales	Bellerocheaceae	<i>Bellerochea</i> asp	D5	Rare
			Chaetocerotales	Chaetocerotaceae	<i>Chaetoceros</i> sp	D6	Occasional
			Rhizosoleniales	Rhizosoleniaceae	<i>Rhizosolenia sp.</i>	D7	Occasional
		Lithodesmiales	Lithodesmiaceae	<i>Ditylum</i> sp	D8	Abundant	
		Bacillariophyceae	Naviculales	Pleurosigma	<i>Pleurosigma</i> sp	D9	Occasional
					<i>Pinnularia</i> sp	D10	Rare
		Fragilariophyceae	Thalassionematales	Thalassionemataceae	<i>Thalassiothrix sp.</i>	D11	Dominant
					<i>Thalassionema sp.</i>	D12	Rare
			Fragilariales	Fragilariaceae	<i>Asterionella sp.</i>	D13	Occasional
					<i>Fragilaria</i> sp	D14	Frequent
<i>Synedra</i> sp	D15				Rare		
DINO FLAGELLATES	Dinoflagellata / Dinozoa	Dinophyceae	Peridinales	Protoperidiniaceae	<i>Protoperidinium sp.</i>	DF1	Rare
			Gonyaulacales	Ceratiaceae	<i>Ceratiumfurca</i>	DF2	Rare
					<i>Ceratiumtripos</i>	DF3	Rare



**TABLE #13 SYSTEMATIC ACCOUNT OF ZOOPLANKTON FROM THE SAMPLING LOCATIONS IN DPT HARBOUR AREA, AND NEAR BY CREEKS DURING SPRING TIDE OF AUGUST,2021**

GROUP	PHYLUM	CLASS	ORDER	FAMILY	GENUS/SPECIES	#	RELATIVE ABUNDANCE
TINTINIDS	PROTOZOA CILIOPHORA	Spirotrichea	Tintinnida	Tintinnidiidae	<i>Leprotintinnus</i> sp.	T1	Frequent
				Codonellidae	<i>Tintinnopsisaccuminata</i>	T2	Rare
					<i>Tintinnopsisfailakkaensis</i>	T3	Rare
					<i>Tintinnopsisgracilis</i>	T4	Occasional
					<i>Tintinnopsisradix</i>	T5	Rare
COPEPODS	ATHROPODA	Crustacea Sub class copepoda	Calanoida	Paracalanidae	<i>Acrocalanus</i> sp.	C1	Frequent
				Temoridae	<i>Temora</i> sp.	C2	Rare
			Cyclopoida	Oithonidae	<i>Oithona</i> sp.	C3	Frequent
			Harpacticoida	Ectinosomatidae	<i>Microsetella</i> sp.	C4	Abundant
				Euterpinidae	<i>Euterpina</i> sp.	C5	Rare
			Poecilostomatoida	Oncaeidae	<i>Oncaea</i> sp.	C6	Rare
CILIATES	CILIOPHORA	Oligohymenophorea	Sessilida	Zoothamniidae	<i>Zoothamnium</i> sp.	CI1	Occasional
CRUSTACEAN LARVAE	ARTHROPODA (CRUSTACEA)	Copepoda			Nauplius larvae of Copepods	L1	Dominant
BIVALVE LARVAE	MOLLUSCA	Pelecypoda			Veliger larvae of Bivalves	L2	Occasional
POLYCHAETE LARVA	ANNELIDA	Polychaeta			Trochophore larvae	L3	Occasional
BARNACLE LARVAE	ATHROPODA CRUSTACEA	Maxillopoda Thecostraca			Cirripede larvae	L4	Rare
MOLLUSCAN LARVAE	MOLLUSCA	Gastropoda Streptoneura			Opisthobranchia larvae	L5	Occasional
BRYOZOA					Cyphonautes larvae	L6	Occasional
FORAMINIFERA	FORAMINIFERA	Globobulimina	Rotaliida	Rotaliidae	<i>Rotalia</i> sp.	F1	Rare

**TABLE # 13 SYSTEMATIC ACCOUNT OF ZOOPLANKTON FROM THE SAMPLING LOCATIONS IN DPT HARBOUR AREA, AND NEAR BY CREEKS DURING NEAP TIDE OF AUGUST,2021**

GROUP	PHYLUM	CLASS	ORDER	FAMILY	GENUS/SPECIES	#	RELATIVE ABUNDANCE
TINTINIDS	PROTOZOA CILIOPHORA	Spirotrichea	Tintinnida	Tintinnidiidae	<i>Leprotintinnussp.</i>	T1	Occasional
				Codonellidae	<i>Tintinnopsisaccuminata</i>	T2	Rare
					<i>Tintinnopsisfailakkaensis</i>	T3	Rare
					<i>Tintinnopsisgracilis</i>	T4	Occasional
					<i>Tintinnopsis radix</i>	T5	Rare
COPEPODS	ATHROPODA	Crustacea Sub class copepoda	Calanoida	Paracalanidae	<i>Acrocalanus sp.</i>	C1	Frequent
				Temoridae	<i>Temora sp.</i>	C2	Rare
			Cyclopoida	Oithonidae	<i>Oithona sp.</i>	C3	Occasional
				Harpacticoida	Ectinosomatidae	<i>Microsetellasp.</i>	C4
			Euterpinidae		<i>Euterpina sp.</i>	C5	Rare
			Poecilostomatatoida		Oncaeiidae	<i>Oncaea sp.</i>	C6
			MYSIDS	ATHROPODA CRUSTACEA	Malacostraca	Mysida, Decapoda	Solenoceridae
Penaeidae	<i>Metapenaeussp.</i>	M2					Rare
CRUSTACEAN LARVAE	ARTHROPODA (CRUSTACEA)	Copepoda			Nauplius larvae of Copepods	L1	Dominant
BIVALVE LARVAE	MOLLUSCA	Pelecypoda			Veliger larvae of Bivalves	L2	Occasional
POLYCHAETE LARVA	ANNELIDA	Polychaeta			Trochophore larvae	L3	Occasional
BARNACLE LARVAE	ATHROPODA CRUSTACEA	Maxillopoda Thecostraca			Cirripede larvae	L4	Rare
MOLLUSCAN LARVAE	MOLLUSCA	Gastropoda Streptoneura			Opisthobranchia larvae	L5	Abundant
CYPHONAUTES LARVAE	BRYOZOA				Cyphonautes larvae	L6	Occasional
ECHINODERMATA larve	ECHINODERMATA	Ophiuroidea			Ophiopluetus larvae	L7	Rare

DCPL/DPT/20-21/16 -AUGUST - 2021

**BENTHIC ORGANISMS:**

Few Benthic organisms were observed in the collected sediments by using the Van-veen grabs during the sampling conducted during spring tide period and Neap tide period from DPT harbour region and nearby creek. The benthic organisms during spring tide were represented by Polychaetes, Nematodes and Amphipods. The polychaetes were represented by *Syllis sp.*, *Polydorasp*, and *Pondodorasp*, during spring tide sampling. The benthic organisms in the collected samples were varying from 0-300 N/M<sup>2</sup> during spring tide and 10-140 NO/M<sup>2</sup> during neap tide sampling

**Table # 14 BENTHIC FAUNA IN THE SAMPLING LOCATIONS IN DPT HARBOUR AREA CREEKS DURING NEAP TIDE IN AUGUST ,2021**

Benthic fauna	ABUNDANCE IN NO/M <sup>2</sup> DIFFERENT SAMPLING STATIONS						
	REPRESENTATION BY GROUP						
	DPT HARBOUR			CREEKS			
POLYCHATES	KPT-1	KPT-2	KPT-3	KPT-4	KPT-5	KPT-6	
Family : Iospilidae <i>Pondodora sp.</i>	0	70	0	0	0	NS	
Family : Spionidae <i>Polydora sp..</i>	10	10	0	20	0	NS	
Family : Syllidae <i>Syllis sp.</i>	0	10	0	10	0	NS	
<b>Total Polychates N/M<sup>2</sup></b>	10	90	0	30	0	NS	
<b>Un identified Nematode worms</b>	40	200	0	10	30	NS	
<b>Amhipods</b>	0	10	0	10	0	NS	
TOTAL Benthic Fauna NUMBER/ M <sup>2</sup>	50	300	0	50	30	NS	

NS : No sample

**Table # 15 BENTHIC FAUNA IN THE SAMPLING LOCATIONS IN DPT HARBOUR AREA CREEKS DURING NEAP TIDE IN AUGUST ,2021**

Benthic fauna	ABUNDANCE IN NO/M <sup>2</sup> DIFFERENT SAMPLING STATIONS REPRESENTATION BY GROUP						
	DPT HARBOUR			CREEKS			
POLYCHATES	KPT-1	KPT-2	KPT-3	KPT-4	KPT-5	KPT-6	
Family : Spionidae <i>Polydora sp..</i>	20	10	0	40	20	NS	
Family : Syllidae <i>Syllis sp.</i>	10	10	0	20	60	NS	
<b>Total Polychates N/M<sup>2</sup></b>	30	20	0	60	80	NS	
<b>Un identified Nematode worms</b>	40	30	10	40	40	NS	
<b>Amhipods</b>	10	10	0	10	20	NS	
<b>TOTAL Benthic Fauna NUMBER/ M<sup>2</sup></b>	80	60	10	110	140	NS	

NS : No sample

## 7. Meteorological Data

Automatic Weather station have been installed in SevaSadan -3 at the Deendayal Port which records the data on Temperature (°C), Humidity (%), Wind (mph), Dew Point (°C), Wind Direction (°), Pressure, Solar radiation, heat Index and UVI.

### Temperature

The mean day time temperature for Deendayal Port was 30.5 °C. The day-time maximum temperature was 34.1 °C. The mean night time temperature was 27.4 °C. The minimum mean night time temperature recorded was 26.1 °C.

### Air Pressure

The mean absolute air pressure for the month of August was 1004.7 hpa, whereas the mean relative pressure was 1001.2 hpa. The maximum absolute air pressure recorded for the month of August was 1008.3 hpa.

### Heat Index

The mean day-time heat index for the month of August was 34.7 °C. The maximum heat index recorded was 42°C.

### Solar Radiation

The mean Solar Radiation in August was 232.4 w/m<sup>2</sup>. The maximum solar radiation recorded in the month of August was 682.8 w/m<sup>2</sup>.

### Humidity

The mean day-time humidity was 73.0 % for the month of August and mean night time humidity was 83.2%. Maximum humidity recorded during day-time was 88.0 % and maximum humidity recorded during night-time was 90.0%.

### Wind Velocity and Wind Direction

The mean wind velocity for the entire month of August was 10.8 km/hour. Maximum wind velocity recorded was 34.9 Km/hr . The wind direction was mostly S to SW.

**Conclusive Summary and Remedial measures Suggested**

- The AAQ monitoring at six locations of Deendayal Port indicates that the mean PM<sub>10</sub> values at four locations viz. Coal storage area, Marine Bhavan and Oil Jetty area were found above the permissible standards (100 µg/m<sup>3</sup>) and PM<sub>2.5</sub> was above permissible limits at Coal storage location (Limit 60 µg/m<sup>3</sup>).
- Drinking water at all the twenty locations was found potable and was within permissible limits of BIS standards (IS 10500).
- Noise quality was also within the set permissible standards of an Industrial Area. The noise level observed during day time was >75 dB (A) and at night time was >70 dB (A) during the entire monitoring period.
- The sewage treated water of Deendayal Port Colony (Gopalpuri) was in line with the standards set by the Gujarat Pollution Control Board. The STP at Deendayal Port is not fully operational and STP at Vadinar Port was found non-operational.

**Reasons for higher Values of PM<sub>10</sub>**

- Large amount of coal is handled at Berth No. 6, 7, 8 and 9. The unloading of coal directly in the truck, using grabs cause coal to spread in air as well as coal dust to fall on ground. This settled coal dust again mixes with the air while trucks travel through it.
- Also, the coal laden trucks are not always covered with tarpaulin sheets and these results in spillage of coal from trucks/dumpers during its transit from vessel to yard or storage site. This also increased PM values around marine Bhavan & Coal storage area.

### **Remedial Measures**

The values of PM<sub>10</sub> during the month of August, 2021 were observed beyond the permissible limit at four locations mentioned above. Given below are the remedial measures suggest to minimize the Air pollution at Deendayal Port.

- Guidelines for Coal Handling by GPCB should be strictly followed. (<http://gpcb.gov.in/pdf/coal-handling-guidelines.pdf>)
- Sewage Treatment Plan at Vadinar Port is not working. Hence, it is recommended to commission the sewage treatment plant at Vadinar immediately.
- Except for the higher values of PM<sub>10</sub> at Coal storage site, Oil Jetty, Tuna Port and Marine Bhavan locations, the monitoring results for the present month suggest that the overall Environment Quality of Deendayal Port is satisfactory.

**SOURCE OF LITERATURE AND ADDITIONAL REFERENCE FOR ECOLOGICAL STUDY**

- 1) ALBERT WEST PHAL (1976) Protozoa Blackwell , London
- 2) BANERJEE R.K. (1989) Heavy metals and Benthic foraminiferal distribution along Bombay coast India. Studies in benthic foraminifera. *Tokyo University Press* Tokyo pp 151-157
- 3) Banse K (1995) Zooplankton: Pivotal role in the control of ocean production: I. Biomass and production. *ICES J Mar Sci* 52: 265–277.
- 4) Beaugrand G, and Ibanez F (2004) Monitoring marine plankton ecosystems. II: long-term changes in North Sea calanoid copepods in relation to hydroclimatic variability. *Inter Res Mar Ecol Prog Ser* 284:35-47.
- 5) DAY F. (1889) The fauna of British India Ceylon and Burma- Fishes Vol-1- Vol-2 *Taylor and Francis* London
- 6) DESIKACHARY V. (1989) Atlas of diatoms, Madras Science Foundation
- 7) DESIKACHARY V. (1959) Cyanophyta ICAP Monographs on Algae *Indian Council of Agricultural research* New Delhi
- 8) FAIZAYOUSIF AL-YAMANI & MARIA A. SABUROVA (2010) illustrative guide on the flagellates of Intertidal soft sediment *Kuwait Institute for scientific Research* Kuwait
- 9) FAIZAYOUSIF AL-YAMANI, VALERIYSKRYABIN, ALEKSANDRA GUBANOVA, SERGEY KHVOROV AND IRINA PRUSOVA (2011), Marine zooplankton Practical guide from North western Arabian gulf Vol-1 and vol-2 *Kuwait Institute for scientific Research* Kuwait
- 10) FAUVEL P. (1953), The fauna of India Annelida - Polychaeta *Indian Press* Allahabad
- 11) Gajbhiye SN, Nair VR, and Desai BN (1984). Diurnal variation of zooplankton in Malad creek, Bombay. *Indian Journal of Marine Science*. 13:75-79.
- 12) HAYWARD P.J AND RYLAND J.S. (1995) Handbook of Marine fauna of north –West Europe *oxford University Press* London
- 13) HIGGINS R.P. HAJAMARTHIEL Eds. (1998) Introduction to the study of Meio Fauna
- 14) HORACE G. BARBER AND ELIZABETH Y. HAWORTH (1981) A guide to the Morphology of DIATOMS FRUSTULES.
- 15) INGRAM HENDEY (1964) An introductory account of smaller Algae of British coastal waters part-V. Bacillariophyceae
- 16) JOHN H. WICKSTEAD (1965) an Introduction to the study of Tropical Plankton . *Hutchinson Tropical Monographs*
- 17) JOYOTHIBABU, R. MADHU, N.V. MAHESHWARAN, P.A., NAIR K.K.C., VENUGOPL, P. BALASUBRAMANIAN T. (2005) Dominance of Dinoflagellates in micro zooplankton communities in the oceanic region Bay of Bengal and Andaman sea *Current science* vol.84. 10<sup>th</sup> May 2003
- 18) KASTURIRANGAN L.R. (1963) A key for the identification of the Common Planktonic Copepoda of Indian Coastal water
- 19) Kusum KK, Vineetha G, Raveendran TV, Nair VR, Muraleedharan KR, Achuthankutty CT and Joseph T (2014) Chaetognath community and their responses to varying environmental factors in the northern Indian ocean. *J Plankton Res* 36(4): 1146- 1152.
- 20) Lalli CM and Parsons TR (1997) Biological Oceanography: An Introduction. DOI <https://doi.org/10.1016/B978-0-7506-3384-0.X5056-7>.
- 21) Madhupratap M (1978) Studies on ecology of zooplankton of Cochin backwaters. *Mahasagar Bull Nat Inst Oceanogr* 11: 45-56.
- 22) Madhupratap M (1979) Distribution, community structure and species succession of copepods from Cochin Backwaters. *Indian J Mar Sci* 8: 1-8.



- 23) Madhupratap M (1987) Status and strategy of zooplankton of tropical Indian estuaries: A review. Bull Plank SocJpn 34: 65-81.
- 24) Madhupratap M (1999) Free living copepods of the Arabian Sea, Distribution and Research Perspectives. I J Mar Sci 146-149.
- 25) Madhupratap M and Haridas P (1986) Epipelagic calanoid copepods of the northern Indian Ocean. OceanologicaActa 9(2):105-117.
- 26) MANAL AL-KANDARI, FAIZA Y. AL-YAMANI , KHOLOOD AL-RIFAIE ( 2009) Marine phytoplankton Atlas of Kuwait's water *Kuwait Institute for scientific Research*
- 27) MPEDA (1998) Commercial Fishes and shell fishes of India
- 28) NEWEL G.E. & NEWELL R.C. (1963) Marine plankton a Practical Guide Hutchinson Educational
- 29) NIGAM R.C. AND CHATURVEDIS.K. (2000) Foraminiferal Study from KharoCreek , Kachchh ( Gujarat) North west coast of *India*. *Indian Journal of marine science* Vol.29 133-189
- 30) OLAV GIERE (1993) Meio benthology , Microscopic Fauna in Aquatic Sediments m Springer London
- 31) PERRAGALLO( 1965) Diatomees marines de franceA. *Asher & Co.* Amsterdam
- 32) Robert P.. Higgins (Eds.), (1985) An introduction to the study of Meio fauna Smithsons Institution press Washington DC
- 33) STERRER W. STERRERC.S Eds. Marine fauna and flora of Bermuda A systematic Guide to the identification of Marine Organisms. *John Wiley and Sons*New York
- 34) Suresh Gandhi. M. (2009) Distribution of certain ecological parameters and Foraminiferaldistribution in the depositional environment of Pak strait east coast of India .*Indian J. of Marine Science* Vol.33 pp 287-295
- 35) Venktaraman (1993) A systematic account of some south Indian diatoms . Proceeding of Indian Academy of Science Vol.X No.6 Sec.B.

\*\*\*\*\*

# ENVIRONMENTAL MONITORING REPORT FOR DEENDAYAL PORT TRUST



REPORT NO. : DCPL/DPT/20-21/17  
Month : September 2021  
Issue No : 01  
Revision No : 00  
Prepared by : DETOX CORPORATION PVT. LTD., SURAT

## TABLE OF CONTENTS

<b>Sr. No.</b>	<b>Particulars</b>	<b>Page No.</b>
<b>1</b>	<b>Ambient Air Quality Monitoring.....</b>	<b>1 - 17</b>
<b>2</b>	<b>Drinking Water Quality Monitoring.....</b>	<b>18 - 28</b>
<b>3</b>	<b>Noise Monitoring.....</b>	<b>29</b>
<b>4</b>	<b>Soil Monitoring.....</b>	<b>30 - 31</b>
<b>5</b>	<b>Sewage Treatment Plant Monitoring.....</b>	<b>32 - 38</b>
<b>6</b>	<b>Marine Water Monitoring.....</b>	<b>39 - 79</b>
<b>7</b>	<b>Meteorological Observations.....</b>	<b>80</b>
<b>8</b>	<b>Conclusive Summary &amp; Remedial Measures .....</b>	<b>81-82</b>
	<b>References.....</b>	<b>83- 84</b>

## **Introduction**

Monitoring of various environmental aspects of the Deendayal port by M/s Detox Corporation Pvt. Ltd. has been carried out through collection of samples, analysis of the same, comparing results with respect to the national standards and any other relevant standards by GBCB/CPCB/MoEF to identify non conformity in the Environment of the Deendayal Port. The results shall address the identified impacts and suggest measures to minimize the environmental impact due to various operations at Deendayal Port.

The environmental monitoring is carried out as per the Environment Management and Monitoring Plan submitted by Detox Corporation Pvt. Ltd.

## **1. Ambient Air Quality Monitoring**

As per the Environmental Monitoring Plan of Deendayal Port Trust, Air monitoring was carried out at six identified locations at Deendayal Port and two locations at Vadinar Port.

### **1.1 Air Quality Monitoring Methodology**

Air quality is measured in all the stations, for 24 hour for Total Suspended Particulate Matter (TSPM), PM<sub>10</sub>, PM<sub>2.5</sub>, SO<sub>2</sub>, NO<sub>x</sub>, NH<sub>3</sub> & Benzene, and Grab-sampling for CO & CO<sub>2</sub> measurements. The Air samplers are operated for a period of 24 hours and after a continuous operation of 8 hours of the sampler, the reagents were replaced to obtain 3 samples per day for each parameter namely, SO<sub>2</sub>, NO<sub>x</sub>. The EPM 2000 filter paper and PTFE Membrane bound filter paper are used for a period of 24 hours to obtain one sample each of TSPM, PM<sub>10</sub> & PM<sub>2.5</sub>.

The AAQ samples are collected twice a week from all the eight locations as per the EMP.

**Environmental Monitoring Report of Deendayal Port Trust, SEPTEMBER-2021**

**1.2 Results**

The ambient air quality monitoring data for six stations, viz. Marine Bhavan, Oil Jetty, Port Colony, Gopalpuri Hospital, Tuna Port and Nr. Coal Storage Area for the month of September 2021 are given in Tables 1A to 6B. The ambient air quality monitoring data for two stations at Vadinar (Nr. Admin Building & Nr. Signal Building) are given in Tables 7A to 8B.

**Location 1: Marine Bhavan (AL1)**

Table 1 : Results of Air Pollutant Concentration at Marine Bhavan										
Parameter	Date	TSPM [µg/m3]	PM10 [µg/m3]	PM2.5 [µg/m3]	SO2 [µg/m3]		NOx [µg/m3]		NH3 [µg/m3]	
					8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
Sampling Period	-	24hr	24hr	24hr						
NAAQMS limit	-	NS	100 µg/m3	60 µg/m3		80 µg/m3		80 µg/m3		400 µg/m3
AL1 – 1	03.09.2021	412	314	76	3.96	5.13	10.80	29.22	5.36	9.53
					6.59		32.39		13.02	
					4.84		44.46		10.21	
AL1 – 2	08.09.2021	673	579	50	3.52	3.81	13.34	26.25	12.25	12.85
					3.08		49.54		12.76	
					4.84		15.88		13.53	
AL1 – 3	10.09.2021	706	552	53	3.08	4.25	11.43	17.78	15.32	16.17
					6.15		17.15		14.55	
					3.52		24.77		18.64	
AL1 – 4	15.09.2021	357	260	82	6.15	4.84	17.78	39.80	18.12	15.06
					3.52		52.72		16.34	
					4.84		48.91		10.72	
AL1 – 5	17.09.2021	297	178	89	1.76	4.40	17.78	28.58	16.34	15.49
					5.28		24.77		15.57	
					6.15		43.19		14.55	
AL1 - 6	22.09.2021	387	309	72	3.52	3.08	40.02	38.53	5.36	9.53
					3.96		45.10		11.23	
					1.76		30.49		12.00	
AL1 - 7	24.09.2021	288	176	67	3.08	3.52	12.70	19.27	20.42	21.95
					4.84		23.50		22.46	
					2.64		21.60		22.98	
AL1 – 8	28.09.2021	471	299	163	17.14	13.48	27.95	27.31	20.68	21.53
					18.90		33.66		19.66	
					4.40		20.33		24.25	
<b>Monthly Average</b>		449	333	81		5.31		28.34		15.26
<b>Standard Deviation</b>		160	153	36		3.37		7.89		4.73

NS: Not Specified

<b>Table 1B : Results of Air Pollutant Concentration at Marine Bhavan</b>					
<b>Parameter</b>	<b>Date</b>	<b>C<sub>6</sub>H<sub>6</sub> [µg/m<sup>3</sup>]</b>	<b>HC* ppm</b>	<b>CO [mg/m<sup>3</sup>]</b>	<b>CO<sub>2</sub> [ppm ]</b>
<b>Sampling Period</b>		<b>8 hr</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>
<b>NAAQMS limit</b>		<b>5.0 µg/m<sup>3</sup></b>	<b>NS</b>	<b>4.0 mg/m<sup>3</sup></b>	<b>NS</b>
AL1 – 1	03.09.2021	1.11	BDL	1.89	492
AL1 – 2	08.09.2021	1.2	BDL	1.92	499
AL1 – 3	10.09.2021	1.3	BDL	1.9	486
AL1 – 4	15.09.2021	1.22	BDL	1.86	496
AL1 – 5	17.09.2021	1.27	BDL	1.88	501
AL1 - 6	22.09.2021	1.16	BDL	1.79	492
AL1 – 7	24.09.2021	1.18	BDL	1.86	496
AL1 – 8	28.09.2021	1.22	BDL	1.92	488
Monthly Average		<b>1.21</b>	-	<b>1.88</b>	<b>494</b>
Standard Deviation		<b>0.06</b>	-	<b>0.04</b>	<b>5</b>

\* NMHC- Non- Methane Hydrocarbons

BDL- Below Detection Limit (Detection Limit – NMHC: 0.5ppm)

NS -Not Specified

At Marine Bhavan, the overall values of TSPM, PM<sub>10</sub>, PM<sub>2.5</sub>, SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> is attributed mainly by motor vehicle emission produced from various types of automobiles (both diesel and petrol driven). Moreover, the loading and unloading of Food Grains and Timber at Jetty no. 1 and 2 also contributes to the high levels of TSPM and PM<sub>10</sub>. The mean TSPM value at Marine Bhavan was 449 µg/m<sup>3</sup>, The mean PM<sub>10</sub> values were 333.0 µg/m<sup>3</sup>, which is above the permissible limit. PM<sub>2.5</sub> values were above the permissible limit (mean = 81 µg/m<sup>3</sup>). The average values of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were within the permissible limit. The average values of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were 5.31 µg/ m<sup>3</sup>, 28.34 µg/ m<sup>3</sup> & 15.26 µg/ m<sup>3</sup> respectively. These were within the permissible limit.

The levels of Benzene, Hydrocarbons (HC) and CO were within the permissible limit at Marine Bhavan. The mean Benzene concentration was 1.21 µg/m<sup>3</sup>, well below the permissible limit of 5.0 µg/m<sup>3</sup>. HC's were below the detectable limit and Carbon Monoxide concentration was 1.88 mg/m<sup>3</sup>, well below the permissible limit of 4.0 mg/m<sup>3</sup>.

**Environmental Monitoring Report of Deendayal Port Trust, SEPTEMBER-2021**

**Location 2: Oil Jetty (AL2)**

Table 2 : Results of Air Pollutant Concentration at Oil Jetty										
Parameters	Date	TSPM [µg/m3]	PM10 [µg/m3]	PM2.5 [µg/m3]	SO2 [µg/m3]		NOx [µg/m3]		NH3 [µg/m3]	
Sampling Period	-	24hr	24hr	24hr	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
NAAQMS limit	-	NS	100 µg/m3	60 µg/m3	-	80 µg/m3	-	80 µg/m3	-	400 µg/m3
AL2 – 1	03.09.2021	645	423	158	6.15	4.84	23.50	16.30	17.87	14.72
					3.52		13.34		16.08	
					4.84		12.07		10.21	
AL2 – 2	08.09.2021	697	594	45	4.40	4.40	12.07	37.26	13.53	13.87
					5.28		45.73		10.21	
					3.52		53.99		17.87	
AL2 – 3	10.09.2021	673	561	62	3.52	4.25	28.58	18.84	9.45	10.81
					4.40		21.60		13.02	
					4.84		6.35		9.96	
AL2 – 4	15.09.2021	604	481	103	2.20	3.52	28.58	42.56	16.59	16.93
					3.08		46.37		17.87	
					5.28		52.72		16.34	
AL2 – 5	17.09.2021	616	571	38	6.15	3.81	46.37	38.53	11.74	8.34
					3.08		55.89		6.13	
					2.20		13.34		7.15	
AL2 – 6	22.09.2021	673	563	102	2.20	4.54	57.16	51.66	10.47	9.02
					5.28		45.10		9.70	
					6.15		52.72		6.89	
AL2 – 7	24.09.2021	245	159	71	6.15	3.52	23.50	38.11	10.47	14.81
					3.08		50.81		15.32	
					1.32		40.02		18.64	
AL2 – 8	28.09.2021	280	178	82	4.40	8.94	15.88	18.42	13.02	15.06
					8.79		13.34		8.42	
					13.63		26.04		23.74	
<b>Monthly Average</b>		554	442	83		4.73		32.71		12.94
<b>Standard Deviation</b>		183	177	39		1.77		13.12		3.14

NS: Not Specified

<b>Table 2B : Results of Air Pollutant Concentration at Oil Jetty</b>					
<b>Parameter</b>	<b>Date</b>	<b>C<sub>6</sub>H<sub>6</sub> [µg/m<sup>3</sup>]</b>	<b>HC* ppm</b>	<b>CO [mg/m<sup>3</sup>]</b>	<b>CO<sub>2</sub> [ppm ]</b>
<b>Sampling Period</b>		<b>8 hr</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>
<b>NAAQMS limit</b>		<b>5.0 µg/m<sup>3</sup></b>	<b>NS</b>	<b>4.0 mg/m<sup>3</sup></b>	<b>NS</b>
AL2 -1	03.09.2021	1.12	BDL	1.56	490
AL2 -2	08.09.2021	1.16	BDL	1.62	488
AL2 -3	10.09.2021	1.06	BDL	1.66	496
AL2 -4	15.09.2021	1.13	BDL	1.72	501
AL2 -5	17.09.2021	1.23	BDL	1.76	490
AL2 -6	22.09.2021	1.06	BDL	1.7	488
AL2 -7	24.09.2021	1.19	BDL	1.68	486
AL2 -8	28.09.2021	1.22	BDL	1.74	493
<b>Monthly Average</b>		<b>1.15</b>	<b>-</b>	<b>1.68</b>	<b>492</b>
<b>Standard Deviation</b>		<b>0.07</b>	<b>-</b>	<b>0.07</b>	<b>5</b>

\* NMHC- Non- Methane Hydrocarbons

BDL- Below Detection Limit (Detection Limit – NMHC : 0.5ppm)

NS- Not Specified

The overall values of TSPM, PM<sub>10</sub>, PM<sub>2.5</sub>, SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> at Oil Jetty area was mainly by motor vehicle emission produced from various types of vehicles Oil Jetty Area. The mean TSPM values at Oil Jetty were 554 µg/m<sup>3</sup>. The mean PM<sub>10</sub> values were 442 µg/m<sup>3</sup>, which is above the permissible limit. PM<sub>2.5</sub> values were above the permissible limit (mean = 83 µg/m<sup>3</sup>). The average values of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were within the permissible limit; The mean concentration of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were 4.73 µg/m<sup>3</sup>, 32.71 µg/m<sup>3</sup> and 12.94 µg/m<sup>3</sup> respectively.

The levels of Benzene, Hydrocarbons (HC) and CO were within the permissible limit at Oil Jetty. The mean Benzene concentration was 1.15 µg/m<sup>3</sup>. Well below the permissible limit of 5.0 µg/m<sup>3</sup>, HC's were below the detectable limit and Carbon Monoxide concentration was 1.68 mg/m<sup>3</sup>, well below the permissible limit of 4.0 mg/m<sup>3</sup>.



**Location 3: Kandla Colony – Estate Office (AL-3)**

Table 3 : Results of Air Pollutant Concentration at Estate Office										
Parameters	Date	TSPM [µg/m <sup>3</sup> ]	PM10 [µg/m <sup>3</sup> ]	PM2.5 [µg/m <sup>3</sup> ]	SO <sub>2</sub> [µg/m <sup>3</sup> ]		NO <sub>x</sub> [µg/m <sup>3</sup> ]		NH <sub>3</sub> [µg/m <sup>3</sup> ]	
Sampling Period	-	24hr	24hr	24hr	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
NAAQMS limit	-	NS	100 µg/m <sup>3</sup>	60 µg/m <sup>3</sup>	-	80 µg/m <sup>3</sup>	-	80 µg/m <sup>3</sup>	-	400 µg/m <sup>3</sup>
AL3 – 1	03.09.2021	239	162	61	3.96	4.54	41.29	37.69	9.45	11.49
					4.40		38.11		11.49	
					5.28		33.66		13.53	
AL3 – 2	08.09.2021	412	288	97	2.64	3.37	20.96	35.36	12.00	12.59
					4.40		40.02		15.57	
					3.08		45.10		10.21	
AL3 – 3	10.09.2021	248	121	41	5.28	4.69	17.78	23.08	15.06	15.57
					5.71		28.58		16.08	
					3.08		22.87		15.57	
AL3 – 4	15.09.2021	195	123	68	4.40	3.08	15.88	17.15	12.00	9.36
					1.76		11.43		7.15	
					3.08		24.14		8.93	
AL3 – 5	17.09.2021	256	194	55	3.08	3.22	32.39	31.55	16.59	14.81
					2.20		30.49		16.85	
					4.40		31.76		10.98	
AL3 – 6	22.09.2021	554	153	39	3.52	3.52	24.14	35.78	18.12	15.06
					3.08		40.02		12.25	
					3.96		43.19		14.81	
AL3 – 7	24.09.2021	467	399	52	2.64	3.08	27.31	34.72	5.36	6.72
					1.76		32.39		6.38	
					4.84		44.46		8.42	
AL3 – 8	28.09.2021	355	253	64	8.79	16.41	7.62	14.19	14.04	10.55
					34.73		14.61		6.13	
					5.71		20.33		11.49	
<b>Monthly Average</b>		341	211	60		5.24		28.69		12.02
<b>Standard Deviation</b>		128	96	18		4.56		9.22		3.11

NS: Not Specified

<b>Table 3B : Results of Air Pollutant Concentration at Kandla Port Colony</b>					
<b>Parameter</b>	<b>Date</b>	<b>C<sub>6</sub>H<sub>6</sub> [µg/m<sup>3</sup>]</b>	<b>HC*</b>	<b>CO [mg/m<sup>3</sup>]</b>	<b>CO<sub>2</sub> [ppm ]</b>
<b>Sampling Period</b>		<b>8 hr</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>
<b>NAAQMS limit</b>		<b>5.0 µg/m<sup>3</sup></b>	<b>NS</b>	<b>4.0 mg/m<sup>3</sup></b>	<b>NS</b>
AL3 -1	03.09.2021	1.1	BDL	1.77	492
AL3 -2	08.09.2021	1.06	BDL	1.82	480
AL3 -3	10.09.2021	1.11	BDL	1.86	479
AL3 -4	15.09.2021	1.16	BDL	1.8	482
AL3 -5	17.09.2021	1.18	BDL	1.92	477
AL3 -6	22.09.2021	1.26	BDL	1.96	486
AL3 -7	24.09.2021	1.22	BDL	1.86	478
AL3 -8	28.09.2021	1.21	BDL	1.78	482
<b>Monthly Average</b>		<b>1.16</b>	-	<b>1.85</b>	<b>482</b>
<b>Standard Deviation</b>		<b>0.07</b>	-	<b>0.07</b>	<b>5</b>

\* NMHC- Non- Methane Hydrocarbons

BDL- Below Detection Limit ( Detection Limit – NMHC : 0.5 ppm)

NS- Not Specified

The overall values of TSPM, PM<sub>10</sub>, PM<sub>2.5</sub>, SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> at Kandla Port Colony was attributed by vehicle emission produced from trucks and heavy duty vehicles that pass through the road outside Kandla Port Colony. The mean TSPM values at Oil Jetty were 341 µg/m<sup>3</sup>, The mean PM<sub>10</sub> values were 211 µg/m<sup>3</sup>, which is above the permissible limit. PM<sub>2.5</sub> values within the permissible limit (mean = 60 µg/m<sup>3</sup>). The average values of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were 5.24 µg/m<sup>3</sup>, 28.69 µg/m<sup>3</sup> and 12.02 µg/m<sup>3</sup> respectively and were all within the permissible limit.

The levels of Benzene, Hydrocarbons (HC) and CO were within the permissible limit at Kandla Port Colony. The mean Benzene concentration was 1.16 µg/m<sup>3</sup>, well below the permissible limit of 5.0 µg/m<sup>3</sup>. HC's were below the detectable limit and Carbon Monoxide concentration was 1.85 mg/m<sup>3</sup>, well below the permissible limit of 4.0 mg/m<sup>3</sup>.

**Location 4: Gopalpuri Hospital (AL-4)**

Table 4 : Results of Air Pollutant Concentration at Gopalpuri Hospital										
Parameter	Date	TSPM [µg/m <sup>3</sup> ]	PM10 [µg/m <sup>3</sup> ]	PM2.5 [µg/m <sup>3</sup> ]	SO <sub>2</sub> [µg/m <sup>3</sup> ]		NO <sub>x</sub> [µg/m <sup>3</sup> ]		NH <sub>3</sub> [µg/m <sup>3</sup> ]	
Sampling Period	-	24hr	24hr	24hr	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
NAAQMS limit	-	NS	100 µg/m <sup>3</sup>	60 µg/m <sup>3</sup>	-	80 µg/m <sup>3</sup>	-	80 µg/m <sup>3</sup>	-	400 µg/m <sup>3</sup>
AL4 -1	03.09.2021	167	118	37	4.40	3.52	13.34	14.40	8.42	6.30
					2.64		23.50		5.36	
					3.52		6.35		5.11	
AL4 -2	08.09.2021	256	178	63	3.08	3.66	13.34	21.17	8.42	8.85
					1.76		36.84		5.36	
					6.15		13.34		12.76	
AL4 -3	10.09.2021	165	122	26	1.32	2.34	30.49	30.06	12.25	9.36
					3.96		36.20		8.17	
					1.76		23.50		7.66	
AL4 -4	15.09.2021	189	124	54	3.08	4.54	48.91	41.29	5.62	8.59
					4.40		40.02		9.45	
					6.15		34.93		10.72	
AL4 -5	17.09.2021	185	104	48	2.20	4.40	11.43	24.56	12.00	11.15
					6.15		22.23		7.91	
					4.84		40.02		13.53	
AL4 -6	22.09.2021	249	101	45	1.76	3.08	17.15	14.40	9.19	8.59
					3.08		12.07		6.89	
					4.40		13.97		9.70	
AL4 -7	24.09.2021	167	116	43	2.20	2.49	24.14	16.94	9.70	11.83
					0.88		15.88		13.53	
					4.40		10.80		12.25	
AL4 -8	28.09.2021	177	122	48	0.88	0.88	5.08	6.14	5.87	5.87
					1.32		5.72		7.15	
					0.44		7.62		4.60	
<b>Monthly Average</b>		194	123	46		3.11		21.12		8.82
<b>Standard Deviation</b>		37	24	11		1.20		10.89		2.07

NS: Not Specified

<b>Table 4B : Results of Air Pollutant Concentration at Gopalpuri Hospital</b>					
<b>Parameter</b>	<b>Date</b>	<b>C<sub>6</sub>H<sub>6</sub> [µg/m<sup>3</sup>]</b>	<b>HC*</b>	<b>CO [mg/m<sup>3</sup>]</b>	<b>CO<sub>2</sub> [ppm ]</b>
<b>Sampling Period</b>		<b>8 hr</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>
<b>NAAQMS limit</b>		<b>5.0 µg/m<sup>3</sup></b>	<b>NS</b>	<b>4.0 mg/m<sup>3</sup></b>	<b>NS</b>
AL4 -1	03.09.2021	1.01	BDL	1.76	485
AL4 -2	08.09.2021	1.1	BDL	1.62	480
AL4 -3	10.09.2021	1.06	BDL	1.7	490
AL4 -4	15.09.2021	1.11	BDL	1.59	494
AL4 -5	17.09.2021	1.18	BDL	1.7	486
AL4 -6	22.09.2021	1.16	BDL	1.81	490
AL4 -7	24.09.2021	1.08	BDL	1.79	487
AL4 -8	28.09.2021	1.06	BDL	1.73	497
<b>Monthly Average</b>		<b>1.10</b>	<b>-</b>	<b>1.71</b>	<b>489</b>
<b>Standard Deviation</b>		<b>0.06</b>	<b>-</b>	<b>0.08</b>	<b>5</b>

\* NMHC- Non- Methane Hydrocarbons

BDL- Below Detection Limit (Detection Limit – NMHC: 0.5 ppm)

NS-Not Specified

The overall values of TSPM, PM<sub>10</sub>, PM<sub>2.5</sub>, SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> at Gopalpuri Hospital was attributed by vehicle emission produced from light motor vehicles of the colony residents. The mean TSPM values at Oil Jetty were 194 µg/m<sup>3</sup>, The mean PM<sub>10</sub> values were 123 µg/m<sup>3</sup>, which is above the permissible limit. PM<sub>2.5</sub> values were in within the permissible limit (mean= 46 µg/m<sup>3</sup>). The average values of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were 3.11 µg/m<sup>3</sup>, 21.12 µg/m<sup>3</sup> and 8.82 µg/m<sup>3</sup> respectively and were all within the permissible limit.

The levels of Benzene, Hydrocarbons (HC) and CO were within the permissible limit at Gopalpuri Hospital. The mean Benzene concentration was 1.10 µg/m<sup>3</sup>, well below the permissible limit of 5.0 µg/m<sup>3</sup>. HC's were below the detectable limit and Carbon Monoxide concentration was 1.71 mg/m<sup>3</sup>, well below the permissible limit of 4.0 mg/m<sup>3</sup>.

**Location 5: Coal Storage Area (AL-5)**

Table 5 : Results of Air Pollutant Concentration at Coal Storage Area										
Parameters	Date	TSPM [µg/m3]	PM10 [µg/m3]	PM2.5 [µg/m3]	SO2 [µg/m3]		NOx [µg/m3]		NH3 [µg/m3]	
Sampling Period	-	24hr	24hr	24hr	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
NAAQMS limit	-	NS	100 µg/m3	60 µg/m3	-	80 µg/m3	-	80 µg/m3	-	400 µg/m3
AL5 – 1	03.09.2021	380	115	86	3.52	4.10	32.39	30.28	13.79	14.30
					3.96		37.47		12.25	
					4.84		20.96		16.85	
AL5 – 2	08.09.2021	275	176	80	2.64	4.40	18.42	30.06	6.38	5.79
					6.59		33.66		5.11	
					3.96		38.11		5.87	
AL5 – 3	10.09.2021	302	225	74	2.20	3.66	52.72	48.91	9.45	10.64
					4.84		31.12		6.38	
					3.96		62.88		16.08	
AL5 – 4	15.09.2021	378	242	97	3.08	4.98	13.34	31.33	9.70	10.72
					5.28		50.81		12.76	
					6.59		29.85		9.70	
AL5 – 5	17.09.2021	210	138	70	4.84	4.10	12.07	27.31	9.70	10.98
					3.52		48.91		10.21	
					3.96		20.96		13.02	
AL5 – 6	22.09.2021	402	305	92	5.28	5.71	19.05	28.37	14.55	12.08
					6.15		26.04		12.25	
					5.71		40.02		9.45	
AL5 – 7	24.09.2021	268	151	73	2.64	4.69	32.39	33.66	18.64	17.61
					4.84		31.76		16.08	
					6.59		36.84		18.12	
AL5 – 8	28.09.2021	375	248	70	5.71	6.15	26.04	18.00	15.32	16.00
					6.15		5.72		19.91	
					6.59		22.23		12.76	
<b>Monthly Average</b>		324	200	80		4.73		30.99		12.26
<b>Standard Deviation</b>		69	65	10		0.96		8.61		3.69

NS: Not Specified

<b>Table 5B : Results of Air Pollutant Concentration at Coal Storage Area</b>					
<b>Parameter</b>	<b>Date</b>	<b>C<sub>6</sub>H<sub>6</sub> [µg/m<sup>3</sup>]</b>	<b>HC*</b>	<b>CO [mg/m<sup>3</sup>]</b>	<b>CO<sub>2</sub> [ppm ]</b>
<b>Sampling Period</b>		<b>8 hr</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>
<b>NAAQMS limit</b>		<b>5.0 µg/m<sup>3</sup></b>	<b>NS</b>	<b>4.0 mg/m<sup>3</sup></b>	<b>NS</b>
AL5 – 1	03.09.2021	1.22	BDL	1.9	489
AL5 – 2	08.09.2021	1.26	BDL	1.86	499
AL5 – 3	10.09.2021	1.3	BDL	1.79	501
AL5 – 4	15.09.2021	1.22	BDL	1.88	486
AL5 – 5	17.09.2021	1.21	BDL	1.86	488
AL5 – 6	22.09.2021	1.35	BDL	1.8	492
AL5 – 7	24.09.2021	1.34	BDL	1.92	496
AL5 – 8	28.09.2021	1.30	BDL	1.93	502
<b>Monthly Average</b>		<b>1.28</b>	<b>-</b>	<b>1.87</b>	<b>494</b>
<b>Standard Deviation</b>		<b>0.06</b>	<b>-</b>	<b>0.05</b>	<b>6</b>

\* NMHC- Non- Methane Hydrocarbons

BDL- Below Detection Limit (Detection Limit – NMHC: 0.5 ppm)

NS-Not Specified

The overall values of TSPM, PM<sub>10</sub>, PM<sub>2.5</sub>, SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> at Coal Storage Area was comparatively highest among all the locations of Air Quality monitoring in Kandla Port. High values of TSPM, PM<sub>10</sub>, PM<sub>2.5</sub>, SO<sub>2</sub>, NO<sub>x</sub> at this location was due to lifting of coal with grab and other coal handling processes near Berth no. 6 & 7. Moreover, the traffic was also heavy around this place for transport of coal thus emissions produced from heavy vehicles. The mean TSPM values at Coal storage were 324 µg/m<sup>3</sup>. The mean PM<sub>10</sub> values were 200 µg/m<sup>3</sup>, which is well above the permissible limit. PM<sub>2.5</sub> values were above the permissible limit (mean = 80 µg/m<sup>3</sup>). The average values of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were 4.73 µg/m<sup>3</sup>, 30.99 µg/m<sup>3</sup> and 12.26 µg/m<sup>3</sup> respectively and were all within the permissible limit.

The levels of Benzene, Hydrocarbons (HC) and CO were within the permissible limit at Coal Storage Area. The mean Benzene concentration was 1.28 µg/m<sup>3</sup>, well below the permissible limit of 5.0 µg/m<sup>3</sup>. HC's were below the detectable limit and Carbon Monoxide concentration was 1.87 mg/m<sup>3</sup>, well below the permissible limit of 4.0 mg/m<sup>3</sup>.

**Environmental Monitoring Report of Deendayal Port Trust, SEPTEMBER-2021**

**Location 6: Tuna Port (AL-6)**

Table 6 : Results of Air Pollutant Concentration at Tuna Port										
Parameters	Date	TSPM [µg/m3]	PM10 [µg/m3]	PM2.5 [µg/m3]	SO2 [µg/m3]		NOx [µg/m3]		NH3 [µg/m3]	
Sampling Period	-	24hr	24hr	24hr	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
NAAQMS limit	-	NS	100 µg/m3	60 µg/m3	-	80 µg/m3	-	80 µg/m3	-	400 µg/m3
AL6 -1	03.09.2021	186	104	52	7.03	5.13	12.70	38.11	14.30	16.34
					4.40		57.16		16.85	
					3.96		44.46		17.87	
AL6 - 2	08.09.2021	253	123	75	4.40	4.69	11.43	17.36	6.38	10.64
					6.15		18.42		14.04	
					3.52		22.23		11.49	
AL6 - 3	10.09.2021	214	128	57	3.52	3.66	25.41	23.29	9.96	13.87
					5.28		32.39		21.70	
					2.20		12.07		9.96	
AL6 - 4	15.09.2021	166	108	49	2.20	4.54	30.49	20.96	9.70	9.70
					4.84		19.05		9.19	
					6.59		13.34		10.21	
AL6 - 5	17.09.2021	253	177	50	3.08	4.69	52.72	45.52	12.25	12.51
					4.84		45.10		15.57	
					6.15		38.74		9.70	
AL6 - 6	22.09.2021	441	135	49	1.32	3.22	27.31	31.97	13.02	13.79
					3.08		38.74		15.57	
					5.28		29.85		12.76	
AL6 - 7	24.09.2021	216	130	46	3.08	3.81	33.66	40.44	16.08	12.00
					5.28		44.46		10.47	
					3.08		43.19		9.45	
AL6 - 8	28.09.2021	179	106	62	1.76	3.37	13.34	7.20	5.87	7.23
					3.96		4.45		5.36	
					4.40		3.81		10.47	
<b>Monthly Average</b>		238	126	55		4.14		28.11		12.01
<b>Standard Deviation</b>		88	24	10		0.71		13.08		2.82

NS: Not Specified

<b>Table 6B : Results of Air Pollutant Concentration at Tuna Port</b>					
<b>Parameter</b>	<b>Date</b>	<b>C<sub>6</sub>H<sub>6</sub> [µg/m<sup>3</sup>]</b>	<b>HC*</b>	<b>CO [mg/m<sup>3</sup>]</b>	<b>CO<sub>2</sub> [ppm ]</b>
<b>Sampling Period</b>		<b>8 hr</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>
<b>NAAQMS limit</b>		<b>5.0 µg/m<sup>3</sup></b>	<b>NS</b>	<b>4.0 mg/m<sup>3</sup></b>	<b>NS</b>
AL6 -1	03.09.2021	1.19	BDL	1.86	478
AL6 – 2	08.09.2021	1.16	BDL	1.92	492
AL6 – 3	10.09.2021	1.21	BDL	1.78	486
AL6 – 4	15.09.2021	1.06	BDL	1.92	482
AL6 – 5	17.09.2021	1.1	BDL	1.86	478
AL6 – 6	22.09.2021	1.02	BDL	1.8	492
AL6 – 7	24.09.2021	1.21	BDL	1.79	488
AL6 – 8	28.09.2021	1.2	BDL	1.86	478
<b>Monthly Average</b>		<b>1.14</b>	<b>-</b>	<b>1.85</b>	<b>484</b>
<b>Standard Deviation</b>		<b>0.07</b>	<b>-</b>	<b>0.05</b>	<b>6</b>

\* NMHC- Non- Methane Hydrocarbons

BDL- Below Detection Limit (Detection Limit – NMHC : 0.5 ppm)

NS- Not Specified

The mean TSPM values at Tuna Port were 238 µg/m<sup>3</sup>, The mean PM<sub>10</sub> values were 126 µg/m<sup>3</sup>, which is above the permissible limit. PM<sub>2.5</sub> values were within the permissible limit (mean = 55 µg/m<sup>3</sup>). The average values of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were 4.14 µg/m<sup>3</sup>, 28.11 µg/m<sup>3</sup> and 12.01 µg/m<sup>3</sup> respectively and were all within the permissible limit.

The levels of Benzene, Hydrocarbons (HC) and CO were within the permissible limit at Tuna Port. The mean Benzene concentration was 1.14 µg/m<sup>3</sup>, well below the permissible limit of 5.0 µg/m<sup>3</sup>. HC's were below the detectable limit and Carbon Monoxide concentration was 1.85 mg/m<sup>3</sup>, well below the permissible limit of 4.0 mg/m<sup>3</sup>.



**Location 7: Signal Building (Vadinar) (AL-7)**

Table 7 : Results of Air Pollutant Concentration at Signal Building										
Parameters	Date	TSPM [µg/m <sup>3</sup> ]	PM10 [µg/m <sup>3</sup> ]	PM2.5 [µg/m <sup>3</sup> ]	SO <sub>2</sub> [µg/m <sup>3</sup> ]		NO <sub>x</sub> [µg/m <sup>3</sup> ]		NH <sub>3</sub> [µg/m <sup>3</sup> ]	
					8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
NAAQMS limit	-	NS	100 µg/m <sup>3</sup>	60 µg/m <sup>3</sup>	-	80 µg/m <sup>3</sup>	-	80 µg/m <sup>3</sup>	-	400 µg/m <sup>3</sup>
AL7 -1	03.09.2021	158	88	43	3.96	12.60	9.53	10.80	5.62	6.21
					30.77		8.89		4.60	
					3.08		13.97		8.42	
AL7 -2	08.09.2021	158	93	59	3.96	3.52	13.97	14.61	9.96	9.28
					4.40		17.78		6.64	
					2.20		12.07		11.23	
AL7 -3	10.09.2021	180	108	53	3.08	3.22	19.05	12.28	5.62	4.51
					3.52		10.80		4.85	
					3.08		6.99		3.06	
AL7 -4	15.09.2021	169	102	33	5.28	4.25	13.97	12.70	6.13	8.85
					3.52		10.16		9.96	
					3.96		13.97		10.47	
AL7 -5	17.09.2021	160	87	27	5.28	3.66	10.80	9.74	8.42	5.87
					2.64		8.26		5.62	
					3.08		10.16		3.57	
AL7 -6	22.09.2021	177	95	64	3.52	4.40	13.97	8.79	10.47	9.87
					3.96		10.80		9.96	
					5.71		1.59		9.19	
AL7 -7	24.09.2021	139	94	32	2.20	3.52	13.97	12.91	6.38	6.72
					4.40		12.70		8.42	
					3.96		12.07		5.36	
AL7 -8	28.09.2021	168	107	43	2.64	3.08	14.61	12.49	8.68	7.40
					3.08		8.89		6.13	
					3.52		13.97		7.40	
<b>Monthly Average</b>		164	97	44		5		12		7
<b>Standard Deviation</b>		13	8	13		3		2		2

NS: Not Specified

<b>Table 7B : Results of Air Pollutant Concentration at Signal Building</b>					
<b>Parameter</b>	<b>Date</b>	<b>C<sub>6</sub>H<sub>6</sub> [µg/m<sup>3</sup>]</b>	<b>HC*</b>	<b>CO [mg/m<sup>3</sup>]</b>	<b>CO<sub>2</sub> [ppm ]</b>
<b>Sampling Period</b>		<b>8 hr</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>
<b>NAAQMS limit</b>		<b>5.0 µg/m<sup>3</sup></b>	<b>NS</b>	<b>4.0 mg/m<sup>3</sup></b>	<b>NS</b>
AL7 -1	03.09.2021	1.12	BDL	1.8	456
AL7 – 2	08.09.2021	1.06	BDL	1.78	462
AL7 – 3	10.09.2021	1.11	BDL	1.86	470
AL7 – 4	15.09.2021	1.18	BDL	1.8	455
AL7 – 5	17.09.2021	1.25	BDL	1.72	469
AL7 – 6	22.09.2021	1.16	BDL	1.68	460
AL7 – 7	24.09.2021	1.2	BDL	1.77	463
AL7 – 8	28.09.2021	1.26	BDL	1.7	460
<b>Monthly Average</b>		<b>1.17</b>	<b>-</b>	<b>1.76</b>	<b>462</b>
<b>Standard Deviation</b>		<b>0.07</b>	<b>-</b>	<b>0.06</b>	<b>5</b>

\* NMHC- Non- Methane Hydrocarbons

BDL- Below Detection Limit (Detection Limit – NMHC : 0.5 ppm)

NS\_ Not Specified

The mean TSPM values at Vadinar Port were 164 µg/m<sup>3</sup>. The mean PM<sub>10</sub> values were 97 µg/m<sup>3</sup>, which is below the permissible limit. PM<sub>2.5</sub> values were also within the permissible limit (mean = 44 µg/m<sup>3</sup>). The average values of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were 5.0 µg/m<sup>3</sup>, 12.0 µg/m<sup>3</sup> and 7.0 µg/m<sup>3</sup> respectively and were all within the permissible limit.

The levels of Benzene, Hydrocarbons (HC) and CO were within the permissible limit at Vadinar Port. The mean Benzene concentration was 1.17 µg/m<sup>3</sup>, well below the permissible limit of 5.0 µg/m<sup>3</sup>. HC's were below the detectable limit and Carbon Monoxide concentration was 1.76 mg/m<sup>3</sup>, well below the permissible limit of 4.0 mg/m<sup>3</sup>.

**Location 8: Admin Building (Vadinar) (AL-8)**

<b>Table 8 : Results of Air Pollutant Concentration at Admin Building</b>										
Parameters	Date	TSPM [µg/m3]	PM10 [µg/m3]	PM2.5 [µg/m3]	SO2 [µg/m3]		NOx [µg/m3]		NH3 [µg/m3]	
Sampling Period	-	24hr	24hr	24hr	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
NAAQMS limit	-	NS	100 µg/m3	60 µg/m3	-	80 µg/m3	-	80 µg/m3	-	400 µg/m3
AL8 -1	03.09.2021	164	83	24	3.96	4.25	13.34	12.28	5.87	4.77
					3.96		12.07		5.87	
					4.84		11.43		2.55	
AL8 -2	08.09.2021	198	130	35	4.40	5.28	20.96	20.54	5.11	5.70
					6.15		17.78		4.85	
					5.28		22.87		7.15	
AL8 -3	10.09.2021	177	86	64	2.64	3.81	8.89	12.91	8.42	8.34
					3.96		13.97		9.19	
					4.84		15.88		7.40	
AL8 -4	15.09.2021	150	78	25	2.64	2.07	20.96	16.94	8.42	6.98
					3.08		17.78		4.08	
					0.48		12.07		8.42	
AL8 -5	17.09.2021	156	84	46	2.20	3.52	247.71	91.46	5.62	6.72
					3.96		12.70		6.89	
					4.40		13.97		7.66	
AL8 -6	22.09.2021	198	123	55	3.52	4.10	11.43	13.97	4.60	4.85
					4.40		14.61		4.34	
					4.40		15.88		5.62	
AL8 -5	24.09.2021	172	101	54	3.08	2.34	9.53	9.10	8.68	7.83
					3.52		6.99		11.23	
					0.44		10.80		3.57	
AL8-6	28.09.2021	135	79	34	4.84	5.57	7.62	9.95	3.57	6.30
					5.71		9.53		5.62	
					6.15		12.70		9.70	
<b>Monthly Average</b>		169	95	42		4		23		6
<b>Standard Deviation</b>		22	20	15		1		28		1

NS: Not Specified

<b>Table 8B : Results of Air Pollutant Concentration at Admin Building</b>					
<b>Parameter</b>	<b>Date</b>	<b>C<sub>6</sub>H<sub>6</sub> [µg/m<sup>3</sup>]</b>	<b>HC*</b>	<b>CO [mg/m<sup>3</sup>]</b>	<b>CO<sub>2</sub> [ppm ]</b>
<b>Sampling Period</b>		<b>8 hr</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>
<b>NAAQMS limit</b>		<b>5.0 µg/m<sup>3</sup></b>	<b>NS</b>	<b>4.0 mg/m<sup>3</sup></b>	<b>NS</b>
AL8 -1	03.09.2021	1.06	BDL	1.78	460
AL8-2	08.09.2021	1.1	BDL	1.8	472
AL8 -3	10.09.2021	1.02	BDL	1.68	460
AL8-4	15.09.2021	1.1	BDL	1.72	461
AL8 -5	17.09.2021	1.17	BDL	1.81	452
AL8-6	22.09.2021	1.06	BDL	1.76	460
AL8-7	24.09.2021	1.1	BDL	1.66	470
AL8-8	28.09.2021	1.11	BDL	1.6	465
Monthly Average		<b>1.09</b>	-	<b>1.73</b>	<b>463</b>
Standard Deviation		<b>0.04</b>	-	<b>0.07</b>	<b>6</b>

\* NMHC- Non- Methane Hydrocarbons

BDL- Below Detection Limit (Detection Limit – NMHC: 0.5 ppm)

NS-Not Specified

The overall values of TSPM, PM<sub>10</sub>, PM<sub>2.5</sub>, SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> at Admin Building Vadinar was comparatively low among all the locations of Air Quality monitoring in Kandla Port and Vadinar Port. The mean TSPM values at Vadinar Port were 169 µg/m<sup>3</sup>. The mean PM<sub>10</sub> values were 95 µg/m<sup>3</sup>, which is below the permissible limit. PM<sub>2.5</sub> values were also within the permissible limit (mean = 42.0 µg/m<sup>3</sup>). The average values of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were 4.0 µg/m<sup>3</sup>, 23.0 µg/m<sup>3</sup> and 6.0 µg/m<sup>3</sup> respectively and were all within the permissible limit.

The levels of Benzene, Hydrocarbons (HC) and CO were within the permissible limit at Admin Building, Vadinar Port. The mean Benzene concentration was 1.09 µg/m<sup>3</sup>, well below the permissible limit of 5.0 µg/m<sup>3</sup>. HC's were below the detectable limit and Carbon Monoxide concentration was 1.73 mg/m<sup>3</sup>, well below the permissible limit of 4.0 mg/m<sup>3</sup>.

#### **1.4 Observations and Conclusion**

During the monitoring period, the overall Ambient Air Quality of the port area was found to be well within the desired levels for various gaseous pollutants. However, Particulate matter as PM<sub>10</sub> and PM<sub>2.5</sub> was found to exceed the limits at locations like Near Coal storage area, Marine Bhavan, Estate Office , Tuna Port and Oil Jetty area.

## 2. Drinking Water Quality Monitoring

Drinking Water Quality Monitoring was carried out at twenty stations at Kandla, Vadinar & Township Area of Deendayal Port.

### 2.1 Drinking Water Monitoring Methodology

Drinking water samples were collected from 20 locations as prescribed in the tender document. Samples for physico-chemical analysis were collected in 1 liter carboys and samples for microbiological parameters were collected in sterilized bottles. These samples were then analyzed in laboratory for various drinking water parameters at Kandla Lab/Surat.

The Sampling and Analysis was done as per standard methods - IS 10500:2012. The water samples were analyzed for various parameters, viz. Color , Odor, Turbidity , Conductivity , pH , Chlorides , TDS, Total Hardness, Iron , Sulphate , Salinity , DO, BOD, Na, K, Ca, Mg, F, NO<sub>3</sub>, NO<sub>2</sub>, Mn, Cr-6, Cu, Cd, As, Hg, Pb, Zn, Bacterial Count (cfu) .

### 2.2 Results

The Drinking Water Quality monitoring data for 20 stations are given in below from table No. 9 to Table No. 15

**Table 9: Drinking Water Quality Monitoring Parameters for Nirman Building 1, P & C building & Main Gate (North) at Kandla**

Sr. No.	Parameter	Unit	Nirman Building 1	P & C building	Main Gate North	Acceptable Limits as per IS 10500 : 2012	Permissible Limits in the absence of Alternate Source as per IS 10500 : 2012
1	pH	pH Unit	7.36	7.31	7.4	6.5 to 8.5	6.5 to 8.5
2	Total Dissolved Solids	mg/l	1343	1312	1350	500	2000
3	Turbidity	NTU	0	1	1	1.0	5.0
4	Odor	-	Odorless	Odorless	Odorless	Agreeable	Agreeable
5	Color	Hazen Units	Colorless	Colorless	Colorless	5.0	15.0
6	Conductivity	µs/cm	2630	2600	2690	NS*	NS*
7	Biochemical Oxygen Demand	mg/l	<2	<2	<2	NS*	NS*
8	Chloride as Cl	mg/l	420.94	365.81	370.82	250.0	1000.0
9	Ca as Ca	mg/l	72.14	56.11	52.10	75.0	200.0
10	Mg as Mg	mg/l	85.05	72.90	65.61	30.0	100.0
11	Total Hardness	mg/l	350	300	270	200.0	600.0
12	Iron as Fe	mg/l	<0.01	<0.01	<0.01	0.3	No Relaxation
13	Fluorides as F	mg/l	0.35	0.54	0.21	1.0	1.5
14	Sulphate as SO <sub>4</sub>	mg/l	228	210	258	200.0	400
15	Nitrite as NO <sub>2</sub>	mg/l	<0.01	<0.01	<0.01	NS*	NS*
16	Nitrate as NO <sub>3</sub>	mg/l	6.27	8.10	13.38	45.0	No Relaxation
17	Salinity	%	0.76	0.66	0.67	NS*	NS*
18	Sodium as Na	mg/l	301	243	265	NS*	NS*
19	Potassium as K	mg/l	3.24	3.68	3.51	NS*	NS*
20	Manganese	mg/l	<0.04	<0.04	<0.04	0.1	0.3
21	Hexavalent Chromium	mg/l	<0.03	<0.03	<0.03	NS*	NS*
22	Copper	mg/l	<0.05	<0.05	<0.05	0.05	1.5
23	Cadmium	mg/l	<0.002	<0.002	<0.002	0.003	0.003
24	Arsenic	mg/l	<0.01	<0.01	<0.01	0.01	0.05
25	Mercury	mg/l	<0.001	<0.001	<0.001	0.001	0.001
26	Lead	mg/l	<0.01	<0.01	<0.01	0.01	0.01
27	Zinc	mg/l	<0.1	<0.1	<0.1	5.0	15.0
28	Bacterial Count	CFU/100ml	Absent	Absent	Absent	Absent	Absent

\*NS: Not Specified

**Environmental Monitoring Report of Deendayal Port Trust, SEPTEMBER-2021**

**Table 10: Drinking Water Quality Monitoring Parameters for Canteen, West Gate – I & Wharf Area at Kandla**

Sr. No.	Parameter	Unit	Canteen	West Gate – I	Wharf Area	Acceptable Limits as per IS 10500 : 2012	Permissible Limits in the absence of Alternate Source as per IS 10500 : 2012
1	pH	pH Unit	7.55	7.6	7.83	6.5 to 8.5	6.5 to 8.5
2	Total Dissolved Solids	mg/l	1390	1360	1500	500	2000
3	Turbidity	NTU	0	0	1	1.0	5.0
4	Odor	-	Odorless	Odorless	Odorless	Agreeable	Agreeable
5	Color	Hazen Units	Colorless	Colorless	Colorless	5.0	15.0
6	Conductivity	µs/cm	2700	2680	2950	NS*	NS*
7	Biochemical Oxygen Demand	mg/l	<2	<2	<2	NS*	NS*
8	Chloride as Cl	mg/l	430.96	360.80	380.85	250.0	1000.0
9	Ca as Ca	mg/l	60.12	56.11	60.12	75.0	200.0
10	Mg as Mg	mg/l	63.18	80.19	85.05	30.0	100.0
11	Total Hardness	mg/l	260	330	350	200.0	600.0
12	Iron as Fe	mg/l	<0.01	<0.01	<0.01	0.3	No Relaxation
13	Fluorides as F	mg/l	0.58	0.79	0.25	1.0	1.5
14	Sulphate as SO4	mg/l	164.4	282	276	200.0	400
15	Nitrite as NO2	mg/l	<0.01	<0.01	<0.01	NS*	NS*
16	Nitrate as NO3	mg/l	8.80	10.42	9.50	45.0	No Relaxation
17	Salinity	%	0.78	0.65	0.69	NS*	NS*
18	Sodium as Na	mg/l	274	251	263	NS*	NS*
19	Potassium as K	mg/l	4.23	3.88	4.21	NS*	NS*
20	Manganese	mg/l	<0.04	<0.04	<0.04	0.1	0.3
21	Hexavalent Chromium	mg/l	<0.03	<0.03	<0.03	NS*	NS*
22	Copper	mg/l	<0.05	<0.05	<0.05	0.05	1.5
23	Cadmium	mg/l	<0.002	<0.002	<0.002	0.003	0.003
24	Arsenic	mg/l	<0.01	<0.01	<0.01	0.01	0.05
25	Mercury	mg/l	<0.001	<0.001	<0.001	0.001	0.001
26	Lead	mg/l	<0.01	<0.01	<0.01	0.01	0.01
27	Zinc	mg/l	<0.1	<0.1	<0.1	5.0	15.0
28	Bacterial Count	CFU/100ml	Absent	Absent	Absent	Absent	Absent

\*NS: Not Specified

**Environmental Monitoring Report of Deendayal Port Trust, SEPTEMBER-2021**

**Table 11: Drinking Water Quality Monitoring Parameters for Sewa sadan – 3, Workshop I & Custom Building at Kandla**

Sr. No.	Parameter	Unit	SewaSadon – 3	Workshop	Custom Building	Acceptable Limits as per IS 10500 : 2012	Permissible Limits in the absence of Alternate Source as per IS 10500 : 2012
1	pH	pH Unit	7.61	7.57	7.45	6.5 to 8.5	6.5 to 8.5
2	Total Dissolved Solids	mg/l	1326	1320	1520	500	2000
3	Turbidity	NTU	2	1	0	1.0	5.0
4	Odor	-	Odorless	Odorless	Odorless	Agreeable	Agreeable
5	Color	Hazen Units	Colorless	Colorless	Colorless	5.0	15.0
6	Conductivity	µs/cm	2650	2610	3010	NS*	NS*
7	Biochemical Oxygen Demand	mg/l	<2	<2	<2	NS*	NS*
8	Chloride	mg/l	410.91	320.71	425.95	250.0	1000.0
9	Ca as Ca	mg/l	48.10	56.11	48.10	75.0	200.0
10	Mg as Mg	mg/l	77.76	82.62	77.76	30.0	100.0
11	Total Hardness	mg/l	320	340	320	200.0	600.0
12	Iron as Fe	mg/l	<0.01	<0.01	<0.01	0.3	No Relaxation
13	Fluorides	mg/l	0.16	0.20	0.77	1.0	1.5
14	Sulphate	mg/l	213.6	195.6	276	200.0	400
15	Nitrite	mg/l	<0.01	<0.01	<0.01	NS*	NS*
16	Nitrate	mg/l	13.73	10.21	12.88	45.0	No Relaxation
17	Salinity	%	0.74	0.58	0.77	NS*	NS*
18	Sodium as Na	mg/l	215	206	166	NS*	NS*
19	Potassium as K	mg/l	3.87	3.73	3.56	NS*	NS*
20	Manganese	mg/l	<0.04	<0.04	<0.04	0.1	0.3
21	Hexavalent Chromium	mg/l	<0.03	<0.03	<0.03	NS*	NS*
22	Copper	mg/l	<0.05	<0.05	<0.05	0.05	1.5
23	Cadmium	mg/l	<0.002	<0.002	<0.002	0.003	0.003
24	Arsenic	mg/l	<0.01	<0.01	<0.01	0.01	0.05
25	Mercury	mg/l	<0.001	<0.001	<0.001	0.001	0.001
26	Lead	mg/l	<0.01	<0.01	<0.01	0.01	0.01
27	Zinc	mg/l	<0.1	<0.1	<0.1	5.0	15.0
28	Bacterial Count	CFU/100ml	Absent	Absent	Absent	Absent	Absent

\*NS: Not Specified



**Table 12: Drinking Water Quality Monitoring Parameters for Port Colony Kandla, Hospital Kandla & A.O. Building at Gandhidham**

Sr. No.	Parameter	Unit	Port Colony Kandla	Hospital Kandla	A.O. Building	Acceptable Limits as per IS 10500 : 2012	Permissible Limits in the absence of Alternate Source as per IS 10500 : 2012
1	pH	pH Unit	7.53	7.56	7.59	6.5 to 8.5	6.5 to 8.5
2	Total Dissolved Solids	mg/l	1370	1350	1450	500	2000
3	Turbidity	NTU	0	0	1	1.0	5.0
4	Odor	-	Odorless	Odorless	Odorless	Agreeable	Agreeable
5	Color	Hazen Units	Colorless	Colorless	Colorless	5.0	15.0
6	Conductivity	µs/cm	2690	2700	2990	NS*	NS*
7	Biochemical Oxygen Demand	mg/l	<2	<2	<2	NS*	NS*
8	Chloride	mg/l	335.75	375.84	821.83	250.0	1000.0
9	Ca as Ca	mg/l	76.15	52.10	72.14	75.0	200.0
10	Mg as Mg	mg/l	87.48	70.47	92.34	30.0	100.0
11	Total Hardness	mg/l	360	290	380	200.0	600.0
12	Iron as Fe	mg/l	<0.01	<0.01	<0.01	0.3	No Relaxation
13	Fluorides	mg/l	0.48	0.14	0.81	1.0	1.5
14	Sulphate	mg/l	336	228	237.6	200.0	400
15	Nitrite	mg/l	<0.01	<0.01	<0.01	NS*	NS*
16	Nitrate	mg/l	9.50	10.21	11.62	45.0	No Relaxation
17	Salinity	%	0.61	0.68	1.48	NS*	NS*
18	Sodium as Na	mg/l	211	196	202	NS*	NS*
19	Potassium as K	mg/l	4.08	4.01	3.99	NS*	NS*
20	Manganese	mg/l	<0.04	<0.04	<0.04	0.1	0.3
21	Hexavalent Chromium	mg/l	<0.03	<0.03	<0.03	NS*	NS*
22	Copper	mg/l	<0.05	<0.05	<0.05	0.05	1.5
23	Cadmium	mg/l	<0.002	<0.002	<0.002	0.003	0.003
24	Arsenic	mg/l	<0.01	<0.01	<0.01	0.01	0.05
25	Mercury	mg/l	<0.001	<0.001	<0.001	0.001	0.001
26	Lead	mg/l	<0.01	<0.01	<0.01	0.01	0.01
27	Zinc	mg/l	<0.1	<0.1	<0.1	5.0	15.0
28	Bacterial Count	CFU/100ml	Absent	Absent	Absent	Absent	Absent

\*NS: Not Specified

**Table 13: Drinking Water Quality Monitoring Parameters for School Gopalpuri, Guest House & E - Type Quarter at Gopalpuri, Gandhidham**

Sr. No.	Parameter	Unit	School Gopalpuri	Guest House	E - Type Quarter	Acceptable Limits as per IS 10500 : 2012	Permissible Limits in the absence of Alternate Source as per IS 10500 : 2012
1	pH	pH Unit	7.64	7.61	7.69	6.5 to 8.5	6.5 to 8.5
2	Total Dissolved Solids	mg/l	1400	1850	1190	500	2000
3	Turbidity	NTU	2	0	1	1.0	5.0
4	Odor	-	Odorless	Odorless	Odorless	Agreeable	Agreeable
5	Color	Hazen Units	Colorless	Colorless	Colorless	5.0	15.0
6	Conductivity	µs/cm	2780	3670	2310	NS*	NS*
7	Biochemical Oxygen Demand	mg/l	<2	<2	<2	NS*	NS*
8	Chloride	mg/l	410.91	471.05	385.86	250.0	1000.0
9	Ca as Ca	mg/l	64.13	76.15	44.09	75.0	200.0
10	Mg as Mg	mg/l	85.05	99.63	80.19	30.0	100.0
11	Total Hardness	mg/l	350	410	330	200.0	600.0
12	Iron as Fe	mg/l	<0.01	<0.01	<0.01	0.3	No Relaxation
13	Fluorides	mg/l	0.34	0.22	0.48	1.0	1.5
14	Sulphate	mg/l	252	284.4	303.6	200.0	400
15	Nitrite	mg/l	<0.01	<0.01	<0.01	NS*	NS*
16	Nitrate	mg/l	9.50	10.70	9.50	45.0	No Relaxation
17	Salinity	%	0.74	0.85	0.70	NS*	NS*
18	Sodium as Na	mg/l	202	184	192	NS*	NS*
19	Potassium as K	mg/l	3.49	3.50	3.56	NS*	NS*
20	Manganese	mg/l	<0.04	<0.04	<0.04	0.1	0.3
21	Hexavalent Chromium	mg/l	<0.03	<0.03	<0.03	NS*	NS*
22	Copper	mg/l	<0.05	<0.05	<0.05	0.05	1.5
23	Cadmium	mg/l	<0.002	<0.002	<0.002	0.003	0.003
24	Arsenic	mg/l	<0.01	<0.01	<0.01	0.01	0.05
25	Mercury	mg/l	<0.001	<0.001	<0.001	0.001	0.001
26	Lead	mg/l	<0.01	<0.01	<0.01	0.01	0.01
27	Zinc	mg/l	<0.1	<0.1	<0.1	5.0	15.0
28	Bacterial Count	CFU/100ml	Absent	Absent	Absent	Absent	Absent

\*NS: Not Specified

**Table 14: Drinking Water Quality Monitoring Parameters for F - Type Quarter, Hospital Gopalpuri & Tuna Port**

Sr. No.	Parameter	Unit	F - Type Quarter	Hospital Gopalpuri	Tuna Port	Acceptable Limits as per IS 10500 : 2012	Permissible Limits in the absence of Alternate Source as per IS 10500 : 2012
1	pH	pH Unit	7.56	7.42	7.32	6.5 to 8.5	6.5 to 8.5
2	Total Dissolved Solids	mg/l	1230	1590	1020	500	2000
3	Turbidity	NTU	1	1	0	1.0	5.0
4	Odor	-	Odorless	Odorless	Odorless	Agreeable	Agreeable
5	Color	Hazen Unit	Colorless	Colorless	Colorless	5.0	15.0
6	Conductivity	µs/cm	2450	3150	2000	NS*	NS*
7	Biochemical Oxygen Demand	mg/l	<2	<2	<2	NS*	NS*
8	Chloride	mg/l	375.84	511.14	604	250.0	1000.0
9	Ca as Ca	mg/l	40.08	60.12	80.16	75.0	200.0
10	Mg as Mg	mg/l	85.05	97.20	60.75	30.0	100.0
11	Total Hardness	mg/l	350	400	330	200.0	600.0
12	Iron as Fe+3	mg/l	<0.01	<0.01	<0.01	0.3	No Relaxation
13	Fluorides	mg/l	0.59	0.36	0.46	1.0	1.5
14	Sulphate	mg/l	260.4	174	180	200.0	400
15	Nitrite	mg/l	<0.01	<0.01	<0.01	NS*	NS*
16	Nitrate	mg/l	10.00	7.96	8.2	45.0	No Relaxation
17	Salinity	%	0.68	0.92	1.09	NS*	NS*
18	Sodium as Na	mg/l	162	206	210	NS*	NS*
19	Potassium as K	mg/l	2.18	4.01	2.6	NS*	NS*
20	Manganese	mg/l	<0.04	<0.04	<0.04	0.1	0.3
21	Hexavalent Chromium	mg/l	<0.03	<0.03	<0.03	NS*	NS*
22	Copper	mg/l	<0.05	<0.05	<0.05	0.05	1.5
23	Cadmium	mg/l	<0.002	<0.002	<0.002	0.003	0.003
24	Arsenic	mg/l	<0.01	<0.01	<0.01	0.01	0.05
25	Mercury	mg/l	<0.001	<0.001	<0.001	0.001	0.001
26	Lead	mg/l	<0.01	<0.01	<0.01	0.01	0.01
27	Zinc	mg/l	<0.1	<0.1	<0.1	5.0	15.0
28	Bacterial Count	CFU/100ml	Absent	Absent	Absent	Absent	Absent

\*NS: Not Specified

**Table 15: Drinking Water Quality Monitoring Parameters for Vadinar Jetty & Port Colony at Vadinar**

Sr. No.	Parameter	Unit	Vadinar Jetty	Port Colony Vadinar	Acceptable Limits as per IS 10500 : 2012	Permissible Limits in the absence of Alternate Source as per IS 10500 : 2012
1	pH	pH Unit	7.5	7.4	6.5 to 8.5	6.5 to 8.5
2	Total Dissolved Solids	mg/l	1160	1150	500	2000
3	Turbidity	NTU	ND	ND	1.0	5.0
4	Odor	-	Odorless	Odorless	Agreeable	Agreeable
5	Color	Hazen Units	Colorless	Colorless	5.0	15.0
6	Conductivity	µs/cm	2390	2300	NS*	NS*
7	Biochemical Oxygen Demand	mg/l	<2	<2	NS*	NS*
8	Chloride	mg/l	425.95	415.92	250.0	1000.0
9	Ca as Ca	mg/l	56.11	64.13	75.0	200.0
10	Mg as Mg	mg/l	75.33	70.47	30.0	100.0
11	Total Hardness	mg/l	310	290	200.0	600.0
12	Iron as Fe+3	mg/l	<0.01	<0.01	0.3	No Relaxation
13	Fluorides	mg/l	0.71	0.62	1.0	1.5
14	Sulphate	mg/l	30.60	28.80	200.0	400
15	Nitrite	mg/l	<0.01	<0.01	NS*	NS*
16	Nitrate	mg/l	9.85	9.71	45.0	No Relaxation
17	Salinity	%	0.76	0.75	NS*	NS*
18	Sodium as Na	mg/l	192.0	183.0	NS*	NS*
19	Potassium as K	mg/l	2.2	2.7	NS*	NS*
20	Manganese	mg/l	<0.04	<0.04	0.1	0.3
21	Hexavalent Chromium	mg/l	<0.03	<0.03	NS*	NS*
22	Copper	mg/l	<0.05	<0.05	0.05	1.5
23	Cadmium	mg/l	<0.002	<0.002	0.003	0.003
24	Arsenic	mg/l	<0.01	<0.01	0.01	0.05
25	Mercury	mg/l	<0.001	<0.001	0.001	0.001
26	Lead	mg/l	<0.01	<0.01	0.01	0.01
27	Zinc	mg/l	<0.1	<0.1	5.0	15.0
28	Bacterial Count	CFU/100ml	Absent	Absent	Absent	Absent

\*NS: Not Specified

### 2.3 Results & Discussion

The colour of all drinking water samples was < 5 Hazen unit and odour of the samples was also agreeable. All parameters are found to be within the specified limit of the Drinking water Standard.

#### pH

The limit of pH value for drinking water is specified as 6.5 to 8.5. pH value in the studied area varied from 7.0 to 8.0 pH unit. All the sampling points showed pH values within the prescribed limit by Indian Standards.

#### Total Dissolved Solids (TDS)

TDS values in the studied area varied between 1000 -1800 mg/l. None of the sampling points showed higher TDS values than the prescribed limit by Indian standards .

#### Conductivity

Electrical Conductivity is the ability of a solution to transfer (conduct) electric current. Conductivity is used to measure the concentration of dissolved solids which have been ionized in a polar solution such as water. The conductivity in the samples collected during the month of September ranged from 2000-3700  $\mu\text{s}/\text{cm}$ . Electrical conductivity standards do not appear in BIS standards for drinking water.

#### BOD

BOD value in the studied area was less than 2.0 mg/L. Indian standards does not show any standard values for BOD in drinking water.

#### Chlorides

Excessive chloride concentration increase rates of corrosion of metals in the distribution system. This can lead to increased concentration of metals in the supply. Chloride value in the studied area varied between 400-900 mg/l and is found to be within the Permissible limit of the Drinking Water Standard.

#### Calcium

Calcium value in the studied area varied between 40 - 80 mg/l and is found to be within the Permissible limit of the Drinking Water Standard. If calcium is present beyond the maximum acceptable limit, it causes incrustation of pipes.

### **Magnesium**

Magnesium value in the studied area varied between 60 – 99 mg/l. All the locations had Magnesium within the prescribed limits of 30-100 mg/L.

### **Total Hardness**

Hardness value in the studied area varied between 260-410 mg/l and is found to be within the Permissible limit of the Drinking Water Standard. The prescribed limit by Indian Standards is 200-600 mg/L.

### **Iron**

Iron value in the studied area was below 0.01mg/L and hence well below the permissible limit as per Indian Standards is 0.3 mg/L. The excess amount of iron causes slight toxicity; gives stringent taste to water.

### **Fluoride**

Fluoride value in the studied area varied between 0.1 – 1.0 mg/l and hence well below the permissible limit as per Indian Standards is 1.0-1.5 mg/L. Moderate amounts lead to dental effects, but long-term ingestion of large amounts can lead to potentially severe skeletal problems.

### **Sulphates**

Sulphate value in the studied area varied between 30 – 350 mg/l. All the sampling points showed sulphate values within the prescribed limits by Indian Standards (200-400 mg/L). Sulphate content in drinking water exceeding the 400 mg/L imparts bitter taste.

### **Nitrites (NO<sub>2</sub>) and Nitrates (NO<sub>3</sub>)**

Nitrite values in all the water samples were <0.1. There are no specified standard values for Nitrites in Drinking water. The mean Nitrate values in drinking water of KPT was 6.27 mg/l which is well within the permissible limit of the Drinking water Standard.

### **Salinity**

Salinity in drinking water in the present samples collected ranged from 0.5 to 1.4 %. There are no prescribed Indian standards for salinity in Drinking water.

### **Sodium and Potassium Salts**

Sodium values in the samples collected ranged from 100 - 300 mg/l and Potassium salts ranged from 2.2 to 4.0 mg/l. There are no prescribed limits of Sodium and Potassium in Indian standards for Drinking water.

### **Heavy Metals in Drinking Water**

In the present study period drinking water samples were analyzed for Mn, Cr, Cu, Cd, As, Hg, Pb and Zn. All these heavy metals were well below the permissible limits prescribed by the Indian Standards.

### **Bacteriological Study**

Analysis of the bacteriological parameter at all location shows that Bacteria is not present and hence Bacterial count is in line with the permissible limit of drinking water. This shows that all the drinking water samples were safe from any bacteriological contamination.

## **2.4 Conclusions**

These results are compared with acceptable limits as prescribed in IS 10500:2012 – Drinking Water Specification. It is seen from the analysis data that during the study period the water was safe for human consumption at all drinking water monitoring stations.

### 3. Noise Level Monitoring

Noise sources in port operations include cargo handling, vehicular traffic, and loading / unloading containers and ships. Noise Monitoring was done at 13 stations at Kandla, Vadinar and Township area.

#### 3.1 Method of Monitoring

Sampling was done at all stations for 24 hour period. Data was recorded using automated sound level meter. The intensity of sound was measured in sound pressure level (SPL) and common unit of measurement is decibel (Db).

#### 3.2 Results

Table 16: Noise Monitoring data for ten locations of Deendayal Port and two locations of Vadinar Port

Sr. No.	Location	Day Time Average Noise Level (SPL) in dB(A)	Night Time Average Noise Level (SPL) in dB(A)
	Sampling Time	6:00 am to 10:00 PM	10:00PM to 6:00 AM
1	Marine Bhavan	58.0	50.4
2	Nirman Building 1	55.3	49.1
3	Tuna Port	52.8	46.5
4	Main Gate North	60.3	55.2
5	West Gate I	67.2	60.6
6	Canteen Area	58.7	50.9
7	Main Road	70.5	59.5
8	ATM Building	69.2	62.3
9	Wharf Area /Jetty Area	73.7	65.4
10	Port & Custom Office	55.2	49.6
	<b>Vadinar Port</b>		
11	Entrance Gate of Vadinar Port	69.6	58.4
12	Nr. Port Colony, Vadinar	61.3	55.8
13	Nr. Vadinar Jetty	68.2	61.5

**3.3 Conclusions-** Noise sources in port operations include cargo handling, vehicular traffic, and loading / unloading containers and ships. The Day Time Average Noise Level (SPL) in all 13 locations at Deendayal Port ranged from 52.0 dB(A) to 73.7 dB(A) and it was within the permissible limits of 75 dB(A) for the industrial area for the daytime. The Night Time Average Noise Level (SPL) in all 13 locations of Deendayal Port ranged from 46.5 dB to 65.4 dB(A) and it was within the permissible limits of 70 dB(A) for the industrial area for the night time.



#### 4. Soil Monitoring

Sampling and analysis of soil samples were undertaken at six locations within the study area (Deendayal Port and Vadinar Port) as a part of EMP. The soil sampling locations are initially decided based on the locations as provided in the tender document of the Deendayal Port.

##### 4.1 Methodology

The soil samples were collected in the month of September 2021. The samples collected from the all locations are homogeneous representative of each location. At random locations were identified at each location and soil was dug from 30 cm below the surface. It was uniformly mixed before homogenizing the soil samples. The samples were filled in polythene bags, labeled in the field with number and site name and sent to laboratory for analysis.

##### 4.2 Results

**Table-17: Chemical Characteristics of Soil in the Study Area**

Sr. No.	Parameter	Unit	Station Name					
			SL1	SL2	SL3	SL4	SL5	SL6
			Tuna Port	IFFCO Plant	Khori Creek	Nakti Creek	KPT Admin Site	KPT Colony
			Near main gate of Port	10 m away from main gate	Sand from creek at low tide		Vadinar	
1	Texture		Sandy Loam	Sandy Loam	Sandy Loam	Sandy Loam	Sandy Loam	Sandy Loam
2	pH	-	8.58	8.16	8.46	8.26	8.02	8.56
3	Electrical Conductivity	µs/cm	18,400.0	25,620.0	17,880.0	16,520.0	523.0	420.0
4	Moisture	%	21.00	22.20	24.10	18.80	8.66	9.02
5	Total Organic Carbon	%	0.48	1.24	0.48	3.93	0.18	0.21
6	Alkalinity	mg/kg	72.07	36.04	190.19	90.09	60.06	100.10
7	Total Nitrogen	%	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001
8	Chloride	mg/kg	1,506.6	6,381.0	1,701.0	1,878.9	52.0	67.8
9	Sulphate	mg/kg	202.0	196.0	112.0	112.0	12.0	18.0
10	Phosphorus	mg/kg	0.89	0.92	1.05	1.10	0.78	0.86
11	Potassium	mg/kg	386.0	820.0	345.0	422.0	110.0	172.0
12	Sodium	mg/kg	1,585.0	3,386.0	2,303.0	1,990.0	990.0	810.0
13	Calcium	mg/kg	228.46	741.50	248.50	468.94	118.00	72.00
14	Copper as Cu	mg/kg	52.2	78.2	46.2	33.8	18.6	28
15	Lead as Pb	mg/kg	4.9	5.6	3.2	4.8	3.2	1.1
16	Nickel as Ni	mg/kg	46.2	28	33.2	26.1	18.2	16.2
17	Zinc as Zn	mg/kg	66.20	41.60	68	49.55	24.00	38.50
18	Cadmium as Cd	mg/kg	BQL	BQL	BQL	BQL	BQL	BQL

### 4.3 Discussion

- The data shows that value of pH ranges from 8.02 at Nakti Creek to 8.58 at Tuna Creek indicating that all soil samples are neutral to slight basic. Iffco Plant samples showed maximum conductivity of 25,620 $\mu$ mhos/cm, while Nakti Creek location showed minimum conductivity of 16,520  $\mu$ mhos/cm. Conductivity at Vadinar Port was 523 and 420  $\mu$ mhos/cm at Admin site and Vadinar Port colony respectively.
- Total organic Carbon ranged from 0.1 % to 3.9 at Deendayal Port. At Vadinar Port, organic carbon content ranged from 0.1 % to 0.2 %.
- The concentration of Phosphorus and Potassium in the soil samples varies from 0.8 to 1.10 mg/kg and 300.0 to 800 mg/kg respectively at Deendayal Port. The mean concentration of Phosphorous at Vadinar site was 0.82 mg/kg and mean concentration of Potassium at Vadinar site was 145 mg/kg.

These differences in NPK in soil at different locations are due to the dissimilar nature of soil at each of the locations. Samples SL3 & SL4 (Khorī Creek & Nakti Creek) are of saline nature as they are coastal soil; where as other locations are inland locations and have different chemical properties.

### Heavy Metals in the Soil

Traces of Copper, Lead, Nickel and Zinc were observed in the soil samples collected from all the four locations of Deendayal Port and two locations of Vadinar Port. Cadmium metal was not detected in the Soil.

### 4.4 Conclusion

The soils of Deendayal Port and Vadinar Port appears to be neutral to basic with varying levels of Chloride, Sulphate, NPK and Calcium. As the nature of soil at different locations are different with respect to its proximity to the sea, the samples showed high degree of variations in their chemical properties.

## 5. Sewage Treatment Plant Monitoring

This involves safe collection of waste water (spent/used water) from wash areas, bathroom, industrial units, etc., waste from toilets of various buildings and its conveyance to the treatment plant and final disposal in conformity with the requirement and guide lines of State Pollution Control Board and other statutory bodies.

### 5.1 Methodology for STP Monitoring

To monitor the working efficiency of Sewage Treatment Plant (STP), STP Inlet and Outlet Samples were collected once a week. Locations selected are namely Gopalpuri Township, Deendayal Port and Vadinar. Samples were collected in 1 lit. Carboys and were analyzed in laboratory for various parameters.

### 5.2 Results

- **Kandla STP**

**Table 18: Sewage Water Monitoring at Kandla STP (1<sup>st</sup> Week)**

Date of Sampling		04.09.2021		
Sr. No.	Parameters	Unit	Results	
			KPT STP I/L	KPT STP O/L
1	pH	pH unit	7.56	7.44
2	Total Suspended Solids	mg/l	64.2	26.6
3	Residual Chlorine	mg/l	<1.0	<0.5
4	COD	mg/l	424.2	103.0
5	BOD @ 27 °C	mg/l	141.0	29.0
6.	Fecal Coliform	MPN Index / 100 ml	-	20.0
<b>Aeration Tank</b>				
7.	MLSS	mg/l	6.0	
8.	MLVSS	%	93.0	

**Table 19: Sewage Water Monitoring at Kandla STP (2<sup>nd</sup> Week)**

Date of Sampling		09.09.2021		
Sr. No.	Parameters	Unit	Results	
			KPT STP I/L	KPT STP O/L
1	pH	pH unit	7.6	7.2
2	Total Suspended Solids	mg/l	152.2	72.4
3	Residual Chlorine	mg/l	<1.0	<0.5
4	COD	mg/l	384	103.0
5	BOD @ 27 °C	mg/l	120.0	24.0
6.	Fecal Coliform	MPN Index / 100 ml	-	31.0
<b>Aeration Tank</b>				
7.	MLSS	mg/l	9.0	
8.	MLVSS	%	89.0	

**Table 20: Sewage Water Monitoring at Kandla STP (3<sup>rd</sup> Week)**

Date of Sampling		16.09.2021		
Sr. No.	Parameters	Unit	Results	
			KPT STP I/L	KPT STP O/L
1	pH	pH unit	7.71	7.18
2	Total Suspended Solids	mg/l	417.8	159.8
3	Residual Chlorine	mg/l	<1.0	<0.5
4	COD	mg/l	383.8	102
5	BOD @ 27 °C	mg/l	128.0	23.0
6.	Fecal Coliform	MPN Index / 100 ml	-	<1.8
<b>Aeration Tank</b>				
7.	MLSS	mg/l	9.0	
8.	MLVSS	%	89.0	

**Table 21: Sewage Water Monitoring at Kandla STP (4<sup>th</sup> Week)**

Date of Sampling		21.09.2021		
Sr. No.	Parameters	Unit	Results	
			KPT STP I/L	KPT STP O/L
1	pH	pH unit	7.53	7.32
2	Total Suspended Solids	mg/l	172.4	75.9
3	Residual Chlorine	mg/l	<1.0	<1.0
4	COD	mg/l	151.5	102.0
5	BOD @ 27 °C	mg/l	106.0	52.0
6.	Fecal Coliform	MPN Index / 100 ml	-	110.0
<b>Aeration Tank</b>				
7.	MLSS	mg/l	16.0	
8	MLVSS	%	82.0	

- **Gopalpuri Colony STP**

**Table 22: Sewage Water Monitoring at Gopalpuri STP (1<sup>st</sup> Week)**

Date of Sampling		04.09.2021		
Sr. No.	Parameters	Unit	Results	
			Gopalpuri STP I/L	Gopalpuri STP O/L
1	pH	pH unit	7.35	7.21
2	Total Suspended Solids	mg/l	108.8	26
3	Residual Chlorine	mg/l	<1.0	<0.5
4	COD	mg/l	316.0	98.0
5	BOD @ 27 °C	mg/l	110.0	28.0
6.	Fecal Coliform	MPN Index / 100 ml	-	>1600.0
<b>Aeration Tank</b>				
7.	MLSS	mg/l	11.0	
8	MLVSS	%	87.0	

**Table 23: Sewage Water Monitoring at Gopalpuri STP (2<sup>nd</sup> Week)**

Date of Sampling		09.09.2021		
Sr. No.	Parameters	Unit	Results	
			Gopalpuri STP I/L	Gopalpuri STP O/L
1	pH	pH unit	7.6	7.41
2	Total Suspended Solids	mg/l	406	107.4
3	Residual Chlorine	mg/l	<1.0	<0.5
4	COD	mg/l	414.1	101
5	BOD @ 27 °C	mg/l	139.0	28.0
6.	Fecal Coliform	MPN Index / 100 ml	-	>1600.0
<b>Aeration Tank</b>				
7.	MLSS	mg/l	14.0	
8	MLVSS	%	90.0	

**Table 24: Sewage Water Monitoring at Gopalpuri STP (3<sup>rd</sup> Week)**

Date of Sampling		16.09.2021		
Sr. No.	Parameters	Unit	Results	
			Gopalpuri STP I/L	Gopalpuri STP O/L
1	pH	pH unit	7.7	7.36
2	Total Suspended Solids	mg/l	276.6	92.1
3	Residual Chlorine	mg/l	<1.0	<0.5
4	COD	mg/l	373.7	104
5	BOD @ 27 °C	mg/l	125.0	28.0
6.	Fecal Coliform	MPN Index / 100 ml	-	>1600.0
<b>Aeration Tank</b>				
7.	MLSS	mg/l	12.0	
8	MLVSS	%	86.0	

**Table 25: Sewage Water Monitoring at Gopalpuri STP (4<sup>th</sup> Week)**

Date of Sampling		05.09.2021		
Sr. No.	Parameters	Unit	Results	
			Gopalpuri STP I/L	Gopalpuri STP O/L
1	pH	pH unit	7.4	7.22
2	Total Suspended Solids	mg/l	182.4	117.8
3	Residual Chlorine	mg/l	<1.0	<0.5
4	COD	mg/l	171.7	101
5	BOD @ 27 °C	mg/l	80.0	23.0
6.	Fecal Coliform	MPN Index / 100 ml	-	920.0
<b>Aeration Tank</b>				
7.	MLSS	mg/l	12.0	
8.	MLVSS	%	88.0	

- **Vadinar STP**

**Table 26: Sewage Water Monitoring at Vadinar STP (1<sup>st</sup> Week)**

Date of Sampling		05.09.2021		
Sr. No.	Parameters	Unit	Results	
			Vadinar STP I/L	Vadinar O/L
1	pH	pH unit	7.22	7.10
2	Total Suspended Solids	mg/l	62	28.0
3	Residual Chlorine	mg/l	<1.0	<0.5
4	COD	mg/l	88.0	56.0
5	BOD @ 27 °C	mg/l	26.0	15.0

**Table 27: Sewage Water Monitoring at Vadinar STP (2<sup>nd</sup> Week)**

<b>Date of Sampling</b>	<b>09.09.2021</b>
-------------------------	-------------------

Sr. No.	Parameters	Unit	Results	
			Vadinar STP I/L	Vadinar O/L
1	pH	pH unit	7.33	7.10
2	Total Suspended Solids	mg/l	72	24.0
3	Residual Chlorine	mg/l	<1.0	<0.5
4	COD	mg/l	88.0	60.0
5	BOD @ 27 °C	mg/l	29.0	18.0

**Table 28: Sewage Water Monitoring at Vadinar STP (3<sup>rd</sup> Week)**

<b>Date of Sampling</b>	<b>16.09.2021</b>
-------------------------	-------------------

Sr. No.	Parameters	Unit	Results	
			Vadinar STP I/L	Vadinar O/L
1	pH	pH unit	7.32	7.12
2	Total Suspended Solids	mg/l	60	58.0
3	Residual Chlorine	mg/l	<1.0	<0.5
4	COD	mg/l	80.0	55.0
5	BOD @ 27 °C	mg/l	26.0	16.0



**Table 29: Sewage Water Monitoring at Vadinar STP (4<sup>th</sup> Week)**

<b>Date of Sampling</b>	<b>21.09.2021</b>
-------------------------	-------------------

Sr. No.	Parameters	Unit	Results	
			Vadinar STP I/L	Vadinar O/L
1	pH	pH unit	7.18	7.10
2	Total Suspended Solids	mg/l	72	42.0
3	Residual Chlorine	mg/l	<1.0	<0.5
4	COD	mg/l	80.0	58.0
5	BOD @ 27 °C	mg/l	26.0	12.0

### 5.3 Conclusions:

The GPCB standards of BOD, TSS and Residual Chlorine for STP outlet are 20 mg/lit, 30 mg/lit & 0.5 mg/lit respectively. It is suggested to do treatment on regular basis to avoid flow of contaminated/polluted water into the sea.

## 6. Marine Water Monitoring

The Forty Second Amendment to the Constitution in 1976 underscored the importance of 'green thinking'. Article 48A enjoins the state to protect and improve the environment and safeguard the forests and wildlife in the country. Further, Article 51A(g) states that the "fundamental duty of every citizen is to protect and improve the natural environment including forests, lakes, rivers and wildlife and to have compassion for living creatures".

Policy Statement for Abatement of Pollution (1992) has suggested developing relevant legislation and regulation, fiscal incentives, voluntary agreements and educational programs and information campaigns. It emphasizes the need for integration by incorporating environmental considerations into decision making at all levels by adopting frameworks namely, pollution prevention at source, application of best practicable solution, ensure polluter pays for control of pollution, focus on heavily polluted areas and river stretches and involve public in decision-making. The National Conservation Strategy and Policy Statement on Environment and Development, (1992) aimed at "integrating environmental concerns with developmental imperatives to meet the challenges by redirecting the thrust of our developmental process so that the basic needs of our people could be fulfilled by making judicious and sustainable use of natural resources." The priorities mentioned in this policy document include the sustainable use of land and water resources, prevention and control of pollution and preservation of biodiversity.

The National Water Policy, (2002) contains provisions for developing, conserving, sustainable utilizing and managing this important water resources and need to be governed by national perspectives.

### Marine Environment

On national and state levels, we have several policies and regulation like Water (Prevention and Control of Pollution) Act, 1974, to regulate pollution discharges and restore water quality of our aquatic resources including the prescription of monitoring activities. One of the important provisions of the Water Act, 1974, is to maintain and restore the 'wholesomeness' of our aquatic resources. Water quality monitoring is one of the first steps required in the rational development and management of water resources. In the field of water quality management, there has been a steady evolution in procedures for designing system to obtain information on the changes of water quality. The monitoring comprises all activities to obtain 'information' with respect to the water system.

## Sampling Stations

The monitoring of marine environment for the study of biological and ecological parameters was carried out on 7<sup>th</sup>& 8<sup>th</sup> September-2021 in harbor regions of KPT and on 7<sup>th</sup> September-2021 at Vadinar during spring tide period of New moon phase of Lunar Cycle. The monitoring of marine environment for the study of biological and ecological parameters was repeated again on 14<sup>th</sup>& 15<sup>th</sup> September 2021 in harbor regions of KPT. 15<sup>th</sup> September -2021 in Vadinar during Neap tide period first quarter of Lunar Cycle..

Plankton samples from sub surface layer was collected both during high tide period and low tide period from 3 water quality monitoring stations of KPT harbour area and two stations in Nakti creek and one station in Khori creek. The same sampling schedule was repeated during consecutive spring tide and neap tide in same month. Plankton samples from sub surface layer was collected both during high tide period and low tide period from 1 water quality monitoring stations near Vadinar jetty area during spring tide and neap tide in this month .Collected water samples were processed for estimation of Chlorophyll- a, Pheophytin- a, qualitative & quantitative evaluation of phytoplankton, qualitative & quantitative evaluation zooplanktons (density and their population).

## Sampling Locations

Offshore monitoring requirement	Number of locations
Offshore Installations	3 in Kandla creek 2 in Nakti creek 1 in Khori creek 1 near Vadinar Jetty 1 near 1 <sup>st</sup> SBM
<b>Total Number of locations</b>	<b>8</b>

### 6.1 Marine Water Quality

Marine water quality of marine waters of Deendayal Port Harbor waters, Khori and Nakti Creeks and two locations of Vadinar are monitored for various physico-chemical parameters during spring and neap tide of each month.

The results of marine water quality and Marine sediments are as below;

**Table 30: Marine Water Quality Monitoring Parameters for location near KPT colony**

Sr. No.	Parameters	Unit	Kandla Creek Near KPT colony (1)			
			23°0'58"N 70°13'22."E			
			Spring Tide		Neap Tide	
			Tide →	High Tide	Low Tide	High Tide
1	pH	pH unit	7.20	7.35	7.31	7.27
2	Color	-	Colorless	Colorless	Colorless	Colorless
3	Odor	-	Odorless	Odorless	Odorless	Odorless
4	Salinity	ppt	31.6	32.0	31.0	31.8
5	Turbidity	NTU	37	35	32	28
6	Total Dissolved Solids	mg/l	59704	58025	34000.0	37060.0
7	Total Suspended Solids	mg/l	282	357	382	303.5
8	Total Solids	mg/l	59986	58382	34382.0	37363.5
9	DO	mg/l	4.5	4.7	4.7	5.3
10	COD	mg/l	78.0	82.0	80.0	86.0
11	BOD	mg/l	<2.0	<2.0	<2.0	<2.0
12	Silica	mg/l	6.09	7.49	0.53	0.42
13	Phosphate	mg/l	0.17	0.16	0.18	0.18
14	Sulphate	mg/l	2640	2280	2808	2568
15	Nitrate	mg/l	2.60	1.43	2.26	2.29
16	Nitrite	mg/l	<0.05	<0.05	<0.05	<0.05
17	Calcium	mg/l	761.52	921.84	521.04	721.44
18	Magnesium	mg/l	1409.4	1263.6	1749.6	1749.6
19	Sodium	mg/l	11280.0	10920.0	11360.0	11062.0
20	Potassium	mg/l	289.0	320.0	296.0	310.0
21	Iron	mg/l	1.95	1.89	1.85	1.79
22	Chromium	mg/l	0.11	0.13	0.11	0.12
23	Copper	mg/l	0.07	0.08	0.08	0.07
24	Arsenic	mg/l	<0.01	<0.01	<0.01	<0.01
25	Cadmium	mg/l	0.04	0.05	0.04	0.05
26	Mercury	mg/l	<0.001	<0.001	<0.001	<0.001
27	Lead	mg/l	0.17	0.18	0.13	0.14
28	Zinc	mg/l	0.05	0.06	0.05	0.07

**Table 31: Marine Water Quality Monitoring Parameters for location near passenger Jetty One at Kandla**

Sr. No.	Parameters	Unit	Near passenger Jetty One (2)			
			23° 0'18 "N 70°13'31"E			
			Spring Tide		Neap Tide	
Tide →		High Tide	Low Tide	High Tide	Low Tide	
1	pH	pH unit	7.40	7.28	7.4	7.43
2	Color	-	Colorless	Colorless	Colorless	Colorless
3	Odor	-	Odorless	Odorless	Odorless	Odorless
4	Salinity	ppt	31.8	31.6	32.5	32.0
5	Turbidity	NTU	26	27	36	27
6	Total Dissolved Solids	mg/l	55555	51116	34060.0	33780.0
7	Total Suspended Solids	mg/l	363	174	242	582.9
8	Total Solids	mg/l	55918	51290	34302.0	34362.9
9	DO	mg/l	4.9	5.2	5.5	4.6
10	COD	mg/l	96.0	90.0	90.0	88.0
11	BOD	mg/l	<2.0	<2.0	<2.0	<2.0
12	Silica	mg/l	5.16	6.84	0.67	0.71
13	Phosphate	mg/l	0.20	0.18	0.16	0.19
14	Sulphate	mg/l	2820	2376	2832	2496
15	Nitrate	mg/l	2.36	2.89	4.00	3.37
16	Nitrite	mg/l	<0.05	<0.05	<0.05	<0.05
17	Calcium	mg/l	721.44	961.92	601.2	681.36
18	Magnesium	mg/l	1409.4	1215	1822.5	1773.9
19	Sodium	mg/l	11862.0	11060.0	11652.0	11110.0
20	Potassium	mg/l	290.0	312.0	299.0	310.0
21	Iron	mg/l	1.96	1.93	1.86	1.93
22	Chromium	mg/l	0.13	0.13	0.13	0.14
23	Copper	mg/l	0.09	0.08	0.06	0.05
24	Arsenic	mg/l	<0.01	<0.01	<0.01	<0.01
25	Cadmium	mg/l	0.06	0.05	0.06	0.04
26	Mercury	mg/l	<0.001	<0.001	<0.001	<0.001
27	Lead	mg/l	0.19	0.16	0.17	0.18
28	Zinc	mg/l	0.08	0.08	0.07	0.06

**Table 32: Marine Water Quality Monitoring Parameters for location Near Coal Berth**

Sr. No.	Parameters	Unit	Near Coal Berth			
			22°59'12"N 70°13'40"E			
			Spring Tide		Neap Tide	
			Tide →	High Tide	Low Tide	High Tide
1	pH	pH unit	7.19	7.38	7.53	7.34
2	Color	-	Colorless	Colorless	Colorless	Colorless
3	Odor	-	Odorless	Odorless	Odorless	Odorless
4	Salinity	ppt	32.5	32.2	32.6	31.6
5	Turbidity	NTU	35	36	34	33
6	Total Dissolved Solids	mg/l	48086	54880	41460.0	39690.0
7	Total Suspended Solids	mg/l	220	220	376.6	359.9
8	Total Solids	mg/l	48306	55100	41836.6	40049.9
9	DO	mg/l	5.0	5.1	4.8	5.2
10	COD	mg/l	89.0	92.0	81.0	78.0
11	BOD	mg/l	<2.0	<2.0	<2.0	<2.0
12	Silica	mg/l	4.67	4.95	0.47	0.71
13	Phosphate	mg/l	0.16	0.21	0.18	0.19
14	Sulphate	mg/l	2376	2964	2376	2352
15	Nitrate	mg/l	2.04	2.26	4.82	4.60
16	Nitrite	mg/l	<0.05	<0.05	<0.05	<0.05
17	Calcium	mg/l	801.60	921.84	440.88	521.04
18	Magnesium	mg/l	1336.5	1287.9	1701	1773.9
19	Sodium	mg/l	12042.0	11910.0	12150.0	11956.0
20	Potassium	mg/l	366.0	372.0	358.0	376.0
21	Iron	mg/l	2.11	2.30	1.96	2.01
22	Chromium	mg/l	0.12	0.15	0.12	0.16
23	Copper	mg/l	0.07	0.09	0.08	0.07
24	Arsenic	mg/l	<0.01	<0.01	<0.01	<0.01
25	Cadmium	mg/l	0.04	0.07	0.05	0.07
26	Mercury	mg/l	<0.001	<0.001	<0.001	<0.001
27	Lead	mg/l	0.19	0.19	0.17	0.2
28	Zinc	mg/l	0.07	0.07	0.05	0.07

**Table 33: Marine Water Quality Monitoring Parameters for location Khori creek at Kandla**

Sr. No.	Parameters	Unit	KPT 4			
			Near 15/16 Berth			
			Spring Tide		Neap Tide	
			Tide →	High Tide	Low Tide	High Tide
1	pH	pH unit	7.40	7.38	7.27	7.22
2	Color	-	Colorless	Colorless	Colorless	Colorless
3	Odor	-	Odorless	Odorless	Odorless	Odorless
4	Salinity	ppt	32.4	31.8	31.8	31.6
5	Turbidity	NTU	27	23	39	45
6	Total Dissolved Solids	mg/l	53390	47930	42746.0	35470.0
7	Total Suspended Solids	mg/l	262	354	561.7	520.9
8	Total Solids	mg/l	53652	48284	43307.7	35990.9
9	DO	mg/l	4.8	4.9	5.5	4.7
10	COD	mg/l	78.0	80.0	86.0	82.0
11	BOD	mg/l	<2.0	<2.0	<2.0	<2.0
12	Silica	mg/l	6.98	6.35	0.98	0.85
13	Phosphate	mg/l	0.23	0.21	0.21	0.19
14	Sulphate	mg/l	2220	2268	2412	2568
15	Nitrate	mg/l	2.87	2.03	2.81	3.32
16	Nitrite	mg/l	<0.05	<0.05	<0.05	<0.05
17	Calcium	mg/l	801.60	881.76	601.2	480.96
18	Magnesium	mg/l	1312.2	1360.8	1773.9	1773.9
19	Sodium	mg/l	12220.0	12052.0	12012.0	12110.0
20	Potassium	mg/l	300.0	278.0	289.0	280.0
21	Iron	mg/l	2.31	2.22	1.88	2.02
22	Chromium	mg/l	0.16	0.14	0.18	0.16
23	Copper	mg/l	0.06	0.08	0.06	0.05
24	Arsenic	mg/l	<0.01	<0.01	<0.01	<0.01
25	Cadmium	mg/l	0.06	0.08	0.06	0.07
26	Mercury	mg/l	<0.001	<0.001	<0.001	<0.001
27	Lead	mg/l	0.19	0.17	0.19	0.16
28	Zinc	mg/l	0.09	0.07	0.06	0.08

**Table 34: Marine Water Quality Monitoring Parameters for location Nakti Creek near Tuna Port**

Sr. No.	Parameters	Unit	Nakti Creek Near Tuna Port			
			22°57'49."N 70° 7'0.67"E			
			Spring Tide		Neap Tide	
			Tide →	High Tide	Low Tide	High Tide
1	pH	pH unit	7.47	7.45	7.22	7.39
2	Color	-	Colorless	Colorless	Colorless	Colorless
3	Odor	-	Odorless	Odorless	Odorless	Odorless
4	Salinity	ppt	31.6	31.9	32.0	31.9
5	Turbidity	NTU	35	37	36	48
6	Total Dissolved Solids	mg/l	52041	55010	35620.0	38755.0
7	Total Suspended Solids	mg/l	360	425	387.9	525.7
8	Total Solids	mg/l	52401	55435	36007.9	39280.7
9	DO	mg/l	4.5	4.7	5.4	5.1
10	COD	mg/l	86.0	82.0	92.0	90.0
11	BOD	mg/l	<2.0	<2.0	<2.0	<2.0
12	Silica	mg/l	8.67	9.40	0.82	0.53
13	Phosphate	mg/l	0.17	0.18	0.22	0.18
14	Sulphate	mg/l	2820	2844	2268	2136
15	Nitrate	mg/l	2.56	1.91	2.42	3.81
16	Nitrite	mg/l	<0.05	<0.05	<0.05	<0.05
17	Calcium	mg/l	761.52	721.44	521.04	601.2
18	Magnesium	mg/l	1409.4	1458	1676.7	1749.6
19	Sodium	mg/l	11958.0	11628.0	11990.0	11558.0
20	Potassium	mg/l	366.0	376.0	360.0	320.0
21	Iron	mg/l	2.35	2.36	2.05	2.10
22	Chromium	mg/l	0.19	0.20	0.18	0.16
23	Copper	mg/l	0.08	0.09	0.05	0.06
24	Arsenic	mg/l	<0.01	<0.01	<0.01	<0.01
25	Cadmium	mg/l	0.06	0.08	0.05	0.08
26	Mercury	mg/l	<0.001	<0.001	<0.001	<0.001
27	Lead	mg/l	0.19	0.21	0.17	0.18
28	Zinc	mg/l	0.09	0.08	0.07	0.08



**Table 35: Marine Water Quality Monitoring Parameters for location Nakti Creek Near NH-8A at Kandla**

Sr. No.	Parameters	Unit	Nakti Creek Near NH-8A			
			23° 02'01"N 70° 09'31"E			
			Spring Tide		Neap Tide	
			Tide →	High Tide	Low Tide	High Tide
1	pH	pH unit	7.36	Sampling not possible during Low Tide	7.39	Sampling not possible during Low Tide
2	Color	-	Colorless		Colorless	
3	Odor	-	Odorless		Odorless	
4	Salinity	ppt	31.6		31.8	
5	Turbidity	NTU	38		37	
6	Total Dissolved Solids	mg/l	54144		35040.0	
7	Total Suspended Solids	mg/l	394		327	
8	Total Solids	mg/l	54538		35367.0	
9	DO	mg/l	4.9		5.6	
10	COD	mg/l	78.0		90.0	
11	BOD	mg/l	<2.0		<2.0	
12	Silica	mg/l	6.96		0.67	
13	Phosphate	mg/l	0.19		0.20	
14	Sulphate	mg/l	2964		2340	
15	Nitrate	mg/l	2.21		25.70	
16	Nitrite	mg/l	<0.05		<0.05	
17	Calcium	mg/l	921.84		641.28	
18	Magnesium	mg/l	1263.6		1725.3	
19	Sodium	mg/l	13125.0		13052.0	
20	Potassium	mg/l	360.0		388.0	
21	Iron	mg/l	2.41		2.30	
22	Chromium	mg/l	0.20		0.19	
23	Copper	mg/l	0.09		0.08	
24	Arsenic	mg/l	<0.01		<0.01	
25	Cadmium	mg/l	0.08		0.07	
26	Mercury	mg/l	<0.001		<0.001	
27	Lead	mg/l	0.18		0.2	
28	Zinc	mg/l	0.09		0.05	

**Table 36: Marine Water Quality Monitoring Parameters for locations Nr. Vadinar Jetty**

Sr. No.	Parameters	Unit	Nr.Vadinar Jetty			
			22°26'25.26"N 69°40'20.41"E			
			Spring Tide		Neap Tide	
			Tide →	High Tide	Low Tide	High Tide
1	pH	pH unit	7.72	7.56	7.4	7.52
2	Color	-	Colorless	Colorless	Colorless	Colorless
3	Odor	-	Odorless	Odorless	Odorless	Odorless
4	Salinity	ppt	31.6	32.0	32.0	32.0
5	Turbidity	NTU	33	34	35	31
6	Total Dissolved Solids	mg/l	41457	45920	34437.0	38630.0
7	Total Suspended Solids	mg/l	299	267	512.6	396
8	Total Solids	mg/l	41756	46187	34949.6	39026.0
9	DO	mg/l	4.7	4.8	5.5	4.8
10	COD	mg/l	90.0	86.0	86.0	89.0
11	BOD	mg/l	<2.0	<2.0	<2.0	<2.0
12	Silica	mg/l	7.27	8.55	0.78	0.77
13	Phosphate	mg/l	0.17	0.18	0.19	0.20
14	Sulphate	mg/l	2316	2388	2388	2532
15	Nitrate	mg/l	3.03	3.15	3.32	2.59
16	Nitrite	mg/l	<0.05	<0.05	<0.05	<0.05
17	Calcium	mg/l	841.68	961.92	561.12	601.2
18	Magnesium	mg/l	1385.1	1263.6	1846.8	1822.5
19	Sodium	mg/l	13820.0	13962.0	13762.0	13888.0
20	Potassium	mg/l	310.0	285.0	316.0	296.0
21	Iron	mg/l	1.96	1.99	1.89	2.00
22	Chromium	mg/l	0.19	0.21	0.17	0.16
23	Copper	mg/l	0.08	0.07	0.07	0.06
24	Arsenic	mg/l	<0.01	<0.01	<0.01	<0.01
25	Cadmium	mg/l	0.07	0.05	0.07	0.06
26	Mercury	mg/l	<0.001	<0.001	<0.001	<0.001
27	Lead	mg/l	0.21	0.16	0.17	0.18
28	Zinc	mg/l	0.08	0.07	0.06	0.05

### **6.1.1 Marine Sediments**

Sediment samples were collected with Van Veen Grab from the six locations in Kandla Port Waters and two locations in Vadinar Port. Samples were collected and preserved in silver foil in ice box to prevent the contamination/decaying of the samples.

## 6.2 Results

The Sediment Quality results are given in below from table no. 34 A & B.

**Table 34A: Results of Analysis of Sediment of Kandla & Vadinar Port (Spring Tide)**

Sr. No.	Parameters	Unit	KPT - 1	KPT - 2	KPT - 4	KPT - 5	Jetty
1	Texture	-	Sandy Loam	Sandy Loam	Sandy Loam	Sandy Loam	Sandy Loam
2	Organic Matter	mg/kg	1.16	1.12	0.83	0.75	0.86
3	Organic Carbon	mg/kg	0.67	0.65	0.48	0.44	0.50
4	Inorganic Phosphate	mg/kg	111.0	126.0	132.0	142.0	175.0
5	Moisture	%	26.00	27.20	42.60	41	28.20
6	Aluminium	mg/kg	ND	ND	ND	ND	ND
7	Silica	mg/kg	18.0	16.0	13.0	16.0	20.0
8	Phosphate	mg/kg	9.20	10.60	10.66	8.40	8.60
9	Sulphate	mg/kg	219.0	253.0	189.0	211.0	186.0
10	Nitrite	mg/kg	0.12	0.13	0.11	0.12	0.13
11	Nitrate	mg/kg	10.20	8.88	9.25	8.69	9.74
12	Calcium	mg/kg	362.0	322.0	410.0	365.0	310.0
13	Magnesium	mg/kg	210.0	192.0	265.0	196.0	188.0
14	Sodium	mg/kg	3824.0	4012.0	2611.0	2978.0	3777.0
15	Potassium	mg/kg	240.0	202.0	160.0	145.0	178.0
16	Chromium	mg/kg	42.5	16	79	19.2	28.7
17	Nickel	mg/kg	24	20.4	16.9	11	19.3
18	Copper	mg/kg	31.8	36.4	34.2	16.8	31.2
19	Zinc	mg/kg	37.10	32.60	28.00	10.20	24.00
20	Cadmium	mg/kg	ND	ND	ND	ND	ND
21	Lead	mg/kg	3.2	3.9	4.8	3	ND
22	Mercury	mg/kg	ND	ND	ND	ND	ND
23	Arsenic	mg/kg	ND	ND	ND	ND	ND

\*Grab samples could not be collected due high at KPT – 3 & Vadinar SBM location.

**Table 34B: Results of Analysis of Sediment of Kandla & Vadinar Port (Neap Tide)**

Sr. No.	Parameters	Unit	KPT - 1	KPT - 2	KPT - 3	KPT - 4	KPT - 5	Jetty
1	Texture	-	Sandy Loam	Sandy Loam	Sandy Loam	Sandy Loam	Sandy Loam	Sandy Loam
2	Organic Matter	mg/kg	1.78	0.90	1.03	2.03	0.81	1.31
3	Organic Carbon	mg/kg	1.03	0.52	0.60	1.18	0.47	0.76
4	Inorganic Phosphate	mg/kg	116.0	136.0	142.0	146.0	149.0	166.0
5	Moisture	%	27.00	19.00	27.0	19.0	27.00	19.00
6	Aluminium	mg/kg	ND	ND	ND	ND	ND	ND
7	Silica	mg/kg	22.20	19.62	18.0	20.2	18.00	16.66
8	Phosphate	mg/kg	7.6	8.2	8.90	10.60	11.20	9.8
9	Sulphate	mg/kg	234.0	268.0	245.0	210.0	265.0	206.0
10	Nitrite	mg/kg	0.11	0.13	0.12	0.1	0.11	0.12
11	Nitrate	mg/kg	8.88	9.20	7.66	9.75	8.88	7.82
12	Calcium	mg/kg	378.0	325.0	389.0	378.0	378.0	296.0
13	Magnesium	mg/kg	216.0	206.0	233.0	186.0	210.0	198.0
14	Sodium	mg/kg	4428.0	3971.0	4554.0	2491.0	3036.0	3798.0
15	Potassium	mg/kg	221.0	152.0	167.0	149.0	116.0	160.30
16	Chromium	mg/kg	38.5	12.1	34.9	77.8	18.7	29.4
17	Nickel	mg/kg	27.3	20.4	36.9	21.6	13.1	19.3
18	Copper	mg/kg	11.8	33.5	40.7	20.2	11	41.2
19	Zinc	mg/kg	47.10	61.00	64.10	38.70	5.20	24.00
20	Cadmium	mg/kg	ND	ND	ND	ND	ND	ND
21	Lead	mg/kg	4.4	4.4	5.6	5.7	2.8	ND
22	Mercury	mg/kg	ND	ND	ND	ND	ND	ND
23	Arsenic	mg/kg	ND	ND	ND	ND	ND	ND

**REPORT**  
**ON**  
**ECOLOGICAL MONITORING**  
**OF MARINE ENVIRONMENT**  
**IN**  
**DPT HARBOUR AREA, NEAR BY CREEKS**  
**For**  
**DEENDAYAL PORT TRUST**

SEPTEMBER, 2021

**INTRODUCTION:**

**Sampling Stations:**

The monitoring of marine environment for the study of biological and ecological Parameters was carried out on 8<sup>th</sup> September 2021 in harbour region of DPT, and on 9<sup>th</sup> September 2021 in creeks near by the port during spring tide. The monitoring of marine environment for the study of biological and ecological parameters was repeated again on 14<sup>th</sup> September 2021 in harbour region of DPT and on 15<sup>th</sup> September 2021 in creeks near by the port during neap tidal condition.

Plankton samples from sub surface layer was collected both during high tide period and low tide period from 3 water quality monitoring stations of DPT harbour area and one stations in Nakti creek and one station in Khori creek. Sampling at second sampling station of Nakti creek was possible only during high tide period. Collected water samples were processed for estimation of Chlorophyll- a, Pheophytin- a, qualitative & quantitative evaluation of phytoplankton, qualitative & quantitative evaluation zooplanktons (density and their population).

**TABLE #1 SAMPLING LOCATIONS**

<b>monitoring requirement</b>	<b>Number of locations</b>
Kandla creek	3 in Kandla creek
Nakti creek	2 in Nakti creek
Khori Creek	1 in Khori creek
<b>Total Number of locations</b>	<b>6</b>

### Sampling methodology adopted:

A marine sampling is an estimation of the body of information in the population. The theory of the sampling design is depending upon the underlying frequency distribution of the population of interest. The requirement for useful water sampling is to collect a representative sample of suitable volume from the specified depth and retain it free from contamination during retrieval.

50 litres of the water sample were collected from Sub surface by using bucket. From the collected water sample 1 litres of water sample were taken in an opaque plastic bottle for chlorophyll estimation, thereafter plankton samples were collected by using filtration assembly with nilyobolt cloth of 20µm mesh size.

### Samples Processing for chlorophyll estimation:

Samples for the chlorophyll estimation were preserved in ice box on board in darkness to avoid degradation in opaque container covered with aluminium foil. Immediately after reaching the shore after sampling, 1 litre of collected water sample was filtered through GF/F filters (pore size 0.45 µm) by using vacuum filtration assembly. After vacuum filtration the glass micro fiber filter paper was grunted in tissue grinder, macerating of glass fiber filter paper along with the filtrate was done in 90% aqueous Acetone in the glass tissue grinder with glass grinding tube. Glass fiber filter paper will assist breaking the cell during grinding and chlorophyll content was extracted with 10 ml of 90% Acetone, under cold dark conditions along with saturated magnesium carbonate solution in glass screw cap tubes. After an extraction period of 24 hours, the samples were transferred to calibrated centrifuge tubes and adjusted the volume to original volume with 90% aqueous acetone solution to make up the evaporation loss. The extract was clarified by using centrifuge in closed tubes. The clarified extracts were then decanted in clean cuvette and optical density was observed at wavelength 664, 665 nm. By using corrected optical density, Chlorophyll-a value was calculated as given in (APHA, 1998).

### PLANKTON:

The entire area open water in the sea is the pelagic realm. Pelagic organisms live in the open sea. In contrast to the pelagic realm, the benthic realm comprises organisms and zone of the bottom of the sea. Vertically the pelagic realm can be dividing into two zones based on light penetration; upper photic or euphotic zone and lower dark water mass, aphotic zone below the photic zone.

The term plankton is general term for organisms have such limited powers of locomotion that they are at the mercy of the prevailing water movement. Plankton is subdivided to phytoplankton and zooplankton. Phytoplankton is free floating organisms that are capable of photosynthesis and zooplankton is the various free floating animals.



Pelagic zone, represents the entire ocean water column from the surface to the deepest depths, is home to a diverse community of organisms. Differences in their locomotive ability categorize the organisms in the pelagic realm into two, *plankton* and *nekton* (Lalli and Parsons, 1997). *Plankton* consists of all organisms drifting in the water and is unable to swim against water currents, whereas *Nekton* includes organisms having strong locomotive power. Ecological studies on the plankton community, which form the base of the aquatic food chain, help in the better understanding of the dynamics and functioning of the marine ecosystem. The term 'Plankton' first coined by Victor Hensen (1887), Plankton, (Greek word: *planktos* meaning "passively drifting or wandering") is defined as drifting or free-floating organisms that inhabit the pelagic zone of water. Based on their mode of nutrition planktonic organisms are categorised into phytoplankton (organisms having an autotrophic mode of nutrition) and zooplankton (organisms having a heterotrophic mode of nutrition).

#### Phytoplankton in the marine environment:

Phytoplankton is free floating unicellular, filamentous and colonial eutrophic organisms that grow in aquatic environments whose movement is more or less dependent upon water currents. These micro flora acts as primary producers as well as the basis of food chain, source of protein, bio purifier and bio indicators of the aquatic ecosystems of which diverse array of the life depends .They are considered as an important component of aquatic flora, play a key role in maintaining equilibrium between abiotic and biotic components of aquatic ecosystem.

The phytoplankton includes a wide range of photosynthetic and phototrophic organisms. Marine phytoplankton is mostly microscopic and unicellular floating flora, which are the primary producers that support the pelagic food-chain. The two most prominent groups of phytoplankton are diatoms (Bacillariophyceae) and dinoflagellates (Dinophyceae).The phytoplankton those normally captured in the net from the Gulf of Kutch is normally dominated by these two major groups; diatoms and dinoflagellates. Phytoplankton also include numerous and diverse collection of extremely small, motile algae which are termed micro flagellates (naked flagellates) as well as and Cyanophytes (blue-green algae).

Algae are an ecologically important group in most aquatic ecosystems and have been an important component of biological monitoring programs. Algae are ideally suited for water quality assessment because they have rapid reproduction rates and very short life cycles, making them valuable indicators of short-term impacts.

Aquatic populations are impacted by anthropogenic stress, resulting in a variety of alterations in the biological integrity of aquatic systems. Algae can serve as an indicator of the degree of deterioration of water quality, and many algal indicators have been used to assess environmental status.

### **Zooplankton in the marine environment:**

Zooplankton includes a taxonomically and morphologically diverse community of heterotrophic organisms that drift in the waters of the world's oceans. Qualitative and quantitative studies on zooplankton community are a prerequisite to delineate the ecological processes active in the marine ecosystem. Zooplankton community plays a pivotal role in the pelagic food web as the primary consumers of phytoplankton and act as the food source for organisms in the higher trophic levels, particularly the economically essential groups such as fish larvae and fishes. They also function in the cycling of elements in the marine ecosystem. The dynamics of the zooplankton community, their reproduction, growth and survival rate are all significant factors determining the recruitment and abundance of fish stocks as they form an essential food for larval, juvenile and adult fishes (Beaugrand et al., 2004). Zooplankton grazing in the marine environment controls the primary production and helps in determining the pelagic ecosystem (Banse, 1995). Through grazing in surface waters and following the production of sinking faecal matters and also by the active transportation of dissolved and particulate matter to deeper waters via vertical migration, they help in the transport of organic carbon to deep ocean layers and thus act as key drivers of biological pump' in the marine ecosystem. Zooplankton grazing and metabolism also, transform particulate organic matter into dissolved forms, promoting primary producer community, microbial demineralization, and particle export to the ocean's interior.

The categorisation of zooplankton into various ecological groups is based on several factors such as duration of planktonic life, size, food preferences and habitat. As they vary significantly in size from microscopic to metazoic forms, the classification of zooplankton based on size has paramount importance in the field of quantitative plankton research.


Based on the duration of planktonic life, zooplankton are categorised into Holoplankton (organisms which complete their entire lifecycle as plankton) and Meroplankton (organisms which are planktonic during the early part of their lives such as the larval stages of benthic and nektonic organisms). Tychoplankton are organisms which live a brief planktonic life, such as the benthic crustaceans (cumaceans, mysids, isopods) which ascend to the water column at night for feeding and certain ectoparasitic copepods, they leave the host and spend their life as plankton during their breeding cycle.

Zooplankton can be subdivided into holoplankton, i.e., permanent members of the plankton (e.g., Calanoid copepods), and meroplankton, i.e., temporary members in the plankton e.g., larvae of fish, shrimp, and crab). The meroplankton group consists of larval and young stages of animals that will adopt a different lifestyle once they mature. In contrast to phytoplankton which consist of a relatively smaller variety of organisms, Zooplankton are extremely divers, consist of a host of larval and adult forms representing many animal phylum.

Among the zooplankton one group always dominate than others; members of sub class copepods (Phylum Athropoda) and Tintinids (Phylum Protozoa) among the net planktons. These small animals are of vital importance in marine ecosystem as one of the primary herbivores animals in the sea, and it is they provide vital link between primary producer (autotrophs) and numerous small and large marine consumers.

As their community structure and function are highly susceptible to changes in the environmental conditions regular monitoring of their distribution as well as their interactions with various physicochemical parameters is inevitable for the sustainable management of the ecosystem (Kusum et al., 2014). Of all the marine zooplankton groups, copepods mainly calanoid copepods are the dominant groups in marine subtropical and tropical waters and exhibit considerable diversity in morphology and habitats they occupy (Madhupratap, 1991;)

It has been well established that potential of pelagic fishes viz. finfishes, crustaceans, molluscs and marine mammals either directly or indirectly depend on zooplankton. The herbivorous zooplankton is efficient grazers of the phytoplankton and is referred to as living machines transforming plant material into animal tissue. Hence they play an essential role as the intermediaries for nutrients/energy transfer between primary and tertiary trophic levels. Due to their large density, shorter lifespan, drifting nature, high group/species diversity and different tolerance to the stress, they used as the indicator organisms for the physical, chemical and biological processes in the aquatic ecosystem (Ghajbhiye, 2002).



### Spatial distribution of Plankton:

A characteristic of plankton population is that they tend to occur in patches, which are varying spatially on a scale of few meters to far as few kilometres in distance. They also vary in time scale, season as well as vertically in the water column. It is this patchiness and its constant changes in time and spot, that has made it so difficult for plankton biologist to learn about the ecology of plankton. The biological factors that causes this patchiness is due to the ability of zooplankton to migrate vertically and graze out the phytoplankton at a rapid rate that can create patchiness. Similarly the active swimming ability by certain zooplankton organisms can cause to aggregate in dense group.

At its most extreme, because the water in which plankton is suspended is constantly moving, each sample taken by the plankton biologists remain a different volume of water, so each sample is unique and replicate does not exist.

Plankton may also exhibit vertical patchiness. Physical factors contribute to this type of patchiness include light intensity, nutrients and density gradients in the water column.

Phytoplankton in particular tends to be unequally distributed vertically, which leads to the existence of different concentration of a chlorophyll value between photic zone and below the photic zone.

**Methodology adopted for Plankton sampling:**

Mixed plankton sample were obtained from the sub surface layer at each sampling locations by towing the net horizontally with the weight .After the tow of about 10-15 minutes, plankton net was pulled up and washed down to the tail and collected the plankton adhered to plankton net in the collection bucket at the bottom by springing outer and inner surface of the net with sea water, while the net was hanging with the mouth upward. For quantitative evaluation 50 L water samples were collected from subsurface layer and filtered through 20µm mesh size net by using bucket and filtration assembly.

**Preservation and storage:**

Both filtered plankton and those collected from the plankton net were preserved with 5% buffered formalin and stored in 1L plastic container for further processing in the laboratory.

**Sample concentration:**

The collected plankton samples were concentrated by using centrifuge and made up to 50 ml with 5% formalin -Glycerine mixture.

**Taxonomic evaluation:**

Before processing, the sample was mixed carefully and a subsample was taken with a calibrated Stempel-pipette. 1 ml of the concentrated plankton samples were transferred on a glass slide with automatic pipette. The plankton sample on the glass slides were stained by using Lugol's iodine and added glycerine to avoid drying while observation. The plankton samples were identified by using Labex triangular Research microscope with photographic attachment. Microphotographs of the plankton samples were taken for record as well as for confirming the identification. The bigger sized zooplankton was observed through dissecting stereomicroscope with magnification of 20-30 x. Plankton organisms in the whole slide were identified to the lowest axon possible. A thorough literature search was conducted for the identification of the different groups of zooplankton that were encountered

**Cell counts by drop count method:**

The common glass slide mounted with a 1ml of concentrated phytoplankton/zooplankton sample in glycerol and covered with cover slip 22x 60mm was placed under the compound microscope provided with a mechanical stage. The plankton was then counted from the microscopic field of the left top corner of the slide. Then slide is moved horizontally along the right side and plankton in each microscopic field was thus counted. When first microscopic field row was finished the next consecutive row was adjusted using the mechanical device of the stage. In this way all the plankton present in entire microscopic field are counted. From this total number in 1ml of the concentrated

plankton, total number of plankton in the original volume of sample filtered was calculated as units/L.

#### **BENTHIC ORGANISMS:**

Benthos is those organisms that are associated with the sea bed or benthic habitats. Epi- benthic organisms live attached to a hard substratum or rooted to a shallow depth below the surface. In fauna organisms live below the sediment–water interface. Interstitial organisms live and move in pore water among sedimentary grains.

Because the benthic organisms are often collected and separated on sieves, a classification based on the overall size is used. Macro benthos include organisms whose shortest dimension is greater than or equal to 0.5 mm. Meio benthos are smaller than 0.5mm but larger than 42 $\mu$  in size.

The terms such as macro fauna and Meio fauna generally have little relevance with taxonomic classification. The terms Meio fauna and macro fauna depend on the size. Meio fauna were considered as good bioassay of community health and rather sensitive indicators of environmental changes

#### **SAMPLING METHODOLOGY ADOPTED FOR SUB TIDAL REGION:**

Van veen sampler (0.09m<sup>2</sup>) was used for sampling bottom sediments. Two sets of sediments were sampled from each location, one for macro fauna and other for Meio fauna. The macro fauna in the sediments were sieved on board to separate out the organisms. The fixation of Meio fauna is normally done by bulk fixation of the sediment sample. The bulk fixation is done by using 10% formalin (Buffered with borate). The organisms were preserved with seawater as diluting agent.

#### **Sample sieving:**

Sediments samples were sieved to extract the organisms. Sieving was performed carefully as possible to avoid any damage to the animals. The large portion of the sediment was split in to smaller portions and mixed with sea water in a bucket. The cohesive lumps were broken down by continuous stirring. The disaggregated sediments were then passed through the sieves.

#### **Sample staining:**

Sorting of the Meio fauna from the sieve is difficult task especially in the preserved material, because organisms are not easily detectable. To facilitate the animal detection the entire sample retained on the sieve after sieving operation were stained by immersing the sieve in a flat bottom tub with 1% Rose Bengal stain; a protein stain. A staining period of 10-30 minutes is sufficient for sample detection.

#### **DIVERSITY INDICES:**

On the whole, diversity indices provide more information about community composition than simply species richness (number of species present); they also, take the relative abundances of different

species into account. Based on this fact, diversity indices therefore depend not only on species richness but on the evenness, or equitability, with which individuals are distributed among the different species (Magurram, A. E. (1988)

A diversity index is a measure of species diversity within a community that consists of co-occurring populations of several (two or more) different species. It includes two components: richness and evenness. Richness is the measure of the number of different species within a sample showing that more the types of species in a community, the higher is the diversity or greater is the richness. Evenness is the measure of relative abundance of the different species with in a community.

The basic idea of diversity index is to obtain a quantitative estimate of biological variability that can be used to compare biological entities composed of discrete components in space and time ( Carol H.R. *etal.* 1998). Biodiversity is commonly expressed through indices based on species richness and species abundances (Whittaker 1972, Lande 1996, Purvis and Hector 2000). Biodiversity indices are a non-parametric tool used to describe the relationship between species number and abundance. The most widely used bio diversity indices are Shannon Weiner index and Simpson's index.

A diversity Index is a single statistic that incorporates in formation on richness and evenness. The diversity measures that incorporate the two concepts may be termed heterogeneity measures (Magurran, 2004).

Any study intended to interpret causes and effect of adverse impact on Biodiversity of communities require suitable measures to evaluate specie richness and Diversity. The former is number of species in community, while latter is a function of relative frequency of different species. Species richness is the iconic measure of biological diversity (Magurran, 2004). Several indices have been created to measure the diversity of species; however, the most widely used in the last decades are the Shannon (1948) and Simpson (1949) (Buzas and Hayek 1996; Gorelick 2006), with the components of diversity: richness (*S*) and evenness (*J*)

### Simpson's diversity index

Simpson's index (**D**) is a measure of diversity, which takes into account both species richness, and evenness of abundance among the species present.. The Simson index is one of the meaningful and robust biodiversity measures available.( Magurran ,2004).

The formula for calculating D is presented as:

$$D = \frac{\sum n_i(n_i - 1)}{N(N - 1)}$$

Where  $n_i$  = the total number of organisms of each individual species

$N$  = the total number of organisms of all species

The value of D ranges from 0 to 1. With this index, 0 represents infinite diversity and, 1, no diversity. When D increases diversity decreases. Simpson's index is therefore usually expressed as  $1-D$  or  $1/D$ . (Magurran, 2004)

Low species diversity suggests:

- relatively few successful species in the habitat
- the environment is quite stressful with relatively few ecological niches and only a few organisms are really well adapted to that environment
- food webs which are relatively simple
- change in the environment would probably have quite serious effects

High species diversity suggests:

- a greater number of successful species and a more stable ecosystem
- more ecological niches are available and the environment is less likely to be hostile complex food webs
- environmental change is less likely to be damaging to the ecosystem as a whole

### **Species richness indices**

The species richness(**S**) is simply the number of species present in an ecosystem. Species richness Indices of species richness are widely used to quantify or monitor the effects of anthropogenic disturbance. A decline in species richness may be concomitant with severe or chronic human-induced perturbation (Fair Fair weather 1990,) Species richness measures have traditionally been the mainstay in assessing the effects of environmental degradation on the biodiversity of natural assemblages of organisms (Clarke & Warwick, 2001)

Species richness is the iconic measure of biological diversity (Magurran, 2004). The species richness(**S**) is simply the number of species present in an ecosystem. This index makes no use of relative abundances. The term species richness was coined by McIntosh (1967) and oldest and most intuitive measure of biological diversity (Magurran, 2004).

Margalef's diversity index is a species richness index. Margalef's Species richness index (d), or indices that describe the evenness of the distribution of the numbers of individuals among species, were derived.

The value of a diversity index increases both when the number of types increases and when evenness increases. For a given number of types, the value of diversity index is maximised when all types are equally abundant (Rosenzweig, M. L. (1995).



**Shannon-Wiener's index:**

An index of diversity commonly used in plankton community analyses is the Shannon-Wiener's index (**H**), which emphasizes not only the number of species (richness or variety), but also the apportionment of the numbers of individuals among the species (Odum 1971 and Reish 1984). Shannon-Wiener's index (**H**) reproduce community parameters to a single number by using an equation.

Shannon and Weiner index represents entropy. It is a diversity index taking into account the number of individuals as well as the number of taxa. It varies from 0 for communities with only single taxa to high values for community with many taxa each with few individuals. This index can also determine the pollution status of a water body. Normal values range from 0 to 4. This index is a combination of species present and the evenness of the species. Examining the diversity in the range of polluted and unpolluted ecosystems, Wilham and Dorris (1968) concluded that the values of the index greater than 3 indicate clean water, values in the range of 1 to 3 are characterized by moderate pollution and values less than 1 are characterized as heavily polluted

$$H' = - \sum_{j=1}^s \frac{n_j}{N} \ln \left( \frac{n_j}{N} \right)$$

**RESULTS:**

**CHLOROPHYLL-a:**

Water Samples for the chlorophyll estimation were collected from sub surface layer during high tide and low tide period of the tidal cycle for each sampling locations and analysed for Chlorophyll -a and after acidification for Pheophytin –a. Chlorophyll- a value was used as algal biomass indicator (APHA,1998) Algal biomass was estimated by converting Chlorophyll value.

In the sub surface water chlorophyll-a was varying from 0.366 -0.613mg/m<sup>3</sup>.in harbour region of DPT during sampling done in spring tide period of September 2021. In the nearby creeks chlorophyll-a was varying from 0.101-0.851mg/m<sup>3</sup>.Pheophytin –a level was below detectable limit- the all the sampling stations during springtide in the harbour region of DPT.

In the sub surface water chlorophyll-a was varying from 0.322 -0.645mg/m<sup>3</sup>.in harbour region of DPT during sampling done in neap tide period of September 2021 . In the nearby creeks chlorophyll-a was varying from 0.291-0.614 mg/m<sup>3</sup>.Pheophytin –a level was below detectable limit- the all the sampling stations during neap tide in the harbour region of DPT.

**TABLE #2 VARIATIONS IN CHLOROPHYLL –a PHEOPHYTIN- a AND ALGAL BIOMASS FROM SAMPLING STATIONS IN DPT HARBOUR AREA ,NEAR BY CREEKS DURING SPRING TIDE IN SEPTEMBER,2021**

Sr. No.	Station	Tide	Chlorophyll-a (mg/m <sup>3</sup> )	Pheophytin- a (mg/m <sup>3</sup> )	Algal Biomass (Chlorophyll method) mg/m <sup>3</sup>
DPT HARBOUR AREA					
1	KPT1	High tide	0.410	BDL	27.47
		Low tide	0.366	BDL	24.52
2	KPT 2	High tide	0.409	BDL	27.40
		Low tide	0.467	BDL	31.29
3	KPT 3	High tide	0.512	BDL	34.30
		Low tide	0.613	BDL	41.07
CREEKS					
4	KPT-4 Khori-I	High tide	0.645	BDL	43.22
		Low tide	0.748	BDL	50.12
5	KPT-5 Nakti-I	High tide	0.818	BDL	54.81
		Low tide	0.851	BDL	57.02
6	KPT-5 Nakti-II	High tide	0.101	BDL	6.76

BDL: Below Detectable Limit.

**TABLE #3 VARIATIONS IN CHLOROPHYLL –a PHEOPHYTIN- a AND ALGAL BIOMASS FROM SAMPLING STATIONS IN DPT HARBOUR AREA ,NEAR BY CREEKS DURING NEAP TIDE IN SEPTEMBER,2021**

Sr. No.	Station	Tide	Chlorophyll-a (mg/m <sup>3</sup> )	Pheophytin- a (mg/m <sup>3</sup> )	Algal Biomass (Chlorophyll method) mg/m <sup>3</sup>
DPT HARBOUR AREA					
1	KPT1	High tide	0.322	BDL	21.57
		Low tide	0.323	BDL	21.64
2	KPT 2	High tide	0.630	BDL	42.21
		Low tide	0.615	BDL	41.21
3	KPT 3	High tide	0.527	BDL	35.31
		Low tide	0.645	BDL	43.22
CREEKS					
4	KPT-4 Khori-I	High tide	0.511	BDL	34.24
		Low tide	0.599	BDL	40.13
5	KPT-5 Nakti-I	High tide	0.529	BDL	35.44
		Low tide	0.614	BDL	41.14
6	KPT-5 Nakti-II	High tide	0.291	BDL	19.50

BDL: Below Detectable Limit.

### PHYTOPLANKTON POPULATION:

For the evaluation of the Phytoplankton population in DPT harbour area and within the immediate surroundings of the port, sampling was conducted from 5 sampling locations (3 in harbour area and two in Nakti creek) during high tide period and low tide period of spring tide and neap tide.

The phytoplankton community of the sub surface water in the harbour and nearby creeks was represented by, Diatoms and blue green algae during spring tide period. Diatoms were represented by 18 genera. Blue green were represented by 3 genera during the sampling conducted in spring tide in September, 2021. Phytoplankton of the sampling stations at sub surface layer in the harbour area and nearby creeks was varying from 85-116 units/ L during high tide period and 103-133 units/ L during low tide of Spring Tide.

The phytoplankton community of the sub surface water in the harbour and nearby creeks was represented by Diatoms and Blue green algae during Neap tide period. Diatoms were represented by 15 genera and Blue green algae were represented 3 genera during the sampling conducted in Neap tide in September, 2021. Phytoplankton of the sampling stations at sub surface layer in the harbour area and nearby creeks was varying from 45 -155 units/ L during high tide period and 131-182 units/ L during low tide of Neap Tide.

### Species Richness Indices and Diversity Indices:

#### Margalef's diversity index (Species Richness) S

At the organismal level, the most widely used biodiversity measures are those based on the number of species present, perhaps adjusted for the number of individuals sampled, Here Margalef's Species richness index (d), or indices that describe the evenness of the distribution of the numbers of individuals among species, are derived.

Margalef's diversity index (Species Richness) S of phytoplankton communities in the stations was varying from 1.801-3.197 with an average of 2.642 during the sampling conducted in High tide period of spring tide. While Margalef's diversity index (Species Richness) S of phytoplankton communities in the Kandla Harbour region and nearby creeks was varying from 12.458-2.904 with an average of 2.697 during the consecutive low tide period .

Margalef's diversity index (Species Richness) S of phytoplankton communities in the stations was varying from 1.808-2.461 with an average of 2.087 during the sampling conducted in High tide period of Neap tide. While Margalef's diversity index (Species Richness) S of phytoplankton communities in the Kandla Harbour region and nearby creeks was varying from. 1.961-2.882 with an average of 2.371 during the consecutive low tide period.

**Shannon-Wiener's index:**

Shannon-Wiener's Index (H) of phytoplankton communities in the sampling stations was in the range of 0.771-0.988 ( $H'(\log_{10})$ ) between selected sampling stations with an average value of 0.934 during high tide period of spring tide. Shannon-Wiener's Index (H) of phytoplankton communities in the sampling stations was in the range of 0.893-0.932 ( $H'(\log_{10})$ ) between selected sampling stations with an average value of 0.916 during consecutive low tide. Shannon-Wiener's Index (H) of phytoplankton communities in the sampling stations was in the range of 0.781-0.911 ( $H'(\log_{10})$ ) between selected sampling stations with an average value of 0.862 during high tide period of neap tide. Shannon-Wiener's Index (H) of phytoplankton communities in the sampling stations was in the range of 0.823-0.969 ( $H'(\log_{10})$ ) between selected sampling stations with an average value of 0.904 during consecutive low tide. Typical values are generally between 1.5 and 3.5 in most ecological studies, and the index is rarely greater than 4. The Shannon-Wiener's index increases as both the richness and the evenness of the community increase. This result indicates that diversity of phytoplankton of Kandla Harbour region and nearby creeks is less but with abundant population of few, with relatively few ecological niches and only very few opportunist organisms are really well adapted to this environment and thrive better than other species.

**Simpson's diversity index:**

Simpson's index (D) is a measure of diversity, which takes into account both species richness, and an evenness of abundance among the species present. The Simpson index is one of the meaningful and robust biodiversity measures available. (Magurran, 2004).

Simpson diversity index (1-D) of phytoplankton communities was below 0.9 at all sampling stations in the Kandla Harbour region and nearby creeks, which was varying from 0.797- 0.882 between selected sampling stations with an average of 0.862 during high tide period of spring tide. Simpson diversity index (1-D) of phytoplankton communities was below 0.9 at all sampling stations in the Kandla Harbour region and nearby creeks, which was varying from 0.845- 0.867 between selected sampling stations with an average of 0.854 during consecutive low tide. Low species diversity suggests a relatively few successful species in this habitat. The environment is quite stressful with relatively few ecological niches and only a few organisms are really well adapted to that environment. Any change in the environment would probably have quite serious effects.

Simpson diversity index (1-D) of phytoplankton communities was below 0.9 at all sampling stations in Kandla Harbour region and nearby creeks, during high tide period and low tide period during neap tide also, which was varying from 0.821-0.857 with an average value of 0.845 between selected sampling stations during high tide period and varying from 0.824-0.870 with an average

**Environmental Monitoring Report of Deendayal Port Trust, SEPTEMBER-2021**

value of 0.853 between selected sampling stations during consecutive low tide period Low species diversity suggests a relatively few successful species in this habitat.

**Table # 4 PHYTOPLANKTON VARIATIONS IN ABUNDANCE AND DIVERSITY IN SUB SURFACE SAMPLING STATIONS IN DPT HARBOUR AREA ,NEAR BY CREEKS DURING SPRING TIDE IN SEPTEMBER,2021**

Tide	Sampling Station	Abundance In units/L	No of Species observed /total species	% of diversity	Margalef's diversity index (Species Richness S)	Shannon Weiner index H (log <sub>10</sub> )	Diversity Index (Simpson's Index) 1-D
HIGH TIDE	1	109	16/21	76.19	3.197	0.9854	0.8739
	2	110	13/21	61.90	2.553	0.9544	0.8754
	3	107	13/21	61.90	2.568	0.9737	0.8824
	4	105	14/21	66.66	2.793	0.9885	0.8811
	5	116	15/21	71.43	2.945	0.9317	0.8627
	6	85	9/21	42.86	1.801	0.7711	0.7978
LOW TIDE	1	103	13/21	61.90	2.589	0.9277	0.8667
	2	132	13/21	61.90	2.458	0.9324	0.8648
	3	124	15/21	71.43	2.904	0.91	0.8451
	4	130	14/21	66.66	2.671	0.8926	0.8458
	5	133	15/21	71.43	2.863	0.9185	0.8479

**Table # 5 PHYTOPLANKTON VARIATIONS IN ABUNDANCE AND DIVERSITY IN SUB SURFACE SAMPLING STATIONS IN DPT HARBOUR AREA ,NEAR BY CREEKS DURING NEAP TIDE IN SEPTEMBER,2021**

Tide	Sampling Station	Abundance In units/L	No of Species observed /total species	% of diversity	Margalef's diversity index (Species Richness S)	Shannon Weiner index H (log <sub>10</sub> )	Diversity Index (Simpson's Index) 1-D
HIGH TIDE	1	131	13/18	72.22	2.461	0.881	0.8452
	2	144	12/18	66.66	2.213	0.8693	0.8503
	3	145	10/18	55.55	1.808	0.8678	0.8511
	4	155	11/18	61.11	1.983	0.8653	0.8484
	5	153	12/18	66.66	2.187	0.911	0.8573
	6	42	8/18	44.44	1.873	0.7809	0.8211
LOW TIDE	1	131	11/18	61.11	2.051	0.8234	0.8243
	2	153	12/18	66.66	2.187	0.8951	0.8542
	3	182	16/18	88.88	2.882	0.9697	0.8703
	4	155	15/18	83.33	2.776	0.9374	0.8618
	5	164	11/18	61.11	1.961	0.8971	0.8564


**Table # 6 ABUNDANCE OF PHYTOPLANKTON SUBSURFACE SAMPLING STATIONS IN DPT HARBOUR AREA, NEAR BY CREEKS DURING SPRING TIDE IN SEPTEMBER, 2021**

Tide	Surface	No of Sampling location	Group of phytoplankton	Phytoplankton Group range Units/L	Genera or species /total Phyto plankton	Taxon Diversity % (Group level)
HIGH TIDE	Sub surface	6	DIATOMS	83-110	18/21	85.71
			BLUE GREEN	2-12	3/21	14.29
			TOTAL PHYTO PLANKTON	<b>85-116</b>	<b>21</b>	-
LOW TIDE	Sub surface	5	DIATOMS	93-129	18/21	85.71
			BLUE GREEN	4-13	3/12	14.29
			TOTAL PHYTO PLANKTON	103-133	<b>21</b>	-


**Table # 7 ABUNDANCE OF PHYTOPLANKTON SUBSURFACE SAMPLING STATIONS IN DPT HARBOUR AREA, NEAR BY CREEKS DURING NEAP TIDE IN SEPTEMBER,2021**

Tide	Surface	No of Sampling location	Group of phytoplankton	Phytoplankton Group range Units/L	Genera or species /total Phyto plankton	Taxon Diversity % (Group level)
HIGH TIDE	Sub surface	6	DIATOMS	38-154	15/18	83.33
			BLUE GREEN	1-4	3/18	16.67
			TOTAL PHYTO PLANKTON	<b>42-155</b>	<b>18</b>	-
LOW TIDE	Sub surface	5	DIATOMS	131-177	15/18	83.33
			BLUE GREEN	0-5	3/18	16.67
			TOTAL PHYTO PLANKTON	<b>131-182</b>	<b>18</b>	-

Taxon Diversity % of Phytoplankton during High tide and Low tide period during spring tide



Taxon Diversity % of Phytoplankton during High tide and Low tide period during Neap tide



### ZOOPLANKTON POPULATION:

For the evaluation of the Zooplankton population in DPT harbour area and within the immediate surroundings of the port sampling was conducted from 6 sampling locations (3 in harbour area and two in Nakti creek and one in Khori creek) during high tide period and low tide period of spring tide and Neap tide in September, 2021. The Zooplankton community of the sub surface water in the harbour and nearby creeks during spring tide was represented by mainly five groups, Tintinids, Copepods, Ciliates, Foraminiferans and larval forms of Crustacea, Molluscs and Polychaetes. The Zooplankton community of the sub surface water in the harbour and nearby creeks during neap tide was represented by mainly seven groups, Tintinids, Copepods, Arrow worms, Ciliates, Mysids, Foraminiferans and larval forms of Crustaceans, Molluscs and Polychaetes.

Zooplankton of the sampling stations at sub surface layer in the DPT harbour area and nearby creek was varying from  $32-86 \times 10^3$  N/  $m^3$  during high tide and  $64-100 \times 10^3$  N/  $m^3$  during low tide of Spring Tide period. Zooplankton of the sampling stations at sub surface layer in the DPT harbour area and nearby creek was varying from  $43-173 \times 10^3$  N/  $m^3$  during high tide and  $115-184 \times 10^3$  N/  $m^3$  during low tide of Neap Tide period.

### Species Richness Indices and Diversity Indices:

#### Margalef's diversity index (Species Richness) S

At the organismal level, the most widely used biodiversity measures are those based on the number of species present, perhaps adjusted for the number of individuals sampled, Here Margalef's Species richness index (d), or indices that describe the evenness of the distribution of the numbers of individuals among species, are derived.

Margalef's diversity index (Species Richness) S of Zooplankton communities in the stations Kandla Harbour region and nearby creeks was varying from 2.825-3.507 with an average of 3.009 during the sampling conducted in High tide period. Margalef's diversity index (Species Richness) S of Zooplankton communities varying from 2.645-3.423 with an average of 3.020 during the sampling conducted in low tide period during Spring tide.

Margalef's diversity index (Species Richness) S of Zooplankton communities in the Kandla Harbour region and nearby creeks sampling stations was varying from 3.722- 4.463 with an average of 4.061 during the sampling conducted in high tide and varying from 4.163-4.647 with an average of 4.458 during the sampling conducted in low tide during Neap tide period **Shannon-Wiener's index:**

Shannon-Wiener's Index (H) of Zooplankton communities in the sampling stations in Kandla Harbour region and nearby creeks was in the range of 0.909-1.014 ( $H'(\log_{10})$ ) between selected sampling stations with an average value of 0.966 ( $H'(\log_{10})$ ) during high tide period of spring tide. Shannon-



Wiener's Index (H) of Zooplankton communities in the sampling stations in Kandla Harbour region and nearby creeks was in the range of 0.929-0.989( $H'(\log_{10})$ ) between selected sampling stations with an average value of 0.963 ( $H'(\log_{10})$ ) during consecutive low tide period .

Shannon-Wiener's Index (H) of Zooplankton communities in the sampling stations in Kandla Harbour region and nearby creeks was in the range of 0.925-1.248 ( $H'(\log_{10})$ ) between selected sampling stations with an average value of 1.155 ( $H'(\log_{10})$ ) during high tide period of Neap tide. Shannon-Wiener's Index (H) of Zooplankton communities in the sampling stations in Kandla Harbour region and nearby creeks was in the range of 1.185-1.254 ( $H'(\log_{10})$ ) between selected sampling stations with an average value of 1.214 ( $H'(\log_{10})$ ) during consecutive low tide period .Typical values are generally between 1.5 and 3.5 in most ecological studies, and the index is rarely greater than 4. The Shannon-Wiener's index increases as both the richness and the evenness of the community increase. This result indicates that diversity of Zooplankton of Kandla Harbour region and nearby creeks stations is slightly high with very minimum diverse population but very few opportunist organisms are really well adapted to this environment and thrive better than other species.

**Simpson's diversity index:**

Simpson's index (D) is a measure of diversity, which takes into account both species richness, and an evenness of abundance among the species present. The Simpson index is one of the meaningful and robust biodiversity measures available. (Magurran, 2004).

Simpson diversity index (1-D) of Zooplankton communities was below 0.9 most of sampling stations in the Kandla Harbour region and nearby creeks during high tide and low tide of spring tide period, which was varying from 0.849-0.889 between selected sampling stations with an average of 0.872 during high tide period and was varying from 0.845- 0.880 with an average value of 0.868 between selected sampling stations during low tide

Simpson diversity index (1-D) of Zooplankton communities was above 0.9 at all sampling stations in the Kandla Harbour region and nearby creeks except few during high tide and low tide period, which was varying from 0.834-0.939 between selected sampling stations with an average of 0.914 during high tide period and was varying from 0.920- 0.939 with an average value of 0.929 between selected sampling stations during consecutive low tide

This high species diversity suggests a relatively more number of successful species in this habitat during September ,2021 sampling.

**Table # 8 ZOOPLANKTON VARIATION IN ABUNDANCE AND DIVERSITY IN SUB SURFACE SAMPLING STATIONS IN DPT HARBOUR AREA ,NEAR BY CREEKS DURING SPRING TIDE IN SEPTEMBER,2021**

Tide	Sampling Station	Abundance In No / m <sup>3</sup>	No of Species/groups observed /total species/group	% of diversity	Margalef's diversity index (Species Richness S)	Shannon Weiner index H (log <sub>10</sub> )	Diversity Index (Simpson's Index) 1-D
HIGH TIDE	1	72 X10 <sup>3</sup>	16/23	69.56	3.507	0.9864	0.8725
	2	70 X10 <sup>3</sup>	13/23	56.52	2.825	0.9094	0.8584
	3	81 X10 <sup>3</sup>	14/23	60.87	2.958	0.9324	0.8494
	4	82 X10 <sup>3</sup>	15/23	65.22	3.177	1.014	0.8871
	5	86 X10 <sup>3</sup>	14/23	60.87	2.918	1.008	0.8892
	6	32 X10 <sup>3</sup>	12/23	52.17	3.174	0.9456	0.875
LOW TIDE	1	80 X10 <sup>3</sup>	16/23	69.56	3.423	0.9692	0.8661
	2	64 X10 <sup>3</sup>	12/23	52.17	2.645	0.9299	0.873
	3	70 X10 <sup>3</sup>	13/23	56.52	2.825	0.9384	0.8451
	4	82 X10 <sup>3</sup>	14/23	60.87	2.95	0.9894	0.8802
	5	100 X10 <sup>3</sup>	16/23	69.56	3.257	0.9872	0.8772

**Table # 9 ZOOPLANKTON VARIATION IN ABUNDANCE AND DIVERSITY IN SUB SURFACE SAMPLING STATIONS IN DPT HARBOUR AREA ,NEAR BY CREEKS DURING NEAP TIDE IN SEPTEMBER,2021**

Tide	Sampling Station	Abundance In No / m <sup>3</sup>	No of Species/groups observed /total species/group	% of diversity	Margalef's diversity index (Species Richness S)	Shannon Weiner index H (log <sub>10</sub> )	Diversity Index (Simpson's Index) 1-D
HIGH TIDE	1	107 X10 <sup>3</sup>	21/29	72.41	4.28	1.16	0.9185
	2	123 X10 <sup>3</sup>	19/29	65.52	3.741	1.19	0.9315
	3	168 X10 <sup>3</sup>	22/29	75.86	4.098	1.205	0.9328
	4	173 X10 <sup>3</sup>	24/29	82.76	4.463	1.248	0.9389
	5	137 X10 <sup>3</sup>	21/29	72.41	4.065	1.203	0.9342
	6	43 X10 <sup>3</sup>	15/29	51.72	3.722	0.9255	0.8339
LOW TIDE	1	115 X10 <sup>3</sup>	22/29	75.86	4.426	1.185	0.9202
	2	122 X10 <sup>3</sup>	21/29	72.41	4.163	1.19	0.9252
	3	175 X10 <sup>3</sup>	25/29	86.21	4.647	1.254	0.9395
	4	184 X10 <sup>3</sup>	25/29	86.21	4.602	1.23	0.9342
	5	140 X10 <sup>3</sup>	23/29	79.31	4.452	1.21	0.9274


**Table # 10 ABUNDANCE OF ZOOPLANKTON IN SUBSURFACE SAMPLING STATIONS IN DPT HARBOUR AREA ,NEAR BY CREEKS DURING SPRING TIDE IN SEPTEMBER,2021**

Tide	Surface	No of Sampling locations	Group of Zooplankton	Abundance of Zooplankton x10 <sup>3</sup> Group Range	Genera or species /total Zooplankton	Taxon Diversity % (Group level)
HIGH TIDE	Sub surface	6	Tintinids	0-8	4/23	17.39
			Copepods	16-51	9/23	39.13
			Ciliates	0-1	1/23	4.35
			Larval forms	14-35	7/23	30.44
			Foraminiferans	0-3	2/23	8.69
			<b>TOTAL ZOOPLANKTON NO/L</b>	<b>32-86</b>	<b>23</b>	<b>23</b>
LOW TIDE	Sub surface	5	Tintinids	2-6	4/23	17.39
			Copepods	29-53	9/23	39.13
			Ciliates	0-1	1/23	4.35
			Larval forms	26-39	7/23	30.44
			Foraminiferans	1-4	2/23	8.69
			<b>TOTAL ZOOPLANKTON NO/M3</b>	<b>64-100</b>	<b>23</b>	<b>23</b>


**Table # 11 ABUNDANCE OF ZOOPLANKTON IN SUBSURFACE SAMPLING STATIONS IN DPT HARBOUR AREA, NEAR BY CREEKS DURING NEAP TIDE IN SEPTEMBER, 2021**

Tide	Surface	No of Sampling locations	Group of Zooplankton	Abundance of Zooplankton x10 <sup>3</sup> Group Range	Genera or species /total Zooplankton	Taxon Diversity % (Group level)
HIGH TIDE	Sub surface	6	Tintinids	1-8	4/29	13.79
			Copepods	17-71	13/29	44.83
			Arrow worms	0-1	1/29	3.45
			Ciliates	1-7	1/29	3.45
			Mysids	0-4	1/29	3.45
			Larval forms	22-87	8/29	27.58
			Foraminiferans	0-2	1/29	3.45
			<b>TOTAL ZOOPLANKTON</b>	<b>43-173</b>	<b>29</b>	<b>-</b>
LOW TIDE	Sub surface	5	Tintinids	2-8	4/29	13.79
			Copepods	38-70	13/29	44.83
			Arrow worms	0-1	1/29	3.45
			Ciliates	3-7	1/29	3.45
			Mysids	1-4	1/29	3.45
			Larval forms	67-106	8/29	27.58
			Foraminiferans	<b>0-1</b>	1/29	3.45
			<b>TOTAL ZOOPLANKTON NO/M3</b>	<b>115-184</b>	<b>29</b>	<b>-</b>

Taxon Diversity % of Zooplankton during High tide and Low tide period of Spring tide



Taxon Diversity % of Zooplankton during High tide and Low tide period of Neap tide



**TABLE # 12 SYSTEMATIC ACCOUNT OF PHYTOPLANKTON IN THE SAMPLING LOCATIONS IN OF DPT HARBOUR AREA AND NEARBY CREEKS DURING SPRING TIDE OF SEPTEMBER 2021**

GROUP	PHYLUM	CLASS	ORDER	FAMILY	GENUS/SPECIES	#	Relative Abundance
BLUE GREEN ALGAE	Cyanophyta	Cyanophyceae	Nostocales	Oscillatoriaceae	<i>Oscillatoria sp.</i>	B1	Occasional
					<i>Arthrospira sp.</i>	B2	Rare
			Stigonematales	Stigonemataceae	<i>Stigonema sp.</i>	B3	Rare
DIATOMS	Bacillariophyta	Coscinodiscophyceae	Thalassiosirales	Thalassiosiraceae	<i>Planktoniellasp</i>	D1	Occasional
			Coscinodiscales	Coscinodiscaceae	<i>Coscinodiscus sp.</i>	D2	Abundant
			Triceratiales	Triceratiaceae	<i>Odontella sp</i>	D3	Frequent
					<i>Triceratium sp.</i>	D4	Rare
			Biddulphiales	Biddulphiaceae	<i>Biddulphiasp</i>	D5	Dominant
			Hemiaulales	Bellerocheaceae	<i>Bellerochea sp</i>	D6	Rare
			Rhizosoleniales	Rhizosoleniaceae	<i>Rhizosolenia sp.</i>	D7	Rare
			Chaetocerotales	Chaetocerotaceae	<i>Chaetoceros sp</i>	D8	Occasional
			Lithodesmiales	Lithodesmiaceae	<i>Ditylum sp</i>	D9	Abundant
		Bacillariophyceae	Naviculales	Pleurosigmaaceae	<i>Pleurosigma sp</i>	D10	Rare
					<i>Navicula sp</i>	D11	Rare
					Surirellales	Surirellaceae	<i>Surirella sp</i>
		Fragilariophyceae	Thalassionematales	Thalassionemataceae	<i>Thalassiothrix sp.</i>	D13	Frequent
					<i>Thalassionema sp.</i>	D14	Rare
			Fragilariales	Fragilariaceae	<i>Asterionellopsis sp.</i>	D15	Rare
					<i>Fragilaria sp</i>	D16	Occasional
					<i>Synedrasp</i>	D17	Rare
					Tabellariales	Tabellariaceae	<i>Tabellaria sp</i>

TABLE # 13 SYSTEMATIC ACCOUNT OF PHYTOPLANKTON IN THE SAMPLING LOCATIONS IN OF DPT HARBOUR AREA AND NEARBY CREEKS DURING AND NEAP TIDE OF SEPTEMBER,2021

GROUP	PHYLUM	CLASS	ORDER	FAMILY	GENUS/SPECIES	#	Relative Abundance
BLUE GREEN ALGAE	Cyanophyta	Cyanophyceae	Nostocales	Oscillatoriaceae	<i>Oscillatoria sp.</i>	B1	Rare
					<i>Arthrospira sp.</i>	B2	Rare
			Stigonematales	Stigonemataceae	<i>Stigonema sp.</i>	B3	Occasional
DIATOMS	Bacillariophyta	Coscinodiscophyceae	Thalassiosirales	Thalassiosiraceae	<i>Planktoniellasp</i>	D1	Rare
			Coscinodiscales	Coscinodiscaceae	<i>Coscinodiscus sp.</i>	D2	Abundant
			Triceratiales	Triceratiaceae	<i>Odontella sp</i>	D3	Frequent
					<i>Triceratium sp.</i>	D4	Occasional
			Biddulphiales	Biddulphiaceae	<i>Biddulphiasp</i>	D5	Dominant
			Hemiaulales	Bellerocheaceae	<i>Bellerochea sp</i>	D6	Occasional
			Chaetocerotales	Chaetocerotaceae	<i>Chaetoceros sp</i>	D7	Rare
		Lithodesmiales	Lithodesmiaceae	<i>Ditylum sp</i>	D8	Abundant	
		Bacillariophyceae	Naviculales	Pleurosigmaaceae	<i>Pleurosigma sp</i>	D9	Occasional
		Fragilariophyceae	Thalassionematales	Thalassionemataceae	<i>Thalassiothrix sp.</i>	D10	Abundant
					<i>Thalassionema sp.</i>	D11	Rare
			Fragilariales	Fragilariaceae	<i>Asterionellopsis sp.</i>	D12	Rare
					<i>Fragilaria sp</i>	D13	Occasional
					<i>Synedrasp</i>	D14	Frequent
			Tabellariales	Tabellariaceae	<i>Tabellaria sp</i>	D15	Rare

**TABLE #14 SYSTEMATIC ACCOUNT OF ZOOPLANKTON FROM THE SAMPLING LOCATIONS IN DPT HARBOUR AREA, AND NEAR BY CREEKS DURING SPRING TIDE OF SEPTEMBER,2021**

GROUP	PHYLUM	CLASS	ORDER	FAMILY	GENUS/SPECIES	#	RELATIVE ABUNDANCE
TINTINIDS	PROTOZOA CILIOPHORA	Spirotrichea	Tintinnida	Tintinnidiidae	<i>Leprotintinnus sp.</i>	T1	Rare
				Codonellidae	<i>Tintinnopsis gracilis</i>	T2	Occasional
					<i>Tintinnopsis radix</i>	T3	Rare
					<i>Tintinnopsis tocaninensis</i>	T4	Rare
COPEPODS	ATHROPODA	Crustacea Sub class copepoda	Calanoida	Paracalanidae	<i>Acrocalanus sp.</i>	C1	Abundant
				Eucalanidae	<i>Pareucalanus sp.</i>	C2	Rare
				Clausocalanidae	<i>Clausocalanus sp.</i>	C3	Occasional
				Acartiidae	<i>Acartia sp.</i>	C4	Rare
				Temoridae	<i>Temora sp.</i>	C5	Occasional
			Cyclopoida	Oithonidae	<i>Oithona sp.</i>	C6	Frequent
			Harpacticoida	Ectinosomatidae	<i>Microsetella sp.</i>	C7	Frequent
				Euterpinidae	<i>Euterpina sp.</i>	C8	Rare
			Poecilostomatatoida	Oncaeidae	<i>Oncaea sp.</i>	C9	Rare
CILIATES	CILIOPHORA	Oligohymenophorea	Sessilida	Zoothamniidae	<i>Zoothamnium sp.</i>	CI1	Rare
CRUSTACEAN LARVAE	ARTHROPODA (CRUSTACEA)	Copepoda			Nauplius larvae of Copepods	L1	Dominant
BIVALVE LARVAE	MOLLUSCA	Pelecypoda			Veliger larvae of Bivalves	L2	Occasional
POLYCHAETE LARVAE	ANNELIDA	Polychaeta			Trochophore larvae	L3	Rare
BARNACLE LARVAE	ARTHROPODA CRUSTACEA	Maxillopoda Thecostraca			Cirripede larvae	L4	Rare
MOLLUSCAN LARVAE	MOLLUSCA	Gastropoda, Streptoneura			Opisthobranchia larvae	L5	Occasional
CYPHONAUTES LARVAE	BRYOZOA				Cyphonautes larvae	L6	Rare
BRACHYURA LARVAE	ARTHROPODA (CRUSTACEA)	Malacostraca Decapoda			Brachyuran Zoea larvae	L7	Rare
FORAMINIFERA	FORAMINIFERA	Globothalamea	Rotaliida	Globigerinidae	<i>Globigerina sp.</i>	F1	Rare
				Rotalliidae	<i>Rotalia sp.</i>	F2	Rare

DCPL/DPT/20-21/17 -SEPTEMBER - 2021

**TABLE # 15 SYSTEMATIC ACCOUNT OF ZOOPLANKTON FROM THE SAMPLING LOCATIONS IN DPT HARBOUR AREA, AND NEAR BY CREEKS DURING NEAP TIDE OF SEPTEMBER,2021**

GROUP	PHYLUM	CLASS	ORDER	FAMILY	GENUS/SPECIES	#	RELATIVE ABUNDANCE	
TINTINIDS	PROTOZOA CILIOPHORA	Spirotrichea	Tintinnida	Tintinnidiidae	<i>Leprotintinnus sp.</i>	T1	Rare	
				Codonellidae	<i>Tintinnopsis gracilis</i>	T2	Occasional	
					<i>Tintinnopsis radix</i>	T3	Rare	
					<i>Tintinnopsis failakkaensis</i>	T4	Rare	
COPEPODS	ATHROPODA	Crustacea Sub class copepoda	Calanoida	Paracalanidae	<i>Acrocalanus sp.</i>	C1	Frequent	
				Eucalanidae	<i>Parvocalanus sp.</i>	C2	Rare	
					<i>Pareucalanus sp.</i>	C3	Rare	
				Subeucalanidae	<i>Subeucalanus sp.</i>	C4	Rare	
					Clausocalanidae	<i>Clausocalanus sp.</i>	C5	Occasional
				Centropagidae	<i>Centropages sp.</i>	C6	Rare	
				Tortanidae	<i>Tortanus sp.</i>	C7	Rare	
				Acartiidae	<i>Acartia sp.</i>	C8	Frequent	
				Temoridae	<i>Temora sp.</i>	C9	Occasional	
				Cyclopoida	Oithonidae	<i>Oithona sp.</i>	C10	Abundant
				Harpacticoida	Ectinosomatidae	<i>Microsetella sp.</i>	C11	Frequent
					Euterpinae	<i>Euterpina sp.</i>	C12	Occasional
				Poecilostomatoida	Oncaeidae	<i>Oncaea sp.</i>	C13	Rare
ARROW WORMS	CHAETOGNATHA	Sagittoidea	Aphragmophora	Sagittidae	<i>Sagitta sp.</i>	A1	Rare	
CILIATES	CILIOPHORA	Oligohymenophorea	Sessilida	Zoothamniidae	<i>Zoothamnium sp.</i>	C11	Occasional	
MYSIDS	ATHROPODA CRUSTACEA	Malacostraca	Mysida, Decapoda	Penaeidae	<i>Penaeus sp.</i>	M1	Occasional	
CRUSTACEAN LARVAE	ARTHROPODA (CRUSTACEA)	Copepoda			Nauplius larvae of Copepods	L1	Dominant	
BRACHYURA LARVAE	ARTHROPODA (CRUSTACEA)	Malacostraca Decapoda			Brachyuran Zoea larvae	L2	Abundant	



**Environmental Monitoring Report of Deendayal Port Trust, SEPTEMBER-2021**

GROUP	PHYLUM	CLASS	ORDER	FAMILY	GENUS/SPECIES	#	RELATIVE ABUNDANCE
BARNACLE LARVAE	ARTHROPODA CRUSTACEA	Maxillopoda Thecostraca			Cirripede larvae	L3	Frequent
CYPHONAUTES LARVAE	BRYOZOA				Cyphonautes larvae	L4	Occasional
ECHINODERMATA LARVAE	ECHINODERMATA				Ophioplutes larvae/ Echinoplutes larvae	L5	Occasional
MOLLUSCAN LARVAE	MOLLUSCA	Gastropoda Streptoneura			Opisthobranchia larvae	L6	Occasional
BIVALVE LARVAE	MOLLUSCA	Pelecypoda			Veliger larvae of Bivalves	L7	Frequent
POLYCHAETE LARVAE	ANNELIDA	Polychaeta			Trochophore larvae	L8	Frequent
FORAMINIFERA	FORAMINIFERA	Globobulimina	Rotaliida	Rotalliidae	<i>Rotalia</i> sp.	F1	Rare

**BENTHIC ORGANISMS:**

Few Benthic organisms were observed in the collected sediments by using the Van-veen grabs during the sampling conducted during spring tide period and while no benthic organisms were observed during sampling conducted in Neap tide period from DPT harbour region and nearby creek except few dead shells. The meiobenthic organisms during spring tide were represented by Polychaetes, and Nematodes. The polychaetes were represented by *Scyphoproctus sp.* and *Branchiicapitelida sps*, during spring tide sampling. The meiobenthic organisms in the collected samples were varying from 0-80N/M<sup>2</sup>.

**Table # 16 BENTHIC FAUNA IN THE SAMPLING LOCATIONS IN DPT HARBOUR AREA CREEKS DURING SPRING TIDE IN SEPTEMBER ,2021**

Benthic fauna	ABUNDANCE IN NO/M <sup>2</sup> DIFFERENT SAMPLING STATIONS REPRESENTATION BY GROUP						
	DPT HARBOUR			CREEKS			
POLYCHAETES	KPT-1	KPT-2	KPT-3	KPT-4	KPT-5	KPT-6	
Family : Capitellidae <i>Scyphoproctus sp.</i>	40	0	0	0	0	NS	
Family : Capitellidae <i>Branchiicapitelida sp.</i>	0	20	0	0	0		
<b>Total Polychates N/M<sup>2</sup></b>	40	20	0	00	0	NS	
<b>Un identified Nematode worms</b>	40	20	0	10	0	NS	
TOTAL Benthic Fauna NUMBER/ M <sup>2</sup>	80	40	0	10	0	-	

NS : No sample

## **7. Meteorological Data**

Automatic Weather station have been installed in SevaSadan -3 at the Deendayal Port which records the data on Temperature (°C), Humidity (%), Wind (mph), Dew Point (°C), Wind Direction (°), Pressure, Solar radiation, heat Index and UVI.

### **Temperature**

The mean day time temperature for Deendayal Port was 29.0 °C. The day-time maximum temperature was 36.2 °C. The mean night time temperature was 26.7 °C. The minimum mean night time temperature recorded was 29.4 °C.

### **Air Pressure**

The mean absolute air pressure for the month of September was 1003.7 hpa, whereas the mean relative pressure was 1002.3 hpa. The maximum absolute air pressure recorded for the month of September was 1010.1 hpa.

### **Heat Index**

The mean day-time heat index for the month of September was 33.5 °C. The maximum heat index recorded was 49°C.

### **Solar Radiation**

The mean Solar Radiation in September was 136.4 w/m<sup>2</sup>. The maximum solar radiation recorded in the month of September was 808.9 w/m<sup>2</sup>.

### **Humidity**

The mean day-time humidity was 83.5 % for the month of September and mean night time humidity was 98.0%. Maximum humidity recorded during day-time was 90.3 % and maximum humidity recorded during night-time was 96.0%.

### **Wind Velocity and Wind Direction**

The mean wind velocity for the entire month of September was 6.88 km/hour. Maximum wind velocity recorded was 43.2 Km/hr . The wind direction was mostly S to SW.

### **Conclusive Summary and Remedial measures Suggested**

- The AAQ monitoring at six locations of Deendayal Port indicates that the mean PM<sub>10</sub> values at four locations viz. Coal storage area, Marine Bhavan and Oil Jetty area were found above the permissible standards (100 µg/m<sup>3</sup>) and PM<sub>2.5</sub> was above permissible limits at Coal storage location (Limit 60 µg/m<sup>3</sup>).
- Drinking water at all the twenty locations was found potable and was within permissible limits of BIS standards (IS 10500).
- Noise quality was also within the set permissible standards of an Industrial Area. The noise level observed during day time was >75 dB (A) and at night time was >70 dB (A) during the entire monitoring period.
- The sewage treated water of Deendayal Port Colony (Gopalpuri) was in line with the standards set by the Gujarat Pollution Control Board. The STP at Deendayal Port is not fully operational and STP at Vadinar Port was found non-operational.

### **Reasons for higher Values of PM<sub>10</sub>**

- Large amount of coal is handled at Berth No. 6, 7, 8 and 9. The unloading of coal directly in the truck, using grabs cause coal to spread in air as well as coal dust to fall on ground. This settled coal dust again mixes with the air while trucks travel through it.
- Also, the coal laden trucks are not always covered with tarpaulin sheets and these results in spillage of coal from trucks/dumpers during its transit from vessel to yard or storage site. This also increased PM values around marine Bhavan & Coal storage area.

### **Remedial Measures**

The values of PM<sub>10</sub> during the month of September, 2021 were observed beyond the permissible limit at four locations mentioned above. Given below are the remedial measures suggest to minimize the Air pollution at Deendayal Port.

- Guidelines for Coal Handling by GPCB should be strictly followed. (<http://gpcb.gov.in/pdf/coal-handling-guidelines.pdf>)
- Except for the higher values of PM<sub>10</sub> at Coal storage site, Oil Jetty, Tuna Port and Marine Bhavan locations, the monitoring results for the present month suggest that the overall Environment Quality of Deendayal Port is satisfactory.

**SOURCE OF LITERATURE AND ADDITIONAL REFERENCE FOR ECOLOGICAL STUDY**

- 1) ALBERT WEST PHAL (1976) Protozoa Blackwell , London
- 2) BANERJEE R.K. (1989) Heavy metals and Benthic foraminiferal distribution along Bombay coast India. Studies in benthic foraminifera. *Tokyo University Press* Tokyo pp 151-157
- 3) Banse K (1995) Zooplankton: Pivotal role in the control of ocean production: I. Biomass and production. *ICES J Mar Sci* 52: 265–277.
- 4) Beaugrand G, and Ibanez F (2004) Monitoring marine plankton ecosystems. II: long-term changes in North Sea calanoid copepods in relation to hydroclimatic variability. *Inter Res Mar Ecol Prog Ser* 284:35-47.
- 5) DAY F. (1889) The fauna of British India Ceylon and Burma- Fishes Vol-1- Vol-2 *Taylor and Francis* London
- 6) DESIKACHARY V. (1989) Atlas of diatoms, Madras Science Foundation
- 7) DESIKACHARY V. (1959) Cyanophyta ICAP Monographs on Algae *Indian Council of Agricultural research* New Delhi
- 8) FAIZAYOUSIF AL-YAMANI & MARIA A. SABUROVA (2010) illustrative guide on the flagellates of Intertidal soft sediment *Kuwait Institute for scientific Research* Kuwait
- 9) FAIZAYOUSIF AL-YAMANI, VALERIYSKRYABIN, ALEKSANDRA GUBANOVA, SERGEY KHVOROV AND IRINA PRUSOVA (2011), Marine zooplankton Practical guide from North western Arabian gulf Vol-1 and vol-2 *Kuwait Institute for scientific Research* Kuwait
- 10) FAUVEL P. (1953), The fauna of India Annelida - Polychaeta *Indian Press* Allahabad
- 11) Gajbhiye SN, Nair VR, and Desai BN (1984). Diurnal variation of zooplankton in Malad creek, Bombay. *Indian Journal of Marine Science*. 13:75-79.
- 12) HAYWARD P.J AND RYLAND J.S. (1995) Handbook of Marine fauna of north –West Europe *oxford University Press* London
- 13) HIGGINS R.P. HAJAMARTHIEL Eds. (1998) Introduction to the study of Meio Fauna
- 14) HORACE G. BARBER AND ELIZABETH Y. HAWORTH (1981) A guide to the Morphology of DIATOMS FRUSTULES.
- 15) INGRAM HENDEY (1964) An introductory account of smaller Algae of British coastal waters part-V. Bacillariophyceae
- 16) JOHN H. WICKSTEAD (1965) an Introduction to the study of Tropical Plankton . *Hutchinson Tropical Monographs*
- 17) JOYOTHIBABU, R. MADHU, N.V. MAHESHWARAN, P.A., NAIR K.K.C., VENUGOPL, P. BALASUBRAMANIAN T. (2005) Dominance of Dinoflagellates in micro zooplankton communities in the oceanic region Bay of Bengal and Andaman sea *Current science* vol.84. 10<sup>th</sup> May 2003
- 18) KASTURIRANGAN L.R. (1963) A key for the identification of the Common Planktonic Copepoda of Indian Coastal water
- 19) Kusum KK, Vineetha G, Raveendran TV, Nair VR, Muraleedharan KR, Achuthankutty CT and Joseph T (2014) Chaetognath community and their responses to varying environmental factors in the northern Indian ocean. *J Plankton Res* 36(4): 1146- 1152.
- 20) Lalli CM and Parsons TR (1997) Biological Oceanography: An Introduction. DOI <https://doi.org/10.1016/B978-0-7506-3384-0.X5056-7>.
- 21) Madhupratap M (1978) Studies on ecology of zooplankton of Cochin backwaters. *Mahasagar Bull Nat Inst Oceanogr* 11: 45-56.
- 22) Madhupratap M (1979) Distribution, community structure and species succession of copepods from Cochin Backwaters. *Indian J Mar Sci* 8: 1-8.

- 23) Madhupratap M (1987) Status and strategy of zooplankton of tropical Indian estuaries: A review. Bull Plank SocJpn 34: 65-81.
- 24) Madhupratap M (1999) Free living copepods of the Arabian Sea, Distribution and Research Perspectives. I J Mar Sci 146-149.
- 25) Madhupratap M and Haridas P (1986) Epipelagic calanoid copepods of the northern Indian Ocean. OceanologicaActa 9(2):105-117.
- 26) MANAL AL-KANDARI, FAIZA Y. AL-YAMANI , KHOLOOD AL-RIFAIE ( 2009) Marine phytoplankton Atlas of Kuwait's water *Kuwait Institute for scientific Research*
- 27) MPEDA (1998) Commercial Fishes and shell fishes of India
- 28) NEWEL G.E. & NEWELL R.C. (1963) Marine plankton a Practical Guide Hutchinson Educational
- 29) NIGAM R.C. AND CHATURVEDIS.K. (2000) Foraminiferal Study from KharoCreek , Kachchh ( Gujarat) North west coast of *India*. *Indian Journal of marine science* Vol.29 133-189
- 30) OLAV GIERE (1993) Meio benthology , Microscopic Fauna in Aquatic Sediments m Springer London
- 31) PERRAGALLO( 1965) Diatomees marines de franceA. *Asher & Co.* Amsterdam
- 32) Robert P.. Higgins (Eds.), (1985) An introduction to the study of Meio fauna Smithsons Institution press Washington DC
- 33) STERRER W. STERRERC.S Eds. Marine fauna and flora of Bermuda A systematic Guide to the identification of Marine Organisms. *John Wiley and Sons*New York
- 34) Suresh Gandhi. M. (2009) Distribution of certain ecological parameters and Foraminiferaldistribution in the depositional environment of Pak strait east coast of India .*Indian J. of Marine Science* Vol.33 pp 287-295
- 35) Venktaraman (1993) A systematic account of some south Indian diatoms . Proceeding of Indian Academy of Science Vol.X No.6 Sec.B.

\*\*\*\*\*

# ENVIRONMENTAL MONITORING REPORT FOR DEENDAYAL PORT TRUST



REPORT NO. : DCPL/DPT/20-21/18  
Month : October 2021  
Issue No : 01  
Revision No : 00  
Prepared by : DETOX CORPORATION PVT. LTD., SURAT



## TABLE OF CONTENTS

<b>Sr. No.</b>	<b>Particulars</b>	<b>Page No.</b>
<b>1</b>	<b>Ambient Air Quality Monitoring.....</b>	<b>1 - 17</b>
<b>2</b>	<b>Drinking Water Quality Monitoring.....</b>	<b>18 - 28</b>
<b>3</b>	<b>Noise Monitoring.....</b>	<b>29</b>
<b>4</b>	<b>Soil Monitoring.....</b>	<b>30 - 31</b>
<b>5</b>	<b>Sewage Treatment Plant Monitoring.....</b>	<b>32 - 38</b>
<b>6</b>	<b>Marine Water Monitoring.....</b>	<b>39 - 92</b>
<b>7</b>	<b>Meteorological Observations.....</b>	<b>93</b>
<b>8</b>	<b>Conclusive Summary &amp; Remedial Measures .....</b>	<b>94-95</b>
	<b>References.....</b>	<b>96- 97</b>

## **Introduction**

Monitoring of various environmental aspects of the Deendayal port by M/s Detox Corporation Pvt. Ltd. has been carried out through collection of samples, analysis of the same, comparing results with respect to the national standards and any other relevant standards by GBCB/CPCB/MoEF to identify non conformity in the Environment of the Deendayal Port. The results shall address the identified impacts and suggest measures to minimize the environmental impact due to various operations at Deendayal Port.

The environmental monitoring is carried out as per the Environment Management and Monitoring Plan submitted by Detox Corporation Pvt. Ltd.

## **1. Ambient Air Quality Monitoring**

As per the Environmental Monitoring Plan of Deendayal Port Trust, Air monitoring was carried out at six identified locations at Deendayal Port and two locations at Vadinar Port.

### **1.1 Air Quality Monitoring Methodology**

Air quality is measured in all the stations, for 24 hour for Total Suspended Particulate Matter (TSPM), PM<sub>10</sub>, PM<sub>2.5</sub>, SO<sub>2</sub>, NO<sub>x</sub>, NH<sub>3</sub> & Benzene, and Grab-sampling for CO & CO<sub>2</sub> measurements. The Air samplers are operated for a period of 24 hours and after a continuous operation of 8 hours of the sampler, the reagents were replaced to obtain 3 samples per day for each parameter namely, SO<sub>2</sub>, NO<sub>x</sub>. The EPM 2000 filter paper and PTFE Membrane bound filter paper are used for a period of 24 hours to obtain one sample each of TSPM, PM<sub>10</sub> & PM<sub>2.5</sub>.

The AAQ samples are collected twice a week from all the eight locations as per the EMP.

**Environmental Monitoring Report of Deendayal Port Trust, OCTOBER-2021**

**1.2 Results**

The ambient air quality monitoring data for six stations, viz. Marine Bhavan, Oil Jetty, Port Colony, Gopalpuri Hospital, Tuna Port and Nr. Coal Storage Area for the month of October 2021 are given in Tables 1A to 6B. The ambient air quality monitoring data for two stations at Vadinar (Nr. Admin Building & Nr. Signal Building) are given in Tables 7A to 8B.

**Location 1: Marine Bhavan (AL1)**

Table 1 : Results of Air Pollutant Concentration at Marine Bhavan										
Parameter	Date	TSPM [µg/m <sup>3</sup> ]	PM10 [µg/m <sup>3</sup> ]	PM2.5 [µg/m <sup>3</sup> ]	SO <sub>2</sub> [µg/m <sup>3</sup> ]		NO <sub>x</sub> [µg/m <sup>3</sup> ]		NH <sub>3</sub> [µg/m <sup>3</sup> ]	
					8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
NAAQMS limit	-	NS	100 µg/m <sup>3</sup>	60 µg/m <sup>3</sup>		80 µg/m <sup>3</sup>		80 µg/m <sup>3</sup>		400 µg/m <sup>3</sup>
AL1 – 1	06.10.2021	349	165	80	2.20	2.64	26.04	25.19	12.76	13.44
					3.96		24.14		12.25	
					1.76		25.41		15.32	
AL1 – 2	08.10.2021	474	229	103	4.40	3.22	15.24	19.27	12.51	13.02
					3.08		16.51		13.02	
					2.20		26.04		13.53	
AL1 – 3	13.10.2021	280	162	58	7.47	7.33	28.58	26.04	10.72	10.81
					8.79		31.12		12.51	
					5.71		18.42		9.19	
AL1 – 4	15.10.2021	404	227	95	3.08	2.49	16.51	15.24	13.79	14.89
					2.64		13.97		15.83	
					1.76		15.24		15.06	
AL1 – 5	20.10.2021	336	156	73	3.52	3.96	18.42	20.54	5.87	9.28
					4.84		20.96		10.72	
					3.52		22.23		11.23	
AL1 - 6	22.10.2021	453	267	85	2.64	3.52	15.88	15.67	10.72	6.47
					5.28		19.69		5.62	
					2.64		11.43		3.06	
AL1 - 7	27.10.2021	338	163	76	3.52	3.37	12.07	16.73	10.47	10.55
					3.96		20.96		11.49	
					2.64		17.15		9.70	
AL1 – 8	29.10.2021	275	152	88	2.64	3.08	24.14	25.19	12.51	9.02
					2.20		29.22		6.64	
					4.40		22.23		7.91	
<b>Monthly Average</b>		364	190	82		3.70		20.48		10.93
<b>Standard Deviation</b>		74	44	14		1.54		4.50		2.75

NS: Not Specified

<b>Table 1B : Results of Air Pollutant Concentration at Marine Bhavan</b>					
<b>Parameter</b>	<b>Date</b>	<b>C<sub>6</sub>H<sub>6</sub> [µg/m<sup>3</sup>]</b>	<b>HC* ppm</b>	<b>CO [mg/m<sup>3</sup>]</b>	<b>CO<sub>2</sub> [ppm ]</b>
<b>Sampling Period</b>		<b>8 hr</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>
<b>NAAQMS limit</b>		<b>5.0 µg/m<sup>3</sup></b>	<b>NS</b>	<b>4.0 mg/m<sup>3</sup></b>	<b>NS</b>
AL1 – 1	06.10.2021	1.12	BQL	1.89	492
AL1 – 2	08.10.2021	1.11	BQL	1.75	489
AL1 – 3	13.10.2021	1.32	BQL	1.82	499
AL1 – 4	15.10.2021	1.15	BQL	1.76	492
AL1 – 5	20.10.2021	1.13	BQL	1.84	493
AL1 - 6	22.10.2021	1.15	BQL	1.86	501
AL1 – 7	27.10.2021	1.21	BQL	1.88	488
AL1 – 8	29.10.2021	1.13	BQL	1.95	511
Monthly Average		<b>1.17</b>	-	<b>1.84</b>	<b>496</b>
Standard Deviation		<b>0.07</b>	-	<b>0.07</b>	<b>8</b>

\* NMHC- Non- Methane Hydrocarbons

BQL- Below Quantification Limit (Quantification Limit – NMHC: 0.5 ppm)

NS -Not Specified

At Marine Bhavan, the overall values of TSPM, PM<sub>10</sub>, PM<sub>2.5</sub>, SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> is attributed mainly by motor vehicle emission produced from various types of automobiles (both diesel and petrol driven). Moreover, the loading and unloading of Food Grains and Timber at Jetty no. 1 and 2 also contributes to the high levels of TSPM and PM<sub>10</sub>. The mean TSPM value at Marine Bhavan was 364 µg/m<sup>3</sup>, The mean PM<sub>10</sub> values were 190.0 µg/m<sup>3</sup>, which is above the permissible limit. PM<sub>2.5</sub> values were above the permissible limit (mean = 82 µg/m<sup>3</sup>). The average values of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were within the permissible limit. The average values of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were 3.70 µg/ m<sup>3</sup>, 20.48 µg/ m<sup>3</sup> & 10.93 µg/ m<sup>3</sup> respectively. These were within the permissible limit.

The levels of Benzene, Hydrocarbons (HC) and CO were within the permissible limit at Marine Bhavan. The mean Benzene concentration was 1.17 µg/m<sup>3</sup>, well below the permissible limit of 5.0 µg/m<sup>3</sup>. HC's were below the detectable limit and Carbon Monoxide concentration was 1.84 mg/m<sup>3</sup>, well below the permissible limit of 4.0 mg/m<sup>3</sup>.

**Location 2: Oil Jetty (AL2)**

**Table 2 : Results of Air Pollutant Concentration at Oil Jetty**

Parameters	Date	TSPM [µg/m3]	PM10 [µg/m3]	PM2.5 [µg/m3]	SO2 [µg/m3]		NOx [µg/m3]		NH3 [µg/m3]	
					8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
NAAQMS limit	-	NS	100 µg/m3	60 µg/m3	-	80 µg/m3	-	80 µg/m3	-	400 µg/m3
AL2 – 1	06.10.2021	380	162	82	4.40	5.28	33.66	33.66	8.93	9.96
					5.28		36.84		10.21	
					6.15		30.49		10.72	
AL2 – 2	08.10.2021	486	284	103	1.76	3.96	8.89	11.01	6.38	9.62
					4.84		9.53		10.98	
					5.28		14.61		11.49	
AL2 – 3	13.10.2021	451	300	89	7.47	9.52	32.39	24.35	3.57	6.30
					9.23		18.42		6.13	
					11.87		22.23		9.19	
AL2 – 4	15.10.2021	480	335	100	2.20	2.05	16.51	14.82	7.40	8.93
					2.64		14.61		10.47	
					1.32		13.34		8.93	
AL2 – 5	20.10.2021	464	190	76	2.64	2.49	20.33	19.48	9.19	7.32
					1.76		24.77		4.85	
					3.08		13.34		7.91	
AL2 – 6	22.10.2021	509	280	100	4.84	5.28	20.96	14.82	5.62	8.00
					7.03		10.16		7.91	
					3.96		13.34		10.47	
AL2 – 7	27.10.2021	448	215	71	1.76	1.76	22.87	19.48	9.19	10.04
					1.32		14.61		12.51	
					2.20		20.96		8.42	
AL2 – 8	29.10.2021	504	204	92	1.32	2.49	14.61	17.36	6.38	9.10
					2.20		22.87		9.96	
					3.96		14.61		10.98	
<b>Monthly Average</b>		465	246	89		4.10		19.37		8.66
<b>Standard Deviation</b>		41	61	12		2.59		7.01		1.34

NS: Not Specified

<b>Table 2B : Results of Air Pollutant Concentration at Oil Jetty</b>					
<b>Parameter</b>	<b>Date</b>	<b>C<sub>6</sub>H<sub>6</sub> [µg/m<sup>3</sup>]</b>	<b>HC* ppm</b>	<b>CO [mg/m<sup>3</sup>]</b>	<b>CO<sub>2</sub> [ppm ]</b>
<b>Sampling Period</b>		<b>8 hr</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>
<b>NAAQMS limit</b>		<b>5.0 µg/m<sup>3</sup></b>	<b>NS</b>	<b>4.0 mg/m<sup>3</sup></b>	<b>NS</b>
AL2 -1	06.10.2021	1.11	BQL	1.88	499
AL2 -2	08.10.2021	1.21	BQL	1.78	495
AL2 -3	13.10.2021	1.26	BQL	1.86	468
AL2 -4	15.10.2021	1.11	BQL	1.83	466
AL2 – 5	20.10.2021	1.22	BQL	1.89	458
AL2 – 6	22.10.2021	1.18	BQL	1.87	488
AL2 -7	27.10.2021	1.01	BQL	1.77	498
AL2 – 8	29.10.2021	1.14	BQL	1.82	501
<b>Monthly Average</b>		<b>1.16</b>	<b>-</b>	<b>1.84</b>	<b>484</b>
<b>Standard Deviation</b>		<b>0.08</b>	<b>-</b>	<b>0.05</b>	<b>17</b>

\* NMHC- Non- Methane Hydrocarbons

BQL- Below Quantification Limit (Quantification Limit – NMHC: 0.5 ppm)

NS- Not Specified

The overall values of TSPM, PM<sub>10</sub>, PM<sub>2.5</sub>, SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> at Oil Jetty area was mainly by motor vehicle emission produced from various types of vehicles Oil Jetty Area. The mean TSPM values at Oil Jetty were 465 µg/m<sup>3</sup>. The mean PM<sub>10</sub> values were 246 µg/m<sup>3</sup>, which is above the permissible limit. PM<sub>2.5</sub> values were above the permissible limit (mean = 89 µg/m<sup>3</sup>). The average values of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were within the permissible limit, The mean concentration of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were 4.10 µg/m<sup>3</sup>, 19.37 µg/m<sup>3</sup> and 8.66 µg/m<sup>3</sup> respectively.

The levels of Benzene, Hydrocarbons (HC) and CO were within the permissible limit at Oil Jetty. The mean Benzene concentration was 1.16 µg/m<sup>3</sup>. Well below the permissible limit of 5.0 µg/m<sup>3</sup>. , HC's were below the detectable limit and Carbon Monoxide concentration was 1.84 mg/m<sup>3</sup>, well below the permissible limit of 4.0 mg/m<sup>3</sup>.

**Location 3: Kandla Colony – Estate Office (AL-3)**

<b>Table 3 : Results of Air Pollutant Concentration at Estate Office</b>										
<b>Parameters</b>	<b>Date</b>	<b>TSPM [µg/m3]</b>	<b>PM10 [µg/m3]</b>	<b>PM2.5 [µg/m3]</b>	<b>SO2 [µg/m3]</b>		<b>NOx [µg/m3]</b>		<b>NH3 [µg/m3]</b>	
<b>Sampling Period</b>	-	<b>24hr</b>	<b>24hr</b>	<b>24hr</b>	<b>8 hr</b>	<b>24hr (Avg.)</b>	<b>8 hr</b>	<b>24hr (Avg.)</b>	<b>8 hr</b>	<b>24hr (Avg.)</b>
<b>NAAQMS limit</b>	-	<b>NS</b>	<b>100 µg/m3</b>	<b>60 µg/m3</b>	-	<b>80 µg/m3</b>	-	<b>80 µg/m3</b>	-	<b>400 µg/m3</b>
<b>AL3 – 1</b>	06.10.2021	355	161	79	4.84	3.37	12.70	23.08	14.30	12.00
					3.52		24.14		12.25	
					1.76		32.39		9.45	
<b>AL3 – 2</b>	08.10.2021	280	121	76	3.96	2.64	13.34	12.49	10.47	9.10
					1.32		9.53		11.49	
					2.64		14.61		5.36	
<b>AL3 – 3</b>	13.10.2021	420	282	98	3.08	3.52	13.97	19.48	7.91	5.87
					4.84		19.69		6.38	
					2.64		24.77		3.32	
<b>AL3 – 4</b>	15.10.2021	530	287	101	4.40	2.93	8.89	8.05	12.25	9.10
					2.64		8.26		9.19	
					1.76		6.99		5.87	
<b>AL3 – 5</b>	20.10.2021	401	239	98	5.28	3.66	18.42	23.50	8.93	9.19
					3.08		32.39		9.70	
					2.64		19.69		8.93	
<b>AL3 – 6</b>	22.10.2021	381	244	93	5.28	4.40	18.42	19.27	10.47	8.25
					1.76		14.61		8.93	
					6.15		24.77		5.36	
<b>AL3 – 7</b>	27.10.2021	466	194	90	4.84	2.93	19.69	17.36	11.23	10.81
					2.64		16.51		10.72	
					1.32		15.88		10.47	
<b>AL3 – 8</b>	29.10.2021	380	222	87	1.76	2.93	15.88	16.94	12.00	9.10
					4.40		15.24		9.70	
					2.64		19.69		5.62	
<b>Monthly Average</b>		402	219	90		3.30		17.52		9.18
<b>Standard Deviation</b>		74	58	9		0.56		5.20		1.79

NS: Not Specified

<b>Table 3B : Results of Air Pollutant Concentration at Kandla Port Colony</b>					
<b>Parameter</b>	<b>Date</b>	<b>C<sub>6</sub>H<sub>6</sub> [µg/m<sup>3</sup>]</b>	<b>HC*</b>	<b>CO [mg/m<sup>3</sup>]</b>	<b>CO<sub>2</sub> [ppm ]</b>
<b>Sampling Period</b>		<b>8 hr</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>
<b>NAAQMS limit</b>		<b>5.0 µg/m<sup>3</sup></b>	<b>NS</b>	<b>4.0 mg/m<sup>3</sup></b>	<b>NS</b>
AL3 -1	06.10.2021	1.01	BQL	1.85	489
AL3 -2	08.10.2021	1.12	BQL	1.98	496
AL3 -3	13.10.2021	1.02	BQL	1.79	488
AL3 -4	15.10.2021	1.11	BQL	1.81	499
AL3 -5	20.10.2021	1.06	BQL	1.88	480
AL3 -6	22.10.2021	1.18	BQL	1.79	485
AL3 -7	27.10.2021	1.26	BQL	1.96	472
AL3 -8	29.10.2021	1.14	BQL	1.88	498
<b>Monthly Average</b>		<b>1.11</b>	<b>-</b>	<b>1.87</b>	<b>488</b>
<b>Standard Deviation</b>		<b>0.08</b>	<b>-</b>	<b>0.07</b>	<b>9</b>

\* NMHC- Non- Methane Hydrocarbons

BQL- Below Quantification Limit (Quantification Limit – NMHC: 0.5 ppm)

NS- Not Specified

The overall values of TSPM, PM<sub>10</sub>, PM<sub>2.5</sub>, SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> at Kandla Port Colony was attributed by vehicle emission produced from trucks and heavy duty vehicles that pass through the road outside Kandla Port Colony. The mean TSPM values at Oil Jetty were 402 µg/m<sup>3</sup>, The mean PM<sub>10</sub> values were 219 µg/m<sup>3</sup>, which is above the permissible limit. PM<sub>2.5</sub> values were above the permissible limit (mean = 90 µg/m<sup>3</sup>). The average values of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were 3.30 µg/m<sup>3</sup>, 17.52 µg/m<sup>3</sup> and 9.18 µg/m<sup>3</sup> respectively and were all within the permissible limit.

The levels of Benzene, Hydrocarbons (HC) and CO were within the permissible limit at Kandla Port Colony. The mean Benzene concentration was 1.11 µg/m<sup>3</sup>, well below the permissible limit of 5.0 µg/m<sup>3</sup>. HC's were below the detectable limit and Carbon Monoxide concentration was 1.87 mg/m<sup>3</sup>, well below the permissible limit of 4.0 mg/m<sup>3</sup>.



**Location 4: Gopalpuri Hospital (AL-4)**

Table 4 : Results of Air Pollutant Concentration at Gopalpuri Hospital										
Parameter	Date	TSPM [µg/m3]	PM10 [µg/m3]	PM2.5 [µg/m3]	SO2 [µg/m3]		NOx [µg/m3]		NH3 [µg/m3]	
Sampling Period	-	24hr	24hr	24hr	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
NAAQMS limit	-	NS	100 µg/m3	60 µg/m3	-	80 µg/m3	-	80 µg/m3	-	400 µg/m3
AL4 -1	06.10.2021	158	96	40	6.15	4.40	18.42	25.83	3.83	5.87
					3.96		25.41		7.40	
					3.08		33.66		6.38	
AL4 -2	08.10.2021	246	115	77	3.96	5.28	12.70	10.37	4.85	5.70
					5.28		9.53		5.11	
					6.59		8.89		7.15	
AL4 -3	13.10.2021	302	129	66	2.20	2.64	10.16	11.22	3.83	4.25
					3.08		12.70		4.85	
					2.64		10.80		4.08	
AL4 -4	15.10.2021	414	267	89	2.20	2.20	10.16	8.68	6.38	5.45
					2.64		9.53		4.60	
					1.76		6.35		5.36	
AL4 -5	20.10.2021	268	128	90	2.64	2.64	14.61	14.61	4.85	6.64
					3.08		9.53		8.42	
					2.20		19.69		6.64	
AL4 -6	22.10.2021	219	114	93	2.64	2.49	13.34	12.49	4.85	8.51
					3.08		9.53		9.19	
					1.76		14.61		11.49	
AL4 -7	27.10.2021	274	132	84	2.64	3.08	17.78	15.88	6.38	6.55
					3.08		13.34		7.91	
					3.52		16.51		5.36	
AL4 -8	29.10.2021	311	142	96	2.20	3.22	13.34	12.70	7.40	8.25
					3.52		13.97		8.42	
					3.96		10.80		8.93	
<b>Monthly Average</b>		274	140	79		3.24		13.97		6.40
<b>Standard Deviation</b>		75	53	18		1.06		5.30		1.43

NS: Not Specified

<b>Table 4B : Results of Air Pollutant Concentration at Gopalpuri Hospital</b>					
<b>Parameter</b>	<b>Date</b>	<b>C<sub>6</sub>H<sub>6</sub> [µg/m<sup>3</sup>]</b>	<b>HC*</b>	<b>CO [mg/m<sup>3</sup>]</b>	<b>CO<sub>2</sub> [ppm ]</b>
<b>Sampling Period</b>		<b>8 hr</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>
<b>NAAQMS limit</b>		<b>5.0 µg/m<sup>3</sup></b>	<b>NS</b>	<b>4.0 mg/m<sup>3</sup></b>	<b>NS</b>
AL4 -1	06.10.2021	1.02	BQL	1.88	496
AL4 -2	08.10.2021	1.11	BQL	1.68	482
AL4 -3	13.10.2021	1.32	BQL	1.65	501
AL4 -4	15.10.2021	1.25	BQL	1.79	499
AL4 -5	20.10.2021	1.52	BQL	1.65	501
AL4 -6	22.10.2021	1.32	BQL	1.82	508
AL4 -7	27.10.2021	1.23	BQL	1.7	487
AL4 -8	29.10.2021	1.58	BQL	1.76	496
<b>Monthly Average</b>		<b>1.29</b>	<b>-</b>	<b>1.74</b>	<b>496</b>
<b>Standard Deviation</b>		<b>0.19</b>	<b>-</b>	<b>0.08</b>	<b>8</b>

\* NMHC- Non- Methane Hydrocarbons

BQL- Below Quantification Limit (Quantification Limit – NMHC: 0.5 ppm)

NS-Not Specified

The overall values of TSPM, PM<sub>10</sub>, PM<sub>2.5</sub>, SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> at Gopalpuri Hospital was attributed by vehicle emission produced from light motor vehicles of the colony residents. The mean TSPM values at Oil Jetty were 274 µg/m<sup>3</sup>, The mean PM<sub>10</sub> values were 140 µg/m<sup>3</sup>, which is above the permissible limit. PM<sub>2.5</sub> values were above the permissible limit (mean= 79 µg/m<sup>3</sup>). The average values of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were 3.24 µg/m<sup>3</sup>, 13.97 µg/m<sup>3</sup> and 6.40 µg/m<sup>3</sup> respectively and were all within the permissible limit.

The levels of Benzene, Hydrocarbons (HC) and CO were within the permissible limit at Gopalpuri Hospital. The mean Benzene concentration was 1.29 µg/m<sup>3</sup>, well below the permissible limit of 5.0 µg/m<sup>3</sup>. HC's were below the detectable limit and Carbon Monoxide concentration was 1.74 mg/m<sup>3</sup>, well below the permissible limit of 4.0 mg/m<sup>3</sup>.

**Location 5: Coal Storage Area (AL-5)**

Table 5 : Results of Air Pollutant Concentration at Coal Storage Area										
Parameters	Date	TSPM [µg/m <sup>3</sup> ]	PM10 [µg/m <sup>3</sup> ]	PM2.5 [µg/m <sup>3</sup> ]	SO <sub>2</sub> [µg/m <sup>3</sup> ]		NO <sub>x</sub> [µg/m <sup>3</sup> ]		NH <sub>3</sub> [µg/m <sup>3</sup> ]	
Sampling Period	-	24hr	24hr	24hr	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
NAAQMS limit	-	NS	100 µg/m <sup>3</sup>	60 µg/m <sup>3</sup>	-	80 µg/m <sup>3</sup>	-	80 µg/m <sup>3</sup>	-	400 µg/m <sup>3</sup>
AL5 – 1	06.10.2021	266	122	92	4.40	5.71	44.46	49.33	15.32	14.04
					6.15		49.54		13.53	
					6.59		53.99		13.27	
AL5 – 2	08.10.2021	360	208	79	3.96	3.96	20.33	18.84	15.57	16.25
					2.64		22.23		17.61	
					5.28		13.97		15.57	
AL5 – 3	13.10.2021	647	226	110	10.11	8.65	22.23	22.87	7.91	7.32
					6.15		26.04		6.13	
					9.67		20.33		7.91	
AL5 – 4	15.10.2021	760	217	118	1.32	2.93	16.51	18.84	12.51	9.62
					3.52		19.69		8.42	
					3.96		20.33		7.91	
AL5 – 5	20.10.2021	597	244	110	4.84	4.25	19.69	20.75	10.72	11.32
					4.40		17.78		10.98	
					3.52		24.77		12.25	
AL5 – 6	22.10.2021	647	206	106	3.52	4.54	14.61	17.15	14.30	14.47
					3.96		15.88		15.06	
					6.15		20.96		14.04	
AL5 – 7	27.10.2021	614	249	107	4.40	4.25	13.34	16.51	9.96	9.36
					4.84		17.78		9.19	
					3.52		18.42		8.93	
AL5 – 8	29.10.2021	324	151	117	4.84	4.10	22.87	25.41	12.51	14.04
					3.96		27.95		14.30	
					3.52		25.41		15.32	
<b>Monthly Average</b>		527	203	105		4.80		23.71		12.05
<b>Standard Deviation</b>		182	44	13		1.73		10.76		3.11

NS: Not Specified

<b>Table 5B : Results of Air Pollutant Concentration at Coal Storage Area</b>					
<b>Parameter</b>	<b>Date</b>	<b>C<sub>6</sub>H<sub>6</sub> [µg/m<sup>3</sup>]</b>	<b>HC*</b>	<b>CO [mg/m<sup>3</sup>]</b>	<b>CO<sub>2</sub> [ppm ]</b>
<b>Sampling Period</b>		<b>8 hr</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>
<b>NAAQMS limit</b>		<b>5.0 µg/m<sup>3</sup></b>	<b>NS</b>	<b>4.0 mg/m<sup>3</sup></b>	<b>NS</b>
AL5 – 1	06.10.2021	1.22	BQL	1.85	498
AL5 – 2	08.10.2021	1.32	BQL	1.86	488
AL5 – 3	13.10.2021	1.22	BQL	1.89	485
AL5 – 4	15.10.2021	1.16	BQL	1.84	501
AL5 – 5	20.10.2021	1.33	BQL	1.86	496
AL5 – 6	22.10.2021	1.24	BQL	1.9	500
AL5 – 7	27.10.2021	1.15	BQL	1.84	490
AL5 – 8	29.10.2021	1.18	BQL	1.98	498
<b>Monthly Average</b>		<b>1.23</b>	<b>-</b>	<b>1.88</b>	<b>495</b>
<b>Standard Deviation</b>		<b>0.07</b>	<b>-</b>	<b>0.05</b>	<b>6</b>

\* NMHC- Non- Methane Hydrocarbons

BQL- Below Quantification Limit (Quantification Limit – NMHC: 0.5 ppm)

NS-Not Specified

The overall values of TSPM, PM<sub>10</sub>, PM<sub>2.5</sub>, SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> at Coal Storage Area was comparatively highest among all the locations of Air Quality monitoring in Kandla Port. High values of TSPM, PM<sub>10</sub>, PM<sub>2.5</sub>, SO<sub>2</sub>, NO<sub>x</sub> at this location was due to lifting of coal with grab and other coal handling processes near Berth no. 6 & 7. Moreover, the traffic was also heavy around this place for transport of coal thus emissions produced from heavy vehicles. The mean TSPM values at Coal storage were 527µg/m<sup>3</sup>. The mean PM<sub>10</sub> values were 203 µg/m<sup>3</sup>, which is well above the permissible limit. PM<sub>2.5</sub> values were above the permissible limit (mean = 105 µg/m<sup>3</sup>). The average values of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were 4.80 µg/m<sup>3</sup>, 23.71 µg/m<sup>3</sup> and 12.05 µg/m<sup>3</sup> respectively and were all within the permissible limit.

The levels of Benzene, Hydrocarbons (HC) and CO were within the permissible limit at Coal Storage Area. The mean Benzene concentration was 1.23 µg/m<sup>3</sup>, well below the permissible limit of 5.0 µg/m<sup>3</sup>. HC's were below the detectable limit and Carbon Monoxide concentration was 1.88 mg/m<sup>3</sup>, well below the permissible limit of 4.0 mg/m<sup>3</sup>.

**Location 6: Tuna Port (AL-6)**

Table 6 : Results of Air Pollutant Concentration at Tuna Port										
Parameters	Date	TSPM [µg/m3]	PM10 [µg/m3]	PM2.5 [µg/m3]	SO2 [µg/m3]		NOx [µg/m3]		NH3 [µg/m3]	
Sampling Period	-	24hr	24hr	24hr	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
NAAQMS limit	-	NS	100 µg/m3	60 µg/m3	-	80 µg/m3	-	80 µg/m3	-	400 µg/m3
AL6 -1	06.10.2021	280	134	98	4.40	4.10	15.88	29.85	5.62	7.40
					6.15		33.66		7.66	
					1.76		40.02		8.93	
AL6 - 2	08.10.2021	293	130	92	1.76	3.22	13.34	12.49	13.53	11.49
					3.08		13.97		12.25	
					4.84		10.16		8.68	
AL6 - 3	13.10.2021	438	251	103	6.15	4.69	16.51	16.94	6.38	5.96
					5.71		20.96		4.08	
					2.20		13.34		7.40	
AL6 - 4	15.10.2021	466	153	100	2.20	1.76	5.08	6.78	5.62	6.47
					1.76		8.26		7.15	
					1.32		6.99		6.64	
AL6 - 5	20.10.2021	480	180	94	1.32	2.64	20.33	16.94	12.25	11.57
					2.64		13.97		11.49	
					3.96		16.51		10.98	
AL6 - 6	22.10.2021	310	123	88	4.84	3.22	32.39	27.31	9.96	12.76
					2.20		20.96		15.57	
					2.64		28.58		12.76	
AL6 - 7	27.10.2021	275	140	93	2.20	2.49	15.24	15.88	9.19	9.36
					1.76		16.51		8.42	
					3.52		15.88		10.47	
AL6 - 8	29.10.2021	352	191	98	2.20	2.93	15.88	15.03	10.72	10.30
					2.64		10.80		8.93	
					3.96		18.42		11.23	
<b>Monthly Average</b>		362	163	96		3.13		17.65		9.41
<b>Standard Deviation</b>		86	43	5		0.92		7.54		2.55

NS: Not Specified

<b>Table 6B : Results of Air Pollutant Concentration at Tuna Port</b>					
<b>Parameter</b>	<b>Date</b>	<b>C<sub>6</sub>H<sub>6</sub> [µg/m<sup>3</sup>]</b>	<b>HC*</b>	<b>CO [mg/m<sup>3</sup>]</b>	<b>CO<sub>2</sub> [ppm ]</b>
<b>Sampling Period</b>		<b>8 hr</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>
<b>NAAQMS limit</b>		<b>5.0 µg/m<sup>3</sup></b>	<b>NS</b>	<b>4.0 mg/m<sup>3</sup></b>	<b>NS</b>
AL6 -1	06.10.2021	1.03	BQL	1.79	510
AL6 – 2	08.10.2021	1.11	BQL	1.84	502
AL6 – 3	13.10.2021	1.14	BQL	1.72	511
AL6 – 4	15.10.2021	1.11	BQL	1.69	496
AL6 – 5	20.10.2021	1.18	BQL	1.88	499
AL6 – 6	22.10.2021	1.06	BQL	1.87	502
AL6 – 7	27.10.2021	1.10	BQL	1.74	506
AL6 – 8	29.10.2021	1.01	BQL	1.7	512
<b>Monthly Average</b>		<b>1.09</b>	<b>-</b>	<b>1.78</b>	<b>505</b>
<b>Standard Deviation</b>		<b>0.06</b>	<b>-</b>	<b>0.08</b>	<b>6</b>

\* NMHC- Non- Methane Hydrocarbons

BQL- Below Quantification Limit (Quantification Limit – NMHC: 0.5 ppm)

NS- Not Specified

The mean TSPM values at Tuna Port were 362 µg/m<sup>3</sup>, The mean PM<sub>10</sub> values were 163 µg/m<sup>3</sup>, which is above the permissible limit. PM<sub>2.5</sub> values were above the permissible limit (mean = 96 µg/m<sup>3</sup>). The average values of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were 3.13 µg/m<sup>3</sup>, 17.65 µg/m<sup>3</sup> and 9.41 µg/m<sup>3</sup> respectively and were all within the permissible limit.

The levels of Benzene, Hydrocarbons (HC) and CO were within the permissible limit at Tuna Port. The mean Benzene concentration was 1.09 µg/m<sup>3</sup>, well below the permissible limit of 5.0 µg/m<sup>3</sup>. HC's were below the detectable limit and Carbon Monoxide concentration was 1.78 mg/m<sup>3</sup>, well below the permissible limit of 4.0 mg/m<sup>3</sup>.

**Environmental Monitoring Report of Deendayal Port Trust, OCTOBER-2021**

**Location 7: Signal Building (Vadinar) (AL-7)**

**Table 7 : Results of Air Pollutant Concentration at Signal Building**

Parameters	Date	TSPM [µg/m <sup>3</sup> ]	PM10 [µg/m <sup>3</sup> ]	PM2.5 [µg/m <sup>3</sup> ]	SO <sub>2</sub> [µg/m <sup>3</sup> ]		NO <sub>x</sub> [µg/m <sup>3</sup> ]		NH <sub>3</sub> [µg/m <sup>3</sup> ]	
					8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
Sampling Period	-	24hr	24hr	24hr	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
NAAQMS limit	-	NS	100 µg/m <sup>3</sup>	60 µg/m <sup>3</sup>	-	80 µg/m <sup>3</sup>	-	80 µg/m <sup>3</sup>	-	400 µg/m <sup>3</sup>
AL7 -1	06.10.2021	115	73	30	4.40	3.52	19.69	18.63	6.89	6.21
					3.52		22.23		6.38	
					2.64		13.97		5.36	
AL7 -2	08.10.2021	111	62	34	3.96	2.78	16.51	16.94	8.42	8.42
					1.76		20.33		6.13	
					2.64		13.97		10.72	
AL7 -3	13.10.2021	198	107	52	3.08	3.08	16.51	16.51	4.60	5.45
					3.96		23.50		5.11	
					2.20		9.53		6.64	
AL7 -4	15.10.2021	146	72	50	3.96	4.40	15.24	12.91	7.91	7.06
					5.28		11.43		9.96	
					3.96		12.07		3.32	
AL7 -5	20.10.2021	171	85	44	3.08	2.64	8.89	9.32	6.89	7.23
					2.20		8.26		8.93	
					2.64		10.80		5.87	
AL7 -6	22.10.2021	178	88	71	3.08	4.54	14.61	12.49	8.42	8.17
					4.84		9.53		8.68	
					5.71		13.34		7.40	
AL7 -7	27.10.2021	160	80	52	3.08	3.81	6.35	11.22	10.98	8.25
					2.64		15.24		5.36	
					5.71		12.07		8.42	
AL7 -8	29.10.2021	177	89	56	2.20	3.52	9.53	10.16	8.42	5.96
					3.96		12.07		3.32	
					4.40		8.89		6.13	
<b>Monthly Average</b>		157	82	49		3.5		13.5		7.1
<b>Standard Deviation</b>		31	14	13		0.7		3.4		1.1

NS: Not Specified

<b>Table 7B : Results of Air Pollutant Concentration at Signal Building</b>					
<b>Parameter</b>	<b>Date</b>	<b>C<sub>6</sub>H<sub>6</sub> [µg/m<sup>3</sup>]</b>	<b>HC*</b>	<b>CO [mg/m<sup>3</sup>]</b>	<b>CO<sub>2</sub> [ppm ]</b>
<b>Sampling Period</b>		<b>8 hr</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>
<b>NAAQMS limit</b>		<b>5.0 µg/m<sup>3</sup></b>	<b>NS</b>	<b>4.0 mg/m<sup>3</sup></b>	<b>NS</b>
AL7 -1	06.10.2021	1.10	BQL	1.71	466
AL7 – 2	08.10.2021	1.13	BQL	1.62	488
AL7 – 3	13.10.2021	1.06	BQL	1.66	479
AL7 – 4	15.10.2021	1.11	BQL	1.72	480
AL7 – 5	20.10.2021	1.16	BQL	1.59	486
AL7 – 6	22.10.2021	1.17	BQL	1.66	477
AL7 – 7	27.10.2021	1.04	BQL	1.79	468
AL7 – 8	29.10.2021	1.10	BQL	1.64	470
<b>Monthly Average</b>		<b>1.11</b>	<b>-</b>	<b>1.67</b>	<b>477</b>
<b>Standard Deviation</b>		<b>0.04</b>	<b>-</b>	<b>0.06</b>	<b>8</b>

\*NMHC- Non- Methane Hydrocarbons

BQL- Below Quantification Limit (Quantification Limit – NMHC: 0.5 ppm)

NS\_ Not Specified

The mean TSPM values at Vadinar Port were 157 µg/m<sup>3</sup>. The mean PM<sub>10</sub> values were 82 µg/m<sup>3</sup>, which is below the permissible limit. PM<sub>2.5</sub> values were also within the permissible limit (mean = 49 µg/m<sup>3</sup>). The average values of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were 3.5 µg/m<sup>3</sup>, 13.5 µg/m<sup>3</sup> and 7.1 µg/m<sup>3</sup> respectively and were all within the permissible limit.

The levels of Benzene, Hydrocarbons (HC) and CO were within the permissible limit at Vadinar Port. The mean Benzene concentration was 1.11 µg/m<sup>3</sup>, well below the permissible limit of 5.0 µg/m<sup>3</sup>. HC's were below the detectable limit and Carbon Monoxide concentration was 1.67 mg/m<sup>3</sup>, well below the permissible limit of 4.0 mg/m<sup>3</sup>.



**Location 8: Admin Building (Vadinar) (AL-8)**

Table 8 : Results of Air Pollutant Concentration at Admin Building										
Parameters	Date	TSPM [µg/m3]	PM10 [µg/m3]	PM2.5 [µg/m3]	SO2 [µg/m3]		NOx [µg/m3]		NH3 [µg/m3]	
Sampling Period	-	24hr	24hr	24hr	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
NAAQMS limit	-	NS	100 µg/m3	60 µg/m3	-	80 µg/m3	-	80 µg/m3	-	400 µg/m3
AL8 -1	06.10.2021	221	113	82	2.64	3.22	8.89	12.28	7.40	5.79
					3.08		14.61		5.87	
					3.96		13.34		4.08	
AL8 -2	08.10.2021	218	126	73	4.40	5.42	27.95	18.21	4.08	6.81
					5.28		15.88		10.72	
					6.59		10.80		5.62	
AL8 -3	13.10.2021	197	104	72	3.08	3.22	10.16	17.57	5.87	9.02
					3.96		26.68		11.74	
					2.64		15.88		9.45	
AL8 -4	15.10.2021	227	111	75	2.20	3.37	20.96	15.24	8.42	6.30
					4.40		14.61		4.08	
					3.52		10.16		6.38	
AL8 -5	20.10.2021	185	88	54	4.40	3.52	15.24	16.73	8.42	6.98
					2.64		20.96		6.64	
					3.52		13.97		5.87	
AL8 -6	22.10.2021	248	121	94	3.96	3.81	8.89	10.37	5.36	5.19
					2.20		8.26		4.08	
					5.28		13.97		6.13	
AL8 -5	27.10.2021	210	138	62	3.08	3.66	13.97	13.76	15.06	10.64
					1.32		6.35		11.49	
					6.59		20.96		5.36	
AL8-6	29.10.2021	186	128	50	3.08	3.23	15.88	11.01	4.08	5.53
					2.20		8.89		5.87	
					4.40		8.26		6.64	
<b>Monthly Average</b>		211	116	70		3.7		14.4		7.0
<b>Standard Deviation</b>		22	16	15		0.7		3.0		1.9

NS: Not Specified

<b>Table 8B : Results of Air Pollutant Concentration at Admin Building</b>					
<b>Parameter</b>	<b>Date</b>	<b>C<sub>6</sub>H<sub>6</sub> [µg/m<sup>3</sup>]</b>	<b>HC*</b>	<b>CO [mg/m<sup>3</sup>]</b>	<b>CO<sub>2</sub> [ppm ]</b>
<b>Sampling Period</b>		<b>8 hr</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>
<b>NAAQMS limit</b>		<b>5.0 µg/m<sup>3</sup></b>	<b>NS</b>	<b>4.0 mg/m<sup>3</sup></b>	<b>NS</b>
AL8 -1	06.10.2021	1.30	BQL	1.86	451
AL8-2	08.10.2021	1.09	BQL	1.79	450
AL8 -3	13.10.2021	1.20	BQL	1.71	462
AL8-4	15.10.2021	1.11	BQL	1.82	455
AL8 -5	20.10.2021	1.16	BQL	1.69	469
AL8-6	22.10.2021	1.06	BQL	1.77	470
AL8-7	27.10.2021	1.30	BQL	1.82	459
AL8-8	29.10.2021	1.10	BQL	1.74	466
Monthly Average		<b>1.17</b>	-	<b>1.78</b>	<b>460</b>
Standard Deviation		<b>0.09</b>	-	<b>0.06</b>	<b>8</b>

\* NMHC- Non- Methane Hydrocarbons

BQL- Below Quantification Limit (Quantification Limit – NMHC: 0.5 ppm)

NS-Not Specified

The overall values of TSPM, PM<sub>10</sub>, PM<sub>2.5</sub>, SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> at Admin Building Vadinar was comparatively low among all the locations of Air Quality monitoring in Kandla Port and Vadinar Port. The mean TSPM values at Vadinar Port were 211 µg/m<sup>3</sup>. The mean PM<sub>10</sub> values were 116 µg/m<sup>3</sup>, which is above the permissible limit. PM<sub>2.5</sub> values were above the permissible limit (mean = 70.0 µg/m<sup>3</sup>). The average values of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were 3.7 µg/m<sup>3</sup>, 14.4 µg/m<sup>3</sup> and 7.0 µg/m<sup>3</sup> respectively and were all within the permissible limit.

The levels of Benzene, Hydrocarbons (HC) and CO were within the permissible limit at Admin Building, Vadinar Port. The mean Benzene concentration was 1.17 µg/m<sup>3</sup>, well below the permissible limit of 5.0 µg/m<sup>3</sup>. HC's were below the detectable limit and Carbon Monoxide concentration was 1.78 mg/m<sup>3</sup>, well below the permissible limit of 4.0 mg/m<sup>3</sup>.

#### **1.4 Observations and Conclusion**

During the monitoring period, the overall Ambient Air Quality of the port area was found to be well within the desired levels for various gaseous pollutants. However, Particulate matter as PM<sub>10</sub> and PM<sub>2.5</sub> was found to exceed the limits at locations like Near Coal storage area, Marine Bhavan, Estate Office, Tuna Port and Oil Jetty area.

## 2. Drinking Water Quality Monitoring

Drinking Water Quality Monitoring was carried out at twenty stations at Kandla, Vadinar & Township Area of Deendayal Port.

### 2.1 Drinking Water Monitoring Methodology

Drinking water samples were collected from 20 locations as prescribed in the tender document. Samples for physico-chemical analysis were collected in 1 liter carboys and samples for microbiological parameters were collected in sterilized bottles. These samples were then analyzed in laboratory for various drinking water parameters at Kandla Lab/Surat.

The Sampling and Analysis was done as per standard methods - IS 10500:2012. The water samples were analyzed for various parameters, viz. Color , Odor, Turbidity , Conductivity , pH , Chlorides , TDS, Total Hardness, Iron , Sulphate , Salinity , DO, BOD, Na, K, Ca, Mg, F, NO<sub>3</sub>, NO<sub>2</sub>, Mn, Cr-6, Cu, Cd, As, Hg, Pb, Zn, Bacterial Count (cfu) .

### 2.2 Results

The Drinking Water Quality monitoring data for 20 stations are given in below from table No. 9 to Table No. 15

**Environmental Monitoring Report of Deendayal Port Trust, OCTOBER-2021**

**Table 9: Drinking Water Quality Monitoring Parameters for Nirman Building 1, P & C building & Main Gate (North) at Kandla**

Sr. No.	Parameter	Unit	Nirman Building 1	P & C building	Main Gate North	Acceptable Limits as per IS 10500 : 2012	Permissible Limits in the absence of Alternate Source as per IS 10500 : 2012
1	pH	pH Unit	7.3	7.4	7.3	6.5 to 8.5	6.5 to 8.5
2	Total Dissolved Solids	mg/l	970	1310	1250	500	2000
3	Turbidity	NTU	0	1	0	1.0	5.0
4	Odor	-	Odorless	Odorless	Odorless	Agreeable	Agreeable
5	Color	Hazen Units	Colorless	Colorless	Colorless	5.0	15.0
6	Conductivity	µs/cm	1860	2560	2430	NS*	NS*
7	Biochemical Oxygen Demand	mg/l	<2	<2	<2	NS*	NS*
8	Chloride as Cl	mg/l	476.06	440.98	506.13	250.0	1000.0
9	Ca as Ca	mg/l	64.13	68.14	72.14	75.0	200.0
10	Mg as Mg	mg/l	63.18	70.47	65.61	30.0	100.0
11	Total Hardness	mg/l	420	460	450	200.0	600.0
12	Iron as Fe	mg/l	<0.01	<0.01	<0.01	0.3	No Relaxation
13	Fluorides as F	mg/l	0.32	0.35	0.17	1.0	1.5
14	Sulphate as SO4	mg/l	286.8	289.2	194.4	200.0	400
15	Nitrite as NO2	mg/l	<0.01	<0.01	<0.01	NS*	NS*
16	Nitrate as NO3	mg/l	6.41	7.88	13.02	45.0	No Relaxation
17	Salinity	%	0.86	0.80	0.91	NS*	NS*
18	Sodium as Na	mg/l	199	193	258	NS*	NS*
19	Potassium as K	mg/l	3.24	3.68	3.51	NS*	NS*
20	Manganese	mg/l	<0.04	<0.04	<0.04	0.1	0.3
21	Hexavalent Chromium	mg/l	<0.03	<0.03	<0.03	NS*	NS*
22	Copper	mg/l	<0.05	<0.05	<0.05	0.05	1.5
23	Cadmium	mg/l	<0.002	<0.002	<0.002	0.003	0.003
24	Arsenic	mg/l	<0.01	<0.01	<0.01	0.01	0.05
25	Mercury	mg/l	<0.001	<0.001	<0.001	0.001	0.001
26	Lead	mg/l	<0.01	<0.01	<0.01	0.01	0.01
27	Zinc	mg/l	<0.1	<0.1	<0.1	5.0	15.0
28	Bacterial Count	CFU/100ml	Absent	Absent	Absent	Absent	Absent

\*NS: Not Specified

**Environmental Monitoring Report of Deendayal Port Trust, OCTOBER-2021**

**Table 10: Drinking Water Quality Monitoring Parameters for Canteen, West Gate – I & Wharf Area at Kandla**

Sr. No.	Parameter	Unit	Canteen	West Gate – I	Wharf Area	Acceptable Limits as per IS 10500 : 2012	Permissible Limits in the absence of Alternate Source as per IS 10500 : 2012
1	pH	pH Unit	7.7	7.4	7.2	6.5 to 8.5	6.5 to 8.5
2	Total Dissolved Solids	mg/l	1340	980	1040	500	2000
3	Turbidity	NTU	0	0	1	1.0	5.0
4	Odor	-	Odorless	Odorless	Odorless	Agreeable	Agreeable
5	Color	Hazen Units	Colorless	Colorless	Colorless	5.0	15.0
6	Conductivity	µs/cm	2600	1940	2040	NS*	NS*
7	Biochemical Oxygen Demand	mg/l	<2	<2	<2	NS*	NS*
8	Chloride as Cl	mg/l	526.17	496.10	481.07	250.0	1000.0
9	Ca as Ca	mg/l	60.12	64.13	56.11	75.0	200.0
10	Mg as Mg	mg/l	70.47	65.61	70.47	30.0	100.0
11	Total Hardness	mg/l	440	430	430	200.0	600.0
12	Iron as Fe	mg/l	<0.01	<0.01	<0.01	0.3	No Relaxation
13	Fluorides as F	mg/l	0.46	0.50	0.52	1.0	1.5
14	Sulphate as SO <sub>4</sub>	mg/l	186	194.4	288	200.0	400
15	Nitrite as NO <sub>2</sub>	mg/l	<0.01	<0.01	<0.01	NS*	NS*
16	Nitrate as NO <sub>3</sub>	mg/l	8.59	10.21	9.22	45.0	No Relaxation
17	Salinity	%	0.95	0.90	0.87	NS*	NS*
18	Sodium as Na	mg/l	215	209	231	NS*	NS*
19	Potassium as K	mg/l	4.23	3.88	4.21	NS*	NS*
20	Manganese	mg/l	<0.04	<0.04	<0.04	0.1	0.3
21	Hexavalent Chromium	mg/l	<0.03	<0.03	<0.03	NS*	NS*
22	Copper	mg/l	<0.05	<0.05	<0.05	0.05	1.5
23	Cadmium	mg/l	<0.002	<0.002	<0.002	0.003	0.003
24	Arsenic	mg/l	<0.01	<0.01	<0.01	0.01	0.05
25	Mercury	mg/l	<0.001	<0.001	<0.001	0.001	0.001
26	Lead	mg/l	<0.01	<0.01	<0.01	0.01	0.01
27	Zinc	mg/l	<0.1	<0.1	<0.1	5.0	15.0
28	Bacterial Count	CFU/100ml	Absent	Absent	Absent	Absent	Absent

\*NS: Not Specified

**Table 11: Drinking Water Quality Monitoring Parameters for Sewa sadan – 3, Workshop I & Custom Building at Kandla**

Sr. No.	Parameter	Unit	SewaSadan – 3	Workshop	Custom Building	Acceptable Limits as per IS 10500 : 2012	Permissible Limits in the absence of Alternate Source as per IS 10500 : 2012
1	pH	pH Unit	7.4	7.8	7.5	6.5 to 8.5	6.5 to 8.5
2	Total Dissolved Solids	mg/l	1190	1420	1160	500	2000
3	Turbidity	NTU	1	1	0	1.0	5.0
4	Odor	-	Odorless	Odorless	Odorless	Agreeable	Agreeable
5	Color	Hazen Units	Colorless	Colorless	Colorless	5.0	15.0
6	Conductivity	µs/cm	1920	2870	2180	NS*	NS*
7	Biochemical Oxygen Demand	mg/l	<2	<2	<2	NS*	NS*
8	Chloride	mg/l	616.37	591.31	491.09	250.0	1000.0
9	Ca as Ca	mg/l	64.13	68.14	60.12	75.0	200.0
10	Mg as Mg	mg/l	68.04	58.32	65.61	30.0	100.0
11	Total Hardness	mg/l	440	410	420	200.0	600.0
12	Iron as Fe	mg/l	<0.01	<0.01	<0.01	0.3	No Relaxation
13	Fluorides	mg/l	0.90	0.57	0.63	1.0	1.5
14	Sulphate	mg/l	217.2	205.2	289.2	200.0	400
15	Nitrite	mg/l	<0.01	<0.01	<0.01	NS*	NS*
16	Nitrate	mg/l	13.52	9.93	12.74	45.0	No Relaxation
17	Salinity	%	1.11	1.07	0.89	NS*	NS*
18	Sodium as Na	mg/l	265	218	323	NS*	NS*
19	Potassium as K	mg/l	3.87	3.73	3.56	NS*	NS*
20	Manganese	mg/l	<0.04	<0.04	<0.04	0.1	0.3
21	Hexavalent Chromium	mg/l	<0.03	<0.03	<0.03	NS*	NS*
22	Copper	mg/l	<0.05	<0.05	<0.05	0.05	1.5
23	Cadmium	mg/l	<0.002	<0.002	<0.002	0.003	0.003
24	Arsenic	mg/l	<0.01	<0.01	<0.01	0.01	0.05
25	Mercury	mg/l	<0.001	<0.001	<0.001	0.001	0.001
26	Lead	mg/l	<0.01	<0.01	<0.01	0.01	0.01
27	Zinc	mg/l	<0.1	<0.1	<0.1	5.0	15.0
28	Bacterial Count	CFU/100ml	Absent	Absent	Absent	Absent	Absent

\*NS: Not Specified

**Environmental Monitoring Report of Deendayal Port Trust, OCTOBER-2021**

**Table 12: Drinking Water Quality Monitoring Parameters for Port Colony Kandla, Hospital Kandla & A.O. Building at Gandhidham**

Sr. No.	Parameter	Unit	Port Colony Kandla	Hospital Kandla	A.O. Building	Acceptable Limits as per IS 10500 : 2012	Permissible Limits in the absence of Alternate Source as per IS 10500 : 2012
1	pH	pH Unit	7.3	7.6	7.2	6.5 to 8.5	6.5 to 8.5
2	Total Dissolved Solids	mg/l	1090	1460	940	500	2000
3	Turbidity	NTU	0	0	1	1.0	5.0
4	Odor	-	Odorless	Odorless	Odorless	Agreeable	Agreeable
5	Color	Hazen Units	Colorless	Colorless	Colorless	5.0	15.0
6	Conductivity	µs/cm	2090	2850	1860	NS*	NS*
7	Biochemical Oxygen Demand	mg/l	<2	<2	<2	NS*	NS*
8	Chloride	mg/l	445.99	466.04	496.10	250.0	1000.0
9	Ca as Ca	mg/l	56.11	68.14	60.12	75.0	200.0
10	Mg as Mg	mg/l	75.33	72.90	80.19	30.0	100.0
11	Total Hardness	mg/l	450	470	480	200.0	600.0
12	Iron as Fe	mg/l	<0.01	<0.01	<0.01	0.3	No Relaxation
13	Fluorides	mg/l	0.87	0.68	0.92	1.0	1.5
14	Sulphate	mg/l	294	318	210	200.0	400
15	Nitrite	mg/l	<0.01	<0.01	<0.01	NS*	NS*
16	Nitrate	mg/l	9.29	10.14	11.69	45.0	No Relaxation
17	Salinity	%	0.81	0.84	0.90	NS*	NS*
18	Sodium as Na	mg/l	101	221	402	NS*	NS*
19	Potassium as K	mg/l	4.08	4.01	3.99	NS*	NS*
20	Manganese	mg/l	<0.04	<0.04	<0.04	0.1	0.3
21	Hexavalent Chromium	mg/l	<0.03	<0.03	<0.03	NS*	NS*
22	Copper	mg/l	<0.05	<0.05	<0.05	0.05	1.5
23	Cadmium	mg/l	<0.002	<0.002	<0.002	0.003	0.003
24	Arsenic	mg/l	<0.01	<0.01	<0.01	0.01	0.05
25	Mercury	mg/l	<0.001	<0.001	<0.001	0.001	0.001
26	Lead	mg/l	<0.01	<0.01	<0.01	0.01	0.01
27	Zinc	mg/l	<0.1	<0.1	<0.1	5.0	15.0
28	Bacterial Count	CFU/100ml	Absent	Absent	Absent	Absent	Absent

\*NS: Not Specified

**Environmental Monitoring Report of Deendayal Port Trust, OCTOBER-2021**

**Table 13: Drinking Water Quality Monitoring Parameters for School Gopalpuri, Guest House & E - Type Quarter at Gopalpuri, Gandhidham**

Sr. No.	Parameter	Unit	School Gopalpuri	Guest House	E - Type Quarter	Acceptable Limits as per IS 10500 : 2012	Permissible Limits in the absence of Alternate Source as per IS 10500 : 2012
1	pH	pH Unit	7.4	7.4	7.3	6.5 to 8.5	6.5 to 8.5
2	Total Dissolved Solids	mg/l	1020	1340	1100	500	2000
3	Turbidity	NTU	1	0	1	1.0	5.0
4	Odor	-	Odorless	Odorless	Odorless	Agreeable	Agreeable
5	Color	Hazen Units	Colorless	Colorless	Colorless	5.0	15.0
6	Conductivity	µs/cm	2010	2660	2140	NS*	NS*
7	Biochemical Oxygen Demand	mg/l	<2	<2	<2	NS*	NS*
8	Chloride	mg/l	466.04	506.13	466.04	250.0	1000.0
9	Ca as Ca	mg/l	52.10	72.14	68.14	75.0	200.0
10	Mg as Mg	mg/l	77.76	55.89	63.18	30.0	100.0
11	Total Hardness	mg/l	450	410	430	200.0	600.0
12	Iron as Fe	mg/l	<0.01	<0.01	<0.01	0.3	No Relaxation
13	Fluorides	mg/l	0.60	0.81	1.08	1.0	1.5
14	Sulphate	mg/l	291.6	294	283.2	200.0	400
15	Nitrite	mg/l	<0.01	<0.01	<0.01	NS*	NS*
16	Nitrate	mg/l	9.29	10.63	9.36	45.0	No Relaxation
17	Salinity	%	0.84	0.91	0.84	NS*	NS*
18	Sodium as Na	mg/l	275	300	130	NS*	NS*
19	Potassium as K	mg/l	3.49	3.50	3.56	NS*	NS*
20	Manganese	mg/l	<0.04	<0.04	<0.04	0.1	0.3
21	Hexavalent Chromium	mg/l	<0.03	<0.03	<0.03	NS*	NS*
22	Copper	mg/l	<0.05	<0.05	<0.05	0.05	1.5
23	Cadmium	mg/l	<0.002	<0.002	<0.002	0.003	0.003
24	Arsenic	mg/l	<0.01	<0.01	<0.01	0.01	0.05
25	Mercury	mg/l	<0.001	<0.001	<0.001	0.001	0.001
26	Lead	mg/l	<0.01	<0.01	<0.01	0.01	0.01
27	Zinc	mg/l	<0.1	<0.1	<0.1	5.0	15.0
28	Bacterial Count	CFU/100ml	Absent	Absent	Absent	Absent	Absent

\*NS: Not Specified



**Environmental Monitoring Report of Deendayal Port Trust, OCTOBER-2021**

**Table 14: Drinking Water Quality Monitoring Parameters for F - Type Quarter, Hospital Gopalpuri & Tuna Port**

Sr. No.	Parameter	Unit	F - Type Quarter	Hospital Gopalpuri	Tuna Port	Acceptable Limits as per IS 10500 : 2012	Permissible Limits in the absence of Alternate Source as per IS 10500 : 2012
1	pH	pH Unit	7.7	7.4	7.38	6.5 to 8.5	6.5 to 8.5
2	Total Dissolved Solids	mg/l	1120	1090	1080	500	2000
3	Turbidity	NTU	0	1	0	1.0	5.0
4	Odor	-	Odorless	Odorless	Odorless	Agreeable	Agreeable
5	Color	Hazen Unit	Colorless	Colorless	Colorless	5.0	15.0
6	Conductivity	µs/cm	2210	2190	2160	NS*	NS*
7	Biochemical Oxygen Demand	mg/l	<2	<2	<2	NS*	NS*
8	Chloride	mg/l	576.28	521.16	520	250.0	1000.0
9	Ca as Ca	mg/l	72.14	76.15	78.56	75.0	200.0
10	Mg as Mg	mg/l	55.89	65.61	54.92	30.0	100.0
11	Total Hardness	mg/l	410	460	422	200.0	600.0
12	Iron as Fe+3	mg/l	<0.01	<0.01	<0.01	0.3	No Relaxation
13	Fluorides	mg/l	1.05	0.93	0.46	1.0	1.5
14	Sulphate	mg/l	265.2	238.8	180	200.0	400
15	Nitrite	mg/l	<0.01	<0.01	<0.01	NS*	NS*
16	Nitrate	mg/l	9.86	7.88	8.2	45.0	No Relaxation
17	Salinity	%	1.04	0.94	0.98	NS*	NS*
18	Sodium as Na	mg/l	235	235	260	NS*	NS*
19	Potassium as K	mg/l	2.18	4.01	2.6	NS*	NS*
20	Manganese	mg/l	<0.04	<0.04	<0.04	0.1	0.3
21	Hexavalent Chromium	mg/l	<0.03	<0.03	<0.03	NS*	NS*
22	Copper	mg/l	<0.05	<0.05	<0.05	0.05	1.5
23	Cadmium	mg/l	<0.002	<0.002	<0.002	0.003	0.003
24	Arsenic	mg/l	<0.01	<0.01	<0.01	0.01	0.05
25	Mercury	mg/l	<0.001	<0.001	<0.001	0.001	0.001
26	Lead	mg/l	<0.01	<0.01	<0.01	0.01	0.01
27	Zinc	mg/l	<0.1	<0.1	<0.1	5.0	15.0
28	Bacterial Count	CFU/100ml	Absent	Absent	Absent	Absent	Absent

\*NS: Not Specified

**Table 15: Drinking Water Quality Monitoring Parameters for Vadinar Jetty & Port Colony at Vadinar**

Sr. No.	Parameter	Unit	Vadinar Jetty	Port Colony Vadinar	Acceptable Limits as per IS 10500 : 2012	Permissible Limits in the absence of Alternate Source as per IS 10500 : 2012
1	pH	pH Unit	7.5	7.6	6.5 to 8.5	6.5 to 8.5
2	Total Dissolved Solids	mg/l	1030	1010	500	2000
3	Turbidity	NTU	ND	ND	1.0	5.0
4	Odor	-	Odorless	Odorless	Agreeable	Agreeable
5	Color	Hazen Units	Colorless	Colorless	5.0	15.0
6	Conductivity	µs/cm	2020	1960	NS*	NS*
7	Biochemical Oxygen Demand	mg/l	<2	<2	NS*	NS*
8	Chloride	mg/l	425.95	415.92	250.0	1000.0
9	Ca as Ca	mg/l	60.12	52.10	75.0	200.0
10	Mg as Mg	mg/l	75.33	94.26	30.0	100.0
11	Total Hardness	mg/l	460	440	200.0	600.0
12	Iron as Fe+3	mg/l	<0.01	<0.01	0.3	No Relaxation
13	Fluorides	mg/l	0.92	0.67	1.0	1.5
14	Sulphate	mg/l	22.44	22.20	200.0	400
15	Nitrite	mg/l	<0.01	<0.01	NS*	NS*
16	Nitrate	mg/l	7.39	9.99	45.0	No Relaxation
17	Salinity	%	0.90	0.92	NS*	NS*
18	Sodium as Na	mg/l	51.1	44.1	NS*	NS*
19	Potassium as K	mg/l	2.2	<2.0	NS*	NS*
20	Manganese	mg/l	<0.04	<0.04	0.1	0.3
21	Hexavalent Chromium	mg/l	<0.03	<0.03	NS*	NS*
22	Copper	mg/l	<0.05	<0.05	0.05	1.5
23	Cadmium	mg/l	<0.002	<0.002	0.003	0.003
24	Arsenic	mg/l	<0.01	<0.01	0.01	0.05
25	Mercury	mg/l	<0.001	<0.001	0.001	0.001
26	Lead	mg/l	<0.01	<0.01	0.01	0.01
27	Zinc	mg/l	<0.1	<0.1	5.0	15.0
28	Bacterial Count	CFU/100ml	Absent	Absent	Absent	Absent

\*NS: Not Specified

### 2.3 Results & Discussion

The colour of all drinking water samples was < 5 Hazen unit and odour of the samples was also agreeable. All parameters are found to be within the specified limit of the Drinking water Standard.

#### **pH**

The limit of pH value for drinking water is specified as 6.5 to 8.5. pH value in the studied area varied from 7.0 to 8.0 pH unit. All the sampling points showed pH values within the prescribed limit by Indian Standards.

#### **Total Dissolved Solids (TDS)**

TDS values in the studied area varied between 900 -1500 mg/l. None of the sampling points showed higher TDS values than the prescribed limit by Indian standards.

#### **Conductivity**

Electrical Conductivity is the ability of a solution to transfer (conduct) electric current. Conductivity is used to measure the concentration of dissolved solids which have been ionized in a polar solution such as water. The conductivity in the samples collected during the month of October ranged from 1800-3000  $\mu\text{s}/\text{cm}$ . Electrical conductivity standards do not appear in BIS standards for drinking water.

#### **BOD**

BOD value in the studied area was less than 2.0 mg/L. Indian standards does not show any standard values for BOD in drinking water.

#### **Chlorides**

Excessive chloride concentration increase rates of corrosion of metals in the distribution system. This can lead to increased concentration of metals in the supply. Chloride value in the studied area varied between 400-650 mg/l and is found to be within the Permissible limit of the Drinking Water Standard.

#### **Calcium**

Calcium value in the studied area varied between 50 - 80 mg/l and is found to be within the Permissible limit of the Drinking Water Standard. If calcium is present beyond the maximum acceptable limit, it causes incrustation of pipes.

### **Magnesium**

Magnesium value in the studied area varied between 50 – 99 mg/l. All the locations had Magnesium within the prescribed limits of 30-100 mg/L.

### **Total Hardness**

Hardness value in the studied area varied between 400-480 mg/l and is found to be within the Permissible limit of the Drinking Water Standard. The prescribed limit by Indian Standards is 200-600 mg/L.

### **Iron**

Iron value in the studied area was below 0.01mg/L and hence well below the permissible limit as per Indian Standards is 0.3 mg/L. The excess amount of iron causes slight toxicity; gives stringent taste to water.

### **Fluoride**

Fluoride value in the studied area varied between 0.1 – 1.0 mg/l and hence well below the permissible limit as per Indian Standards is 1.0-1.5 mg/L. Moderate amounts lead to dental effects, but long-term ingestion of large amounts can lead to potentially severe skeletal problems.

### **Sulphates**

Sulphate value in the studied area varied between 20 – 350 mg/l. All the sampling points showed sulphate values within the prescribed limits by Indian Standards (200-400 mg/L). Sulphate content in drinking water exceeding the 400 mg/L imparts bitter taste.

### **Nitrites (NO<sub>2</sub>) and Nitrates (NO<sub>3</sub>)**

Nitrite values in all the water samples were <0.1. There are no specified standard values for Nitrites in Drinking water. The mean Nitrate values in drinking water of KPT was 6.41 mg/l which is well within the permissible limit of the Drinking water Standard.

### **Salinity**

Salinity in drinking water in the present samples collected ranged from 0.5 to 1.4 %. There are no prescribed Indian standards for salinity in Drinking water.

### **Sodium and Potassium Salts**

Sodium values in the samples collected ranged from 50 - 400 mg/l and Potassium salts ranged from 2.0 to 4.5 mg/l. There are no prescribed limits of Sodium and Potassium in Indian standards for Drinking water.

### **Heavy Metals in Drinking Water**

In the present study period drinking water samples were analyzed for Mn, Cr, Cu, Cd, As, Hg, Pb and Zn. All these heavy metals were well below the permissible limits prescribed by the Indian Standards.

### **Bacteriological Study**

Analysis of the bacteriological parameter at all location shows that Bacteria is not present and hence Bacterial count is in line with the permissible limit of drinking water. This shows that all the drinking water samples were safe from any bacteriological contamination.

## **2.4 Conclusions**

These results are compared with acceptable limits as prescribed in IS 10500:2012 – Drinking Water Specification. It is seen from the analysis data that during the study period the water was safe for human consumption at all drinking water monitoring stations.

### 3. Noise Level Monitoring

Noise sources in port operations include cargo handling, vehicular traffic, and loading / unloading containers and ships. Noise Monitoring was done at 13 stations at Kandla, Vadinar and Township area.

#### 3.1 Method of Monitoring

Sampling was done at all stations for 24 hour period. Data was recorded using automated sound level meter. The intensity of sound was measured in sound pressure level (SPL) and common unit of measurement is decibel (Db).

#### 3.2 Results

Table 16: Noise Monitoring data for ten locations of Deendayal Port and two locations of Vadinar Port

Sr. No.	Location	Day Time Average Noise Level (SPL) in dB(A)	Night Time Average Noise Level (SPL) in dB(A)
	Sampling Time	6:00 am to 10:00 PM	10:00PM to 6:00 AM
1	Marine Bhavan	73.9	53.4
2	Nirman Building 1	62.3	51.0
3	Tuna Port	57.2	50.9
4	Main Gate North	67.0	61.8
5	West Gate I	70.5	65.1
6	Canteen Area	64.5	51.0
7	Main Road	68.4	51.5
8	ATM Building	80.4	57.3
9	Wharf Area /Jetty Area	72.9	68.1
10	Port & Custom Office	67.8	41.8
<b>Vadinar Port</b>			
11	Entrance Gate of Vadinar Port	66.4	53.2
12	Nr. Port Colony, Vadinar	60.7	54.1
13	Nr. Vadinar Jetty	72.4	66.5

**3.3 Conclusions-** Noise sources in port operations include cargo handling, vehicular traffic, and loading / unloading containers and ships. The Day Time Average Noise Level (SPL) in all 13 locations at Deendayal Port ranged from 57.0 dB(A) to 73.9 dB(A) and it was within the permissible limits of 75 dB(A) for the industrial area for the daytime. The Night Time Average Noise Level (SPL) in all 13 locations of Deendayal Port ranged from 41.8 dB to 68.4 dB(A) and it was within the permissible limits of 70 dB(A) for the industrial area for the night time.

#### 4. Soil Monitoring

Sampling and analysis of soil samples were undertaken at six locations within the study area (Deendayal Port and Vadinar Port) as a part of EMP. The soil sampling locations are initially decided based on the locations as provided in the tender document of the Deendayal Port.

##### 4.1 Methodology

The soil samples were collected in the month of October 2021. The samples collected from the all locations are homogeneous representative of each location. At random locations were identified at each location and soil was dug from 30 cm below the surface. It was uniformly mixed before homogenizing the soil samples. The samples were filled in polythene bags, labeled in the field with number and site name and sent to laboratory for analysis.

##### 4.2 Results

**Table-17: Chemical Characteristics of Soil in the Study Area**

Sr. No.	Parameter	Unit	Station Name					
			SL1	SL2	SL3	SL4	SL5	SL6
			Tuna Port	IFFCO Plant	Khori Creek	Nakti Creek	KPT Admin Site	KPT Colony
			Near main gate of Port	10 m away from main gate	Sand from creek at low tide		Vadinar	
1	Texture		Sandy Loam	Sandy Loam	Sandy Loam	Sandy Loam	Sandy Loam	Sandy Loam
2	pH	-	8.62	8.10	8.75	8.33	8.10	8.22
3	Electrical Conductivity	µs/cm	16,200.0	26,820.0	16,252.0	17,520.0	560.0	480.0
4	Moisture	%	17.00	18.20	19.10	20.22	7.26	8.22
5	Total Organic Carbon	%	0.52	1.02	0.62	3.10	0.12	0.20
6	Alkalinity	mg/kg	60.06	80.44	140.20	80.44	60.06	80.44
7	Total Nitrogen	%	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001
8	Chloride	mg/kg	1,620.0	5,380.0	1,820.0	2,078.0	62.0	77.0
9	Sulphate	mg/kg	230.0	198.0	120.0	118.0	16.0	20.0
10	Phosphorus	mg/kg	0.90	0.82	0.96	1.02	0.80	0.72
11	Potassium	mg/kg	396.0	810.0	366.0	460.0	120.0	160.0
12	Sodium	mg/kg	1,620.0	3,400.0	2,122.0	2,012.0	910.0	888.0
13	Calcium	mg/kg	230.32	722.20	252.00	470.42	110.00	82.00
14	Copper as Cu	mg/kg	17.40	38.80	21.20	35.10	16.6	17.0
15	Lead as Pb	mg/kg	6.40	7.90	29.10	7.60	4.8	2.0
16	Nickel as Ni	mg/kg	33.50	13.90	34.50	13.20	13.2	12.2
17	Zinc as Zn	mg/kg	55.90	91.90	77.9	81.90	28.00	36.22
18	Cadmium as Cd	mg/kg	BQL	BQL	BQL	BQL	BQL	BQL

### 4.3 Discussion

- The data shows that value of pH ranges from 8.10 at IFFCO Plant to 8.75 at Khori Creek indicating that all soil samples are neutral to slight basic. Iffco Plant samples showed maximum conductivity of 26,820  $\mu\text{mhos/cm}$ , while Tuna Port location showed minimum conductivity of 16,200  $\mu\text{mhos/cm}$ . Conductivity at Vadinar Port was 560 and 480  $\mu\text{mhos/cm}$  at Admin site and Vadinar Port colony respectively.
- Total organic Carbon ranged from 0.5 % to 3.1 at Deendayal Port. At Vadinar Port, organic carbon content ranged from 0.1 % to 0.2 %.
- The concentration of Phosphorus and Potassium in the soil samples varies from 0.7 to 1.0 mg/kg and 300.0 to 800 mg/kg respectively at Deendayal Port. The mean concentration of Phosphorous at Vadinar site was 0.76 mg/kg and mean concentration of Potassium at Vadinar site was 140 mg/kg.

These differences in NPK in soil at different locations are due to the dissimilar nature of soil at each of the locations. Samples SL3 & SL4 (Khori Creek & Nakti Creek) are of saline nature as they are coastal soil; where as other locations are inland locations and have different chemical properties.

### Heavy Metals in the Soil

Traces of Copper, Lead, Nickel and Zinc were observed in the soil samples collected from all the four locations of Deendayal Port and two locations of Vadinar Port. Cadmium metal was not detected in the Soil.

### 4.4 Conclusion

The soils of Deendayal Port and Vadinar Port appears to be neutral to basic with varying levels of Chloride, Sulphate, NPK and Calcium. As the nature of soil at different locations are different with respect to its proximity to the sea, the samples showed high degree of variations in their chemical properties.



## 5. Sewage Treatment Plant Monitoring

This involves safe collection of waste water (spent/used water) from wash areas, bathroom, industrial units, etc., waste from toilets of various buildings and its conveyance to the treatment plant and final disposal in conformity with the requirement and guide lines of State Pollution Control Board and other statutory bodies.

### 5.1 Methodology for STP Monitoring

To monitor the working efficiency of Sewage Treatment Plant (STP), STP Inlet and Outlet Samples were collected once a week. Locations selected are namely Gopalpuri Township, Deendayal Port and Vadinar. Samples were collected in 1 lit. Carboys and were analyzed in laboratory for various parameters.

### 5.2 Results

- **Kandla STP**

**Table 18: Sewage Water Monitoring at Kandla STP (1<sup>st</sup> Week)**

Date of Sampling		08.10.2021		
Sr. No.	Parameters	Unit	Results	
			KPT STP I/L	KPT STP O/L
1	pH	pH unit	7.62	7.4
2	Total Suspended Solids	mg/l	99.2	64.7
3	Residual Chlorine	mg/l	-	<0.5
4	COD	mg/l	424.2	98.0
5	BOD @ 27 °C	mg/l	141.0	23.0
6.	Fecal Coliform	MPN Index / 100 ml	-	79.0
<b>Aeration Tank</b>				
7.	MLSS	mg/l	6.0	
8.	MLVSS	%	93.0	

**Table 19: Sewage Water Monitoring at Kandla STP (2<sup>nd</sup> Week)**

Date of Sampling		12.10.2021		
Sr. No.	Parameters	Unit	Results	
			KPT STP I/L	KPT STP O/L
1	pH	pH unit	7.6	7.2
2	Total Suspended Solids	mg/l	152.2	72.4
3	Residual Chlorine	mg/l	-	<0.5
4	COD	mg/l	384	103.0
5	BOD @ 27 °C	mg/l	120.0	24.0
6.	Fecal Coliform	MPN Index / 100 ml	-	8.0
<b>Aeration Tank</b>				
7.	MLSS	mg/l	12.0	
8.	MLVSS	%	84.0	

**Table 20: Sewage Water Monitoring at Kandla STP (3<sup>rd</sup> Week)**

Date of Sampling		21.10.2021		
Sr. No.	Parameters	Unit	Results	
			KPT STP I/L	KPT STP O/L
1	pH	pH unit	7.65	7.41
2	Total Suspended Solids	mg/l	223.4	99.8
3	Residual Chlorine	mg/l	-	<0.5
4	COD	mg/l	181.8	101
5	BOD @ 27 °C	mg/l	68.0	24.0
6.	Fecal Coliform	MPN Index / 100 ml	-	11.0
<b>Aeration Tank</b>				
7.	MLSS	mg/l	9.0	
8.	MLVSS	%	87.0	

**Table 21: Sewage Water Monitoring at Kandla STP (4<sup>th</sup> Week)**

Date of Sampling		25.10.2021		
Sr. No.	Parameters	Unit	Results	
			KPT STP I/L	KPT STP O/L
1	pH	pH unit	7.72	7.5
2	Total Suspended Solids	mg/l	284.6	113.6
3	Residual Chlorine	mg/l	-	<0.5
4	COD	mg/l	212	101.0
5	BOD @ 27 °C	mg/l	98.0	26.0
6.	Fecal Coliform	MPN Index / 100 ml	-	3.6
<b>Aeration Tank</b>				
7.	MLSS	mg/l	14.0	
8	MLVSS	%	84.0	

- **Gopalpuri Colony STP**

**Table 22: Sewage Water Monitoring at Gopalpuri STP (1<sup>st</sup> Week)**

Date of Sampling		08.10.2021		
Sr. No.	Parameters	Unit	Results	
			Gopalpuri STP I/L	Gopalpuri STP O/L
1	pH	pH unit	7.6	7.4
2	Total Suspended Solids	mg/l	195.6	84.0
3	Residual Chlorine	mg/l	-	<0.5
4	COD	mg/l	304.0	104.0
5	BOD @ 27 °C	mg/l	120.0	26.0
6.	Fecal Coliform	MPN Index / 100 ml	-	49.0
<b>Aeration Tank</b>				
7.	MLSS	mg/l	10.0	
8	MLVSS	%	87.0	

**Table 23: Sewage Water Monitoring at Gopalpuri STP (2<sup>nd</sup> Week)**

Date of Sampling		12.10.2021		
Sr. No.	Parameters	Unit	Results	
			Gopalpuri STP I/L	Gopalpuri STP O/L
1	pH	pH unit	7.7	7.5
2	Total Suspended Solids	mg/l	388	131.8
3	Residual Chlorine	mg/l	-	<0.5
4	COD	mg/l	414.1	106.00
5	BOD @ 27 °C	mg/l	139.0	28.0
6.	Fecal Coliform	MPN Index / 100 ml	-	6.0
<b>Aeration Tank</b>				
7.	MLSS	mg/l	14.0	
8	MLVSS	%	90.0	

**Table 24: Sewage Water Monitoring at Gopalpuri STP (3<sup>rd</sup> Week)**

Date of Sampling		21.10.2021		
Sr. No.	Parameters	Unit	Results	
			Gopalpuri STP I/L	Gopalpuri STP O/L
1	pH	pH unit	7.52	7.21
2	Total Suspended Solids	mg/l	354.2	103.3
3	Residual Chlorine	mg/l	-	<0.5
4	COD	mg/l	161.6	90.9
5	BOD @ 27 °C	mg/l	80.0	26.0
6.	Fecal Coliform	MPN Index / 100 ml	-	46.0
<b>Aeration Tank</b>				
7.	MLSS	mg/l	14.0	
8	MLVSS	%	84.0	

**Table 25: Sewage Water Monitoring at Gopalpuri STP (4<sup>th</sup> Week)**

Date of Sampling		25.10.2021		
Sr. No.	Parameters	Unit	Results	
			Gopalpuri STP I/L	Gopalpuri STP O/L
1	pH	pH unit	7.45	7.19
2	Total Suspended Solids	mg/l	345.8	105
3	Residual Chlorine	mg/l	-	<0.5
4	COD	mg/l	232	106
5	BOD @ 27 °C	mg/l	82.0	28.0
6.	Fecal Coliform	MPN Index / 100 ml	-	32.0
<b>Aeration Tank</b>				
7.	MLSS	mg/l	10.0	
8.	MLVSS	%	89.0	

- **Vadinar STP**

**Table 26: Sewage Water Monitoring at Vadinar STP (1<sup>st</sup> Week)**

Date of Sampling		08.10.2021		
Sr. No.	Parameters	Unit	Results	
			Vadinar STP I/L	Vadinar O/L
1	pH	pH unit	7.62	7.4
2	Total Suspended Solids	mg/l	99.2	64.7
3	Residual Chlorine	mg/l	-	<0.5
4	COD	mg/l	88.0	60.0
5	BOD @ 27 °C	mg/l	32.0	16.0

**Table 27: Sewage Water Monitoring at Vadinar STP (2<sup>nd</sup> Week)**

<b>Date of Sampling</b>	<b>12.10.2021</b>
-------------------------	-------------------

Sr. No.	Parameters	Unit	Results	
			Vadinar STP I/L	Vadinar O/L
1	pH	pH unit	7.7	7.5
2	Total Suspended Solids	mg/l	133.5	59.9
3	Residual Chlorine	mg/l	-	<0.5
4	COD	mg/l	188.0	60.0
5	BOD @ 27 °C	mg/l	60.0	16.0

**Table 28: Sewage Water Monitoring at Vadinar STP (3<sup>rd</sup> Week)**

<b>Date of Sampling</b>	<b>21.10.2021</b>
-------------------------	-------------------

Sr. No.	Parameters	Unit	Results	
			Vadinar STP I/L	Vadinar O/L
1	pH	pH unit	7.7	7.5
2	Total Suspended Solids	mg/l	105	38
3	Residual Chlorine	mg/l	-	<0.5
4	COD	mg/l	161.6	70.7
5	BOD @ 27 °C	mg/l	62.0	20.0

**Table 29: Sewage Water Monitoring at Vadinar STP (4<sup>th</sup> Week)**

<b>Date of Sampling</b>	<b>25.10.2021</b>
-------------------------	-------------------

Sr. No.	Parameters	Unit	Results	
			Vadinar STP I/L	Vadinar O/L
1	pH	pH unit	7.7	7.2
2	Total Suspended Solids	mg/l	105	58.8
3	Residual Chlorine	mg/l	-	<0.5
4	COD	mg/l	202	80.8
5	BOD @ 27 °C	mg/l	60.0	20.0

**5.3 Conclusions:**

The GPCB standards of BOD, TSS and Residual Chlorine for STP outlet are 20 mg/lit, 30 mg/lit & 0.5 mg/lit respectively. It is suggested to do treatment on regular basis to avoid flow of contaminated/polluted water into the sea.

## 6. Marine Water Monitoring

The Forty Second Amendment to the Constitution in 1976 underscored the importance of 'green thinking'. Article 48A enjoins the state to protect and improve the environment and safeguard the forests and wildlife in the country. Further, Article 51A(g) states that the "fundamental duty of every citizen is to protect and improve the natural environment including forests, lakes, rivers and wildlife and to have compassion for living creatures".

Policy Statement for Abatement of Pollution (1992) has suggested developing relevant legislation and regulation, fiscal incentives, voluntary agreements and educational programs and information campaigns. It emphasizes the need for integration by incorporating environmental considerations into decision making at all levels by adopting frameworks namely, pollution prevention at source, application of best practicable solution, ensure polluter pays for control of pollution, focus on heavily polluted areas and river stretches and involve public in decision-making. The National Conservation Strategy and Policy Statement on Environment and Development, (1992) aimed at "integrating environmental concerns with developmental imperatives to meet the challenges by redirecting the thrust of our developmental process so that the basic needs of our people could be fulfilled by making judicious and sustainable use of natural resources." The priorities mentioned in this policy document include the sustainable use of land and water resources, prevention and control of pollution and preservation of biodiversity.

The National Water Policy, (2002) contains provisions for developing, conserving, sustainable utilizing and managing this important water resources and need to be governed by national perspectives.

### Marine Environment

On national and state levels, we have several policies and regulation like Water (Prevention and Control of Pollution) Act, 1974, to regulate pollution discharges and restore water quality of our aquatic resources including the prescription of monitoring activities. One of the important provisions of the Water Act, 1974, is to maintain and restore the 'wholesomeness' of our aquatic resources. Water quality monitoring is one of the first steps required in the rational development and management of water resources. In the field of water quality management, there has been a steady evolution in procedures for designing system to obtain information on the changes of water quality. The monitoring comprises all activities to obtain 'information' with respect to the water system.



## Sampling Stations

The monitoring of marine environment for the study of biological and ecological parameters was carried out on 6<sup>th</sup>& 7<sup>th</sup> October-2021 in harbor regions of KPT and on 7<sup>th</sup> October-2021 at Vadinar during spring tide period of New moon phase of Lunar Cycle. The monitoring of marine environment for the study of biological and ecological parameters was repeated again on 13<sup>th</sup>& 14<sup>th</sup> October 2021 in harbor regions of KPT. 14<sup>th</sup> October -2021 in Vadinar during Neap tide period first quarter of Lunar Cycle..

Plankton samples from sub surface layer was collected both during high tide period and low tide period from 3 water quality monitoring stations of KPT harbour area and two stations in Nakti creek and one station in Khori creek. The same sampling schedule was repeated during consecutive spring tide and neap tide in same month. Plankton samples from sub surface layer was collected both during high tide period and low tide period from 1 water quality monitoring stations near Vadinar jetty area during spring tide and neap tide in this month .Collected water samples were processed for estimation of Chlorophyll- a, Pheophytin- a, qualitative & quantitative evaluation of phytoplankton, qualitative & quantitative evaluation zooplanktons (density and their population).

### Sampling Locations

Offshore monitoring requirement	Number of locations
Offshore Installations	3 in Kandla creek 2 in Nakti creek 1 in Khori creek 1 near Vadinar Jetty 1 near 1 <sup>st</sup> SBM
<b>Total Number of locations</b>	<b>8</b>

#### 6.1 Marine Water Quality

Marine water quality of marine waters of Deendayal Port Harbor waters, Khori and Nakti Creeks and two locations of Vadinar are monitored for various physico-chemical parameters during spring and neap tide of each month.

The results of marine water quality and Marine sediments are as below;

**Table 30: Marine Water Quality Monitoring Parameters for location near KPT colony**

Sr. No.	Parameters	Unit	Kandla Creek Near KPT colony (1)			
			23°0'58"N 70°13'22."E			
			Spring Tide		Neap Tide	
Tide →		High Tide	Low Tide	High Tide	Low Tide	
1	pH	pH unit	7.14	7.16	7.42	7.36
2	Color	-	Colorless	Colorless	Colorless	Colorless
3	Odor	-	Odorless	Odorless	Odorless	Odorless
4	Salinity	ppt	32.5	32.0	32.6	32.0
5	Turbidity	NTU	38	29	36	31
6	Total Dissolved Solids	mg/l	42450	39030	42122.0	41187.0
7	Total Suspended Solids	mg/l	685	950	764.9	558.1
8	Total Solids	mg/l	43135	39980	42886.9	41745.1
9	DO	mg/l	4.1	4	4.3	4.2
10	COD	mg/l	78.0	80.0	80.0	78.0
11	BOD	mg/l	<2.0	<2.0	<2.0	<2.0
12	Silica	mg/l	0.75	0.64	0.65	0.53
13	Phosphate	mg/l	0.36	0.26	0.16	0.18
14	Sulphate	mg/l	3060	2892	2256	2532
15	Nitrate	mg/l	2.89	2.46	2.50	3.48
16	Nitrite	mg/l	<0.05	<0.05	<0.05	<0.05
17	Calcium	mg/l	601.20	521.04	561.12	521.04
18	Magnesium	mg/l	1628.1	1603.8	0	0
19	Sodium	mg/l	9473.0	8438.0	9368	8523
20	Potassium	mg/l	362.1	314.0	360.8	302.8
21	Iron	mg/l	1.63	1.34	1.35	1.24
22	Chromium	mg/l	0.11	0.13	0.12	0.13
23	Copper	mg/l	0.06	0.05	0.06	0.06
24	Arsenic	mg/l	<0.01	<0.01	<0.01	<0.01
25	Cadmium	mg/l	0.04	0.05	0.04	0.04
26	Mercury	mg/l	<0.001	<0.001	<0.001	<0.001
27	Lead	mg/l	0.09	0.11	0.11	0.09
28	Zinc	mg/l	0.07	0.06	0.06	0.05

**Table 31: Marine Water Quality Monitoring Parameters for location near passenger Jetty One at Kandla**

Sr. No.	Parameters	Unit	Near passenger Jetty One (2)			
			23° 0'18 "N 70°13'31"E			
			Spring Tide		Neap Tide	
			Tide →	High Tide	Low Tide	High Tide
1	pH	pH unit	7.25	7.21	7.51	7.1
2	Color	-	Colorless	Colorless	Colorless	Colorless
3	Odor	-	Odorless	Odorless	Odorless	Odorless
4	Salinity	ppt	31.8	31.9	31.8	31.9
5	Turbidity	NTU	36	25	39	45
6	Total Dissolved Solids	mg/l	33930	47550	40323.0	40031.0
7	Total Suspended Solids	mg/l	658	769	569.8	528.6
8	Total Solids	mg/l	34588	48319	40892.8	40559.6
9	DO	mg/l	4.4	3.8	4.4	4.1
10	COD	mg/l	82.0	86.0	88.0	80.0
11	BOD	mg/l	<2.0	<2.0	<2.0	<2.0
12	Silica	mg/l	0.51	0.60	0.80	0.75
13	Phosphate	mg/l	0.23	0.26	0.18	0.19
14	Sulphate	mg/l	2784	3252	2388	2652
15	Nitrate	mg/l	3.03	3.59	2.89	4.04
16	Nitrite	mg/l	<0.05	<0.05	<0.05	<0.05
17	Calcium	mg/l	561.12	641.28	480.96	561.12
18	Magnesium	mg/l	1555.2	1628.1	0	0
19	Sodium	mg/l	9670.0	9156.0	9686	9192
20	Potassium	mg/l	380.0	326.1	354.2	278.2
21	Iron	mg/l	1.90	1.73	1.68	1.33
22	Chromium	mg/l	0.13	0.11	0.11	0.15
23	Copper	mg/l	0.07	0.05	0.07	0.07
24	Arsenic	mg/l	<0.01	<0.01	<0.01	<0.01
25	Cadmium	mg/l	0.05	0.04	0.04	0.05
26	Mercury	mg/l	<0.001	<0.001	<0.001	<0.001
27	Lead	mg/l	0.12	0.12	0.09	0.11
28	Zinc	mg/l	0.05	0.06	0.08	0.09

**Table 32: Marine Water Quality Monitoring Parameters for location Near Coal Berth**

Sr. No.	Parameters	Unit	Near Coal Berth			
			22°59'12"N 70°13'40"E			
			Spring Tide		Neap Tide	
			Tide →	High Tide	Low Tide	High Tide
1	pH	pH unit	7.40	7.52	7.2	7.41
2	Color	-	Colorless	Colorless	Colorless	Colorless
3	Odor	-	Odorless	Odorless	Odorless	Odorless
4	Salinity	ppt	32.2	32.0	32.0	31.7
5	Turbidity	NTU	36	28	33	41
6	Total Dissolved Solids	mg/l	45010	41120	40162.0	42404.0
7	Total Suspended Solids	mg/l	586	838	492.9	627.8
8	Total Solids	mg/l	45596	41958	40654.9	43031.8
9	DO	mg/l	4.1	5	4.5	5.2
10	COD	mg/l	88.0	90.0	79.0	74.0
11	BOD	mg/l	<2.0	<2.0	<2.0	<2.0
12	Silica	mg/l	0.69	0.76	0.53	0.71
13	Phosphate	mg/l	0.27	0.37	0.16	0.19
14	Sulphate	mg/l	3300	1872	2688	2256
15	Nitrate	mg/l	3.87	4.36	2.96	2.59
16	Nitrite	mg/l	<0.05	<0.05	<0.05	<0.05
17	Calcium	mg/l	601.20	681.36	521.04	480.96
18	Magnesium	mg/l	1676.7	1652.4	0	0
19	Sodium	mg/l	9421.0	8958.0	9328	8688
20	Potassium	mg/l	354.2	343.7	283.8	332.6
21	Iron	mg/l	1.56	1.88	1.93	1.57
22	Chromium	mg/l	0.14	0.14	0.14	0.11
23	Copper	mg/l	0.06	0.07	0.06	0.06
24	Arsenic	mg/l	<0.01	<0.01	<0.01	<0.01
25	Cadmium	mg/l	0.05	0.04	0.06	0.05
26	Mercury	mg/l	<0.001	<0.001	<0.001	<0.001
27	Lead	mg/l	0.13	0.10	0.08	0.10
28	Zinc	mg/l	0.07	0.08	0.06	0.07

**Table 33: Marine Water Quality Monitoring Parameters for location Khori creek at Kandla**

Sr. No.	Parameters	Unit	KPT 4			
			Near 15/16 Berth			
			Spring Tide		Neap Tide	
			Tide →	High Tide	Low Tide	High Tide
1	pH	pH unit	7.28	7.40	7.3	7.4
2	Color	-	Colorless	Colorless	Colorless	Colorless
3	Odor	-	Odorless	Odorless	Odorless	Odorless
4	Salinity	ppt	31.5	32.6	32.4	32.0
5	Turbidity	NTU	46	36	50	30
6	Total Dissolved Solids	mg/l	42910	48900	40963.0	41643.0
7	Total Suspended Solids	mg/l	660	562	711.1	509.7
8	Total Solids	mg/l	43570	49462	41674.1	42152.7
9	DO	mg/l	4.7	4.6	5.8	5.6
10	COD	mg/l	72.0	76.0	82.0	78.0
11	BOD	mg/l	<2.0	<2.0	<2.0	<2.0
12	Silica	mg/l	0.73	0.62	0.65	0.49
13	Phosphate	mg/l	0.28	0.24	0.22	0.19
14	Sulphate	mg/l	1500	3336	2412	2124
15	Nitrate	mg/l	1.76	2.89	2.78	2.02
16	Nitrite	mg/l	<0.05	<0.05	<0.05	<0.05
17	Calcium	mg/l	521.04	601.20	440.88	601.2
18	Magnesium	mg/l	1676.7	1555.2	0	0
19	Sodium	mg/l	9979.0	9708.0	9808	9629
20	Potassium	mg/l	373.2	343.9	327.8	305.6
21	Iron	mg/l	1.73	1.67	1.02	1.68
22	Chromium	mg/l	0.15	0.17	0.09	0.09
23	Copper	mg/l	0.05	0.08	0.07	0.07
24	Arsenic	mg/l	<0.01	<0.01	<0.01	<0.01
25	Cadmium	mg/l	0.03	0.06	0.07	0.04
26	Mercury	mg/l	<0.001	<0.001	<0.001	<0.001
27	Lead	mg/l	0.09	0.01	0.09	0.08
28	Zinc	mg/l	0.06	0.08	0.07	0.05

**Table 34: Marine Water Quality Monitoring Parameters for location Nakti Creek near Tuna Port**

Sr. No.	Parameters	Unit	Nakti Creek Near Tuna Port			
			22°57'49."N 70° 7'0.67"E			
			Spring Tide		Neap Tide	
			Tide →	High Tide	Low Tide	High Tide
1	pH	pH unit	7.20	7.40	7.2	7.4
2	Color	-	Colorless	Colorless	Colorless	Colorless
3	Odor	-	Odorless	Odorless	Odorless	Odorless
4	Salinity	ppt	32.1	31.9	31.6	31.2
5	Turbidity	NTU	34	46	33	29
6	Total Dissolved Solids	mg/l	48700	49900	40306.0	38117.0
7	Total Suspended Solids	mg/l	867	848	441.1	513
8	Total Solids	mg/l	49567	50748	40747.1	38630.0
9	DO	mg/l	4.2	4.8	4.8	4.7
10	COD	mg/l	96.0	98.0	90.0	92.0
11	BOD	mg/l	<2.0	<2.0	<2.0	<2.0
12	Silica	mg/l	0.53	0.84	0.73	0.60
13	Phosphate	mg/l	0.25	0.24	0.20	0.17
14	Sulphate	mg/l	3504	3780	2772	2364
15	Nitrate	mg/l	3.24	3.59	2.74	4.60
16	Nitrite	mg/l	<0.05	<0.05	<0.05	<0.05
17	Calcium	mg/l	521.04	601.20	480.96	521.04
18	Magnesium	mg/l	1652.4	1676.7	0	0
19	Sodium	mg/l	10156.0	10254.0	10268	10438
20	Potassium	mg/l	336.9	336.4	278.6	297.8
21	Iron	mg/l	1.54	1.83	1.55	1.50
22	Chromium	mg/l	0.12	0.11	0.14	0.16
23	Copper	mg/l	0.08	0.07	0.05	0.05
24	Arsenic	mg/l	<0.01	<0.01	<0.01	<0.01
25	Cadmium	mg/l	0.04	0.06	0.06	0.04
26	Mercury	mg/l	<0.001	<0.001	<0.001	<0.001
27	Lead	mg/l	0.12	0.13	0.11	0.13
28	Zinc	mg/l	0.08	0.09	0.05	0.09

**Table 35: Marine Water Quality Monitoring Parameters for location Nakti Creek Near NH-8A at Kandla**

Sr. No.	Parameters	Unit	Nakti Creek Near NH-8A			
			23° 02'01"N 70° 09'31"E			
			Spring Tide		Neap Tide	
			Tide →		High Tide	Low Tide
1	pH	pH unit	7.50	Sampling not possible during Low Tide	7.5	Sampling not possible during Low Tide
2	Color	-	Colorless		Colorless	
3	Odor	-	Odorless		Odorless	
4	Salinity	ppt	31.8		31.0	
5	Turbidity	NTU	34		34	
6	Total Dissolved Solids	mg/l	43730		40522.0	
7	Total Suspended Solids	mg/l	635		432.9	
8	Total Solids	mg/l	44365		40954.9	
9	DO	mg/l	4.6		5.2	
10	COD	mg/l	98.0		90.0	
11	BOD	mg/l	<2.0		<2.0	
12	Silica	mg/l	1.00		0.51	
13	Phosphate	mg/l	0.24		0.17	
14	Sulphate	mg/l	3576		2352	
15	Nitrate	mg/l	3.03		3.37	
16	Nitrite	mg/l	<0.05		<0.05	
17	Calcium	mg/l	561.12		601.2	
18	Magnesium	mg/l	1725.3		0	
19	Sodium	mg/l	10760.0		10536	
20	Potassium	mg/l	335.1		335.1	
21	Iron	mg/l	1.80		1.33	
22	Chromium	mg/l	0.16		0.10	
23	Copper	mg/l	0.07		0.06	
24	Arsenic	mg/l	<0.01		<0.01	
25	Cadmium	mg/l	0.03		0.05	
26	Mercury	mg/l	<0.001		<0.001	
27	Lead	mg/l	0.12		0.11	
28	Zinc	mg/l	0.09		0.07	

**Table 36: Marine Water Quality Monitoring Parameters for locations Nr. Vadinar Jetty**

Sr. No.	Parameters	Unit	Nr.Vadinar Jetty			
			22°26'25.26"N 69°40'20.41"E			
			Spring Tide		Neap Tide	
			Tide →	High Tide	Low Tide	High Tide
1	pH	pH unit	7.50	7.80	7.54	7.45
2	Color	-	Colorless	Colorless	Colorless	Colorless
3	Odor	-	Odorless	Odorless	Odorless	Odorless
4	Salinity	ppt	32.0	32.0	32.1	31.9
5	Turbidity	NTU	46	36	46	42
6	Total Dissolved Solids	mg/l	47700	46610	37421.0	38258.0
7	Total Suspended Solids	mg/l	483	476	553.6	490
8	Total Solids	mg/l	48183	47086	37974.6	38748.0
9	DO	mg/l	4.3	4.5	4.6	4.8
10	COD	mg/l	86.0	88.0	88.0	86.0
11	BOD	mg/l	<2.0	<2.0	<2.0	<2.0
12	Silica	mg/l	0.85	0.93	0.71	0.89
13	Phosphate	mg/l	0.23	0.25	0.18	0.17
14	Sulphate	mg/l	2784	2556	2532	2448
15	Nitrate	mg/l	2.46	2.89	3.60	3.37
16	Nitrite	mg/l	<0.05	<0.05	<0.05	<0.05
17	Calcium	mg/l	641.28	561.12	561.12	521.04
18	Magnesium	mg/l	1506.6	1579.5	0	0
19	Sodium	mg/l	10233.0	10490.0	10860.0	10880.0
20	Potassium	mg/l	331.3	363.9	336.0	333.0
21	Iron	mg/l	1.38	1.69	1.34	1.2
22	Chromium	mg/l	0.15	0.19	0.11	0.12
23	Copper	mg/l	0.06	0.08	0.06	0.06
24	Arsenic	mg/l	<0.01	<0.01	<0.01	<0.01
25	Cadmium	mg/l	0.04	0.04	0.06	0.05
26	Mercury	mg/l	<0.001	<0.001	<0.001	<0.001
27	Lead	mg/l	0.11	0.11	0.09	0.09
28	Zinc	mg/l	0.06	0.08	0.06	0.05



**Table 36 (a): Marine Water Quality Monitoring Parameters for locations Nr. Vadinar SPM**

Sr. No.	Parameters	Unit	Nr.Vadinar SPM			
			22°30'56.15"N 69°42'12.07"E			
			Spring Tide		Neap Tide	
			Tide →	High Tide	Low Tide	High Tide
1	pH	pH unit	7.70	7.50	7.42	7.36
2	Color	-	Colorless	Colorless	Colorless	Colorless
3	Odor	-	Odorless	Odorless	Odorless	Odorless
4	Salinity	ppt	31.5	32.6	32.0	31.8
5	Turbidity	NTU	34	32	38	44
6	Total Dissolved Solids	mg/l	40230	39870	40119.0	40011.0
7	Total Suspended Solids	mg/l	470	447	496.1	457.4
8	Total Solids	mg/l	40700	40317	40615.1	40468.4
9	DO	mg/l	4.2	4.1	4.4	4.5
10	COD	mg/l	88.0	90	88.0	86
11	BOD	mg/l	<2.0	<2.0	<2.0	<2.0
12	Silica	mg/l	0.82	0.93	0.62	0.76
13	Phosphate	mg/l	0.24	0.25	0.17	0.17
14	Sulphate	mg/l	2688	2340	2352	2472
15	Nitrate	mg/l	2.68	2.82	3.27	3.60
16	Nitrite	mg/l	<0.05	<0.05	<0.05	<0.05
17	Calcium	mg/l	601.20	641.28	0	0
18	Magnesium	mg/l	1579.5	1652.4	0	0
19	Sodium	mg/l	10575	10639	10936.0	10886.0
20	Potassium	mg/l	370.1	367.9	331	402
21	Iron	mg/l	1.57	1.81	1.07	1.63
22	Chromium	mg/l	0.17	0.18	0.09	0.11
23	Copper	mg/l	0.06	0.08	0.06	0.04
24	Arsenic	mg/l	<0.01	<0.01	<0.01	<0.01
25	Cadmium	mg/l	0.04	0.05	0.04	0.06
26	Mercury	mg/l	<0.001	<0.001	<0.001	<0.001
27	Lead	mg/l	0.13	0.12	0.08	0.08
28	Zinc	mg/l	0.06	0.07	0.07	0.08

### 6.1.1 Marine Sediments

Sediment samples were collected with Van Veen Grab from the six locations in Kandla Port Waters and two locations in Vadinar Port. Samples were collected and preserved in silver foil in ice box to prevent the contamination/decaying of the samples.

## 6.2 Results

The Sediment Quality results are given in below from table no. 34 A & B.

**Table 34A: Results of Analysis of Sediment of Kandla & Vadinar Port (Spring Tide)**

Sr. No.	Parameters	Unit	KPT - 1	KPT - 2	KPT - 3	KPT - 4	KPT - 5	Jetty	SPM
1	Texture	-	Sandy Loam	Sandy Loam	Sandy Loam	Sandy Loam	Sandy Loam	Sandy Loam	Sandy Loam
2	Organic Matter	mg/kg	1.10	1.08	1.01	1.02	0.98	2.03	0.81
3	Organic Carbon	mg/kg	0.65	0.52	0.60	0.56	0.56	1.08	0.47
4	Inorganic Phosphate	mg/kg	110.0	131.0	132.0	145.0	145.0	132.0	149.0
5	Moisture	%	18.96	19.65	21.0	22.10	23.5	17.7	27.00
6	Aluminium	mg/kg	ND	ND	ND	ND	ND	ND	ND
7	Silica	mg/kg	16.5	15.6	12.0	16.3	15.5	18.7	18.00
8	Phosphate	mg/kg	8.02	10.99	7.22	11.00	7.65	9.52	11.20
9	Sulphate	mg/kg	205.0	265.0	266.0	198.0	221.0	197.6	265.0
10	Nitrite	mg/kg	0.1	0.15	0.11	0.13	0.13	0.09	0.11
11	Nitrate	mg/kg	9.35	7.32	7.06	8.65	8.99	7.65	8.88
12	Calcium	mg/kg	325.0	306.0	396.0	388.0	324.0	324.0	378.0
13	Magnesium	mg/kg	195.0	185.0	243.0	244.0	188.0	175.0	210.0
14	Sodium	mg/kg	3745.0	3945.0	4660.0	2566.0	2899.0	2253.0	2854.0
15	Potassium	mg/kg	238.0	194.0	186.0	178.0	138.0	152.0	110.0
16	Chromium	mg/kg	8.1	48.3	30.7	40.3	23.3	36.4	6.6
17	Nickel	mg/kg	16.4	31.8	22.9	25.8	14	46.6	3.7
18	Copper	mg/kg	27.7	36.9	8.7	14.3	4.2	19.2	1.9
19	Zinc	mg/kg	32.40	40.50	35.10	36.20	21.20	21.30	5.00
20	Cadmium	mg/kg	ND	ND	ND	ND	ND	ND	ND
21	Lead	mg/kg	3.8	5.8	3.3	4.4	4.9	ND	1.2
22	Mercury	mg/kg	ND	ND	ND	ND	ND	ND	ND
23	Arsenic	mg/kg	ND	ND	ND	ND	ND	ND	ND

\*ND - Not Detected

**Table 34B: Results of Analysis of Sediment of Kandla & Vadinar Port (Neap Tide)**

Sr. No.	Parameters	Unit	KPT – 1	KPT - 2	KPT - 4	SPM
1	Texture	-	Sandy Loam	Sandy Loam	Sandy Loam	Sandy Loam
2	Organic Matter	mg/kg	1.99	0.85	1.11	1.08
3	Organic Carbon	mg/kg	1.12	0.41	0.65	0.65
4	Inorganic Phosphate	mg/kg	120.0	130.0	148.0	152.0
5	Moisture	%	19.20	18.50	17.7	17.56
6	Aluminium	mg/kg	ND	ND	ND	ND
7	Silica	mg/kg	21.88	18.00	17.5	14.72
8	Phosphate	mg/kg	5.62	8	7.65	8.65
9	Sulphate	mg/kg	225.0	240.0	211.5	195.0
10	Nitrite	mg/kg	0.1	0.11	0.09	0.1
11	Nitrate	mg/kg	7.66	8.11	6.65	6.85
12	Calcium	mg/kg	321.0	310.0	345.0	265.0
13	Magnesium	mg/kg	205.0	197.0	202.0	169.0
14	Sodium	mg/kg	4120.0	3842.0	4465.0	3589.0
15	Potassium	mg/kg	201.0	147.0	154.0	154.00
16	Chromium	mg/kg	13.3	10.5	13	16.8
17	Nickel	mg/kg	8.8	6	8.2	10.5
18	Copper	mg/kg	4.3	2	2.2	5.5
19	Zinc	mg/kg	18.20	9.90	10.30	12.10
20	Cadmium	mg/kg	ND	ND	ND	ND
21	Lead	mg/kg	2.1	2.5	2.8	1.8
22	Mercury	mg/kg	ND	ND	ND	ND
23	Arsenic	mg/kg	ND	ND	ND	ND

\*Grab samples could not be collected due high current at KPT – 3,KPT – 5 Location.

\*ND - Not Detected

**REPORT**  
**ON**  
**ECOLOGICAL MONITORING**  
**OF MARINE ENVIRONMENT**  
**IN**  
**DPT HARBOUR AREA, NEAR BY CREEKS**  
**AND**  
**VADINAR JETTY AND SPM**  
**FOR**  
**DEENDAYAL PORT TRUST**

OCTOBER, 2021

**INTRODUCTION:**

**Sampling Stations:**

The monitoring of marine environment for the study of biological and ecological Parameters was carried out on 6<sup>th</sup> October, 2021 in harbour region of DPT at Kandla Creek, and on 7<sup>th</sup> October, 2021 in creeks near by the port during spring tide. The monitoring of marine environment for the study of biological and ecological parameters was repeated again on 13<sup>th</sup> October, 2021 in harbour region of DPT at Kandla Creek and on 14<sup>th</sup> October 2021 in creeks near by the port during neap tidal condition.

Plankton samples from sub surface layer was collected both during high tide period and low tide period from 3 water quality monitoring stations of KPT harbour area and one stations in Nakti creek and one station in Khori creek. Sampling at second sampling station of Nakti creek was possible only during high tide period. The same sampling schedule was repeated during consecutive Neap tide and spring tide in same month.

Plankton samples from sub surface layer were collected during high tide period and low tide period from monitoring station near Vadinar jetty at Path Finder Creek during spring tide period and during neap tide. Sampling was conducted at only during Neap tide period near SPM both during High tide period and low tide period. Collected water samples were processed for estimation of Chlorophyll-a, Pheophytin-a, qualitative & quantitative evaluation of phytoplankton, qualitative & quantitative evaluation zooplanktons (density and their population).

**TABLE #1 SAMPLING LOCATIONS**

<b>monitoring requirement</b>	<b>Number of locations</b>
Kandla creek	3 in Kandla creek
Nakti creek	2 in Nakti creek
Khori Creek	1 in Khori creek
Vadinar jetty	1 near Vadinar Jetty
SPM	1 near I <sup>st</sup> SPM
<b>Total Number of locations</b>	<b>8</b>

### Sampling methodology adopted:

A marine sampling is an estimation of the body of information in the population. The theory of the sampling design is depending upon the underlying frequency distribution of the population of interest. The requirement for useful water sampling is to collect a representative sample of suitable volume from the specified depth and retain it free from contamination during retrieval.

50 litres of the water sample were collected from Sub surface by using bucket. From the collected water sample 1 litres of water sample were taken in an opaque plastic bottle for chlorophyll estimation, thereafter plankton samples were collected by using filtration assembly with nilyobolt cloth of 20 $\mu$ m mesh size.

### Samples Processing for chlorophyll estimation:

Samples for the chlorophyll estimation were preserved in ice box on board in darkness to avoid degradation in opaque container covered with aluminium foil. Immediately after reaching the shore after sampling, 1 litre of collected water sample was filtered through GF/F filters (pore size 0.45  $\mu$ m) by using vacuum filtration assembly. After vacuum filtration the glass micro fiber filter paper was grunted in tissue grinder, macerating of glass fiber filter paper along with the filtrate was done in 90% aqueous Acetone in the glass tissue grinder with glass grinding tube. Glass fiber filter paper will assist breaking the cell during grinding and chlorophyll content was extracted with 10 ml of 90% Acetone, under cold dark conditions along with saturated magnesium carbonate solution in glass screw cap tubes. After an extraction period of 24 hours, the samples were transferred to calibrated centrifuge tubes and adjusted the volume to original volume with 90% aqueous acetone solution to make up the evaporation loss. The extract was clarified by using centrifuge in closed tubes. The clarified extracts were then decanted in clean cuvette and optical density was observed at wavelength 664, 665 nm. By using corrected optical density, Chlorophyll-a value was calculated as given in (APHA, 1998).

### PLANKTON:

The entire area open water in the sea is the pelagic realm. Pelagic organisms live in the open sea. In contrast to the pelagic realm, the benthic realm comprises organisms and zone of the bottom of the sea. Vertically the pelagic realm can be dividing into two zones based on light penetration; upper photic or euphotic zone and lower dark water mass, aphotic zone below the photic zone.

The term plankton is general term for organisms have such limited powers of locomotion that they are at the mercy of the prevailing water movement. Plankton is subdivided to phytoplankton and zooplankton. Phytoplankton is free floating organisms that are capable of photosynthesis and zooplankton is the various free floating animals.

Pelagic zone, represents the entire ocean water column from the surface to the deepest depths, is home to a diverse community of organisms. Differences in their locomotive ability categorize the organisms in the pelagic realm into two, **plankton** and **nekton** (Lalli and Parsons, 1997). **Plankton** consists of all organisms drifting in the water and is unable to swim against water currents, whereas **Nekton** includes organisms having strong locomotive power. Ecological studies on the plankton community, which form the base of the aquatic food chain, help in the better understanding of the dynamics and functioning of the marine ecosystem. The term 'Plankton' first coined by Victor Hensen (1887), Plankton, (Greek word: *planktos* meaning "passively drifting or wandering") is defined as drifting or free-floating organisms that inhabit the pelagic zone of water. Based on their mode of nutrition planktonic organisms are categorised into phytoplankton (organisms having an autotrophic mode of nutrition) and zooplankton (organisms having a heterotrophic mode of nutrition).

#### Phytoplankton in the marine environment:

Phytoplankton is free floating unicellular, filamentous and colonial eutrophic organisms that grow in aquatic environments whose movement is more or less dependent upon water currents. These micro flora acts as primary producers as well as the basis of food chain, source of protein, bio purifier and bio indicators of the aquatic ecosystems of which diverse array of the life depends .They are considered as an important component of aquatic flora, play a key role in maintaining equilibrium between abiotic and biotic components of aquatic ecosystem.

The phytoplankton includes a wide range of photosynthetic and phototrophic organisms. Marine phytoplankton is mostly microscopic and unicellular floating flora, which are the primary producers that support the pelagic food-chain. The two most prominent groups of phytoplankton are diatoms (Bacillariophyceae) and dinoflagellates (Dinophyceae).The phytoplankton those normally captured in the net from the Gulf of Kutch is normally dominated by these two major groups; diatoms and dinoflagellates. Phytoplankton also include numerous and diverse collection of extremely small, motile algae which are termed micro flagellates (naked flagellates) as well as and Cyanophytes (blue-green algae).

Algae are an ecologically important group in most aquatic ecosystems and have been an important component of biological monitoring programs. Algae are ideally suited for water quality assessment because they have rapid reproduction rates and very short life cycles, making them valuable indicators of short-term impacts.



Aquatic populations are impacted by anthropogenic stress, resulting in a variety of alterations in the biological integrity of aquatic systems. Algae can serve as an indicator of the degree of deterioration of water quality, and many algal indicators have been used to assess environmental status.

#### **Zooplankton in the marine environment:**

Zooplankton includes a taxonomically and morphologically diverse community of heterotrophic organisms that drift in the waters of the world's oceans. Qualitative and quantitative studies on zooplankton community are a prerequisite to delineate the ecological processes active in the marine ecosystem. Zooplankton community plays a pivotal role in the pelagic food web as the primary consumers of phytoplankton and act as the food source for organisms in the higher trophic levels, particularly the economically essential groups such as fish larvae and fishes. They also function in the cycling of elements in the marine ecosystem. The dynamics of the zooplankton community, their reproduction, and growth and survival rate are all significant factors determining the recruitment and abundance of fish stocks as they form an essential food for larval, juvenile and adult fishes (Beaugrand et al., 2004). Zooplankton grazing in the marine environment controls the primary production and helps in determining the pelagic ecosystem (Banse, 1995). Through grazing in surface waters and following the production of sinking faecal matters and also by the active transportation of dissolved and particulate matter to deeper waters via vertical migration, they help in the transport of organic carbon to deep ocean layers and thus act as key drivers of biological pump' in the marine ecosystem. Zooplankton grazing and metabolism also, transform particulate organic matter into dissolved forms, promoting primary producer community, microbial demineralization, and particle export to the ocean's interior.

The categorisation of zooplankton into various ecological groups is based on several factors such as duration of planktonic life, size, food preferences and habitat. As they vary significantly in size from microscopic to metazoic forms, the classification of zooplankton based on size has paramount importance in the field of quantitative plankton research.

Based on the duration of planktonic life, zooplankton are categorised into Holoplankton (organisms which complete their entire lifecycle as plankton) and Meroplankton (organisms which are planktonic during the early part of their lives such as the larval stages of benthic and nektonic organisms). Tychoplankton are organisms which live a brief planktonic life, such as the benthic crustaceans (Cumaceans, mysids, isopods) which ascend to the water column at night for feeding and certain ectoparasitic copepods, they leave the host and spend their life as plankton during their breeding cycle.


Zooplankton can be subdivided into holoplankton, i.e., permanent members of the plankton (e.g., Calanoid copepods), and meroplankton, i.e., temporary members in the plankton e.g., larvae of fish,

shrimp, and crab). The meroplankton group consists of larval and young stages of animals that will adopt a different lifestyle once they mature. In contrast to phytoplankton which consist of a relatively smaller variety of organisms, Zooplankton are extremely divers, consist of a host of larval and adult forms representing many animal phylum.

Among the zooplankton one group always dominate than others; members of sub class copepods (Phylum Athropoda) and Tintinids (Phylum Protozoa) among the net planktons. These small animals are of vital importance in marine ecosystem as one of the primary herbivores animals in the sea, and it is they provide vital link between primary producer (autotrophs) and numerous small and large marine consumers.

As their community structure and function are highly susceptible to changes in the environmental conditions regular monitoring of their distribution as well as their interactions with various physicochemical parameters is inevitable for the sustainable management of the ecosystem (Kusum et al., 2014). Of all the marine zooplankton groups, copepods mainly Calanoid copepods are the dominant groups in marine subtropical and tropical waters and exhibit considerable diversity in morphology and habitats they occupy (Madhupratap, 1991;)

It has been well established that potential of pelagic fishes viz. finfishes, crustaceans, molluscs and marine mammals either directly or indirectly depend on zooplankton. The herbivorous zooplankton is efficient grazers of the phytoplankton and is referred to as living machines transforming plant material into animal tissue. Hence they play an essential role as the intermediaries for nutrients/energy transfer between primary and tertiary trophic levels. Due to their large density, shorter lifespan, drifting nature, high group/species diversity and different tolerance to the stress, they used as the indicator organisms for the physical, chemical and biological processes in the aquatic ecosystem (Ghajibhiye, 2002).



**Spatial distribution of Plankton:**

A characteristic of plankton population is that they tend to occur in patches, which are varying spatially on a scale of few meters to far as few kilometres in distance. They also vary in time scale, season as well as vertically in the water column. It is this patchiness and its constant changes in time and spot, that has made it so difficult for plankton biologist to learn about the ecology of plankton. The biological factors that causes this patchiness is due to the ability of zooplankton to migrate vertically and graze out the phytoplankton at a rapid rate that can create patchiness. Similarly the active swimming ability by certain zooplankton organisms can cause to aggregate in dense group.

At its most extreme, because the water in which plankton is suspended is constantly moving, each sample taken by the plankton biologists remain a different volume of water, so each sample is unique and replicate does not exist.

Plankton may also exhibit vertical patchiness. Physical factors contribute to this type of patchiness include light intensity, nutrients and density gradients in the water column.

Phytoplankton in particular tends to be unequally distributed vertically, which leads to the existence of different concentration of a chlorophyll value between photic zone and below the photic zone.

**Methodology adopted for Plankton sampling:**

Mixed plankton sample were obtained from the sub surface layer at each sampling locations by towing the net horizontally with the weight .After the tow of about 15-30minutes, plankton net was pulled up and washed down to the tail and collected the plankton adhered to plankton net in the collection bucket at the bottom by springing outer and inner surface of the net with sea water, while the net was hanging with the mouth upward. For quantitative evaluation 50 L water samples were collected from subsurface layer and filtered through 20µm mesh size net by using bucket and filtration assembly.

**Preservation and storage:**

Both filtered plankton and those collected from the plankton net were preserved with 5% buffered formalin and stored in 1L plastic container for further processing in the laboratory.

**Sample concentration:**

The collected plankton samples were concentrated by using centrifuge and made up to 50 ml with 5% formalin -Glycerine mixture.

**Taxonomic evaluation:**

Before processing, the sample was mixed carefully and a subsample was taken with a calibrated Stempel-pipette. 1 ml of the concentrated plankton samples were transferred on a glass slide with automatic pipette. The plankton sample on the glass slides were stained by using Lugol's iodine and added glycerine to avoid drying while observation. The plankton samples were identified by using Labex triangular Research microscope with photographic attachment. Microphotographs of the plankton samples were taken for record as well as for confirming the identification. The bigger sized zooplankton was observed through dissecting stereomicroscope with magnification of 20-30 x. Plankton organisms in the whole slide were identified to the lowest axon possible. A thorough literature search was conducted for the identification of the different groups of zooplankton that were encountered

**Cell counts by drop count method:**

The common glass slide mounted with a 1ml of concentrated phytoplankton/zooplankton sample in glycerol and covered with cover slip 22x 60mm was placed under the compound microscope provided with a mechanical stage. The plankton was then counted from the microscopic field of the left top corner of the slide. Then slide is moved horizontally along the right side and plankton in each microscopic field was thus counted. When first microscopic field row was finished the next consecutive row was adjusted using the mechanical device of the stage. In this way all the plankton present in entire microscopic field are counted. From this total number in 1ml of the concentrated

plankton, total number of plankton in the original volume of sample filtered was calculated as units/L.

#### **BENTHIC ORGANISMS:**

Benthos is those organisms that are associated with the sea bed or benthic habitats. Epi- benthic organisms live attached to a hard substratum or rooted to a shallow depth below the surface. In fauna organisms live below the sediment–water interface. Interstitial organisms live and move in pore water among sedimentary grains.

Because the benthic organisms are often collected and separated on sieves, a classification based on the overall size is used. Macro benthos include organisms whose shortest dimension is greater than or equal to 0.5 mm. Meio benthos are smaller than 0.5mm but larger than 42 $\mu$  in size.

The terms such as macro fauna and Meio fauna generally have little relevance with taxonomic classification. The terms Meio fauna and macro fauna depend on the size. Meio fauna were considered as good bioassay of community health and rather sensitive indicators of environmental changes

#### **SAMPLING METHODOLOGY ADOPTED FOR SUB TIDAL REGION:**

Van veen sampler (0.09m<sup>2</sup>) was used for sampling bottom sediments. Two sets of sediments were sampled from each location, one for macro fauna and other for Meio fauna. The macro fauna in the sediments were sieved on board to separate out the organisms. The fixation of Meio fauna is normally done by bulk fixation of the sediment sample. The bulk fixation is done by using 10% formalin (Buffered with borate). The organisms were preserved with seawater as diluting agent.

#### **Sample sieving:**

Sediments samples were sieved to extract the organisms. Sieving was performed carefully as possible to avoid any damage to the animals. The large portion of the sediment was split in to smaller portions and mixed with sea water in a bucket. The cohesive lumps were broken down by continuous stirring. The disaggregated sediments were then passed through the sieves.

#### **Sample staining:**

Sorting of the Meio fauna from the sieve is difficult task especially in the preserved material, because organisms are not easily detectable. To facilitate the animal detection the entire sample retained on the sieve after sieving operation were stained by immersing the sieve in a flat bottom tub with 1% Rose Bengal stain; a protein stain. A staining period of 10-30 minutes is sufficient for sample detection.

#### **DIVERSITY INDICES:**

On the whole, diversity indices provide more information about community composition than simply species richness (number of species present); they also, take the relative abundances of different

species into account. Based on this fact, diversity indices therefore depend not only on species richness but on the evenness, or equitability, with which individuals are distributed among the different species (Magurram, A. E. (1988)

A diversity index is a measure of species diversity within a community that consists of co-occurring populations of several (two or more) different species. It includes two components: richness and evenness. Richness is the measure of the number of different species within a sample showing that more the types of species in a community, the higher is the diversity or greater is the richness. Evenness is the measure of relative abundance of the different species with in a community.

The basic idea of diversity index is to obtain a quantitative estimate of biological variability that can be used to compare biological entities composed of discrete components in space and time (Carol H.R. *etal.* 1998). Biodiversity is commonly expressed through indices based on species richness and species abundances (Whittaker 1972, Lande 1996, Purvis and Hector 2000). Biodiversity indices are a non-parametric tool used to describe the relationship between species number and abundance. The most widely used bio diversity indices are Shannon Weiner index and Simpson's index.

A diversity Index is a single statistic that incorporates in formation on richness and evenness. The diversity measures that incorporate the two concepts may be termed heterogeneity measures (Magurran, 2004).

Any study intended to interpret causes and effect of adverse impact on Biodiversity of communities require suitable measures to evaluate specie richness and Diversity. The former is number of species in community, while latter is a function of relative frequency of different species. Species richness is the iconic measure of biological diversity (Magurran, 2004). Several indices have been created to measure the diversity of species; however, the most widely used in the last decades are the Shannon (1948) and Simpson (1949) (Buzas and Hayek 1996; Gorelick 2006), with the components of diversity: richness (*S*) and evenness (*J*)

### Simpson's diversity index

Simpson's index (**D**) is a measure of diversity, which takes into account both species richness, and evenness of abundance among the species present.. The Simson index is one of the meaningful and robust biodiversity measures available.(Magurran ,2004).

The formula for calculating D is presented as:

$$D = \frac{\sum n_i(n_i - 1)}{N(N - 1)}$$

Where  $n_i$  = the total number of organisms of each individual species

$N$  = the total number of organisms of all species

The value of D ranges from 0 to 1. With this index, 0 represents infinite diversity and, 1, no diversity. When D increases diversity decreases. Simpson's index is therefore usually expressed as  $1-D$  or  $1/D$ . (Magurran, 2004)

Low species diversity suggests:

- relatively few successful species in the habitat
- the environment is quite stressful with relatively few ecological niches and only a few organisms are really well adapted to that environment
- food webs which are relatively simple
- change in the environment would probably have quite serious effects

High species diversity suggests:

- a greater number of successful species and a more stable ecosystem
- more ecological niches are available and the environment is less likely to be hostile complex food webs
- environmental change is less likely to be damaging to the ecosystem as a whole

### Species richness indices

The species richness (**S**) is simply the number of species present in an ecosystem. Species richness Indices of species richness are widely used to quantify or monitor the effects of anthropogenic disturbance. A decline in species richness may be concomitant with severe or chronic human-induced perturbation (Fair Fair weather 1990,) Species richness measures have traditionally been the mainstay in assessing the effects of environmental degradation on the biodiversity of natural assemblages of organisms (Clarke & Warwick, 2001)

Species richness is the iconic measure of biological diversity (Magurran, 2004). The species richness (**S**) is simply the number of species present in an ecosystem. This index makes no use of relative abundances. The term species richness was coined by McIntosh (1967) and oldest and most intuitive measure of biological diversity (Magurran, 2004).

Margalef's diversity index is a species richness index. Margalef's Species richness index (d), or indices that describe the evenness of the distribution of the numbers of individuals among species, were derived.

The value of a diversity index increases both when the number of types increases and when evenness increases. For a given number of types, the value of diversity index is maximised when all types are equally abundant (Rosenzweig, M. L. (1995).

**Shannon-Wiener's index:**

An index of diversity commonly used in plankton community analyses is the Shannon-Wiener's index (**H**), which emphasizes not only the number of species (richness or variety), but also the apportionment of the numbers of individuals among the species (Odum 1971 and Reish 1984). Shannon-Wiener's index (**H**) reproduces community parameters to a single number by using an equation.

Shannon and Weiner index represents entropy. It is a diversity index taking into account the number of individuals as well as the number of taxa. It varies from 0 for communities with only single taxa to high values for community with many taxa each with few individuals. This index can also determine the pollution status of a water body. Normal values range from 0 to 4. This index is a combination of species present and the evenness of the species. Examining the diversity in the range of polluted and unpolluted ecosystems, Wilham and Dorris (1968) concluded that the values of the index greater than 3 indicate clean water, values in the range of 1 to 3 are characterized by moderate pollution and values less than 1 are characterized as heavily polluted

$$H' = - \sum_{j=1}^s \frac{n_j}{N} \ln \left( \frac{n_j}{N} \right)$$

**RESULTS:**

**CHLOROPHYLL-a:**

Water Samples for the chlorophyll estimation were collected from sub surface layer during high tide and low tide period of the tidal cycle for each sampling locations and analysed for Chlorophyll -a and after acidification for Pheophytin -a. Chlorophyll- a value was used as algal biomass indicator (APHA,1998) Algal biomass was estimated by converting Chlorophyll value.

In the sub surface water chlorophyll-a was varying from 0.615 -1.459mg/m<sup>3</sup>.in harbour region of DPT in Kandla Creek during sampling done in spring tide period of October, 2021. In the nearby creeks chlorophyll-a was varying from 0.153 -1.497mg/m<sup>3</sup>.Pheophytin -a level was below detectable limit- the all the sampling stations during springtide except at KPT-I in the harbour region of DPT.

In the sub surface water chlorophyll-a was varying from 0.204 -0.749mg/m<sup>3</sup>.in harbour region of DPT in Kandla Creek during sampling done in neap tide period of October , 2021 . In the nearby creeks chlorophyll-a was varying from 0.184-0.610 mg/m<sup>3</sup>.Pheophytin -a level was below detectable limit- the all the sampling stations during neap tide in the harbour region of DPT.



**Environmental Monitoring Report of Deendayal Port Trust, OCTOBER-2021**

In the sub surface water chlorophyll-a was varying from 0.527 -0.733 mg/m<sup>3</sup>.in harbour region of DPT OOT in path finder Creek during sampling done in spring tide period of October, 2021. In the sub surface water chlorophyll-a was varying from 0.750 -1.175 mg/m<sup>3</sup>.in harbour region of DPT OOT in path finder Creek during sampling done in Neap Tide period of October, 2021

**TABLE #2 VARIATIONS IN CHLOROPHYLL –a PHEOPHYTIN- a AND ALGAL BIOMASS FROM SAMPLING STATIONS IN DPT HARBOUR AREA IN KANDLA CREEK ,NEAR BY CREEKS AND DPT OOT JETTY IN PATH FINDER CREEK AND SPM NEAR VADINAR DURING SPRING TIDE IN OCTOBER,2021**

Sr.No.	Station	Tide	Chlorophyll-a (mg/m <sup>3</sup> )	Pheophytin- a (mg/m <sup>3</sup> )	Algal Biomass (Chlorophyll method) mg/m <sup>3</sup>
DPT HARBOUR AREA KANDLA CREEK					
1	KPT1	High tide	1.459	0.821	97.75
		Low tide	1.187	0.661	79.53
2	KPT 2	High tide	0.765	BQL	51.25
		Low tide	0.852	BQL	57.08
3	KPT 3	High tide	0.968	BQL	64.85
		Low tide	0.615	BQL	41.20
CREEKS					
4	KPT-4 Khori-I	High tide	1.056	BQL	70.75
		Low tide	1.497	BQL	100.2
5	KPT-5 Nakti-I	High tide	0.764	BQL	51.19
		Low tide	0.612	BQL	41.00
6	KPT-5 Nakti-II	High tide	0.153	BQL	10.24
PATHFINDER CREEK VADINAR					
7	VADINAR-I jetty	Low tide	0.527	BQL	35.31
8		High tide	0.733	BQL	49.11
9	SPM	High tide	No sample	--	--
10	SPM	Low tide	No sample	-	--

BDL: Below Quantification Limit.

**TABLE #3 VARIATIONS IN CHLOROPHYLL –a PHEOPHYTIN- a AND ALGAL BIOMASS FROM SAMPLING STATIONS IN DPT HARBOUR AREA ,NEAR BY CREEKS AND DPT OOT JETTY IN PATH FINDER CREEK AND SPM NEAR VADINAR DURING NEAP TIDE IN OCTOBER,2021**

Sr.No.	Station	Tide	Chlorophyll-a (mg/m <sup>3</sup> )	Pheophytin- a (mg/m <sup>3</sup> )	Algal Biomass (Chlorophyll method) mg/m <sup>3</sup>
DPTHARBOUR AREA KANDLA CREEK					
1	KPT1	High tide	0.307	BQL	20.57
		Low tide	0.529	BQL	35.44
2	KPT 2	High tide	0.749	BQL	50.18
		Low tide	0.614	BQL	41.14
3	KPT 3	High tide	0.204	BQL	13.67
		Low tide	0.542	BQL	36.31
CREEKS					
4	KPT-4 Khori-I	High tide	0.441	BQL	29.54
		Low tide	0.426	BQL	28.54
5	KPT-5 Nakti-I	High tide	0.610	BQL	40.87
		Low tide	0.441	BQL	29.55
6	KPT-5 Nakti-II	High tide	0.184	BQL	12.33
PATHFINDER CREEK VADINAR					
7	VADINAR-I jetty	Low tide	0.750	0.435	50.25
8		High tide	0.820	0.484	54.94
9	SPM	High tide	1.071	0.610	71.76
10	SPM	Low tide	1.175	0.680	78.73

BDL: Below Quantification Limit.

**PHYTOPLANKTON POPULATION:**

For the evaluation of the Phytoplankton population in DPT harbour area and within the immediate surroundings of the port, sampling was conducted from 5 sampling locations (3 in harbour area and two in Nakti creek) during high tide period and low tide period of spring tide and neap tide.

The phytoplankton community of the sub surface water in the harbour and nearby creeks was represented by, Diatoms blue green algae and dinoflagellates during spring tide period. Diatoms were represented by 19 genera. Blue green were represented by 3 genera and dinoflagellates were represented by two genera during the sampling conducted in spring tide in OCTOBER, 2021. Phytoplankton of the sampling stations at sub surface layer in the harbour area and nearby creeks was varying from 46-209 units/ L during high tide period and 183-229 units/ L during low tide of Spring Tide.

The phytoplankton community of the sub surface water in the harbour and nearby creeks was represented by Diatoms. Blue green algae and Dinoflagellates during Neap tide period. Diatoms were represented by 20 genera. Blue green algae were represented 3 genera and dinoflagellates with two genera during the sampling conducted in Neap tide in October, 2021. Phytoplankton of the sampling

stations at sub surface layer in the harbour area and nearby creeks was varying from 88-170 units/ L during high tide period and 120-157 units/ L during low tide of Neap Tide.

For the evaluation of the Phytoplankton population in DPT OOT jetty area in Path Finder creek sampling was conducted from one sampling locations ; jetty area during high tide period and low tide of spring tide. For the evaluation of the Phytoplankton population in DPT OOT jetty area in Path Finder creek in Vadinar sampling was conducted from 2 sampling locations ; jetty area and one near SPM during high tide period and low tide of Neap tide.

The phytoplankton community of the sub surface water in the path finder creeks was represented by only Diatoms spring tide period. Diatoms were represented by 9 genera during the sampling conducted in spring tide in October, 2021. Phytoplankton of the sampling stations at sub surface layer in the harbour area and nearby creeks was varying from 131 units/ L during high tide period and 147 units/ L during low tide of Spring Tide.

The phytoplankton community of the sub surface water in the path finder creeks was represented by Diatoms, Blue green algae and Dinoflagellates during Neap tide period. Diatoms were represented by 10 genera Blue green algae were represented single genera and dinoflagellates by four genera during the sampling conducted in Neap tide in October, 2021. Phytoplankton of the sampling stations at sub surface path finder creek was varying from 203-427 units/ L during high tide period and 544-744 units/ L during low tide of Neap Tide.

#### **Species Richness Indices and Diversity Indices:**

##### **Margalef's diversity index (Species Richness) S**

At the organismal level, the most widely used biodiversity measures are those based on the number of species present, perhaps adjusted for the number of individuals sampled, Here Margalef's Species richness index (d), or indices that describe the evenness of the distribution of the numbers of individuals among species, are derived.

Margalef's diversity index (Species Richness) S of phytoplankton communities in the Kandla creek and nearby creeks sampling stations was varying from 2.059-3.212 with an average of 2.632 during the sampling conducted in High tide period of spring tide. While Margalef's diversity index (Species Richness) S of phytoplankton communities in the Kandla creek region and nearby creeks was varying from 2.687-3.144 with an average of 2.923 during the consecutive low tide period.

Margalef's diversity index (Species Richness) S of phytoplankton communities in the stations in Kandla creek and nearby creeks was varying from 1.582-3.384 with an average of 2.477 during the sampling conducted in High tide period of Neap tide. While Margalef's diversity index (Species Richness) S of phytoplankton communities in the Kandla creek region and nearby creeks was varying from 1.582-3.040 with an average of 2.141 during consecutive low tide.

Margalef's diversity index (Species Richness) S of phytoplankton communities in the stations was 1.641 during the sampling conducted in High tide period of spring tide at Path Finder Creek, Vadinar . While Margalef's diversity index (Species Richness) S of phytoplankton communities in the path finder creek was 1.603 during the consecutive low tide period at Path Finder Creek, Vadinar .

Margalef's diversity index (Species Richness) S of phytoplankton communities in the stations was varying from 1.981-2.064 during the sampling conducted in consecutive High tide period and low tide period of Neap Tide at jetty area of Path finder Creek. While Margalef's diversity index (Species Richness) S of phytoplankton communities near the SPM was varying from 2.259- 1.512 during the consecutive high tide and low tide period of Neap tide.

#### **Shannon-Wiener's index:**

Shannon-Wiener's Index (H) of phytoplankton communities in the sampling stations was in the range of 0.802- 0.935 ( $H'(\log_{10})$ ) between selected sampling stations with an average value of 0.860 during high tide period of spring tide at Kandla creek and nearby creeks. Shannon-Wiener's Index (H) of phytoplankton communities in the sampling stations was in the range of 0.858-0.979 ( $H'(\log_{10})$ ) between selected sampling stations with an average value of 0.909 during consecutive low tide at Kandla creek and nearby creeks. Shannon-Wiener's Index (H) of phytoplankton communities in the sampling stations was in the range of 0.774 -0.934 ( $H'(\log_{10})$ ) between selected sampling stations with an average value of 0.868 during high tide period of neap tide at Kandla creek and nearby creeks. Shannon-Wiener's Index (H) of phytoplankton communities in the sampling stations was in the range of 0.773 -0.927 ( $H'(\log_{10})$ ) between selected sampling stations with an average value of 0.842 during consecutive low tide at Kandla creek and nearby creeks.

Shannon-Wiener's Index (H) of phytoplankton communities in the sampling stations in the stations was 0.684 during the sampling conducted in High tide period of spring tide at Path Finder Creek, Vadinar. Shannon-Wiener's Index (H) of phytoplankton communities in the path finder creek was 0.695 during the consecutive low tide period at Path Finder Creek, Vadinar .

Shannon-Wiener's Index (H) of phytoplankton communities in the stations was varying from 0.356- 0.255 during the sampling conducted in consecutive High tide period and low tide period of Neap Tide at jetty area of Path finder Creek. While Shannon-Wiener's Index (H) of phytoplankton communities near the SPM was varying from 0.508-0.234 during the consecutive high tide and low tide period of Neap tide.

Typical values are generally between 1.5 and 3.5 in most ecological studies, and the index is rarely greater than 4. The Shannon-Wiener's index increases as both the richness and the evenness of the community increase. This result indicates that diversity of phytoplankton of Kandla Harbour region

and nearby creeks is less but with abundant population of few, with relatively few ecological niches and only very few opportunist organisms are really well adapted to this environment and thrive better than other species.

**Simpson's diversity index:**

Simpson's index (D) is a measure of diversity, which takes into account both species richness, and an evenness of abundance among the species present. The Simpson index is one of the meaningful and robust biodiversity measures available. (Magurran,2004).

Simpson diversity index (1-D) of phytoplankton communities was below 0.9 at all sampling stations in the Kandla Harbour region and nearby creeks, which was varying from 0.809- 0.852 between selected sampling stations with an average of 0.827 during high tide period of spring tide. Simpson diversity index (1-D) of phytoplankton communities was below 0.9 at all sampling stations in the Kandla Harbour region and nearby creeks, which was varying from 0.819- 0.865 between selected sampling stations with an average of 0.840 during consecutive low tide .

Simpson diversity index (1-D) of phytoplankton communities was below 0.9 at all sampling stations in Kandla Harbour region and nearby creeks, during high tide period and low tide period during neap tide also, which was varying from 0.809-0.853 with an average value of 0.835 between selected sampling stations during high tide period and varying from 0.809-0.852 with an average value of 0.831 between selected sampling stations during consecutive low tide period Low species diversity suggests a relatively few successful species in this habitat.

Low species diversity suggests a relatively few successful species in this habitat. The environment is quite stressful with relatively few ecological niches and only a few organisms are really well adapted to that environment. Any change in the environment would probably have quite serious effects.

Simpson diversity index (1-D) of phytoplankton communities was below 0.9 at all sampling stations in path finder Creek, which was 0.720 during high tide period and 0.751 during low tide period of spring tide at Jetty region . Simpson diversity index (1-D) of phytoplankton communities was below 0.9 at all sampling stations in the in path finder Creek, which was varying from 0.720-0.751 at jetty region of path finder creek during consecutive high tide and low tide period of Spring Tide and recorded below 9 at SPM during consecutive high tide and low tide period of Neap tide also , 0.329 - 0.499 during high tide and 0.218-0.201 during low tide

**Table # 4 PHYTOPLANKTON VARIATIONS IN ABUNDANCE AND DIVERSITY IN SUB SURFACE SAMPLING STATIONS IN DPT HARBOUR AREA AT KANDLA CREEK AND ,NEAR BY CREEKS DURING SPRING TIDE IN OCTOBER,2021**

Tide	Sampling Station	Abundance In units/L	No of Species observed /total species	% of diversity	Margalef's diversity index (Species Richness S)	Shannon Weiner index H (log <sub>10</sub> )	Diversity Index (Simpson's Index) 1-D
HIGH TIDE	1	199	18/24	75	3.212	0.9355	0.8519
	2	209	12/24	50	2.059	0.8018	0.8091
	3	204	14/24	58.3	2.444	0.8451	0.8244
	4	204	17/24	70.83	3.009	0.9095	0.837
	5	199	14/24	58.3	2.456	0.8497	0.8167
	6	46	11/24	45.83	2.612	0.8208	0.8232
LOW TIDE	1	223	18/24	75	3.144	0.9797	0.8653
	2	214	17/24	70.83	2.982	0.9126	0.8446
	3	193	17/24	70.83	3.04	0.919	0.8416
	4	229	16/24	66.66	2.761	0.8806	0.8312
	5	183	15/24	62.5	2.687	0.858	0.8193

**Table # 5 PHYTOPLANKTON VARIATIONS IN ABUNDANCE AND DIVERSITY IN SUB SURFACE SAMPLING STATIONS IN DPT HARBOUR AREA AT KANDLA CREEK AND ,NEAR BY CREEKS DURING NEAP TIDE IN OCTOBER,2021**

Tide	Sampling Station	Abundance In units/L	No of Species observed /total species	% of diversity	Margalef's diversity index (Species Richness S)	Shannon Weiner index H (log <sub>10</sub> )	Diversity Index (Simpson's Index) 1-D
HIGH TIDE	1	170	18/25	72	3.31	0.9214	0.8451
	2	133	14/25	56	2.658	0.8621	0.8308
	3	142	11/25	44	2.018	0.8693	0.847
	4	152	18/25	72	3.384	0.9173	0.8463
	5	128	12/25	48	2.267	0.8398	0.828
	6	88	14/25	56	2.904	0.9345	0.8406
LOW TIDE	1	143	11/25	44	2.015	0.8355	0.832
	2	149	13/25	52	2.398	0.8822	0.842
	3	139	16/25	64	3.04	0.9274	0.8526
	4	157	9/25	36	1.582	0.7937	0.8176
	5	120	9/25	36	1.671	0.7738	0.8094

**Table # 6 ABUNDANCE OF PHYTOPLANKTON SUBSURFACE SAMPLING STATIONS IN DPT HARBOUR AREA AT KANDLA CREEK AND , NEAR BY CREEKS DURING SPRING TIDE IN OCTOBER, 2021**

Tide	Surface	No of Sampling location	Group of phytoplankton	Phytoplankton Group range Units/L	Genera or species /total Phyto plankton	Taxon Diversity % (Group level)
HIGH TIDE	Sub surface	6	DIATOMS	39-204	19/24	79.2
			BLUE GREEN	3-8	3/24	12.5
			DINOFLAGELLATES	2-9	2/24	8.3
			TOTAL PHYTO PLANKTON	<b>46-209</b>	<b>24</b>	-
LOW TIDE	Sub surface	5	DIATOMS	176-222	19/24	79.2
			BLUE GREEN	3-8	3/24	12.5
			DINOFLAGELLATES	2-6	2/24	8.3
			TOTAL PHYTO PLANKTON	183-229	<b>24</b>	-

**Table # 7 ABUNDANCE OF PHYTOPLANKTON SUBSURFACE SAMPLING STATIONS IN DPT HARBOUR AREA AT KANDLA CREEK AND , NEAR BY CREEKS DURING NEAP TIDE IN OCTOBER,2021**

Tide	Surface	No of Sampling location	Group of phytoplankton	Phytoplankton Group range Units/L	Genera or species /total Phyto plankton	Taxon Diversity % (Group level)
HIGH TIDE	Sub surface	6	DIATOMS	81-164	20/25	80
			BLUE GREEN	1-6	3/25	12
			DINOFLAGELLATES	0-4	2/25	8
			TOTAL PHYTO PLANKTON	<b>88-170</b>		
LOW TIDE	Sub surface	5	DIATOMS	117-157	20/25	80
			BLUE GREEN	0-7	3/25	12
			DINOFLAGELLATES	0-4	2/25	8
			TOTAL PHYTO PLANKTON	<b>120-157</b>		

**Table # 8 PHYTOPLANKTON VARIATIONS IN ABUNDANCE AND DIVERSITY IN SUB SURFACE SAMPLING STATIONS IN DPT OOTAT PATH FINDER CREEK , VADINAR & NEAR BY SPM, DURING SPRING TIDE IN OCTOBER, 2021**

Tide	Sampling Station	Abundance In units/L	No of Species observed /total species	% of diversity	Margalef's diversity index (Species Richness S)	Shannon Weiner index H (log <sub>10</sub> )	Diversity Index (Simpson's Index) 1-D
HIGH TIDE	jetty	131	9/9	100	1.641	0.6874	0.7201
LOW TIDE	jetty	147	9/9	100	1.603	0.6956	0.7508

**Table # 9 PHYTOPLANKTON VARIATIONS IN ABUNDANCE AND DIVERSITY IN SUB SURFACE SAMPLING STATIONS IN DPT OOTAT PATH FINDER CREEK , VADINAR & NEAR BY SPM, DURING NEAP TIDE IN OCTOBER, 2021**

Tide	Sampling Station	Abundance In units/L	No of Species observed /total species	% of diversity	Margalef's diversity index (Species Richness S)	Shannon Weiner index H (log <sub>10</sub> )	Diversity Index (Simpson's Index) 1-D
HIGH TIDE	jetty	427	13/15	86.67	1.981	0.3559	0.3291
	SPM	203	13/15	86.67	2.259	0.5079	0.499
LOW TIDE	jetty	544	14/15	93.33	2.064	0.2555	0.2188
	SPM	744	11/15	73.33	1.512	0.2343	0.2011

**Table # 10 ABUNDANCE OF PHYTOPLANKTON SUBSURFACE SAMPLING STATIONS IN DPT DPTOOT AT PATH FINDER CREEK , VADINAR & NEAR BY SPM, DURING SPRING TIDE IN OCTOBER, 2021**


Tide	Surface	No of Sampling location	Group of phytoplankton	Phytoplankton Group range Units/L	Genera or species /total Phyto plankton	Taxon Diversity % (Group level)
HIGH TIDE	Sub surface	1	DIATOMS	131	9/9	100
			TOTAL PHYTO PLANKTON	<b>131</b>	<b>9</b>	
LOW TIDE	Sub surface	1	DIATOMS	147	9/9	100
			TOTAL PHYTO PLANKTON	147	<b>9</b>	

**Table # 11 ABUNDANCE OF PHYTOPLANKTON SUBSURFACE SAMPLING STATIONS IN DPT DPTOOT AT PATH FINDER CREEK , VADINAR & NEAR BY SPM, DURING NEAP TIDE IN OCTOBER, 2021**


Tide	Surface	No of Sampling location	Group of phytoplankton	Phytoplankton Group range Units/L	Genera or species /total Phyto plankton	Taxon Diversity % (Group level)
HIGH TIDE	Sub surface	2	DIATOMS	189-424	10/15	66.5
			BLUE GREEN	1-4	1/15	7.5
			DINOFLAGELLATES	4-10	4/15	26.0
			TOTAL PHYTO PLANKTON	<b>203-427</b>		
LOW TIDE	Sub surface	2	DIATOMS	480-664		
			BLUE GREEN	2	10/15	66.5
			DINOFLAGELLATES	5-10	1/15	7.5
			TOTAL PHYTO PLANKTON	544-744	4/15	26.0




Taxon Diversity % of Phytoplankton during High tide and Low tide period during spring tide in Kandala creek and nearby creeks




Taxon Diversity % of Phytoplankton during High tide and Low tide period during Neap tide in Kandala creek and nearby creeks



**Taxon Diversity % of Phytoplankton during High tide and Low tide period during spring tide in Path Finder Creek, Vadinar**



**Taxon Diversity % of Phytoplankton during High tide and Low tide period during Neap tide in Path Finder Creek, Vadinar**



**ZOOPLANKTON POPULATION:**

For the evaluation of the Zooplankton population in DPT harbour area and within the immediate surroundings of the port sampling was conducted from 6 sampling locations (3 in harbour area and two in Nakti creek and one in Khoricreek) during high tide period and low tide period of spring tide DCPL/DPT/20-21/18 -OCTOBER - 2021

and Neap tide in October, 2021. The Zooplankton community of the sub surface water in the harbour and nearby creeks during spring tide was represented by mainly 9 groups, and 7 larval forms; Tintinids, Copepods, Rotifers, Arrow worms, Mysids, Urochordates, Ciliates, Unidentified medusa and Foraminiferans. Larval forms represented from the group of Crustacea, Molluscs and Polychaetes. The Zooplankton community of the sub surface water in the harbour and nearby creeks during neap tide was represented by mainly 9 groups, and Eight larval forms. The major zooplankton group was Tintinids, Copepods, Rotifers, Arrow worms, Mysids, Urochordata, Ciliates, Medusa and Foraminiferans. Larval forms were represented from the group of Crustaceans, Echinodermata, Bryozoans, Molluscs and Polychaetes,

Zooplankton of the sampling stations at sub surface layer in the DPT harbour area and nearby creek was varying from  $33-132 \times 10^3$  N/ m<sup>3</sup> during high tide and  $81-107 \times 10^3$  N/ m<sup>3</sup> during low tide of Spring Tide period. Zooplankton of the sampling stations at sub surface layer in the DPT harbour area and nearby creek was varying from  $47-167 \times 10^3$  N/ m<sup>3</sup> during high tide and  $9-112 \times 10^3$  N/ m<sup>3</sup> during low tide of Neap Tide period.

For the evaluation of the Zoo plankton population in DPT OOT jetty area in Path Finder creek and SPM in Vadinar selected 2 sampling locations (1 in jetty area and one near SPM) During spring tide sampling plankton sample were collected only at Jetty area during consecutive high tide period and low tide period. During Neap tide sampling Plankton samples were collected from jetty area and SPM during consecutive high tide period and low tide period.

The Zooplankton community of the sub surface water in the path finder creek creeks during spring tide was represented by mainly two groups, Tintinids, Copepods, and larval forms of Crustacea and Molluscs. The Zooplankton community of the sub surface water in the path Finder creeks at Jetty region and SPM during neap tide was represented by mainly Five groups, Tintinids, Copepods, Arrow worms, Urochordata and Mysids. Larval forms were represented from the major group of Crustaceans, Molluscs, Echinodermata and Polychaetes,

Zooplankton of the sampling stations at sub surface layer in the DPT OOT jetty area of path finder creek was  $76 \times 10^3$  N/ m<sup>3</sup> during high tide and  $74 \times 10^3$  N/ m<sup>3</sup> during low tide of Spring Tide period. Zooplankton of the sampling stations at sub surface layer in the DPT OOT jetty area in path finder creek was recorded  $54 \times 10^3$  N/ m<sup>3</sup> during high tide and  $73 \times 10^3$  N/ m<sup>3</sup> during consecutive low tide period of Neap. Zooplankton of the sampling stations at sub surface layer in the DPT SPM area in path finder creek was recorded  $78 \times 10^3$  N/ m<sup>3</sup> during high tide and  $82 \times 10^3$  N/ m<sup>3</sup> during consecutive low tide period of Neap Tide.

### Species Richness Indices and Diversity Indices:

#### Margalef's diversity index (Species Richness) S

At the organismal level, the most widely used biodiversity measures are those based on the number of species present, perhaps adjusted for the number of individuals sampled, Here Margalef's Species richness index (d), or indices that describe the evenness of the distribution of the numbers of individuals among species, are derived.

Margalef's diversity index (Species Richness) S of Zooplankton communities in the stations Kandla creek region and nearby creeks was varying from 3.146-4.804 with an average of 3.874 during the sampling conducted in High tide period. Margalef's diversity index (Species Richness) S of Zooplankton communities varying from 2.842-4.280 with an average of 3.393 during the sampling conducted in low tide period during Spring tide.

Margalef's diversity index (Species Richness) S of Zooplankton communities in the Kandla creek region and nearby creeks sampling stations was varying from 3.117- 6.839 with an average of 5.036 during the sampling conducted in high tide and varying from 3.034 -4.570 with an average of 3.769 during the sampling conducted in low tide during Neap tide period

Margalef's diversity index (Species Richness) S of Zooplankton communities in the sampling station near jetty at Path Finder Creek, Vadinar during the sampling conducted in consecutive High tide period and low tide of spring tide was recorded as 1.847 and 1.859 respectively..

Margalef's diversity index (Species Richness) S of Zooplankton communities in the two stations at Path finder creek was varying from 3.259-3.443 during the sampling conducted in High tide period of Neap tide. While Margalef's diversity index (Species Richness) S of Zooplankton communities in the two stations at Path finder creek was varying from 2.797-4.085 during the consecutive low tide period.

#### Shannon-Wiener's index:

Shannon-Wiener's Index (H) of Zooplankton communities in the sampling stations in Kandla Harbour region and nearby creeks was in the range of 0.912 -1.060 ( $H'(\log_{10})$ ) between selected sampling stations with an average value of 0.992 ( $H'(\log_{10})$ ) during high tide period of spring tide. Shannon-Wiener's Index (H) of Zooplankton communities in the sampling stations in Kandla Harbour region and nearby creeks was in the range of 0.844-0.965 ( $H'(\log_{10})$ ) between selected sampling stations with an average value of 0.894 ( $H'(\log_{10})$ ) during consecutive low tide period .

Shannon-Wiener's Index (H) of Zooplankton communities in the sampling stations in Kandla Harbour region and nearby creeks was in the range of 0.834 -1.336 ( $H'(\log_{10})$ ) between selected sampling stations with an average value of 1.146 ( $H'(\log_{10})$ ) during high tide period of Neap tide. Shannon-Wiener's Index (H) of Zooplankton communities in the sampling stations in Kandla Harbour region

and nearby creeks was in the range of 0.949-1.139 ( $H'(\log_{10})$ ) between selected sampling stations with an average value of 1.043 ( $H'(\log_{10})$ ) during consecutive low tide period.

Shannon-Wiener's Index (H) of Zooplankton communities in the sampling station near jetty at Path Finder Creek, Vadinar during the sampling conducted in consecutive High tide period and low tide of spring tide was recorded as 0.775 and 0.822 respectively..

Shannon-Wiener's Index (H) of Zooplankton communities in the two stations at Path finder creek was varying from 0.991-0.927 during the sampling conducted in High tide period of Neap tide. While Shannon-Wiener's Index (H) of Zooplankton communities in the two stations at Path finder creek was varying from 0.815-1.004 during the consecutive low tide period.

Typical values are generally between 1.5 and 3.5 in most ecological studies, and the index is rarely greater than 4. The Shannon-Wiener's index increases as both the richness and the evenness of the community increase. This result indicates that diversity of Zooplankton of Kandla Harbour region and nearby creeks stations is slightly high with very minimum diverse population but very few opportunist organisms are really well adapted to this environment and thrive better than other species.

#### **Simpson's diversity index:**

Simpson's index (D) is a measure of diversity, which takes into account both species richness, and an evenness of abundance among the species present. The Simpson index is one of the meaningful and robust biodiversity measures available. (Magurran, 2004).

Simpson diversity index (1-D) of Zooplankton communities was below 0.9 most of sampling stations in the Kandla Harbour region and nearby creeks during high tide and low tide of spring tide period, which was varying from 0.848-0.881 between selected sampling stations with an average of 0.865 during high tide period and was varying from 0.812- 0.845 with an average value of 0.827 between selected sampling stations during low tide

Simpson diversity index (1-D) of Zooplankton communities was below 0.9 at all sampling stations in the Kandla Harbour region and nearby creeks except few during high tide and low tide period, which was varying from 0.793- 0.937 between selected sampling stations with an average of 0.894 during high tide period and was varying from 0.861- 0.910 with an average value of 0.884 between selected sampling stations during consecutive low tide

This high species diversity suggests a relatively more number of successful species in this habitat during October, 2021 sampling.

Simpson diversity index (1-D) of Zooplankton communities in the sampling station near jetty at Path Finder Creek, Vadinar during the sampling conducted in consecutive High tide period and low tide of spring tide was recorded as 0.800 and 0.827 respectively..

**Environmental Monitoring Report of Deendayal Port Trust, OCTOBER-2021**

Simpson diversity index (1-D) of Zooplankton communities in the two stations at Path finder creek was varying from 0.884 -0.838 during the sampling conducted in High tide period of Neap tide. While Shannon-Wiener's Index (H) of Zooplankton communities in the two stations at Path finder creek was varying from 0.816 -0.863 during the consecutive low tide period.

**Table # 12 ZOOPLANKTON VARIATION IN ABUNDANCE AND DIVERSITY IN SUB SURFACE SAMPLING STATIONS IN DPT HARBOUR AREA AT KANDLA CREEK AND NEAR BY CREEKS DURING SPRING TIDE IN OCTOBER,2021**

Tide	Sampling Station	Abundance In N / m <sup>3</sup>	No of Species/groups observed /total species/group	% of diversity	Margalef's diversity index (Species Richness S)	Shannon Weiner index H (log <sub>10</sub> )	Diversity Index (Simpson's Index) 1-D
HIGH TIDE	1	120 x10 <sup>3</sup>	24/32	75	4.804	1.06	0.8695
	2	105 x10 <sup>3</sup>	21/32	65.62	4.297	1.045	0.8753
	3	132 x10 <sup>3</sup>	18/32	56.25	3.482	0.9555	0.8523
	4	117 x10 <sup>3</sup>	18/32	56.25	3.57	1.032	0.8811
	5	123 x10 <sup>3</sup>	20/32	62.50	3.948	0.9513	0.8481
	6	33 x10 <sup>3</sup>	12/32	37.5	3.146	0.9125	0.8636
LOW TIDE	1	97 x10 <sup>3</sup>	14/32	43.75	2.842	0.8439	0.8174
	2	107 x10 <sup>3</sup>	21/32	65.63	4.28	0.9654	0.8455
	3	81 x10 <sup>3</sup>	15/32	46.87	3.186	0.9009	0.8315
	4	93 x10 <sup>3</sup>	16/32	50	3.309	0.8669	0.8125
	5	88 x10 <sup>3</sup>	16/32	50	3.35	0.8971	0.8325

**Table # 13 ZOOPLANKTON VARIATION IN ABUNDANCE AND DIVERSITY IN SUB SURFACE SAMPLING STATIONS IN DPT HARBOUR AREA AT KANDLA CREEK AND NEAR BY CREEKS DURING NEAP TIDE IN OCTOBER,2021**

Tide	Sampling Station	Abundance In No / m <sup>3</sup>	No of Species/groups observed /total species/group	% of diversity	Margalef's diversity index (Species Richness S)	Shannon Weiner index H (log <sub>10</sub> )	Diversity Index (Simpson's Index) 1-D
HIGH TIDE	1	167 x10 <sup>3</sup>	36/37	97.30	6.839	1.336	0.9367
	2	124 x10 <sup>3</sup>	28/37	75.68	5.601	1.282	0.9373
	3	133 x10 <sup>3</sup>	24/37	64.86	4.703	1.163	0.9108
	4	144 x10 <sup>3</sup>	29/37	78.38	5.634	1.19	0.8993
	5	102 x10 <sup>3</sup>	21/37	56.75	4.324	1.075	0.8893
	6	47 x10 <sup>3</sup>	13/37	35.13	3.117	0.834	0.7928
LOW TIDE	1	112 x10 <sup>3</sup>	18/37	48.65	3.603	0.9973	0.8637
	2	90 x10 <sup>3</sup>	19/37	51.35	4	1.065	0.8974
	3	99 x10 <sup>3</sup>	22/37	59.46	4.57	1.139	0.9105
	4	107 x10 <sup>3</sup>	18/37	48.65	3.638	1.068	0.8889
	5	101 x10 <sup>3</sup>	15/37	40.54	3.034	0.9498	0.861

**Table # 14 ABUNDANCE OF ZOOPLANKTON IN SUBSURFACE SAMPLING STATIONS IN DPT HARBOUR AREA ATKANDLA CREEK AND ,NEAR BY CREEKS DURING SPRING TIDE IN OCTOBER,2021**

Tide	Surface	No of Sampling locations	Group of Zooplankton	Abundance of Zooplankton $\times 10^3$ Group Range	Genera or species /total Zooplankton	Taxon Diversity % (Group level)
HIGH TIDE	Sub surface	6	Tintinids	3-21	6/32	18.75
			Copepods	13-68	10/32	31.25
			Rotifers	0-4	1/32	3.13
			Arrow worms	0-2	1/32	3.13
			Mysids	0-2	2/32	6.25
			Urochordata	0-1	1/32	3.12
			Ciliates	1-7	1/32	3.12
			Medusa	0-1	1/32	3.12
			Larval forms	11-43	7/32	21.88
			Foraminiferans	0-1	2/32	6.25
			<b>TOTAL ZOOPLANKTON N/ M<sup>3</sup></b>	<b>33-132</b>	<b>32</b>	
LOW TIDE	Sub surface	5	Tintinids	4-15	6/32	18.75
			Copepods	45-59	10/32	31.25
			Rotifers	0-1	1/32	3.13
			Arrow worms	0-1	1/32	3.13
			Mysids	0-1	2/32	6.25
			Urochordata	0-1	1/32	3.12
			Ciliates	1-6	1/32	3.12
			Medusa	0	1/32	3.12
			Larval forms	22-33	7/32	21.88
			Foraminiferans	0-2	2/32	6.25
			<b>TOTAL ZOOPLANKTON N/M<sup>3</sup></b>	<b>81-107</b>	<b>32</b>	

**Table # 15 ABUNDANCE OF ZOOPLANKTON IN SUBSURFACE SAMPLING STATIONS IN DPT HARBOUR AREA IN KANDLA CREEK AND , NEAR BY CREEKS DURING NEAP TIDE IN OCTOBER, 2021**

Tide	Surface	No of Sampling locations	Group of Zooplankton	Abundance of Zooplankton $\times 10^3$ Group Range	Genera or species /total Zooplankton	Taxon Diversity % (Group level)
HIGH TIDE	Sub surface	6	Tintinids	2-26	6/37	16.22
			Copepods	18-76	11/37	29.73
			Rotifers	0-2	1/37	2.70
			Mysids	0-6	4/37	10.81
			Arrow worms	0-4	1/37	2.70
			Urochordata	0-4	1/37	2.70
			Ciliates	0-8	1/37	2.70
			Medusa	0-4	1/37	2.70
			Larval forms	20-46	9/37	24.32
			Foraminiferans	0-6	2/37	5.42
			<b>TOTAL ZOOPLANKTON N/M<sup>3</sup></b>	<b>47-165</b>		
LOW TIDE	Sub surface	5	Tintinids	13-22	6/37	16.22
			Copepods	42-47	11/37	29.73
			Rotifers	0	1/37	2.70
			Mysids	0-4	4/37	10.81
			Arrow worms	0-1	1/37	2.70
			Urochordata	0-2	1/37	2.70
			Ciliates	1-5	1/37	2.70
			Medusa	0	1/37	2.70
			Larval forms	21-40	9/37	24.32
			Foraminiferans	0-5	2/37	5.42
			<b>TOTAL ZOOPLANKTON N/M<sup>3</sup></b>	<b>89-112</b>		

**Table # 16 ZOOPLANKTON VARIATION IN ABUNDANCE AND DIVERSITY IN SUB SURFACE SAMPLING STATIONS IN DPT OOT AREA AT PATH FINDER CREEK AND NEAR BY SPM DURING SPRING TIDE IN OCTOBER,2021**

Tide	Sampling Station	Abundance In N / m <sup>3</sup>	No of Species/groups observed /total species/group	% of diversity	Margalef's diversity index (Species Richness S)	Shannon Weiner index H (log <sub>10</sub> )	Diversity Index (Simpson's Index) 1-D
HIGH TIDE	Jetty	76 $\times 10^3$	9/9	100	1.847	0.7749	0.8004
LOW TIDE	Jetty	74 $\times 10^3$	9/9	100	1.859	0.8222	0.8278



**Table # 17 ZOOPLANKTON VARIATION IN ABUNDANCE AND DIVERSITY IN SUB SURFACE SAMPLING STATIONS IN DPT OOT AREA AT PATH FINDER CREEK AND NEAR BY SPM DURING NEAP TIDE IN OCTOBER,2021**

Tide	Sampling Station	Abundance In N / m <sup>3</sup>	No of Species/groups observed /total species/group	% of diversity	Margalef's diversity index (Species Richness S)	Shannon Weiner index H (log <sub>10</sub> )	Diversity Index (Simpson's Index) 1-D
HIGH TIDE	Jetty	54 x10 <sup>3</sup>	14/23	60.87	3.259	0.9911	0.884
	SPM	78 x10 <sup>3</sup>	16/23	69.56	3.443	0.9276	0.8385
LOW TIDE	Jetty	73 x10 <sup>3</sup>	13/23	56.52	2.797	0.8156	0.8166
	SPM	82 x10 <sup>3</sup>	19/23	82.61	4.085	1.004	0.863


**Table # 18 ABUNDANCE OF ZOOPLANKTON IN SUBSURFACE SAMPLING STATIONS IN DPT OOT AREA AT PATH FINDER CREEK AND NEAR BY SPM DURING SPRING TIDE IN OCTOBER,2021**

Tide	Surface	No of Sampling locations	Group of Zooplankton	Abundance of Zooplankton x10 <sup>3</sup> Group Range	Genera or species /total Zooplankton	Taxon Diversity % (Group level)
HIGH TIDE	Sub surface	1	Tintinids	40	5/9	55.56
			Copepods	18	2/9	22.22
			Larval forms	18	2/9	22.22
			TOTAL ZOOPLANKTON NO/L	<b>76</b>	<b>9</b>	
LOW TIDE	Sub surface	1	Tintinids	38	5/9	55.56
			Copepods	16	2/9	22.22
			Larval forms	20	2/9	22.22
			TOTAL ZOOPLANKTON NO/M3	<b>74</b>	<b>9</b>	


**Table # 19 ABUNDANCE OF ZOOPLANKTON IN SUBSURFACE SAMPLING STATIONS IN DPT OOT AREA AT PATH FINDER CREEK AND NEAR BY SPM DURING NEAP TIDE IN OCTOBER, 2021**

Tide	Surface	No of Sampling locations	Group of Zooplankton	Abundance of Zooplankton $\times 10^3$ Group Range	Genera or species /total Zooplankton	Taxon Diversity % (Group level)
HIGH TIDE	Sub surface	2	Tintinids	1-2	2/23	8.70
			Copepods	30-45	8/23	34.78
			Arrow worms	0-1	1/23	4.35
			Mysids	4-9	4/23	17.39
			Urochordata	2-4	1/23	4.35
			Larval forms	12-22	7/23	30.43
			<b>TOTAL ZOOPLANKTON</b>	<b>41-60</b>		
LOW TIDE	Sub surface	2	Tintinids	1-2	2/23	8.70
			Copepods	49-53	8/23	34.78
			Arrow worms	1	1/23	4.35
			Mysids	2-5	4/23	17.39
			Urochordata	1	1/23	4.35
			Larval forms	19-20	7/23	30.43
			<b>TOTAL ZOOPLANKTON NO/M3</b>	<b>73-82</b>		


**Taxon Diversity % of Zooplankton during High tide and Low tide period of Spring tide In Kandla Creek and nearby Creeks**




**Taxon Diversity % of Zooplankton during High tide and Low tide period of Neap tide In Kandla Creek and nearby Creeks**



**Taxon Diversity % of Zooplankton during High tide and Low tide period of Spring tide In Path Finder Creek and near Jetty**



**Taxon Diversity % of Zooplankton during High tide and Low tide period of Neap tide In Path Finder Creek near jetty and nearby SPM**



**TABLE # 20 SYSTEMATIC ACCOUNT OF PHYTOPLANKTON IN THE SAMPLING LOCATIONS OF DPT HARBOUR AREA AT KANDLA CREEK AND NEARBY CREEKS DURING SPRING TIDE OF OCTOBER 2021**

GROUP	PHYLUM	CLASS	ORDER	FAMILY	GENUS/SPECIES	#	Relative Abundance
BLUE GREEN ALGAE	Cyanophyta	Cyanophyceae	Nostocales	Oscillatoriaceae	<i>Oscillatoria sp.</i>	B1	Occasional
					<i>Arthrospira sp.</i>	B2	Rare
			Stigonematales	Stigonemataceae	<i>Stigonema sp.</i>	B3	Occasional
DIATOMS	Bacillariophyta	Coscinodiscophyceae	Thalassiosirales	Thalassiosiraceae	<i>Planktoniellasp</i>	D1	Occasional
			Coscinodiscales	Coscinodiscaceae	<i>Coscinodiscus sp.</i>	D2	Dominant
			Triceratiales	Triceratiaceae	<i>Odontellasp</i>	D3	Occasional
					<i>Triceratiumsp.</i>	D4	Rare
			Biddulphiales	Biddulphiaceae	<i>Biddulphiasp</i>	D5	Abundant
			Hemiaulales	Bellerocheaceae	<i>Bellerocheasp</i>	D6	Rare
			Rhizosoleniales	Rhizosoleniaceae	<i>Rhizosolenia sp.</i>	D7	Rare
			Chaetocerotales	Chaetocerotaceae	<i>Chaetocerossp</i>	D8	Frequent
			Leptocylindrales	Leptocylindraceae	<i>Leptocylindrussp</i>	D9	Occasional
			Lithodesmiales	Lithodesmiaceae	<i>Ditylumsp</i>	D10	Dominant
		Melosirales	Melosiraceae	<i>Melosirasp</i>	D11	Rare	
		Bacillariophyceae	Naviculales	Pleurosigmaaceae	<i>Pleurosigmasp</i>	D12	Rare
			Surirellales	Surirellaceae	<i>Surirellasp</i>	D13	Rare
		Fragilariophyceae	Thalassionematales	Thalassionemataceae	<i>Thalassiothrix sp.</i>	D14	Abundant
					<i>Thalassionema sp.</i>	D15	Rare
			Fragilariales	Fragilariaceae	<i>Asterionellopsis sp.</i>	D16	Occasional
					<i>Fragilariasp</i>	D17	Frequent
					<i>Synedrasp</i>	D18	Rare
		Tabellariales	Tabellariaceae	<i>Tabellariasp</i>	D19	Rare	
DINO FLAGELLATES	Dinoflagellata / Dinozoa	Dinophyceae	Peridinales	Protopteridiniaceae	<i>Protopteridium sp.</i>	DF1	Rare
			Gonyaulacales	Ceratiaceae	<i>Ceratiumfurca</i>	DF2	Occasional

**TABLE # 21 SYSTEMATIC ACCOUNT OF PHYTOPLANKTON IN THE SAMPLING LOCATIONS IN OF DPT HARBOUR AREA AT KANDLA CREEK AND NEARBY CREEKS DURING AND NEAP TIDE OF OCTOBER,2021**

GROUP	PHYLUM	CLASS	ORDER	FAMILY	GENUS/SPECIES	#	Relative Abundance
BLUE GREEN ALGAE	Cyanophyta	Cyanophyceae	Nostocales	Oscillatoriaceae	<i>Oscillatoria sp.</i>	B1	Rare
					<i>Arthrospira sp.</i>	B2	Rare
			Stigonematales	Stigonemataceae	<i>Stigonema sp.</i>	B3	Rare
DIATOMS	Bacillariophyta	Coccinodiscophyceae	Thalassiosirales	Thalassiosiraceae	<i>Planktoniellasp</i>	D1	Occasional
					<i>Thalassiosirasp</i>	D2	Rare
			Coccinodiscales	Coccinodiscaceae	<i>Coccinodiscus sp.</i>	D3	Dominant
			Triceratiales	Triceratiaceae	<i>Odontellasp</i>	D4	Occasional
					<i>Triceratiumsp.</i>	D5	Rare
			Biddulphiales	Biddulphiaceae	<i>Biddulphiasp</i>	D6	Abundant
			Hemiaulales	Belleracheaceae	<i>Belleracheasp</i>	D7	Rare
			Rhizosoleniales	Rhizosoleniaceae	<i>Rhizosolenia sp.</i>	D8	Rare
			Chaetocerotales	Chaetocerotaceae	<i>Chaetocerossp</i>	D9	Frequent
			Leptocylindrales	Leptocylindraceae	<i>Leptocylindrussp</i>	D10	Occasional
		Lithodesmiales	Lithodesmiaceae	<i>Ditylumsp</i>	D11	Dominant	
		Bacillariophyceae	Naviculales	Naviculaceae	<i>Naviculasp</i>	D12	Rare
				Pleurosigmaaceae	<i>Gyrosigma sp.</i>	D13	Rare
					<i>Pleurosigma sp.</i>	D14	Rare
		Surirellales	Surirellaceae	<i>Surirellasp</i>	D15	Rare	
		Fragilariophyceae	Thalassionematales	Thalassionemataceae	<i>Thalassiothrix sp.</i>	D16	Abundant
					<i>Thalassionema sp.</i>	D17	Rare
			Fragilariales	Fragilariaceae	<i>Fragilariasp</i>	D18	Frequent
					<i>Synedrassp</i>	D19	Rare
		Tabellariales	Tabellariaceae	<i>Tabellariasp</i>	D20	Rare	
DINO FLAGELLATES	Dinoflagellata / Dinozoa	Dinophyceae	Peridinales	Protoperidiniaceae	<i>Protoperidinium sp.</i>	DF1	Rare
			Gonyaulacales	Ceratiaceae	<i>Ceratiumfurca</i>	DF2	Occasional

**TABLE # 22 SYSTEMATIC ACCOUNT OF PHYTOPLANKTON IN THE SAMPLING LOCATIONS IN OF DPT OOT AREA AT PATH FINDER CREEK AND NEARBY SPM AT VADINAR DURING SPRING TIDE OF OCTOBER 2021**

GROUP	PHYLUM	CLASS	ORDER	FAMILY	GENUS/SPECIES	#	Relative Abundance
DIATOMS	Bacillariophyta	Coscinodiscophyceae	Coscinodiscales	Coscinodiscaceae	<i>Coscinodiscus sp.</i>	D1	Occasional
			Triceratiales	Triceratiaceae	<i>Triceratiumsp.</i>	D2	Rare
			Biddulphiales	Biddulphiaceae	<i>Biddulphiasp</i>	D3	Rare
			Hemiaulales	Bellercheaceae	<i>Bellercheasp</i>	D4	Rare
			Rhizosoleniales	Rhizosoleniaceae	<i>Rhizosolenia sp.</i>	D5	Rare
			Chaetocerotales	Chaetocerotaceae	<i>Chaetocerossp</i>	D6	Dominant
			Lithodesmiales	Lithodesmiaceae	<i>Ditylumsp</i>	D7	Occasional
			Bacillariales	Bacillariaceae	<i>Pseudo-Nitzschiasp</i>	D8	Abundant
		Fragilariophyceae	Thalassionematales	Thalassionemataceae	<i>Thalassiothrix sp.</i>	D9	Frequent

**TABLE # 23 SYSTEMATIC ACCOUNT OF PHYTOPLANKTON IN THE SAMPLING LOCATIONS IN OF DPT OOT AREA AT PATH FINDER CREEK AND NEARBY SPM AT VADINAR DURING AND NEAP TIDE OF OCTOBER, 2021**

GROUP	PHYLUM	CLASS	ORDER	FAMILY	GENUS/SPECIES	#	Relative Abundance
BLUE GREEN ALGAE	Cyanophyta	Cyanophyceae	Nostocales	Oscillatoriaceae	<i>Oscillatoria sp.</i>	B1	Rare
DIATOMS	Bacillariophyta	Coscinodiscophyceae	Thalassiosirales	Thalassiosiraceae	<i>Planktoniellasp</i>	D1	Occasional
				Lauderiaceae	<i>Lauderiasp</i>	D2	Rare
			Coscinodiscales	Coscinodiscaceae	<i>Coscinodiscus sp.</i>	D3	Abundant
			Hemiaulales	Bellercheaceae	<i>Bellercheasp</i>	D4	Occasional
			Rhizosoleniales	Rhizosoleniaceae	<i>Rhizosolenia sp.</i>	D5	Rare
			Chaetocerotales	Chaetocerotaceae	<i>Chaetocerossp</i>	D6	Dominant
			Lithodesmiales	Lithodesmiaceae	<i>Ditylumsp</i>	D7	Occasional
			Naviculales	Pleurosigmataceae	<i>Pleurosigmasp</i>	D8	Rare
		Bacillariales	Bacillariaceae	<i>Pseudo-Nitzschiasp</i>	D9	Frequent	
		Fragilariales	Fragilariaceae	<i>Synedrasp</i>	D10	Rare	

**Environmental Monitoring Report of Deendayal Port Trust, OCTOBER-2021**

GROUP	PHYLUM	CLASS	ORDER	FAMILY	GENUS/SPECIES	#	Relative Abundance
DINO FLAGELLATES	Dinoflagellata / Dinozoa	Dinophyceae	Peridiniales	Protoperidiniaceae	<i>Protoperidinium sp.</i>	DF1	Rare
			Gonyaulacales	Ceratiaceae	<i>Ceratiumfurca</i>	DF2	Occasional
					<i>Ceratiumfusus</i>	DF3	Rare
					<i>Ceratiumtripos</i>	DF4	Rare

**TABLE #24 SYSTEMATIC ACCOUNT OF ZOOPLANKTON FROM THE SAMPLING LOCATIONS OF DPT HARBOUR AREA AT KANDLA CREEK AND NEARBY CREEKS DURING SPRING TIDE OF OCTOBER, 2021**

GROUP	PHYLUM	CLASS	ORDER	FAMILY	GENUS/SPECIES	#	RELATIVE ABUNDANCE
TINTINIDS	PROTOZOA CILIOPHORA	Spirotrichea	Tintinnida	Tintinnidiidae	<i>Leprotintinnus sp.</i>	T1	Occasional
				Codonellidae	<i>Tintinnopsis dadayi</i>	T2	Rare
					<i>Tintinnopsis gracilis</i>	T3	Occasional
					<i>Tintinnopsis radix</i>	T4	Rare
					<i>Tintinnopsis failakkaensis</i>	T5	Occasional
Xystonellidae	<i>Favella sp.</i>	T6	Rare				
COPEPODS	ATHROPODA	Crustacea Subclass: Copepoda	Calanoida	Paracalanidae	<i>Acrocalanus sp.</i>	C1	Abundant
					<i>Bestiolina sp.</i>	C2	Rare
				Eucalanidae	<i>Subeucalanus sp.</i>	C3	Rare
				Clausocalanidae	<i>Clausocalanus sp.</i>	C4	Occasional
				Centropagidae	<i>Centropages sp.</i>	C5	Rare
			Acartiidae	<i>Acartia sp.</i>	C6	Rare	
			Cyclopoida	Oithonidae	<i>Oithona sp.</i>	C7	Dominant
			Harpacticoida	Ectinosomatidae	<i>Microsetella sp.</i>	C8	Rare
				Euterpinae	<i>Euterpina sp.</i>	C9	Frequent
				Poecilostomatoida	Oncaidae	<i>Oncaea sp.</i>	C10
ROTIFERS	ROTIFERA	Rotifera Subclass: Eurotatoria	Superorder: Monogononta Order: Ploimida	Brachionidae	<i>Brachionus plicatilis</i>	R1	Rare
ARROW WORMS	CHAETOGNATHA	Sagittioidea	Aphragmophora	Sagittidae	<i>Sagitta sp.</i>	A1	Rare
MYSIDS	ATHROPODA CRUSTACEA	Malacostraca	Mysida, Decapoda	Penaeidae	<i>Metapenaeus sp.</i>	M1	Rare
					<i>Penaeus sp.</i>	M2	Rare

**Environmental Monitoring Report of Deendayal Port Trust, OCTOBER-2021**

GROUP	PHYLUM	CLASS	ORDER	FAMILY	GENUS/SPECIES	#	RELATIVE ABUNDANCE
UROCHORDATA	CHORDATA SUB PHYLUM UROCHORDATA	Appendicularia		Oikopleuridae	<i>Oikopleura sp.</i>	U1	Rare
CILIATES	CILIOPHORA	Oligohymenophorea	Sessilida	Zoothamniidae	<i>Zoothamniumsp.</i>	CI1	Occasional
MEDUSA	PHYLUM CNIDARIA	Hydrozoa			Unidentified medusa	ME1	Rare
CRUSTACEAN LARVAE	ARTHROPODA (CRUSTACEA)	Copepoda			Nauplius larvae of Copepods	L1	Dominant
BRACHYURA LARVAE	ARTHROPODA (CRUSTACEA)	Malacostraca Decapoda			Brachyuran Zoea larvae	L2	Rare
BARNACLE LARVAE	ARTHROPODA CRUSTACEA	Maxillopoda Thecostraca			Cirripede larvae	L3	Rare
MOLLUSCAN LARVAE	MOLLUSCA	Gastropoda Streptoneura			Opisthobranchia larvae	L4	Rare
CHORDATA	VERTEBRATA	Pisces			Fish larvae	L5	Rare
POLYCHAETE LARVAE	ANNELIDA	Polychaeta			Trochophore larvae	L6	Occasional
BIVALVE LARVAE	MOLLUSCA	Pelecypoda			Veliger larvae of Bivalves	L7	Rare
FORAMINIFERA	FORAMINIFERA	Globothalamea	Rotaliida	Globigerinidae	<i>Globigerina sp.</i>	F1	Rare
				Rotalliidae	<i>Rotalia sp.</i>	F2	Rare



**TABLE # 25 SYSTEMATIC ACCOUNT OF ZOOPLANKTON FROM THE SAMPLING OF DPT HARBOUR AREA AT KANDLA CREEK AND NEARBY CREEKS DURING NEAP TIDE OF OCTOBER, 2021**

GROUP	PHYLUM	CLASS	ORDER	FAMILY	GENUS/SPECIES	#	RELATIVE ABUNDANCE
TINTINIDS	PROTOZOA CILIOPHORA	Spirotrichea	Tintinnida	Tintinnidiidae	<i>Leprotintinnus</i> sp.	T1	Frequent
				Codonellidae	<i>Tintinnopsis dadayi</i>	T2	Rare
					<i>Tintinnopsis gracilis</i>	T3	Rare
					<i>Tintinnopsis radix</i>	T4	Occasional
					<i>Tintinnopsis failakkaensis</i>	T5	Rare
				Codonellopsidae	<i>Codonellopsis</i> sp.	T6	Rare
COPEPODS	ATHROPODA	Crustacea Subclass: Copepoda	Calanoida	Paracalanidae	<i>Acrocalanus</i> sp.	C1	Abundant
				Eucalanidae	<i>Pareucalanus</i> sp.	C2	Rare
					<i>Subeucalanus</i> sp.	C3	Rare
					Clausocalanidae	<i>Clausocalanus</i> sp.	C4
				Centropagidae	<i>Centropages</i> sp.	C5	Rare
				Temoridae	<i>Temora</i> sp.	C6	Rare
				Acartiidae	<i>Acartia</i> sp.	C7	Occasional
			Cyclopoida	Oithonidae	<i>Oithona</i> sp.	C8	Abundant
			Harpacticoida	Ectinosomatidae	<i>Microsetella</i> sp.	C9	Occasional
				Euterpinae	<i>Euterpina</i> sp.	C10	Frequent
			Poecilostomatoida	Oncaeidae	<i>Oncaea</i> sp.	C11	Rare
ROTIFERS	ROTIFERA	Rotifera Subclass: Eurotatoria	Superorder: Monogononta Order: Ploimida	Brachionidae	<i>Brachionus plicatilis</i>	R1	Rare
ARROW WORMS	CHAETOGNATHA	Sagittoidea	Aphragmophora	Sagittidae	<i>Sagitta</i> sp.	A1	Rare
MYSIDS	ATHROPODA CRUSTACEA	Malacostraca	Mysida, Decapoda	Solenoceridae	<i>Solenocera</i> sp.	M1	Rare
				Penaeidae	<i>Metapenaeus</i> sp.	M2	Rare
					<i>Penaeus</i> sp.	M3	Rare
				Luciferidae	<i>Lucifer</i> sp.	M4	Rare
UROCHORDATA	CHORDATA SUB PHYLUM UROCHORDATA	Appendicularia		Oikopleuridae	<i>Oikopleura</i> sp.	U1	Rare
CILIATES	CILIOPHORA	Oligohymenophorea	Sessilida	Zoothamniidae	<i>Zoothamnium</i> sp.	CI1	Occasional

DCPL/DPT/20-21/18 -OCTOBER - 2021

**Environmental Monitoring Report of Deendayal Port Trust, OCTOBER-2021**

GROUP	PHYLUM	CLASS	ORDER	FAMILY	GENUS/SPECIES	#	RELATIVE ABUNDANCE
MEDUSA	PHYLUM CNIDARIA	Hydrozoa			Unidentified medusa	ME1	Rare
CRUSTACEAN LARVAE	ARTHROPODA (CRUSTACEA)	Copepoda			Nauplius larvae of Copepods	L1	Dominant
BRACHYURA LARVAE	ARTHROPODA (CRUSTACEA)	Malacostraca Decapoda			Brachyuran Zoea larvae	L2	Occasional
BARNACLE LARVAE	ARTHROPODA CRUSTACEA	Maxillopoda Thecostraca			Cirripede larvae	L3	Rare
CYPHONAUTES LARVAE	BRYOZOA				Cyphonautes larvae	L4	Occasional
MOLLUSCAN LARVAE	MOLLUSCA	Gastropoda Streptoneura			Opisthobranchia larvae	L5	Rare
ECHINODERMATA LARVAE	ECHINODERMATA				Ophioplutes larvae/ Echinoplutes larvae	L6	Rare
CHORDATA	VERTEBRATA	Pisces			Fish larvae	L7	Rare
POLYCHAETE LARVAE	ANNELIDA	Polychaeta			Trochophore larvae	L8	Occasional
BIVALVE LARVAE	MOLLUSCA	Pelecypoda			Veliger larvae of Bivalves	L9	Rare
FORAMINIFERA	FORAMINIFERA	Globothalamea	Rotalliida	Globigerinidae	<i>Globigerina</i> sp.	F1	Occasional
				Rotalliidae	<i>Rotalia</i> sp.	F2	Rare

**TABLE # 26 SYSTEMATIC ACCOUNT OF ZOOPLANKTON FROM THE SAMPLING LOCATIONS OF DPT OOT AREA AT PATH FINDER CREEK AND NEARBY SPM AT VADINAR DURING SPRING TIDE OF OCTOBER,2021**

GROUP	PHYLUM	CLASS	ORDER	FAMILY	GENUS/SPECIES	#	RELATIVE ABUNDANCE
TINTINIDS	PROTOZOA CILIOPHORA	Spirotrichea	Tintinnida	Tintinnidiidae	<i>Leprotintinnussp.</i>	T1	Dominant
				Codonellidae	<i>Tintinnopsisgracilis</i>	T2	Occasional
					<i>Tintinnopsis radix</i>	T3	Occasional
					<i>Tintinnopsistocantinensis</i>	T4	Occasional
				Xystonellidae	<i>Favella sp.</i>	T5	Rare
COPEPODS	ATHROPODA	Crustacea Subclass: Copepoda	Calanoida	Paracalanidae	<i>Acrocalanus sp.</i>	C1	Occasional
			Cyclopoida	Oithonidae	<i>Oithona sp.</i>	C2	Frequent
CRUSTACEAN LARVAE	ARTHROPODA (CRUSTACEA)	Copepoda			Nauplius larvae of Copepods	L1	Abundant
BIVALVE LARVAE	MOLLUSCA	Pelecypoda			Veliger larvae of Bivalves	L2	Rare

**TABLE # 27 SYSTEMATIC ACCOUNT OF ZOOPLANKTON FROM THE SAMPLING LOCATIONS OF DPT OOT AREA AT PATH FINDER CREEK AND NEARBY SPM AT VADINAR DURING NEAP TIDE OF OCTOBER,2021**

GROUP	PHYLUM	CLASS	ORDER	FAMILY	GENUS/SPECIES	#	RELATIVE ABUNDANCE
TINTINIDS	PROTOZOA CILIOPHORA	Spirotrichea	Tintinnida	Tintinnidiidae	<i>Leprotintinnussp.</i>	T1	Rare
				Xystonellidae	<i>Favella sp.</i>	T2	Rare
COPEPODS	ATHROPODA	Crustacea Subclass: Copepoda	Calanoida	Paracalanidae	<i>Acrocalanus sp.</i>	C1	Dominant
				Eucalanidae	<i>Pareucalanus sp.</i>	C2	Rare
					<i>Subeucalanus sp.</i>	C3	Rare
				Clausocalanidae	<i>Clausocalanus sp.</i>	C4	Occasional
			Tortanidae	<i>Tortanus sp.</i>	C5	Rare	
			Cyclopoida	Oithonidae	<i>Oithona sp.</i>	C6	Abundant
			Harpacticoida	Euterpinidae	<i>Euterpina sp.</i>	C7	Frequent
Poecilostomatatoida	Corycaeidae	<i>Corycaeus sp.</i>	C8	Rare			

**Environmental Monitoring Report of Deendayal Port Trust, OCTOBER-2021**

GROUP	PHYLUM	CLASS	ORDER	FAMILY	GENUS/SPECIES	#	RELATIVE ABUNDANCE
ARROW WORMS	CHAETOGNATHA	Sagittoidea	Aphragmophora	Sagittidae	<i>Sagitta sp.</i>	A1	Rare
MYSIDS	ARTHROPODA CRUSTACEA	Malacostraca	Mysida, Decapoda	Solenoceridae	<i>Solenocerasp.</i>	M1	Rare
				Penaeidae	<i>Metapenaeussp.</i>	M2	Rare
					<i>Penaeussp.</i>	M3	Rare
				Luciferidae	<i>Lucifer sp.</i>	M4	Rare
UROCHORDATA	CHORDATA SUB PHYLUM UROCHORDATA	Appendicularia		Oikopleuridae	<i>Oikopleura sp.</i>	U1	Rare
CRUSTACEAN LARVAE	ARTHROPODA (CRUSTACEA)	Copepoda			Nauplius larvae of Copepods	L1	Frequent
BRACHYURA LARVAE	ARTHROPODA (CRUSTACEA)	Malacostraca Decapoda			Brachyuran Zoea larvae	L2	Rare
BARNACLE LARVAE	ARTHROPODA CRUSTACEA	Maxillopoda Thecostraca			Cirripede larvae	L3	Rare
MOLLUSCAN LARVAE	MOLLUSCA	Gastropoda Streptoneura			Opisthobranchia larvae	L4	Rare
ECHINODERMATA LARVAE	ECHINODERMATA				Ophioplutes larvae/ Echinoplutes larvae	L5	Rare
POLYCHAETE LARVAE	ANNELIDA	Polychaeta			Trochophore larvae	L6	Occasional
BIVALVE LARVAE	MOLLUSCA	Pelecypoda			Veliger larvae of Bivalves	L7	Rare

## Environmental Monitoring Report Of Deendayal Port Trust, OCTOBER-2021

### BENTHIC ORGANISMS:

Few Benthic organisms were observed in the collected sediments by using the Van-veen grabs during the sampling conducted during spring tide period and while no Benthic organisms were observed during sampling conducted in Neap tide period from DPT harbour region and nearby creek except few dead shells. The meiobenthic organisms during spring tide were represented by Polychaetes, and Nematodes. The polychaetes were represented by *Scyphoproctus sp.*, during spring tide sampling. The meiobenthic organisms in the collected samples were varying from 40-60 N/M<sup>2</sup> during spring tide

**Table # 28 BENTHIC FAUNA IN THE SAMPLING LOCATIONS IN DPT HARBOUR AREA CREEKS DURING NEAP TIDE IN OCTOBER, 2021**

Benthic fauna	ABUNDANCE IN NO/M <sup>2</sup> DIFFERENT SAMPLING STATIONS REPRESENTATION BY GROUP						
	DPT HARBOUR			CREEKS			
POLYCHAETES	KPT-1	KPT-2	KPT-3	KPT-4	KPT-5	KPT-6	
Family : Capitellidae <i>Scyphoproctus sp.</i>	20	40	20	20	20		NS
<b>Total Polychaetes N/M<sup>2</sup></b>	20	40	20	20	20	NS	
<b>Un identified Nematode worms</b>	40	20	20	40	20	NS	
TOTAL Benthic Fauna NUMBER/ M <sup>2</sup>	60	60	40	60	40	-	

NS : No sample

## 7. Meteorological Data

Automatic Weather station have been installed in Seva Sadan -3 at the Deendayal Port which records the data on Temperature (°C), Humidity (%), Wind (mph), Dew Point (°C), Wind Direction (°), Pressure, Solar radiation, heat Index and UVI.

### Temperature

The mean day time temperature for Deendayal Port was 30.3 °C. The day-time maximum temperature was 38.6 °C. The mean night time temperature was 26.5 °C. The minimum mean night time temperature recorded was 30.6 °C.

### Air Pressure

The mean absolute air pressure for the month of October was 1009.9 hpa, whereas the mean relative pressure was 1009.3 hpa. The maximum absolute air pressure recorded for the month of October was 1016.5 hpa.

### Heat Index

The mean day-time heat index for the month of October was 33.8 °C. The maximum heat index recorded was 55°C.

### Solar Radiation

The mean Solar Radiation in October was 252.2 w/m<sup>2</sup>. The maximum solar radiation recorded in the month of October was 746.6 w/m<sup>2</sup>.

### Humidity

The mean day-time humidity was 60.0 % for the month of October and mean night time humidity was 72.4%. Maximum humidity recorded during day-time was 94.0 % and maximum humidity recorded during night-time was 93.0%.

### Wind Velocity and Wind Direction

The mean wind velocity for the entire month of October was 4.6 km/hour. Maximum wind velocity recorded was 29.2 Km/hr . The wind direction was mostly S to N.

### **Conclusive Summary and Remedial measures Suggested**

- The AAQ monitoring at six locations of Deendayal Port indicates that the mean PM<sub>10</sub> values at four locations viz. Coal storage area, Marine Bhavan and Oil Jetty area were found above the permissible standards (100 µg/m<sup>3</sup>) and PM<sub>2.5</sub> was above permissible limits at Coal storage location (Limit 60 µg/m<sup>3</sup>).
- Drinking water at all the twenty locations was found potable and was within permissible limits of BIS standards (IS 10500).
- Noise quality was also within the set permissible standards of an Industrial Area. The noise level observed during day time was >75 dB (A) and at night time was >70 dB (A) during the entire monitoring period.
- The sewage treated water of Deendayal Port Colony (Gopalpuri) was in line with the standards set by the Gujarat Pollution Control Board. The STP at Deendayal Port is not fully operational and STP at Vadinar Port was found non-operational.

### **Reasons for higher Values of PM<sub>10</sub>**

- Large amount of coal is handled at Berth No. 6, 7, 8 and 9. The unloading of coal directly in the truck, using grabs cause coal to spread in air as well as coal dust to fall on ground. This settled coal dust again mixes with the air while trucks travel through it.
- Also, the coal laden trucks are not always covered with tarpaulin sheets and these results in spillage of coal from trucks/dumpers during its transit from vessel to yard or storage site. This also increased PM values around marine Bhavan & Coal storage area.

### **Remedial Measures**

The values of PM<sub>10</sub> during the month of October, 2021 were observed beyond the permissible limit at four locations mentioned above. Given below are the remedial measures suggest to minimize the Air pollution at Deendayal Port.

- Guidelines for Coal Handling by GPCB should be strictly followed. (<http://gpcb.gov.in/pdf/coal-handling-guidelines.pdf>)
- Except for the higher values of PM<sub>10</sub> at Coal storage site, Oil Jetty, Tuna Port and Marine Bhavan locations, the monitoring results for the present month suggest that the overall Environment Quality of Deendayal Port is satisfactory.



**SOURCE OF LITERATURE AND ADDITIONAL REFERENCE FOR ECOLOGICAL STUDY**

- 1) ALBERT WEST PHAL (1976) Protozoa Blackwell , London
- 2) BANERJEE R.K. (1989) Heavy metals and Benthic foraminiferal distribution along Bombay coast India. Studies in benthic foraminifera. *Tokyo University Press* Tokyo pp 151-157
- 3) Banse K (1995) Zooplankton: Pivotal role in the control of ocean production: I. Biomass and production. *ICES J Mar Sci* 52: 265–277.
- 4) Beaugrand G, and Ibanez F (2004) Monitoring marine plankton ecosystems. II: long-term changes in North Sea calanoid copepods in relation to hydroclimatic variability. *Inter Res Mar Ecol Prog Ser* 284:35-47.
- 5) DAY F. (1889) The fauna of British India Ceylon and Burma- Fishes Vol-1- Vol-2 *Taylor and Francis* London
- 6) DESIKACHARY V. (1989) Atlas of diatoms, Madras Science Foundation
- 7) DESIKACHARY V. (1959) Cyanophyta ICAP Monographs on Algae *Indian Council of Agricultural research* New Delhi
- 8) FAIZAYOUSIF AL-YAMANI & MARIA A. SABUROVA (2010) illustrative guide on the flagellates of Intertidal soft sediment *Kuwait Institute for scientific Research* Kuwait
- 9) FAIZAYOUSIF AL-YAMANI, VALERIYSKRYABIN, ALEKSANDRA GUBANOVA, SERGEY KHVOROV AND IRINA PRUSOVA (2011), Marine zooplankton Practical guide from North western Arabian gulf Vol-1 and vol-2 *Kuwait Institute for scientific Research* Kuwait
- 10) FAUVEL P. (1953), The fauna of India Annelida - Polychaeta *Indian Press* Allahabad
- 11) Gajbhiye SN, Nair VR, and Desai BN (1984). Diurnal variation of zooplankton in Malad creek, Bombay. *Indian Journal of Marine Science*. 13:75-79.
- 12) HAYWARD P.J AND RYLAND J.S. (1995) Handbook of Marine fauna of north –West Europe *oxford University Press* London
- 13) HIGGINS R.P. HAJAMARTHIEL Eds. (1998) Introduction to the study of Meio Fauna
- 14) HORACE G. BARBER AND ELIZABETH Y. HAWORTH (1981) A guide to the Morphology of DIATOMS FRUSTULES.
- 15) INGRAM HENDEY (1964) An introductory account of smaller Algae of British coastal waters part-V. Bacillariophyceae
- 16) JOHN H. WICKSTEAD (1965) an Introduction to the study of Tropical Plankton .Hutchinson Tropical Monographs
- 17) JOYOTHIBABU, R. MADHU, N.V. MAHESHWARAN, P.A., NAIR K.K.C., VENUGOPL, P. BALASUBRAMANIAN T. (2005) Dominance of Dinoflagellates in micro zooplankton communities in the oceanic region Bay of Bengal and Andaman sea *Current science* vol.84. 10<sup>th</sup> May 2003
- 18) KASTURIRANGAN L.R. (1963) A key for the identification of the Common Planktonic Copepoda of Indian Coastal water
- 19) Kusum KK, Vineetha G, Raveendran TV, Nair VR, Muraleedharan KR, Achuthankutty CT and Joseph T (2014) Chaetognath community and their responses to varying environmental factors in the northern Indian ocean. *J Plankton Res* 36(4): 1146- 1152.
- 20) Lalli CM and Parsons TR (1997) Biological Oceanography: An Introduction. DOI <https://doi.org/10.1016/B978-0-7506-3384-0.X5056-7>.
- 21) Madhupratap M (1978) Studies on ecology of zooplankton of Cochin backwaters. *Mahasagar Bull Nat Inst Oceanogr* 11: 45-56.
- 22) Madhupratap M (1979) Distribution, community structure and species succession of copepods from Cochin Backwaters. *Indian J Mar Sci* 8: 1-8.

- 23) Madhupratap M (1987) Status and strategy of zooplankton of tropical Indian estuaries: A review. Bull Plank SocJpn 34: 65-81.
- 24) Madhupratap M (1999) Free living copepods of the Arabian Sea, Distribution and Research Perspectives. I J Mar Sci 146-149.
- 25) Madhupratap M and Haridas P (1986) Epipelagic calanoid copepods of the northern Indian Ocean. OceanologicaActa 9(2):105-117.
- 26) MANAL AL-KANDARI, FAIZA Y. AL-YAMANI , KHOLOOD AL-RIFAIE ( 2009) Marine phytoplankton Atlas of Kuwait's water *Kuwait Institute for scientific Research*
- 27) MPEDA (1998) Commercial Fishes and shell fishes of India
- 28) NEWEL G.E. & NEWELL R.C. (1963) Marine plankton a Practical Guide Hutchinson Educational
- 29) NIGAM R.C. AND CHATURVEDIS.K. (2000) Foraminiferal Study from KharoCreek , Kachchh ( Gujarat) North west coast of *India*. *Indian Journal of marine science* Vol.29 133-189
- 30) OLAV GIERE (1993) Meio benthology , Microscopic Fauna in Aquatic Sediments m Springer London
- 31) PERRAGALLO( 1965) Diatomees marines de franceA. *Asher & Co.* Amsterdam
- 32) Robert P.. Higgins (Eds.), (1985) An introduction to the study of Meio fauna Smithsons Institution press Washington DC
- 33) STERRER W. STERRERC.S Eds. Marine fauna and flora of Bermuda A systematic Guide to the identification of Marine Organisms. *John Wiley and Sons*New York
- 34) Suresh Gandhi. M. (2009) Distribution of certain ecological parameters and Foraminiferaldistribution in the depositional environment of Pak strait east coast of India .*Indian J. of Marine Science* Vol.33 pp 287-295
- 35) Venktaraman (1993) A systematic account of some south Indian diatoms . Proceeding of Indian Academy of Science Vol.X No.6 Sec.B.

\*\*\*\*\*

# ENVIRONMENTAL MONITORING REPORT FOR DEENDAYAL PORT TRUST



**REPORT** : **DCPL/DPT/20-21/19**  
**Mont** : **November**  
**Issue** : **01**  
**Revision** : **00**  
**Prepare** : **DETOX CORPORATION PVT. LTD., SURAT**

**TABLE OF CONTENTS**

<b>Sr. No.</b>	<b>Particulars</b>	<b>Page No.</b>
<b>1</b>	<b>Ambient Air Qualit Monitoring .....</b>	<b>1 - 18</b>
<b>2</b>	<b>Drinking Water Quality Monitoring .....</b>	<b>19 - 28</b>
<b>3</b>	<b>Noise Monitoring .....</b>	<b>29</b>
<b>4</b>	<b>Soil Monitoring .....</b>	<b>30 - 31</b>
<b>5</b>	<b>Sewage Treatment Plant Monitoring .....</b>	<b>32 - 38</b>
<b>6</b>	<b>Marine Water Monitoring .....</b>	<b>39 - 98</b>
<b>7</b>	<b>Meteorological Observations .....</b>	<b>99</b>
<b>8</b>	<b>Conclusive Summary &amp; Remedial Measures .....</b>	<b>100</b>
	<b>References .....</b>	<b>102- 103</b>

## **Introduction**

Monitoring of various environmental aspects of the Deendayal port by M/s Detox Corporation Pvt. Ltd. has been carried out through collection of samples, analysis of the same, comparing results with respect to the national standards and any other relevant standards by GBCB/CPCB/MoEF to identify non conformity in the Environment of the Deendayal Port. The results shall address the identified impacts and suggest measures to minimize the environmental impact due to various operations at Deendayal Port.

The environmental monitoring is carried out as per the Environment Management and Monitoring Plan submitted by Detox Corporation Pvt. Ltd.

## **1. Ambient Air Quality Monitoring**

As per the Environmental Monitoring Plan of Deendayal Port Trust, Air monitoring was carried out at six identified locations at Deendayal Port and two locations at Vadinar Port.

### **1.1 Air Quality Monitoring Methodology**

Air quality is measured in all the stations, for 24 hour for Total Suspended Particulate Matter (TSPM), PM<sub>10</sub>, PM<sub>2.5</sub>, SO<sub>2</sub>, NO<sub>x</sub>, NH<sub>3</sub> & Benzene, and Grab-sampling for CO & CO<sub>2</sub> measurements. The Air samplers are operated for a period of 24 hours and after a continuous operation of 8 hours of the sampler, the reagents were replaced to obtain 3 samples per day for each parameter namely, SO<sub>2</sub>, NO<sub>x</sub>. The EPM 2000 filter paper and PTFE Membrane bound filter paper are used for a period of 24 hours to obtain one sample each of TSPM, PM<sub>10</sub> & PM<sub>2.5</sub>. The AAQ samples are collected twice a week from all the eight locations as per the EMP.

### **1.2 Results**

The ambient air quality monitoring data for six stations, viz. Marine Bhavan, Oil Jetty, Port Colony, Gopalpuri Hospital, Tuna Port and Nr. Coal Storage Area for the month of November 2021 are given in Tables 1A to 6B. The ambient air quality monitoring data for two stations at Vadinar (Nr. Admin Building & Nr. Signal Building) are given in Tables 7A to 8B.

### **Location 1: Marine Bhavan (AL1)**

**Environmental Monitoring Report of Deendayal Port Trust,  
November-2021**

**Table 1 : Results of Air Pollutant Concentration at Marine Bhavan**

Parameter	Date	TSPM [µg/m <sup>3</sup> ]	PM10 [µg/m <sup>3</sup> ]	PM2.5 [µg/m <sup>3</sup> ]	SO <sub>2</sub> [µg/m <sup>3</sup> ]		NO <sub>x</sub> [µg/m <sup>3</sup> ]		NH <sub>3</sub> [µg/m <sup>3</sup> ]	
					8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
<b>Sampling Period</b>	-	<b>24hr</b>	<b>24hr</b>	<b>24hr</b>						
<b>NAAQMS limit</b>	-	<b>NS</b>	<b>100 µg/m<sup>3</sup></b>	<b>60 µg/m<sup>3</sup></b>		<b>80 µg/m<sup>3</sup></b>		<b>80 µg/m<sup>3</sup></b>		<b>400 µg/m<sup>3</sup></b>
<b>AL1 - 1</b>	01.11.21	326	193	86	3.52	3.96	22.23	21.60	9.70	10.21
					4.84		19.69		9.96	
					3.52		22.87		10.98	
<b>AL1 - 2</b>	09.11.21	303	156	90	5.71	6.15	15.88	20.54	13.02	13.36
					6.15		17.15		12.76	
					6.59		28.58		14.30	
<b>AL1 - 3</b>	12.11.21	402	191	96	7.47	6.74	28.58	27.31	11.49	11.49
					7.03		32.39		13.27	
					5.71		20.96		9.70	
<b>AL1 - 4</b>	17.11.21	438	180	90	3.08	2.64	19.69	16.73	15.57	16.00
					2.20		14.61		18.12	
					2.64		15.88		14.30	
<b>AL1 - 5</b>	19.11.21	530	156	88	4.40	4.40	20.96	20.54	5.62	9.19
					5.28		18.42		11.49	
					3.52		22.23		10.47	
<b>AL1 - 6</b>	24.11.21	468	182	90	2.64	3.52	14.61	16.30	10.98	7.49
					5.28		20.96		6.64	
					2.64		13.34		4.85	
<b>AL1 - 7</b>	26.11.21	597	274	92	3.52	2.93	14.61	17.78	14.30	10.89
					3.08		19.69		9.96	
					2.20		19.05		8.42	
<b>AL1 - 8</b>	29.11.21	613	210	90	2.20	2.78	26.04	24.98	10.47	8.00
					2.64		29.22		6.38	
					3.52		19.69		7.15	
<b>Monthly Average</b>		460	193	90		4.14		20.72		10.83
<b>Standard Deviation</b>		116	38	3		1.55		3.90		2.82

NS: Not Specified

**DCPL/DPT/20-21/19 -NOVEMBER - 2021**

**Detox Corporation Pvt. Ltd., Surat**

<b>Table 1B : Results of Air Pollutant Concentration at Marine Bhavan</b>					
<b>Parameter</b>	<b>Date</b>	<b>C<sub>6</sub>H<sub>6</sub> [µg/m<sup>3</sup>]</b>	<b>HC* ppm</b>	<b>CO [mg/m<sup>3</sup>]</b>	<b>CO<sub>2</sub> [ppm ]</b>
<b>Sampling Period</b>		<b>8 hr</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>
<b>NAAQMS limit</b>		<b>5.0 µg/m<sup>3</sup></b>	<b>NS</b>	<b>4.0 mg/m<sup>3</sup></b>	<b>NS</b>
AL1 - 1	1.11.21	1.27	BQL	2.15	581
AL1 - 2	9.11.21	1.12	BQL	1.9	497
AL1 - 3	12.11.21	1.11	BQL	1.94	361
AL1 - 4	17.11.21	1.05	BQL	2.04	357
AL1 - 5	19.11.21	1.12	BQL	1.85	541
AL1 - 6	24.11.21	1.38	BQL	1.91	561
AL1 - 7	26.11.21	1.07	BQL	1.98	567
AL1 - 8	29.11.21	1.14	BQL	2.21	541
Monthly Average		<b>1.16</b>	-	<b>2.00</b>	<b>501</b>
Standard Deviation		<b>0.11</b>	-	<b>0.13</b>	<b>91</b>

\* NMHC- Non- Methane Hydrocarbons

BQL- Below Quantification Limit (Quantification Limit - NMHC: 0.5 ppm)

NS -Not Specified

At Marine Bhavan, the overall values of TSPM, PM<sub>10</sub>, PM<sub>2.5</sub>, SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> is attributed mainly by motor vehicle emission produced from various types of automobiles (both diesel and petrol driven). Moreover, the loading and unloading of Food Grains and Timber at Jetty no. 1 and 2 also contributes to the high levels of TSPM and PM<sub>10</sub>. The mean TSPM value at Marine Bhavan was 460 µg/m<sup>3</sup>, The mean PM<sub>10</sub> values were 193.0 µg/m<sup>3</sup>, which is above the permissible limit. PM<sub>2.5</sub> values were above the permissible limit (mean 90.0 µg/m<sup>3</sup>). The average values of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were within the permissible limit. The average values of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were 4.14 µg/ m<sup>3</sup>, 20.72 µg/ m<sup>3</sup> & 10.83 µg/ m<sup>3</sup> respectively. These were within the permissible limit.

The levels of Benzene, Hydrocarbons (HC) and CO were within the permissible limit at Marine Bhavan. The mean Benzene concentration was 1.16 µg/m<sup>3</sup>, well below the permissible limit of 5.0 µg/m<sup>3</sup>. HC's were below the detectable limit and Carbon Monoxide concentration was 2.0 mg/m<sup>3</sup>, well below the permissible limit of 4.0 mg/m<sup>3</sup>.

**DCPL/DPT/20-21/19 -NOVEMBER - 2021**

**Detox Corporation Pvt. Ltd., Surat**

**Environmental Monitoring Report of Deendayal Port Trust,  
November-2021**

**Location 2: Oil Jetty (AL2)**

**Table 2 : Results of Air Pollutant Concentration at Oil Jetty**

Parameter s	Date	TSPM [µg/m <sup>3</sup> ]	PM10 [µg/m <sup>3</sup> ]	PM2.5 [µg/m <sup>3</sup> ]	SO <sub>2</sub> [µg/m <sup>3</sup> ]		NO <sub>x</sub> [µg/m <sup>3</sup> ]		NH <sub>3</sub> [µg/m <sup>3</sup> ]	
					8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
Sampling Period	-	24hr	24hr	24hr	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
NAAQMS limit	-	NS	100 µg/m <sup>3</sup>	60 µg/m <sup>3</sup>	-	80 µg/m <sup>3</sup>	-	80 µg/m <sup>3</sup>	-	400 µg/m <sup>3</sup>
AL2 - 1	01.11.21	421	151	92	3.96	3.66	17.15	15.67	13.79	14.81
					4.40		13.34		15.06	
					2.64		16.51		15.57	
AL2 - 2	09.11.21	532	176	89	1.76	2.93	13.34	12.49	6.13	9.70
					2.64		11.43		11.23	
					4.40		12.70		11.74	
AL2 - 3	12.11.21	539	180	96	7.03	8.79	32.39	25.19	4.85	7.32
					9.23		20.96		7.91	
					10.11		22.23		9.19	
AL2 - 4	17.11.21	510	200	101	3.96	2.93	15.88	15.24	7.91	9.79
					2.20		16.51		11.49	
					2.64		13.34		9.96	
AL2 - 5	19.11.21	407	234	98	3.52	2.49	17.78	19.48	9.19	8.00
					2.20		24.77		5.87	
					1.76		15.88		8.93	
AL2 - 6	24.11.21	520	152	100	7.03	6.45	20.96	15.88	5.87	8.42
					8.35		11.43		8.93	
					3.96		15.24		10.47	
AL2 - 7	26.11.21	434	150	98	1.32	1.90	22.87	19.69	9.19	10.04
					1.76		15.24		13.02	
					2.64		20.96		7.91	
AL2 - 8	29.11.21	551	278	100	2.20	2.93	16.51	18.42	6.64	9.02
					2.64		22.87		9.45	
					3.96		15.88		10.98	
<b>Monthly Average</b>		489	190	97		4.01		17.76		9.64
<b>Standard Deviation</b>		59	46	4		2.37		3.86		2.29

NS: Not Specified

**DCPL/DPT/20-21/19 -NOVEMBER - 2021**

**Detox Corporation Pvt. Ltd., Surat**



<b>Table 2B : Results of Air Pollutant Concentration at Oil Jetty</b>					
<b>Parameter</b>	<b>Date</b>	<b>C<sub>6</sub>H<sub>6</sub> [µg/m<sup>3</sup>]</b>	<b>HC* ppm</b>	<b>CO [mg/m<sup>3</sup>]</b>	<b>CO<sub>2</sub> [ppm ]</b>
<b>Sampling Period</b>		<b>8 hr</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>
<b>NAAQMS limit</b>		<b>5.0 µg/m<sup>3</sup></b>	<b>NS</b>	<b>4.0 mg/m<sup>3</sup></b>	<b>NS</b>
AL2 -1	1.11.21	1.12	BQL	2	618
AL2 -2	9.11.21	1.09	BQL	1.91	583
AL2 -3	12.11.21	1.07	BQL	2.08	509
AL2 -4	17.11.21	1.19	BQL	2.04	487
AL2 - 5	19.11.21	1.27	BQL	2.07	590
AL2 - 6	24.11.21	1.16	BQL	2.05	549
AL2 -7	26.11.21	1.17	BQL	1.99	578
AL2 - 8	29.11.21	1.09	BQL	1.98	624
<b>Monthly Average</b>		<b>1.15</b>	<b>-</b>	<b>2.02</b>	<b>567</b>
<b>Standard Deviation</b>		<b>0.07</b>	<b>-</b>	<b>0.06</b>	<b>49</b>

\* NMHC- Non- Methane Hydrocarbons

BQL- Below Quantification Limit (Quantification Limit – NMHC: 0.5 ppm)

NS- Not Specified

The overall values of TSPM, PM<sub>10</sub>, PM<sub>2.5</sub>, SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> at Oil Jetty area was mainly by motor vehicle emission produced from various types of vehicles Oil Jetty Area. The mean TSPM values at Oil Jetty were 489 µg/m<sup>3</sup>. The mean PM<sub>10</sub> values were 190 µg/m<sup>3</sup>, which is above the permissible limit. PM<sub>2.5</sub> values were above the permissible limit (mean = 97 µg/m<sup>3</sup>). The average values of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were within the permissible limit, The mean concentration of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were 4.01 µg/m<sup>3</sup>, 17.76 µg/m<sup>3</sup> and 9.79 µg/m<sup>3</sup> respectively.

The levels of Benzene, Hydrocarbons (HC) and CO were within the permissible limit at Oil Jetty. The mean Benzene concentration was 1.15 µg/m<sup>3</sup>. Well below the permissible limit of 5.0 µg/m<sup>3</sup>. , HC's were below the detectable limit and Carbon Monoxide concentration was 2.02 mg/m<sup>3</sup>, well below the permissible limit of 4.0 mg/m<sup>3</sup>.

**DCPL/DPT/20-21/19 -NOVEMBER - 2021**

**Detox Corporation Pvt. Ltd., Surat**

**Location 3: Kandla Colony - Estate Office (AL-3)**

**Table 3 : Results of Air Pollutant Concentration at Estate Office**

Parameters	Date	TSPM [µg/m <sup>3</sup> ]	PM10 [µg/m <sup>3</sup> ]	PM2.5 [µg/m <sup>3</sup> ]	SO <sub>2</sub> [µg/m <sup>3</sup> ]		NO <sub>x</sub> [µg/m <sup>3</sup> ]		NH <sub>3</sub> [µg/m <sup>3</sup> ]	
					8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
<b>NAAQMS limit</b>		<b>NS</b>	<b>100 µg/m<sup>3</sup></b>	<b>60 µg/m<sup>3</sup></b>	<b>-</b>	<b>80 µg/m<sup>3</sup></b>	<b>-</b>	<b>80 µg/m<sup>3</sup></b>	<b>-</b>	<b>400 µg/m<sup>3</sup></b>
<b>AL3 - 1</b>	01.11.21	205	114	80	2.20	2.64	22.87	22.23	13.79	16.34
					2.64		30.49		16.85	
					3.08		13.34		18.38	
<b>AL3 - 2</b>	09.11.21	302	134	90	3.08	3.08	13.97	13.97	9.70	7.49
					2.64		11.43		7.40	
					3.52		16.51		5.36	
<b>AL3 - 3</b>	12.11.21	422	215	92	3.52	3.81	13.34	19.27	8.93	6.98
					4.84		19.69		7.40	
					3.08		24.77		4.60	
<b>AL3 - 4</b>	17.11.21	610	270	108	5.28	3.08	10.80	9.32	13.02	10.21
					2.64		10.16		10.47	
					1.32		6.99		7.15	
<b>AL3 - 5</b>	19.11.21	459	269	100	5.28	3.96	26.04	25.41	8.93	9.19
					3.96		33.66		9.96	
					2.64		16.51		8.68	
<b>AL3 - 6</b>	24.11.21	736	363	102	5.71	4.84	19.69	19.48	10.47	8.93
					2.64		14.61		9.70	
					6.15		24.14		6.64	
<b>AL3 - 7</b>	26.11.21	483	180	98	5.28	3.81	20.96	17.57	11.49	10.98
					3.96		15.24		10.98	
					2.20		16.51		10.47	
<b>AL3 - 8</b>	29.11.21	677	189	105	2.20	3.22	15.88	16.73	12.25	9.02
					4.84		14.61		8.93	
					2.64		19.69		5.87	
<b>Monthly Average</b>		487	217	97		3.55		18.00		9.89
<b>Standard Deviation</b>		182	81	9		0.69		4.93		2.91

NS: Not Specified

<b>Table 3B : Results of Air Pollutant Concentration at Kandla Port Colony</b>					
<b>Parameter</b>	<b>Date</b>	<b>C<sub>6</sub>H<sub>6</sub> [µg/m<sup>3</sup>]</b>	<b>HC*</b>	<b>CO [mg/m<sup>3</sup>]</b>	<b>CO<sub>2</sub> [ppm]</b>
<b>Sampling Period</b>		<b>8 hr</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>
<b>NAAQMS limit</b>		<b>5.0 µg/m<sup>3</sup></b>	<b>NS</b>	<b>4.0 mg/m<sup>3</sup></b>	<b>NS</b>
AL3 -1	1.11.21	1.07	BQL	2.07	577
AL3 -2	9.11.21	1.1	BQL	1.91	583
AL3 -3	12.11.21	1.19	BQL	1.8	510
AL3 -4	17.11.21	1.11	BQL	1.91	480
AL3 - 5	19.11.21	1	BQL	2.1	652
AL3 - 6	24.11.21	1.1	BQL	1.84	672
AL3 - 7	26.11.21	1.26	BQL	2.12	364
AL3 - 8	29.11.21	1.26	BQL	2.01	426
<b>Monthly Average</b>		<b>1.14</b>	<b>-</b>	<b>1.97</b>	<b>533</b>
<b>Standard Deviation</b>		<b>0.09</b>	<b>-</b>	<b>0.12</b>	<b>108</b>

\* NMHC- Non- Methane Hydrocarbons

BQL- Below Quantification Limit (Quantification Limit - NMHC: 0.5 ppm)

NS- Not Specified

The overall values of TSPM, PM<sub>10</sub>, PM<sub>2.5</sub>, SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> at Kandla Port Colony was attributed by vehicle emission produced from trucks and heavy duty vehicles that pass through the road outside Kandla Port Colony. The mean TSPM values at Oil Jetty were 487 µg/m<sup>3</sup>, The mean PM<sub>10</sub> values were 217 µg/m<sup>3</sup>, which is above the permissible limit. PM<sub>2.5</sub> values were above the permissible limit (mean = 97 µg/m<sup>3</sup>). The average values of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were 3.55 µg/m<sup>3</sup>, 18.0 µg/m<sup>3</sup> and 9.89 µg/m<sup>3</sup> respectively and were all within the permissible limit.

The levels of Benzene, Hydrocarbons (HC) and CO were within the permissible limit at Kandla Port Colony. The mean Benzene concentration was 1.14 µg/m<sup>3</sup>, well below the permissible limit of 5.0 µg/m<sup>3</sup>. HC's were below the detectable limit and Carbon Monoxide concentration was 1.97 mg/m<sup>3</sup>, well below the permissible limit of 4.0 mg/m<sup>3</sup>.

**Environmental Monitoring Report of Deendayal Port Trust,  
November-2021**

**Location 4: Gopalpuri Hospital (AL-4)**

<b>Table 4 : Results of Air Pollutant Concentration at Gopalpuri Hospital</b>										
<b>Parameter</b>	<b>Date</b>	<b>TSPM [µg/m3]</b>	<b>PM10 [µg/m3]</b>	<b>PM2.5 [µg/m3]</b>	<b>SO2 [µg/m3]</b>		<b>NOx [µg/m3]</b>		<b>NH3 [µg/m3]</b>	
<b>Sampling Period</b>		<b>24hr</b>	<b>24hr</b>	<b>24hr</b>	<b>8 hr</b>	<b>24hr (Avg.)</b>	<b>8 hr</b>	<b>24hr (Avg.)</b>	<b>8 hr</b>	<b>24hr (Avg.)</b>
<b>NAAQMS limit</b>		<b>NS</b>	<b>100 µg/m3</b>	<b>60 µg/m3</b>	<b>-</b>	<b>80 µg/m3</b>	<b>-</b>	<b>80 µg/m3</b>	<b>-</b>	<b>400 µg/m 3</b>
<b>AL4 -1</b>	01.11.21	145	81	36	3.52	2.78	8.26	12.91	9.19	8.93
					2.20		15.88		10.47	
					2.64		14.61		7.15	
<b>AL4 -2</b>	09.11.21	254	132	89	2.64	2.05	9.53	12.28	5.87	5.36
					1.32		13.34		5.36	
					2.20		13.97		4.85	
<b>AL4 -3</b>	12.11.21	309	136	92	2.20	2.78	10.16	9.95	3.83	5.02
					3.52		8.26		5.87	
					2.64		11.43		5.36	
<b>AL4 -4</b>	17.11.21	474	249	101	3.52	2.78	9.53	10.16	6.64	5.70
					2.64		11.43		4.60	
					2.20		9.53		5.87	
<b>AL4 -5</b>	19.11.21	298	127	90	2.64	2.64	16.51	15.46	5.87	6.81
					3.52		10.16		8.42	
					1.76		19.69		6.13	
<b>AL4 -6</b>	24.11.21	351	170	98	3.08	2.49	15.88	13.13	5.87	8.85
					2.64		10.16		9.19	
					1.76		13.34		11.49	
<b>AL4 -7</b>	26.11.21	285	132	87	2.64	3.22	17.78	16.73	6.38	6.55
					3.96		16.51		7.91	
					3.08		15.88		5.36	
<b>AL4 -8</b>	29.11.21	738	469	104	3.52	3.37	13.34	13.34	8.17	8.34
					3.96		14.61		8.93	
					2.64		12.07		7.91	
<b>Monthly Average</b>		357	187	87		2.77		12.99		6.95
<b>Standard Deviation</b>		180	124	21		0.41		2.33		1.58

NS: Not Specified

<b>Table 4B : Results of Air Pollutant Concentration at Gopalpuri Hospital</b>					
<b>Parameter</b>	<b>Date</b>	<b>C<sub>6</sub>H<sub>6</sub> [µg/m<sup>3</sup>]</b>	<b>HC*</b>	<b>CO [mg/m<sup>3</sup>]</b>	<b>CO<sub>2</sub> [ppm ]</b>
<b>Sampling Period</b>		<b>8 hr</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>
<b>NAAQMS limit</b>		<b>5.0 µg/m<sup>3</sup></b>	<b>NS</b>	<b>4.0 mg/m<sup>3</sup></b>	<b>NS</b>
AL4 -1	1.11.21	1.02	BQL	2.01	609
AL4 -2	9.11.21	1.15	BQL	2.04	509
AL4 -3	12.11.21	1.26	BQL	1.94	487
AL4 -4	17.11.21	1.09	BQL	1.73	450
AL4 - 5	19.11.21	1.08	BQL	1.88	544
AL4 - 6	24.11.21	1.2	BQL	2.13	580
AL4 - 7	26.11.21	1.18	BQL	1.91	559
AL4 - 8	29.11.21	1.14	BQL	2.2	505
<b>Monthly Average</b>		<b>1.14</b>	<b>-</b>	<b>1.98</b>	<b>530</b>
<b>Standard Deviation</b>		<b>0.08</b>	<b>-</b>	<b>0.15</b>	<b>52</b>

\* NMHC- Non- Methane Hydrocarbons

BQL- Below Quantification Limit (Quantification Limit - NMHC: 0.5 ppm)

NS-Not Specified

The overall values of TSPM, PM<sub>10</sub>, PM<sub>2.5</sub>, SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> at Gopalpuri Hospital was attributed by vehicle emission produced from light motor vehicles of the colony residents. The mean TSPM values at Oil Jetty were 357 µg/m<sup>3</sup>, The mean PM<sub>10</sub> values were 187 µg/m<sup>3</sup>, which is above the permissible limit. PM<sub>2.5</sub> values were above the permissible limit (mean= 87 µg/m<sup>3</sup>). The average values of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were 2.77 µg/m<sup>3</sup>, 12.99 µg/m<sup>3</sup> and 6.95 µg/m<sup>3</sup> respectively and were all within the permissible limit.

The levels of Benzene, Hydrocarbons (HC) and CO were within the permissible limit at Gopalpuri Hospital. The mean Benzene concentration was 1.14 µg/m<sup>3</sup>, well below the permissible limit of 5.0 µg/m<sup>3</sup>. HC's were below the detectable limit and Carbon Monoxide concentration was 1.98 mg/m<sup>3</sup>, well below the permissible limit of 4.0 mg/m<sup>3</sup>.

**Environmental Monitoring Report of Deendayal Port Trust,  
November-2021**

**Location 5: Coal Storage Area (AL-5)**

**Table 5 : Results of Air Pollutant Concentration at Coal Storage Area**

Parameters	Date	TSPM [µg/m <sup>3</sup> ]	PM10 [µg/m <sup>3</sup> ]	PM2.5 [µg/m <sup>3</sup> ]	SO <sub>2</sub> [µg/m <sup>3</sup> ]		NO <sub>x</sub> [µg/m <sup>3</sup> ]		NH <sub>3</sub> [µg/m <sup>3</sup> ]	
					8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
<b>NAAQMS limit</b>	-	<b>NS</b>	<b>100 µg/m<sup>3</sup></b>	<b>60 µg/m<sup>3</sup></b>	-	<b>80 µg/m<sup>3</sup></b>	-	<b>80 µg/m<sup>3</sup></b>	-	<b>400 µg/m<sup>3</sup></b>
<b>AL5 - 1</b>	01.11.21	306	154	89	2.20	3.22	20.96	20.33	8.93	9.79
					3.08		21.60		7.40	
					4.40		18.42		13.02	
<b>AL5 - 2</b>	09.11.21	590	288	106	9.67	6.15	19.69	19.69	15.83	16.34
					4.84		23.50		17.61	
					3.96		15.88		15.57	
<b>AL5 - 3</b>	12.11.21	680	351	104	10.5 5	9.08	22.87	26.68	8.68	7.83
					7.03		26.04		6.64	
					9.67		31.12		8.17	
<b>AL5 - 4</b>	17.11.21	712	355	108	2.20	2.49	19.69	20.96	13.53	10.21
					1.76		20.96		9.19	
					3.52		22.23		7.91	
<b>AL5 - 5</b>	19.11.21	696	380	110	5.28	5.13	20.96	22.44	10.72	11.23
					6.15		19.69		9.96	
					3.96		26.68		13.02	
<b>AL5 - 6</b>	24.11.21	622	352	101	3.52	4.40	14.61	17.57	15.83	14.64
					3.96		18.42		15.06	
					5.71		19.69		13.02	
<b>AL5 - 7</b>	26.11.21	578	218	106	3.52	3.96	16.51	18.63	10.98	9.36
					4.84		20.96		9.19	
					3.52		18.42		7.91	
<b>AL5 - 8</b>	29.11.21	596	241	108	5.28	3.37	22.23	26.25	10.98	12.76
					2.64		27.31		13.02	
					2.20		29.22		14.30	
<b>Monthly Average</b>		598	292	104		4.73		21.57		11.52
<b>Standard Deviation</b>		128	81	7		2.10		3.36		2.87

NS: Not Specified

**DCPL/DPT/20-21/19 -NOVEMBER - 2021**

**Detox Corporation Pvt. Ltd., Surat**

<b>Table 5B : Results of Air Pollutant Concentration at Coal Storage Area</b>					
<b>Parameter</b>	<b>Date</b>	<b>C<sub>6</sub>H<sub>6</sub> [µg/m<sup>3</sup>]</b>	<b>HC*</b>	<b>CO [mg/m<sup>3</sup>]</b>	<b>CO<sub>2</sub> [ppm ]</b>
<b>Sampling Period</b>		<b>8 hr</b>	<b>Grab Samplin g</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>
<b>NAAQMS limit</b>		<b>5.0 µg/m<sup>3</sup></b>	<b>NS</b>	<b>4.0 mg/m<sup>3</sup></b>	<b>NS</b>
AL5 - 1	1.11.21	1.1	BQL	1.97	593
AL5 - 2	9.11.21	1.28	BQL	1.56	582
AL5 - 3	12.11.21	1.07	BQL	2.1	453
AL5 - 4	17.11.21	1.02	BQL	1.98	494
AL5 - 5	19.11.21	1.17	BQL	2.07	544
AL5 - 6	24.11.21	1	BQL	1.91	571
AL5 - 7	26.11.21	1.23	BQL	1.78	636
AL5 - 8	29.11.21	1.20	BQL	2.07	511
<b>Monthly Average</b>		<b>1.13</b>	<b>-</b>	<b>1.93</b>	<b>548</b>
<b>Standard Deviation</b>		<b>0.10</b>	<b>-</b>	<b>0.18</b>	<b>59</b>

\* NMHC- Non- Methane Hydrocarbons

BQL- Below Quantification Limit (Quantification Limit - NMHC: 0.5 ppm)

NS-Not Specified

The overall values of TSPM, PM<sub>10</sub>, PM<sub>2.5</sub>, SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> at Coal Storage Area was comparatively highest among all the locations of Air Quality monitoring in Kandla Port. High values of TSPM, PM<sub>10</sub>, PM<sub>2.5</sub>, SO<sub>2</sub>, NO<sub>x</sub> at this location was due to lifting of coal with grab and other coal handling processes near Berth no. 6 & 7. Moreover, the traffic was also heavy around this place for transport of coal thus emissions produced from heavy vehicles. The mean TSPM values at Coal storage were 598µg/m<sup>3</sup>. The mean PM<sub>10</sub> values were 292 µg/m<sup>3</sup>, which is well above the permissible limit. PM<sub>2.5</sub> values were above the permissible limit (mean = 104 µg/m<sup>3</sup>). The average values of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were 4.73 µg/m<sup>3</sup>, 21.57 µg/m<sup>3</sup> and 11.52 µg/m<sup>3</sup> respectively and were all within the permissible limit.

The levels of Benzene, Hydrocarbons (HC) and CO were within the permissible limit at Coal Storage Area. The mean Benzene concentration was 1.13 µg/m<sup>3</sup>, well below the permissible limit of 5.0 µg/m<sup>3</sup>. HC's were below the detectable limit and Carbon Monoxide concentration was 1.93 mg/m<sup>3</sup>, well below the permissible limit of 4.0 mg/m<sup>3</sup>.

**Environmental Monitoring Report of Deendayal Port Trust,  
November-2021**

**Location 6: Tuna Port (AL-6)**

**Table 6 : Results of Air Pollutant Concentration at Tuna Port**

Parameters	Date	TSPM [µg/m3]	PM10 [µg/m3]	PM2.5 [µg/m3]	SO2 [µg/m3]		NOx [µg/m3]		NH3 [µg/m3]	
					8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
Sampling Period	-	24hr	24hr	24hr	-	80 µg/m3	-	80 µg/m3	-	400 µg/m3
NAAQMS limit	-	NS	100 µg/m3	60 µg/m3	-	80 µg/m3	-	80 µg/m3	-	400 µg/m3
AL6 - 1	01.11.21	214	97	61	3.52	2.64	20.96	14.82	14.04	12.08
					2.64		12.07		12.51	
					1.76		11.43		9.70	
AL6 - 2	09.11.21	314	149	102	1.76	1.90	10.16	10.59	6.38	7.15
					2.64		11.43		6.64	
					1.32		10.16		8.42	
AL6 - 3	12.11.21	425	208	104	4.84	4.25	16.51	17.57	6.64	6.55
					5.28		20.96		4.85	
					2.64		15.24		8.17	
AL6 - 4	17.11.21	504	280	110	2.20	2.34	5.08	7.20	6.13	7.06
					1.76		7.62		8.42	
					3.08		8.89		6.64	
AL6 - 5	19.11.21	432	242	106	2.64	2.20	22.87	18.21	13.02	11.74
					2.20		13.34		11.74	
					1.76		18.42		10.47	
AL6 - 6	24.11.21	315	149	96	5.28	3.66	30.49	26.89	9.96	12.76
					1.76		22.23		15.57	
					3.96		27.95		12.76	
AL6 - 7	26.11.21	326	140	98	2.20	2.93	13.34	15.03	9.19	9.53
					3.08		15.24		8.93	
					3.52		16.51		10.47	
AL6 - 8	29.11.21	569	298	104	2.20	2.93	15.88	15.88	10.72	10.30
					2.64		13.34		8.93	
					3.96		18.42		11.23	
<b>Monthly Average</b>		387	195	98		2.86		15.77		9.65
<b>Standard Deviation</b>		116	73	15		0.78		5.79		2.48

NS: Not Specified



<b>Table 6B : Results of Air Pollutant Concentration at Tuna Port</b>					
<b>Parameter</b>	<b>Date</b>	<b>C<sub>6</sub>H<sub>6</sub> [µg/m<sup>3</sup>]</b>	<b>HC*</b>	<b>CO [mg/m<sup>3</sup>]</b>	<b>CO<sub>2</sub> [ppm]</b>
<b>Sampling Period</b>		<b>8 hr</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>
<b>NAAQMS limit</b>		<b>5.0 µg/m<sup>3</sup></b>	<b>NS</b>	<b>4.0 mg/m<sup>3</sup></b>	<b>NS</b>
AL6 -1	01.11.21	1.2	BQL	1.91	586
AL6 - 2	09.11.21	1.02	BQL	2.1	600
AL6 - 3	12.11.21	1.08	1.91	1.91	452
AL6 - 4	17.11.21	1.26	BQL	1.94	507
AL6 - 5	19.11.21	1.21	BQL	2.08	593
AL6 - 6	24.11.21	1.01	BQL	1.98	568
AL6 - 7	26.11.21	1.203	BQL	2.08	577
AL6 - 8	29.11.21	1.19	BQL	2.04	517
<b>Monthly Average</b>		<b>1.15</b>	<b>-</b>	<b>2.01</b>	<b>550</b>
<b>Standard Deviation</b>		<b>0.10</b>	<b>-</b>	<b>0.08</b>	<b>52</b>

\* NMHC- Non- Methane Hydrocarbons

BQL- Below Quantification Limit (Quantification Limit - NMHC: 0.5 ppm)

NS- Not Specified

The mean TSPM values at Tuna Port were 387 µg/m<sup>3</sup>, The mean PM<sub>10</sub> values were 195 µg/m<sup>3</sup>, which is above the permissible limit. PM<sub>2.5</sub> values were above the permissible limit (mean = 98 µg/m<sup>3</sup>). The average values of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were 2.86 µg/m<sup>3</sup>, 15.77 µg/m<sup>3</sup> and 9.65 µg/m<sup>3</sup> respectively and were all within the permissible limit.

The levels of Benzene, Hydrocarbons (HC) and CO were within the permissible limit at Tuna Port. The mean Benzene concentration was 1.15 µg/m<sup>3</sup>, well below the permissible limit of 5.0 µg/m<sup>3</sup>. HC's were below the detectable limit and Carbon Monoxide concentration was 2.01 mg/m<sup>3</sup>, well below the permissible limit of 4.0 mg/m<sup>3</sup>.

**Environmental Monitoring Report of Deendayal Port Trust,  
November-2021**

**Location 7: Signal Building (Vadinar) (AL-7)**

Table 7 : Results of Air Pollutant Concentration at Signal Building										
Parameters	Date	TSPM [µg/m <sup>3</sup> ]	PM10 [µg/m <sup>3</sup> ]	PM2.5 [µg/m <sup>3</sup> ]	SO <sub>2</sub> [µg/m <sup>3</sup> ]		NO <sub>x</sub> [µg/m <sup>3</sup> ]		NH <sub>3</sub> [µg/m <sup>3</sup> ]	
Sampling Period	-	24hr	24hr	24hr	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
NAAQMS limit	-	NS	100 µg/m <sup>3</sup>	60 µg/m <sup>3</sup>	-	80 µg/m <sup>3</sup>	-	80 µg/m <sup>3</sup>	-	400 µg/m <sup>3</sup>
AL7 -1	01.11.21	251	137	109	2.20	3.22	7.62	11.86	6.89	5.70
					3.96		13.34		6.13	
					3.52		14.61		4.08	
AL7 -2	09.11.21	215	115	84	5.71	6.30	26.04	17.36	4.60	6.98
					6.15		14.61		10.47	
					7.03		11.43		5.87	
AL7 -3	12.11.21	202	104	76	3.52	3.96	10.16	16.51	6.38	8.68
					4.84		26.04		10.72	
					3.52		13.34		8.93	
AL7 -4	17.11.21	200	103	84	2.64	3.96	19.69	14.40	7.91	6.13
					5.28		13.34		4.60	
					3.96		10.16		5.87	
AL7 -5	19.11.21	224	104	94	5.71	3.96	13.97	16.30	9.19	7.66
					3.52		19.69		7.15	
					2.64		15.24		6.64	
AL7 -6	24.11.21	238	118	77	4.40	4.40	10.16	10.59	5.87	5.96
					2.64		6.99		4.60	
					6.15		14.61		7.40	
AL7 -7	26.11.21	213	120	64	2.64	3.52	15.88	14.40	14.04	10.38
					2.20		7.62		10.72	
					5.71		19.69		6.38	
AL7 -8	29.11.21	207	115	84	5.71	3.22	10.16	13.34	8.17	7.66
					2.64		15.24		7.91	
					1.32		14.61		6.89	
<b>Monthly Average</b>		219	115	84		4		14		7
<b>Standard Deviation</b>		18	11	13		1		2		2

NS: Not Specified

**DCPL/DPT/20-21/19 -NOVEMBER - 2021**

**Detox Corporation Pvt. Ltd., Surat**

<b>Table 7B : Results of Air Pollutant Concentration at Signal Building</b>					
<b>Parameter</b>	<b>Date</b>	<b>C<sub>6</sub>H<sub>6</sub> [µg/m<sup>3</sup>]</b>	<b>HC*</b>	<b>CO [mg/m<sup>3</sup>]</b>	<b>CO<sub>2</sub> [ppm ]</b>
<b>Sampling Period</b>		<b>8 hr</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>
<b>NAAQMS limit</b>		<b>5.0 µg/m<sup>3</sup></b>	<b>NS</b>	<b>4.0 mg/m<sup>3</sup></b>	<b>NS</b>
AL7 -1	01.11.21	1.03	BQL	1.75	569
AL7 - 2	09.11.21	1.16	BQL	1.85	629
AL7 - 3	15.11.2021	1.35	BQL	1.78	501
AL7 - 4	18.11.2021	1.09	BQL	2	449
AL7 - 5	19.11.2021	1	BQL	1.89	458
AL7 - 6	22.11.2021	1.22	BQL	1.87	510
AL7 - 7	25.11.2021	1.08	BQL	1.99	541
AL7 - 8	29.11.2021	1.18	BQL	1.88	565
<b>Monthly Average</b>		<b>1.14</b>	<b>-</b>	<b>1.88</b>	<b>528</b>
<b>Standard Deviation</b>		<b>0.11</b>	<b>-</b>	<b>0.09</b>	<b>60</b>

\*NMHC- Non- Methane Hydrocarbons

BQL- Below Quantification Limit (Quantification Limit - NMHC: 0.5 ppm)

NS\_ Not Specified

The mean TSPM values at Vadinar Port were 219 µg/m<sup>3</sup>. The mean PM<sub>10</sub> values were 115 µg/m<sup>3</sup>, which is below the permissible limit. PM<sub>2.5</sub> values were also within the permissible limit (mean = 84 µg/m<sup>3</sup>). The average values of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were 4.0 µg/m<sup>3</sup>, 14.0 µg/m<sup>3</sup> and 7.0µg/m<sup>3</sup> respectively and were all within the permissible limit.

The levels of Benzene, Hydrocarbons (HC) and CO were within the permissible limit at Vadinar Port. The mean Benzene concentration was 1.14 µg/m<sup>3</sup>, well below the permissible limit of 5.0 µg/m<sup>3</sup>. HC's were below the detectable limit and Carbon Monoxide concentration was 1.88 mg/m<sup>3</sup>, well below the permissible limit of 4.0 mg/m<sup>3</sup>.

**Environmental Monitoring Report of Deendayal Port Trust,  
November-2021**

**Location 8: Admin Building (Vadinar) (AL-8)**

**Table 8 : Results of Air Pollutant Concentration at Admin Building**

Parameters	Date	TSPM [µg/m <sup>3</sup> ]	PM10 [µg/m <sup>3</sup> ]	PM2.5 [µg/m <sup>3</sup> ]	SO <sub>2</sub> [µg/m <sup>3</sup> ]		NO <sub>x</sub> [µg/m <sup>3</sup> ]		NH <sub>3</sub> [µg/m <sup>3</sup> ]	
					8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
Sampling Period	-	24hr	24hr	24hr	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
NAAQMS limit	-	NS	100 µg/m <sup>3</sup>	60 µg/m <sup>3</sup>	-	80 µg/m <sup>3</sup>	-	80 µg/m <sup>3</sup>	-	400 µg/m <sup>3</sup>
AL8 -1	01.11.21	204	83	73	4.84	3.22	20.96	18.63	6.38	5.96
					2.64		19.69		5.87	
					2.20		15.24		5.62	
AL8 -2	09.11.21	193	86	75	4.40	3.08	13.34	15.67	8.93	8.85
					2.64		20.33		6.64	
					2.20		13.34		10.98	
AL8 -3	12.11.21	241	126	107	2.64	2.49	15.88	16.09	4.85	5.96
					3.08		22.87		5.62	
					1.76		9.53		7.40	
AL8 -4	17.11.21	167	100	53	3.52	3.81	17.78	12.91	8.42	7.57
					4.84		10.16		10.47	
					3.08		10.80		3.83	
AL8 -5	19.11.21	183	151	72	2.20	2.49	10.16	10.59	5.36	6.55
					1.76		9.53		8.17	
					3.52		12.07		6.13	
AL8 -6	24.11.21	197	104	80	3.52	5.28	15.88	13.55	8.93	7.66
					5.71		10.16		7.91	
					6.59		14.61		6.13	
AL8 -5	26.11.21	226	111	88	3.52	3.37	10.16	11.43	11.74	8.85
					1.76		13.34		5.87	
					4.84		10.80		8.93	
AL8-6	29.11.21	226	104	106	2.64	2.78	10.16	13.13	9.19	7.40
					3.52		20.96		5.87	
					2.20		8.26		7.15	
<b>Monthly Average</b>		205	108	82		3		14		7
<b>Standard Deviation</b>		25	22	18		1		3		1

NS: Not Specified

**DCPL/DPT/20-21/19 -NOVEMBER - 2021**

**Detox Corporation Pvt. Ltd., Surat**

<b>Table 8B : Results of Air Pollutant Concentration at Admin Building</b>					
<b>Parameter</b>	<b>Date</b>	<b>C<sub>6</sub>H<sub>6</sub> [µg/m<sup>3</sup>]</b>	<b>HC*</b>	<b>CO [mg/m<sup>3</sup>]</b>	<b>CO<sub>2</sub> [ppm]</b>
<b>Sampling Period</b>		<b>8 hr</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>	<b>Grab Sampling</b>
<b>NAAQMS limit</b>		<b>5.0 µg/m<sup>3</sup></b>	<b>NS</b>	<b>4.0 mg/m<sup>3</sup></b>	<b>NS</b>
AL8 -1	01.11.21	1.23	BDL	1.88	571
AL8-2	09.11.21	1.09	BDL	2.04	581
AL8 -3	15.11.2021	1.29	BDL	2.17	465
AL8-4	18.11.2021	1.05	BDL	1.82	452
AL8 -5	19.11.2021	1.01	BDL	1.92	482
AL8-6	22.11.2021	1.28	BDL	1.73	496
AL8-7	25.11.2021	1.15	BDL	1.85	524
AL8-8	29.11.2021	1.13	BDL	2.02	561
Monthly Average		<b>1.15</b>	-	<b>1.93</b>	<b>517</b>
Standard Deviation		<b>0.10</b>	-	<b>0.14</b>	<b>50</b>

\* NMHC- Non- Methane Hydrocarbons

BQL- Below Quantification Limit (Quantification Limit - NMHC: 0.5 ppm)

NS-Not Specified

The overall values of TSPM, PM<sub>10</sub>, PM<sub>2.5</sub>, SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> at Admin Building Vadinar was comparatively low among all the locations of Air Quality monitoring in Kandla Port and Vadinar Port. The mean TSPM values at Vadinar Port were 205 µg/m<sup>3</sup>. The mean PM<sub>10</sub> values were 108 µg/m<sup>3</sup>, which is above the permissible limit. PM<sub>2.5</sub> values were above the permissible limit (mean = 82.0 µg/m<sup>3</sup>). The average values of SO<sub>2</sub>, NO<sub>x</sub> and NH<sub>3</sub> were 3.0 µg/m<sup>3</sup>, 14.0 µg/m<sup>3</sup> and 7.0 µg/m<sup>3</sup> respectively and were all within the permissible limit.

The levels of Benzene, Hydrocarbons (HC) and CO were within the permissible limit at Admin Building, Vadinar Port. The mean Benzene concentration was 1.15 µg/m<sup>3</sup>, well below the permissible limit of 5.0 µg/m<sup>3</sup>. HC's were below the detectable limit and Carbon Monoxide concentration was 1.93 mg/m<sup>3</sup>, well below the permissible limit of 4.0 mg/m<sup>3</sup>.

#### **1.4 Observations and Conclusion**

During the monitoring period, the overall Ambient Air Quality of the port area was found to be well within the desired levels for various gaseous pollutants. However, Particulate matter as PM<sub>10</sub> and PM<sub>2.5</sub> was found to exceed the limits at locations like Near Coal storage area, Marine Bhavan, Estate Office, Tuna Port and Oil Jetty area.

## **2. Drinking Water Quality Monitoring**

Drinking Water Quality Monitoring was carried out at twenty stations at Kandla, Vadinar & Township Area of Deendayal Port.

### **2.1 Drinking Water Monitoring Methodology**

Drinking water samples were collected from 20 locations as prescribed in the tender document. Samples for physico-chemical analysis were collected in 1 liter carboys and samples for microbiological parameters were collected in sterilized bottles. These samples were then analyzed in laboratory for various drinking water parameters at Kandla Lab/Surat.

The Sampling and Analysis was done as per standard methods - IS 10500:2012. The water samples were analyzed for various parameters, viz. Color , Odor, Turbidity , Conductivity , pH , Chlorides , TDS, Total Hardness, Iron , Sulphate , Salinity , DO, BOD, Na, K, Ca, Mg, F, NO<sub>3</sub>, NO<sub>2</sub>, Mn, Cr-6, Cu, Cd, As, Hg, Pb, Zn, Bacterial Count (cfu) .

### **2.2 Results**

The Drinking Water Quality monitoring data for 20 stations are given in below from table No. 9 to Table No. 15

**Table 9: Drinking Water Quality Monitoring Parameters for Nirman Building 1, P & C building & Main Gate (North) at Kandla**

Sr. No.	Parameter	Unit	Nirman Building 1	P & C building	Main Gate North	Acceptable Limits as per IS 10500 : 2012	Permissible Limits in the absence of Alternate Source as per IS 10500 : 2012
1	pH	pH Unit	7.3	7.4	7.2	6.5 to 8.5	6.5 to 8.5
2	Total Dissolved Solids	mg/l	990	1280	1310	500	2000
3	Turbidity	NTU	0	1	0	1.0	5.0
4	Odor	-	Odorless	Odorless	Odorless	Agreeable	Agreeable
5	Color	Hazen Units	Colorless	Colorless	Colorless	5.0	15.0
6	Conductivity	µs/cm	1860	2430	2540	NS*	NS*
7	Biochemical Oxygen Demand	mg/l	BQL	BQL	BQL	NS*	NS*
8	Chloride as Cl	mg/l	491.09	461.02	516.15	250.0	1000.0
9	Ca as Ca	mg/l	64.13	68.14	60.12	75.0	200.0
10	Mg as Mg	mg/l	63.18	70.47	68.04	30.0	100.0
11	Total Hardness	mg/l	420	460	430	200.0	600.0
12	Iron as Fe	mg/l	BQL	BQL	BQL	0.3	No Relaxation
13	Fluorides as F	mg/l	0.32	0.35	0.34	1.0	1.5
14	Sulphate as SO <sub>4</sub>	mg/l	286.8	289.2	283.2	200.0	400
15	Nitrite as NO <sub>2</sub>	mg/l	<0.1	<0.1	<0.1	NS*	NS*
16	Nitrate as NO <sub>3</sub>	mg/l	6.41	7.88	6.20	45.0	No Relaxation
17	Salinity	%	0.86	0.80	0.89	NS*	NS*
18	Sodium as Na	mg/l	202	225	277	NS*	NS*
19	Potassium as K	mg/l	5.08	3.26	5.26	NS*	NS*
20	Manganese	mg/l	BQL	BQL	BQL	0.1	0.3
21	Hexavalent Chromium	mg/l	BQL	BQL	BQL	NS*	NS*
22	Copper	mg/l	BQL	BQL	BQL	0.05	1.5
23	Cadmium	mg/l	BQL	BQL	BQL	0.003	0.003
24	Arsenic	mg/l	BQL	BQL	BQL	0.01	0.05
25	Mercury	mg/l	BQL	BQL	BQL	0.001	0.001
26	Lead	mg/l	BQL	BQL	BQL	0.01	0.01
27	Zinc	mg/l	BQL	BQL	BQL	5.0	15.0
28	Bacterial Count	CFU/100 ml	Absent	Absent	Absent	Absent	Absent

\*NS: Not Specified

BQL- Below Quantification Limit, (BOD-2.0 mg/l, Fe-0.009 mg/l, Mn- 0.01 mg/l, Cr<sup>+6</sup>- 0.03 mg/l, Cu-0.004 mg/l, Cd-0.003 mg/l, As-0.003mg/l, Hg-0.001 mg/l, Pb-0.006mg/l, Zinc-0.021 mg/l).

**Environmental Monitoring Report of Deendayal Port Trust,  
November-2021**

**Table 10: Drinking Water Quality Monitoring Parameters for Canteen, West Gate - I & Wharf Area at Kandla**

Sr. No	Parameter	Unit	Canteen	West Gate - I	Wharf Area	Acceptable Limits as per IS 10500 : 2012	Permissible Limits in the absence of Alternate Source as per IS 10500 : 2012
1	pH	pH Unit	7.5	7.3	7.2	6.5 to 8.5	6.5 to 8.5
2	Total Dissolved Solids	mg/l	1410	1350	1420	500	2000
3	Turbidity	NTU	0	0	1	1.0	5.0
4	Odor	-	Odorless	Odorless	Odorless	Agreeable	Agreeable
5	Color	Hazen Units	Colorless	Colorless	Colorless	5.0	15.0
6	Conductivity	µs/cm	2710	2560	2730	NS*	NS*
7	Biochemical Oxygen Demand	mg/l	BQL	BQL	BQL	NS*	NS*
8	Chloride as Cl	mg/l	541.20	486.08	491.09	250.0	1000.0
9	Ca as Ca	mg/l	64.13	56.11	64.13	75.0	200.0
10	Mg as Mg	mg/l	68.04	68.04	72.90	30.0	100.0
11	Total Hardness	mg/l	440	420	460	200.0	600.0
12	Iron as Fe	mg/l	BQL	BQL	BQL	0.3	No Relaxation
13	Fluorides as F	mg/l	0.41	0.29	0.34	1.0	1.5
14	Sulphate as SO <sub>4</sub>	mg/l	291.6	204.0	194.4	200.0	400
15	Nitrite as NO <sub>2</sub>	mg/l	<0.1	<0.1	<0.1	NS*	NS*
16	Nitrate as NO <sub>3</sub>	mg/l	8.10	12.25	8.87	45.0	No Relaxation
17	Salinity	%	0.83	0.93	0.98	NS*	NS*
18	Sodium as Na	mg/l	201	195	279	NS*	NS*
19	Potassium as K	mg/l	4.28	4.08	4.69	NS*	NS*
20	Manganese	mg/l	BQL	BQL	BQL	0.1	0.3
21	Hexavalent Chromium	mg/l	BQL	BQL	BQL	NS*	NS*
22	Copper	mg/l	BQL	BQL	BQL	0.05	1.5
23	Cadmium	mg/l	BQL	BQL	BQL	0.003	0.003
24	Arsenic	mg/l	BQL	BQL	BQL	0.01	0.05
25	Mercury	mg/l	BQL	BQL	BQL	0.001	0.001
26	Lead	mg/l	BQL	BQL	BQL	0.01	0.01
27	Zinc	mg/l	BQL	BQL	BQL	5.0	15.0
28	Bacterial Count	CFU/100 ml	Absent	Absent	Absent	Absent	Absent

\*NS: Not Specified

BQL- Below Quantification Limit, (BOD-2.0 mg/l, Fe-0.009 mg/l, Mn- 0.01 mg/l, Cr<sup>+6</sup>- 0.03 mg/l, Cu- 0.004 mg/l, Cd-0.003 mg/l, As-0.003mg/l, Hg-0.001 mg/l, Pb-0.006mg/l, Zinc-0.021 mg/l).



**Environmental Monitoring Report of Deendayal Port Trust,  
November-2021**

**Table 11: Drinking Water Quality Monitoring Parameters for Sewa sadan - 3, Workshop I & Custom Building at Kandla**

Sr. No	Parameter	Unit	Sewa Sadan - 3	Workshop	Custom Building	Acceptable Limits as per IS 10500 : 2012	Permissible Limits in the absence of Alternate Source as per IS 10500 : 2012
1	pH	pH Unit	7.4	7.7	7.5	6.5 to 8.5	6.5 to 8.5
2	Total Dissolved Solids	mg/l	1360	1325	1430	500	2000
3	Turbidity	NTU	1	1	0	1.0	5.0
4	Odor	-	Odorless	Odorless	Odorless	Agreeable	Agreeable
5	Color	Hazen Units	Colorless	Colorless	Colorless	5.0	15.0
6	Conductivity	µs/cm	2530	2480	2680	NS*	NS*
7	Biochemical Oxygen Demand	mg/l	BQL	BQL	BQL	NS*	NS*
8	Chloride	mg/l	656.46	611.36	516.15	250.0	1000.0
9	Ca as Ca	mg/l	56.11	60.12	64.13	75.0	200.0
10	Mg as Mg	mg/l	75.33	65.61	72.90	30.0	100.0
11	Total Hardness	mg/l	450	420	460	200.0	600.0
12	Iron as Fe	mg/l	BQL	BQL	BQL	0.3	No Relaxation
13	Fluorides	mg/l	0.23	0.45	0.46	1.0	1.5
14	Sulphate	mg/l	198.0	290.4	230.4	200.0	400
15	Nitrite	mg/l	<0.1	<0.1	<0.1	NS*	NS*
16	Nitrate	mg/l	10.42	9.57	13.94	45.0	No Relaxation
17	Salinity	%	0.88	0.89	1.19	NS*	NS*
18	Sodium as Na	mg/l	303	248	327	NS*	NS*
19	Potassium as K	mg/l	4.30	5.61	8.26	NS*	NS*
20	Manganese	mg/l	BQL	BQL	BQL	0.1	0.3
21	Hexavalent Chromium	mg/l	BQL	BQL	BQL	NS*	NS*
22	Copper	mg/l	BQL	BQL	BQL	0.05	1.5
23	Cadmium	mg/l	BQL	BQL	BQL	0.003	0.003
24	Arsenic	mg/l	BQL	BQL	BQL	0.01	0.05
25	Mercury	mg/l	BQL	BQL	BQL	0.001	0.001
26	Lead	mg/l	BQL	BQL	BQL	0.01	0.01
27	Zinc	mg/l	BQL	BQL	BQL	5.0	15.0
28	Bacterial Count	CFU/100ml	Absent	Absent	Absent	Absent	Absent

\*NS: Not Specified

BQL- Below Quantification Limit, (BOD-2.0 mg/l, Fe-0.009 mg/l, Mn- 0.01 mg/l, Cr<sup>+6</sup>- 0.03 mg/l, Cu-0.004 mg/l, Cd-0.003 mg/l, As-0.003mg/l, Hg-0.001 mg/l, Pb-0.006mg/l, Zinc-0.021 mg/l).

**Table 12: Drinking Water Quality Monitoring Parameters for Port Colony Kandla, Hospital Kandla & A.O. Building at Gandhidham**

Sr No	Parameter	Unit	Port Colony Kandla	Hospital Kandla	A.O. Building	Acceptable Limits as per IS 10500 : 2012	Permissible Limits in the absence of Alternate Source as per IS 10500 : 2012
1	pH	pH Unit	7.6	7.3	7.2	6.5 to 8.5	6.5 to 8.5
2	Total Dissolved Solids	mg/l	1310	1410	1430	500	2000
3	Turbidity	NTU	0	0	1	1.0	5.0
4	Odor	-	Odorless	Odorless	Odorless	Agreeable	Agreeable
5	Color	Hazen Units	Colorless	Colorless	Colorless	5.0	15.0
6	Conductivity	µs/cm	2540	2690	2740	NS*	NS*
7	Biochemical Oxygen Demand	mg/l	<0.1	<0.1	<0.1	NS*	NS*
8	Chloride	mg/l	481.07	531.18	516.15	250.0	1000.0
9	Ca as Ca	mg/l	72.14	76.15	64.13	75.0	200.0
10	Mg as Mg	mg/l	55.89	58.32	68.04	30.0	100.0
11	Total Hardness	mg/l	410	430	440	200.0	600.0
12	Iron as Fe	mg/l	BQL	BQL	BQL	0.3	No Relaxation
13	Fluorides	mg/l	0.48	0.85	0.52	1.0	1.5
14	Sulphate	mg/l	210.0	291.6	301.2	200.0	400
15	Nitrite	mg/l	<0.1	<0.1	<0.1	NS*	NS*
16	Nitrate	mg/l	10.28	13.24	9.79	45.0	No Relaxation
17	Salinity	%	1.10	0.93	0.87	NS*	NS*
18	Sodium as Na	mg/l	154	384	218	NS*	NS*
19	Potassium as K	mg/l	3.26	4.69	4.03	NS*	NS*
20	Manganese	mg/l	BQL	BQL	BQL	0.1	0.3
21	Hexavalent Chromium	mg/l	BQL	BQL	BQL	NS*	NS*
22	Copper	mg/l	BQL	BQL	BQL	0.05	1.5
23	Cadmium	mg/l	BQL	BQL	BQL	0.003	0.003
24	Arsenic	mg/l	BQL	BQL	BQL	0.01	0.05
25	Mercury	mg/l	BQL	BQL	BQL	0.001	0.001
26	Lead	mg/l	BQL	BQL	BQL	0.01	0.01
27	Zinc	mg/l	BQL	BQL	BQL	5.0	15.0
28	Bacterial Count	CFU/100 ml	Absent	Absent	Absent	Absent	Absent

\*NS: Not Specified

BQL- Below Quantification Limit, (BOD-2.0 mg/l, Fe-0.009 mg/l, Mn- 0.01 mg/l, Cr<sup>+6</sup>- 0.03 mg/l, Cu-0.004 mg/l, Cd-0.003 mg/l, As-0.003mg/l, Hg-0.001 mg/l, Pb-0.006mg/l, Zinc-0.021 mg/l).

**Table 13: Drinking Water Quality Monitoring Parameters for School Gopalpuri, Guest House & E - Type Quarter at Gopalpuri, Gandhidham**

Sr No	Parameter	Unit	School Gopalpuri	Guest House	E - Type Quarter	Acceptable Limits as per IS 10500 : 2012	Permissible Limits in the absence of Alternate Source as per IS 10500 : 2012
1	pH	pH Unit	7.4	7.6	7.5	6.5 to 8.5	6.5 to 8.5
2	Total Dissolved Solids	mg/l	1400	1720	1090	500	2000
3	Turbidity	NTU	1	0	1	1.0	5.0
4	Odor	-	Odorless	Odorless	Odorless	Agreeable	Agreeable
5	Color	Hazen Units	Colorless	Colorless	Colorless	5.0	15.0
6	Conductivity	µs/cm	2640	2730	2130	NS*	NS*
7	Biochemical Oxygen Demand	mg/l	BQL	BQL	BQL	NS*	NS*
8	Chloride	mg/l	496.10	526.17	481.07	250.0	1000.0
9	Ca as Ca	mg/l	60.12	56.11	68.14	75.0	200.0
10	Mg as Mg	mg/l	80.19	77.76	65.61	30.0	100.0
11	Total Hardness	mg/l	480	460	440	200.0	600.0
12	Iron as Fe	mg/l	BQL	BQL	BQL	0.3	No Relaxation
13	Fluorides	mg/l	0.60	0.81	0.63	1.0	1.5
14	Sulphate	mg/l	314.4	214.8	289.2	200.0	400
15	Nitrite	mg/l	<0.1	<0.1	<0.1	NS*	NS*
16	Nitrate	mg/l	10.91	12.39	10.00	45.0	No Relaxation
17	Salinity	%	0.96	0.93	0.90	NS*	NS*
18	Sodium as Na	mg/l	287	106	246	NS*	NS*
19	Potassium as K	mg/l	5.28	6.29	2.25	NS*	NS*
20	Manganese	mg/l	BQL	BQL	BQL	0.1	0.3
21	Hexavalent Chromium	mg/l	BQL	BQL	BQL	NS*	NS*
22	Copper	mg/l	BQL	BQL	BQL	0.05	1.5
23	Cadmium	mg/l	BQL	BQL	BQL	0.003	0.003
24	Arsenic	mg/l	BQL	BQL	BQL	0.01	0.05
25	Mercury	mg/l	BQL	BQL	BQL	0.001	0.001
26	Lead	mg/l	BQL	BQL	BQL	0.01	0.01
27	Zinc	mg/l	BQL	BQL	BQL	5.0	15.0
28	Bacterial Count	CFU/100 ml	Absent	Absent	Absent	Absent	Absent

\*NS: Not Specified

BQL- Below Quantification Limit, (BOD-2.0 mg/l, Fe-0.009 mg/l, Mn- 0.01 mg/l, Cr<sup>+6</sup>- 0.03 mg/l, Cu- 0.004 mg/l, Cd-0.003 mg/l, As-0.003mg/l, Hg-0.001 mg/l, Pb-0.006mg/l, Zinc-0.021 mg/l).

**Table 14: Drinking Water Quality Monitoring Parameters for F - Type  
Quarter, Hospital Gopalpuri & Tuna Port**

Sr. No	Parameter	Unit	F - Type Quarter	Hospital Gopalpuri	Tuna Port	Acceptable Limits as per IS 10500 : 2012	Permissible Limits in the absence of Alternate Source as per IS 10500 : 2012
1	pH	pH Unit	7.6	7.3	7.42	6.5 to 8.5	6.5 to 8.5
2	Total Dissolved Solids	mg/l	1020	1250	1150	500	2000
3	Turbidity	NTU	0	1	0	1.0	5.0
4	Odor	-	Odorless	Odorless	Odorless	Agreeable	Agreeable
5	Color	Hazen Unit	Colorless	Colorless	Colorless	5.0	15.0
6	Conductivity	µs/cm	1950	2380	2000	NS*	NS*
7	Biochemical Oxygen Demand	mg/l	BQL	BQL	BQL	NS*	NS*
8	Chloride	mg/l	611.36	576.28	520	250.0	1000.0
9	Ca as Ca	mg/l	64.13	60.12	76.15	75.0	200.0
10	Mg as Mg	mg/l	55.89	72.90	55.89	30.0	100.0
11	Total Hardness	mg/l	390	450	420	200.0	600.0
12	Iron as Fe+3	mg/l	BQL	BQL	BQL	0.3	No Relaxation
13	Fluorides	mg/l	0.86	0.57	0.75	1.0	1.5
14	Sulphate	mg/l	301.2	285.6	274.8	200.0	400
15	Nitrite	mg/l	<0.1	<0.1	<0.1	NS*	NS*
16	Nitrate	mg/l	10.91	10.07	9.93	45.0	No Relaxation
17	Salinity	%	0.95	0.87	1.10	NS*	NS*
18	Sodium as Na	mg/l	235	235	248	NS*	NS*
19	Potassium as K	mg/l	3.98	5.54	4.8	NS*	NS*
20	Manganese	mg/l	BQL	BQL	BQL	0.1	0.3
21	Hexavalent Chromium	mg/l	BQL	BQL	BQL	NS*	NS*
22	Copper	mg/l	BQL	BQL	BQL	0.05	1.5
23	Cadmium	mg/l	BQL	BQL	BQL	0.003	0.003
24	Arsenic	mg/l	BQL	BQL	BQL	0.01	0.05
25	Mercury	mg/l	BQL	BQL	BQL	0.001	0.001
26	Lead	mg/l	BQL	BQL	BQL	0.01	0.01
27	Zinc	mg/l	BQL	BQL	BQL	5.0	15.0
28	Bacterial Count	CFU/100 ml	Absent	Absent	Absent	Absent	Absent

\*NS: Not Specified

BQL- Below Quantification Limit, (BOD-2.0 mg/l, Fe-0.009 mg/l, Mn- 0.01 mg/l, Cr<sup>+6</sup>- 0.03 mg/l, Cu- 0.004 mg/l, Cd-0.003 mg/l, As-0.003mg/l, Hg-0.001 mg/l, Pb-0.006mg/l, Zinc-0.021 mg/l).

**Table 15: Drinking Water Quality Monitoring Parameters for Vadinar Jetty & Port Colony at Vadinar**

Sr. No	Parameter	Unit	Vadinar Jetty	Port Colony Vadinar	Acceptable Limits as per IS 10500 : 2012	Permissible Limits in the absence of Alternate Source as per IS 10500 : 2012
1	pH	pH Unit	7.5	7.3	6.5 to 8.5	6.5 to 8.5
2	Total Dissolved Solids	mg/l	1060	1120	500	2000
3	Turbidity	NTU	ND	ND	1.0	5.0
4	Odor	-	Odorless	Odorless	Agreeable	Agreeable
5	Color	Hazen Units	Colorless	Colorless	5.0	15.0
6	Conductivity	µs/cm	1960	2150	NS*	NS*
7	Biochemical Oxygen Demand	mg/l	BQL	BQL	NS*	NS*
8	Chloride	mg/l	486.08	521.16	250.0	1000.0
9	Ca as Ca	mg/l	68.14	60.12	75.0	200.0
10	Mg as Mg	mg/l	68.04	68.04	30.0	100.0
11	Total Hardness	mg/l	450	430	200.0	600.0
12	Iron as Fe+3	mg/l	BQL	BQL	0.3	No Relaxation
13	Fluorides	mg/l	0.86	0.69	1.0	1.5
14	Sulphate	mg/l	23.04	22.56	200.0	400
15	Nitrite	mg/l	<0.1	<0.1	NS*	NS*
16	Nitrate	mg/l	7.88	10.63	45.0	No Relaxation
17	Salinity	%	0.88	0.94	NS*	NS*
18	Sodium as Na	mg/l	52.8	40.2	NS*	NS*
19	Potassium as K	mg/l	3.3	2.1	NS*	NS*
20	Manganese	mg/l	BQL	BQL	0.1	0.3
21	Hexavalent Chromium	mg/l	BQL	BQL	NS*	NS*
22	Copper	mg/l	BQL	BQL	0.05	1.5
23	Cadmium	mg/l	BQL	BQL	0.003	0.003
24	Arsenic	mg/l	BQL	BQL	0.01	0.05
25	Mercury	mg/l	BQL	BQL	0.001	0.001
26	Lead	mg/l	BQL	BQL	0.01	0.01
27	Zinc	mg/l	BQL	BQL	5.0	15.0
28	Bacterial Count	CFU/10 Oml	Absent	Absent	Absent	Absent

\*NS: Not Specified

BQL- Below Quantification Limit, (BOD-2.0 mg/l, Fe-0.009 mg/l, Mn- 0.01 mg/l, Cr<sup>+6</sup>- 0.03 mg/l, Cu-0.004 mg/l, Cd-0.003 mg/l, As-0.003mg/l, Hg-0.001 mg/l, Pb-0.006mg/l, Zinc-0.021 mg/l).

### **2.3 Results & Discussion**

The colour of all drinking water samples was < 5 Hazen unit and odour of the samples was also agreeable. All parameters are found to be within the specified limit of the Drinking water Standard.

#### **pH**

The limit of pH value for drinking water is specified as 6.5 to 8.5. pH value in the studied area varied from 7.1 to 7.7 pH unit. All the sampling points showed pH values within the prescribed limit by Indian Standards.

#### **Total Dissolved Solids (TDS)**

TDS values in the studied area varied between 900 -1800 mg/l. None of the sampling points showed higher TDS values than the prescribed limit by Indian standards.

#### **Conductivity**

Electrical Conductivity is the ability of a solution to transfer (conduct) electric current. Conductivity is used to measure the concentration of dissolved solids which have been ionized in a polar solution such as water. The conductivity in the samples collected during the month of November ranged from 1800-3000  $\mu\text{s}/\text{cm}$ . Electrical conductivity standards do not appear in BIS standards for drinking water.

#### **BOD**

BOD value in the studied area was found Below Quantification Limit (2.0 mg/l). Indian standards does not show any standard values for BOD in drinking water.

#### **Chlorides**

Excessive chloride concentration increase rates of corrosion of metals in the distribution system. This can lead to increased concentration of metals in the supply. Chloride value in the studied area varied between 400-700 mg/l and is found to be within the Permissible limit of the Drinking Water Standard.

#### **Calcium**

Calcium value in the studied area varied between 50 - 80 mg/l and is found to be within the Permissible limit of the Drinking Water Standard. If calcium is present beyond the maximum acceptable limit, it causes incrustation of pipes.

### **Magnesium**

Magnesium value in the studied area varied between 50-85 mg/l. All the locations had Magnesium within the prescribed limits of 30-100 mg/L.

### **Total Hardness**

Hardness value in the studied area varied between 350-480 mg/l and is found to be within the Permissible limit of the Drinking Water Standard. The prescribed limit by Indian Standards is 200-600 mg/L.

### **Iron**

Iron value in the studied area was found Below Quantification Limit (0.009 mg/l) and hence well below the permissible limit as per Indian Standards is 0.3 mg/L. The excess amount of iron causes slight toxicity; gives stringent taste to water.

### **Fluoride**

Fluoride value in the studied area varied between 0.1 – 1.0 mg/l and hence well below the permissible limit as per Indian Standards is 1.0-1.5 mg/L. Moderate amounts lead to dental effects, but long-term ingestion of large amounts can lead to potentially severe skeletal problems.

### **Sulphates**

Sulphate value in the studied area varied between 20 – 350 mg/l. All the sampling points showed sulphate values within the prescribed limits by Indian Standards (200-400 mg/L). Sulphate content in drinking water exceeding the 400 mg/L imparts bitter taste.

### **Nitrites (NO<sub>2</sub>) and Nitrates (NO<sub>3</sub>)**

Nitrite values in all the water samples were found Below Quantification Limit (0.1 mg/l). There are no specified standard values for Nitrites in Drinking water. The minimum Nitrate value in drinking water of KPT was 6.20 mg/l which is well within the permissible limit of the Drinking water Standard.

### **Salinity**

Salinity in drinking water in the present samples collected ranged from 0.5 to 1.4 %. There are no prescribed Indian standards for salinity in Drinking water.

### **Sodium and Potassium Salts**

Sodium values in the samples collected ranged from 40 - 400 mg/l and Potassium salts ranged from 2.0 to 8.5 mg/l. There are no prescribed limits of Sodium and Potassium in Indian standards for Drinking water.

### **Heavy Metals in Drinking Water**

In the present study period drinking water samples were analyzed for Mn, Cr, Cu, Cd, As, Hg, Pb and Zn. All these heavy metals were well Below the Quantification limits prescribed by the Indian Standards.

### **Bacteriological Study**

Analysis of the bacteriological parameter at all location shows that Bacteria is not present and hence Bacterial count is in line with the permissible limit of drinking water. This shows that all the drinking water samples were safe from any bacteriological contamination.

## **2.4 Conclusions**

These results are compared with acceptable limits as prescribed in IS 10500:2012 - Drinking Water Specification. It is seen from the analysis data that during the study period the water was safe for human consumption at all drinking water monitoring stations.

## **3. Noise Level Monitoring**

Noise sources in port operations include cargo handling, vehicular traffic, and loading / unloading containers and ships. Noise Monitoring was done at 13 stations at Kandla, Vadinar and Township area.

### **3.1 Method of Monitoring**

Sampling was done at all stations for 24 hour period. Data was recorded using automated sound level meter. The intensity of sound was measured in sound pressure level (SPL) and common unit of measurement is decibel (dB).

### **3.2 Results**

Table 16: Noise Monitoring data for ten locations of Deendayal Port and two locations of Vadinar Port

<b>Sr. No.</b>	<b>Location</b>	<b>Day Time Average Noise Level (SPL) in dB(A)</b>	<b>Night Time Average Noise Level (SPL) in dB(A)</b>
	<b>Sampling Time</b>	<b>6:00 am to 10:00 PM</b>	<b>10:00PM to 6:00 AM</b>
1	Marine Bhavan	73.9	53.4
2	Nirman Building 1	62.3	51.0
3	Tuna Port	57.2	50.9
4	Main Gate North	67.0	61.8
5	West Gate I	70.5	65.1

**DCPL/DPT/20-21/19 -NOVEMBER - 2021**

---

**Detox Corporation Pvt. Ltd., Surat**



6	Canteen Area	64.5	51.0
7	Main Road	68.4	51.5
8	ATM Building	74.4	57.3
9	Wharf Area /Jetty Area	72.9	68.1
10	Port & Custom Office	67.8	41.8
<b>Vadinar Port</b>			
11	Entrance Gate of Vadinar Port	66.4	53.2
12	Nr. Port Colony, Vadinar	60.7	54.1
13	Nr. Vadinar Jetty	72.4	66.5

**3.3 Conclusions-** Noise sources in port operations include cargo handling, vehicular traffic, and loading / unloading containers and ships. The Day Time Average Noise Level (SPL) in all 13 locations at Deendayal Port ranged from 57.2 dB(A) to 74.4 dB(A) and it was within the permissible limits of 75 dB(A) for the industrial area for the daytime. The Night Time Average Noise Level (SPL) in all 13 locations of Deendayal Port ranged from 41.8 dB to 68.4 dB(A) and it was within the permissible limits of 70 dB(A) for the industrial area for the night time.

#### **4. Soil Monitoring**

Sampling and analysis of soil samples were undertaken at six locations within the study area (Deendayal Port and Vadinar Port) as a part of EMP. The soil sampling locations are initially decided based on the locations as provided in the tender document of the Deendayal Port.

##### **4.1 Methodology**

The soil samples were collected in the month of November 2021. The samples collected from the all locations are homogeneous representative of each location. At random locations were identified at each location and soil was dug from 30 cm below the surface. It was uniformly mixed before homogenizing the soil samples. The samples were filled in polythene bags, labeled in the field with number and site name and sent to laboratory for analysis.

## 4.2 Results

**Table-17: Chemical Characteristics of Soil in the Study Area**

Sr. No	Parameter	Unit	Station Name					
			SL1	SL2	SL3	SL4	SL5	SL6
			Tuna Port	IFFCO Plant	Khori Creek	Nakti Creek	KPT Admin Site	KPT Colony
			Near main gate of Port	10 m away from main gate	Sand from creek at low tide		Vadinar	
1	Texture		Sandy Loam	Sandy Loam	Sandy Loam	Sandy Loam	Sandy Loam	Sandy Loam
2	pH	-	8.42	7.92	8.44	8.23	7.79	8.43
3	Electrical Conductivity	µs/cm	14,070.0	16,210.0	13,680.0	9,240.0	387.0	314.0
4	Moisture	%	18.17	9.01	21.39	21.08	3.46	3.95
5	Total Organic Carbon	%	0.20	0.49	0.20	0.72	0.85	0.43
6	Alkalinity	mg/kg	80.08	120.12	60.06	100.10	60.06	80.08
7	Total Nitrogen	%	BQL	BQL	BQL	BQL	BQL	BQL
8	Chloride	mg/kg	1,956.8	4,112.2	1,800.9	514.7	21.7	113.4
9	Sulphate	mg/kg	212.0	279.0	93.3	165.1	44.7	27.7
10	Phosphorus	mg/kg	2.20	1.89	1.41	2.15	BQL	1.74
11	Potassium	mg/kg	539.0	327.4	409.2	667.6	70.4	62.0
12	Sodium	mg/kg	5,752.0	4,061.6	3,954.0	1,477.0	72.8	65.9
13	Calcium	mg/kg	200.40	488.98	252.00	470.42	436.87	256.51
14	Copper as Cu	mg/kg	14.90	29.50	9.80	27.60	88.4	48.4
15	Lead as Pb	mg/kg	5.80	6.40	3.50	8.20	BQL	4.2
16	Nickel as Ni	mg/kg	35.30	16.60	23.50	37.70	33.8	27.3
17	Zinc as Zn	mg/kg	40.60	104.80	25.4	55.20	66.00	30.50
18	Cadmium as Cd	mg/kg	BQL	BQL	BQL	BQL	BQL	BQL

BQL- Below Quantification Limit, ( TN:0.001%, Cd: 1.0mg/kg).

**DCPL/DPT/20-21/19 -NOVEMBER - 2021**

**Detox Corporation Pvt. Ltd., Surat**

### **4.3 Discussion**

The data shows that value of pH ranges from 7.92 at IFFCO Plant to 8.44 at Khori Creek indicating that all soil samples are neutral to slight basic. Iffco Plant samples showed maximum conductivity of 16,210.0 µmhos/cm, while Tuna Port location showed minimum conductivity of 14,070.0 µmhos/cm. Conductivity at Vadinar Port was 387 and 314 µmhos/cm at Admin site and Vadinar Port colony respectively.

Total organic Carbon ranged from 0.2 % to 0.72 at Deendayal Port. At Vadinar Port, organic carbon content ranged from 0.85 % to 0.43 %.

The concentration of Phosphorus and Potassium in the soil samples varies from 1.41 to 2.15 mg/kg and 327.0 to 670.0 mg/kg respectively at Deendayal Port. The mean concentration of Phosphorous at Vadinar site was 1.74 mg/kg and mean concentration of Potassium at Vadinar site was 132 mg/kg.

These differences in NPK in soil at different locations are due to the dissimilar nature of soil at each of the locations. Samples SL3 & SL4 (Khori Creek & Nakti Creek) are of saline nature as they are coastal soil; where as other locations are inland locations and have different chemical properties.

### **Heavy Metals in the Soil**

Traces of Copper, Lead, Nickel and Zinc were observed in the soil samples collected from all the four locations of Deendayal Port and two locations of Vadinar Port. Cadmium metal was below detection limit in the Soil.

### **4.4 Conclusion**

The soils of Deendayal Port and Vadinar Port appears to be neutral to basic with varying levels of Chloride, Sulphate, NPK and Calcium. As the nature of soil at different locations are different with respect to its proximity to the sea, the samples showed high degree of variations in their chemical properties.

## **5. Sewage Treatment Plant Monitoring**

This involves safe collection of waste water (spent/used water) from wash areas, bathroom, industrial units, etc., waste from toilets of various buildings and its conveyance to the treatment plant and final disposal in conformity with the requirement and guide lines of State Pollution Control Board and other statutory bodies.

### **5.1 Methodology for STP Monitoring**

To monitor the working efficiency of Sewage Treatment Plant (STP), STP Inlet and Outlet Samples were collected once a week. Locations selected are namely Gopalpuri Township, Deendayal Port and Vadinar. Samples were collected in 1 lit. Carboys and were analyzed in laboratory for various parameters.

## 5.2 Results

### Kandla STP

**Table 18: Sewage Water Monitoring at Kandla STP (1<sup>st</sup> Week)**

<b>Date of Sampling</b>	<b>02.11.2021</b>
-------------------------	-------------------

Sr. No.	Parameters	Unit	Results	
			KPT STP I/L	KPT STP O/L
1	pH	pH unit	7.82	7.46
2	Total Suspended Solids	mg/l	206	116.1
3	Residual Chlorine	mg/l	-	<0.5
4	COD	mg/l	393.0	152.0
5	BOD @ 27 °C	mg/l	110.0	53.0
6.	Fecal Coliform	MPN Index / 100 ml	-	>1600.0
<b>Aeration Tank</b>				
7.	MLSS	mg/l	35.0	
8.	MLVSS	%	6.0	

**Table 19: Sewage Water Monitoring at Kandla STP (2<sup>nd</sup> Week)**

<b>Date of Sampling</b>	11.11.2021
-------------------------	------------

Sr. No.	Parameters	Unit	Results	
			KPT STP I/L	KPT STP O/L
1	pH	pH unit	7.6	7.2
2	Total Suspended Solids	mg/l	152.2	72.4
3	Residual Chlorine	mg/l	-	<0.5
4	COD	mg/l	384	103.0
5	BOD @ 27 °C	mg/l	120.0	24.0
6.	Fecal Coliform	MPN Index / 100 ml	-	9.2
<b>Aeration Tank</b>				
7.	MLSS	mg/l	7.0	
8.	MLVSS	%	90.0	

**Table 20: Sewage Water Monitoring at Kandla STP (3<sup>rd</sup> Week)**

<b>Date of Sampling</b>	17.11.2021
-------------------------	------------

Sr. No.	Parameters	Unit	Results	
			KPT STP I/L	KPT STP O/L
1	pH	pH unit	7.35	7.14
2	Total Suspended Solids	mg/l	204	144
3	Residual Chlorine	mg/l	-	<0.5
4	COD	mg/l	162	71
5	BOD @ 27 °C	mg/l	60.0	20.0
6.	Fecal Coliform	MPN Index / 100 ml	-	24.0
<b>Aeration Tank</b>				
7.	MLSS	mg/l	12.0	
8.	MLVSS	%	93.0	

**Table 21: Sewage Water Monitoring at Kandla STP (4<sup>th</sup> Week)**

<b>Date of Sampling</b>	22.11.2021
-------------------------	------------

Sr. No.	Parameters	Unit	Results	
			KPT STP I/L	KPT STP O/L
1	pH	pH unit	7.82	7.46
2	Total Suspended Solids	mg/l	306	116
3	Residual Chlorine	mg/l	-	<0.5
4	COD	mg/l	242	103.0
5	BOD @ 27 °C	mg/l	86.0	26.0
6.	Fecal Coliform	MPN Index / 100 ml	-	170.0
<b>Aeration Tank</b>				
7.	MLSS	mg/l	9.0	
8	MLVSS	%	98.0	

**Gopalpuri Colony STP**

**Table 22: Sewage Water Monitoring at Gopalpuri STP (1<sup>st</sup> Week)**

<b>Date of Sampling</b>	02.11.2021
-------------------------	------------

Sr. No.	Parameters	Unit	Results	
			Gopalpuri STP I/L	Gopalpuri STP O/L
1	pH	pH unit	7.50	7.20
2	Total Suspended Solids	mg/l	210	120
3	Residual Chlorine	mg/l	-	<0.5
4	COD	mg/l	414.0	142.0
5	BOD @ 27 °C	mg/l	122.0	53.0
6.	Fecal Coliform	MPN Index / 100 ml	-	>1600.0
<b>Aeration Tank</b>				
7.	MLSS	mg/l	86.0	
8	MLVSS	%	97.0	

**Table 23: Sewage Water Monitoring at Gopalpuri STP (2<sup>nd</sup> Week)**

<b>Date of Sampling</b>	11.11.2021
-------------------------	------------

Sr. No.	Parameters	Unit	Results	
			Gopalpuri STP I/L	Gopalpuri STP O/L
1	pH	pH unit	7.46	7.2
2	Total Suspended Solids	mg/l	379.2	118
3	Residual Chlorine	mg/l	-	<0.5
4	COD	mg/l	454	163.00
5	BOD @ 27 °C	mg/l	148.0	56.0
6.	Fecal Coliform	MPN Index / 100 ml	-	21.0
<b>Aeration Tank</b>				
7.	MLSS	mg/l		94.0
8	MLVSS	%		92.0

**Table 24: Sewage Water Monitoring at Gopalpuri STP (3<sup>rd</sup> Week)**

<b>Date of Sampling</b>	17.11.2021
-------------------------	------------

Sr. No.	Parameters	Unit	Results	
			Gopalpuri STP I/L	Gopalpuri STP O/L
1	pH	pH unit	7.35	7.14
2	Total Suspended Solids	mg/l	204	144
3	Residual Chlorine	mg/l	-	<0.5
4	COD	mg/l	162	71
5	BOD @ 27 °C	mg/l	60.0	20.0
6.	Fecal Coliform	MPN Index / 100 ml	-	24.0
<b>Aeration Tank</b>				
7.	MLSS	mg/l		12.0
8	MLVSS	%		93.0

**Table 25: Sewage Water Monitoring at Gopalpuri STP (4<sup>th</sup> Week)**

**DCPL/DPT/20-21/19 -NOVEMBER - 2021**

**Environmental Monitoring Report of Deendayal Port Trust,  
November-2021**

<b>Date of Sampling</b>	22.11.2021
-------------------------	------------

Sr. No.	Parameters	Unit	Results	
			Gopalpuri STP I/L	Gopalpuri STP O/L
1	pH	pH unit	7.82	7.46
2	Total Suspended Solids	mg/l	306	116
3	Residual Chlorine	mg/l	-	<0.5
4	COD	mg/l	242	103.0
5	BOD @ 27 °C	mg/l	86.0	26.0
6.	Fecal Coliform	MPN Index / 100 ml	-	170.0
<b>Aeration Tank</b>				
7.	MLSS	mg/l	9.0	
8.	MLVSS	%	98.0	

**Vadinar STP**

**Table 26: Sewage Water Monitoring at Vadinar STP (1<sup>st</sup> Week)**

<b>Date of Sampling</b>	02.11.2021
-------------------------	------------

Sr. No.	Parameters	Unit	Results	
			Vadinar STP I/L	Vadinar O/L
1	pH	pH unit	7.62	7.41
2	Total Suspended Solids	mg/l	121	69
3	Residual Chlorine	mg/l	-	<0.5
4	COD	mg/l	89.0	72.0
5	BOD @ 27 °C	mg/l	34.0	15.0



**Table 27: Sewage Water Monitoring at Vadinar STP (2<sup>nd</sup> Week)**

<b>Date of Sampling</b>	11.11.2021
-------------------------	------------

<b>Sr. No.</b>	<b>Parameters</b>	<b>Unit</b>	<b>Results</b>	
			<b>Vadinar STP I/L</b>	<b>Vadinar O/L</b>
1	pH	pH unit	7.5	7.2
2	Total Suspended Solids	mg/l	109	31
3	Residual Chlorine	mg/l	-	<0.5
4	COD	mg/l	173.0	62.0
5	BOD @ 27 °C	mg/l	48.0	20.0

**Table 28: Sewage Water Monitoring at Vadinar STP (3<sup>rd</sup> Week)**

<b>Date of Sampling</b>	17.11.2021
-------------------------	------------

<b>Sr. No.</b>	<b>Parameters</b>	<b>Unit</b>	<b>Results</b>	
			<b>Vadinar STP I/L</b>	<b>Vadinar O/L</b>
1	pH	pH unit	7.7	7.5
2	Total Suspended Solids	mg/l	105	38
3	Residual Chlorine	mg/l	-	<0.5
4	COD	mg/l	150	62
5	BOD @ 27 °C	mg/l	48.0	18.0

**Table 29: Sewage Water Monitoring at Vadinar STP (4<sup>th</sup> Week)**

<b>Date of Sampling</b>	<b>25.10.2021</b>
-------------------------	-------------------

Sr. No.	Parameters	Unit	Results	
			Vadinar STP I/L	Vadinar O/L
1	pH	pH unit	7.5	7.3
2	Total Suspended Solids	mg/l	117	69
3	Residual Chlorine	mg/l	-	<0.5
4	COD	mg/l	192	101
5	BOD @ 27 °C	mg/l	60.0	24.0

### **5.3 Conclusions:**

The GPCB standards of BOD, TSS and Residual Chlorine for STP outlet are 20 mg/lit, 30 mg/lit & 0.5 mg/lit respectively. It is suggested to do treatment on regular basis to avoid flow of contaminated/polluted water into the sea.

### **Marine Water Monitoring**

The Forty Second Amendment to the Constitution in 1976 underscored the importance of 'green thinking'. Article 48A enjoins the state to protect and improve the environment and safeguard the forests and wildlife in the country. Further, Article 51A(g) states that the "fundamental duty of every citizen is to protect and improve the natural environment including forests, lakes, rivers and wildlife and to have compassion for living creatures".

Policy Statement for Abatement of Pollution (1992) has suggested developing relevant legislation and regulation, fiscal incentives, voluntary agreements and educational programs and information campaigns. It emphasizes the need for integration by incorporating environmental considerations into decision making at all levels by adopting frameworks namely, pollution prevention at source, application of best practicable solution, ensure polluter pays for control of pollution, focus on heavily polluted areas and river stretches and involve public in decision-making. The National Conservation Strategy and Policy Statement on Environment and Development, (1992) aimed at

“integrating environmental concerns with developmental imperatives to meet the challenges by redirecting the thrust of our developmental process so that the basic needs of our people could be fulfilled by making judicious and sustainable use of natural resources.” The priorities mentioned in this policy document include the sustainable use of land and water resources, prevention and control of pollution and preservation of biodiversity.

The National Water Policy, (2002) contains provisions for developing, conserving, sustainable utilizing and managing this important water resources and need to be governed by national perspectives.

### **Marine Environment**

On national and state levels, we have several policies and regulation like Water (Prevention and Control of Pollution) Act, 1974, to regulate pollution discharges and restore water quality of our aquatic resources including the prescription of monitoring activities. One of the important provisions of the Water Act, 1974, is to maintain and restore the ‘wholesomeness’ of our aquatic resources. Water quality monitoring is one of the first steps required in the rational development and management of water resources. In the field of water quality management, there has been a steady evolution in procedures for designing system to obtain information on the changes of water quality. The monitoring comprises all activities to obtain ‘information’ with respect to the water system.

### **Sampling Stations**

The monitoring of marine environment for the study of biological and ecological parameters was carried out on 19<sup>th</sup>& 20<sup>th</sup> November-2021 in harbor regions of KPT and on 19<sup>th</sup> November-2021 at Vadinar during spring tide period of New moon phase of Lunar Cycle. The monitoring of marine environment for the study of biological and ecological parameters was repeated again on 26<sup>th</sup>& 27<sup>th</sup> November 2021 in harbor regions of KPT. 26<sup>th</sup> November -2021 in Vadinar during Neap tide period first quarter of Lunar Cycle..

Plankton samples from sub surface layer was collected both during high tide period and low tide period from 3 water quality monitoring stations of KPT harbour area and two stations in Nakti creek and one station in Khori creek. The same sampling schedule was repeated during consecutive spring tide and neap tide in same month. Plankton samples from sub surface layer was

collected both during high tide period and low tide period from 1 water quality monitoring stations near Vadinar jetty area during spring tide and neap tide in this month .Collected water samples were processed for estimation of Chlorophyll- a, Pheophytin- a, qualitative & quantitative evaluation of phytoplankton, qualitative & quantitative evaluation zooplanktons (density and their population).

### Sampling Locations

Offshore monitoring requirement	Number of locations
Offshore Installations	3 in Kandla creek 2 in Nakti creek 1 in Khori creek 1 near Vadinar Jetty 1 near 1 <sup>st</sup> SBM
<b>Total Number of locations</b>	<b>8</b>

### 5.4 Marine Water Quality

Marine water quality of marine waters of Deendayal Port Harbor waters, Khori and Nakti Creeks and two locations of Vadinar are monitored for various physico-chemical parameters during spring and neap tide of each month.

The results of marine water quality and Marine sediments are as below;

**Table 30: Marine Water Quality Monitoring Parameters for location near KPT colony**

Sr. No.	Parameters	Unit	Kandla Creek Near KPT colony (1)			
			23°0'58"N 70°13'22."E			
			Spring Tide		Neap Tide	
Tide	High Tide	Low Tide	High Tide	Low Tide		
1	pH	pH unit	7.21	7.26	7.45	7.26
2	Color	-	Colorless	Colorless	Colorless	Colorless
3	Odor	-	Odorless	Odorless	Odorless	Odorless
4	Salinity	ppt	31.6	31.9	32.0	31.5
5	Turbidity	NTU	36	31	33	32
6	Total Dissolved Solids	mg/l	41592	42007	41300.0	41443.0

**DCPL/DPT/20-21/19 -NOVEMBER - 2021**

**Detox Corporation Pvt. Ltd., Surat**

**Environmental Monitoring Report of Deendayal Port Trust,  
November-2021**

Sr. No.	Parameters	Unit	Kandla Creek Near KPT colony (1)			
			23°0'58"N 70°13'22."E			
	Tide	High Tide	Low Tide	High Tide	Low Tide	
7	Total Suspended Solids	mg/l	655	870	754.2	571.1
8	Total Solids	mg/l	42247	42877	42054.2	42014.1
9	DO	mg/l	4	4.2	4.1	4.3
10	COD	mg/l	82.0	90.0	80.0	78.0
11	BOD	mg/l	BQL	BQL	BQL	BQL
12	Silica	mg/l	0.65	0.60	0.76	0.65
13	Phosphate	mg/l	0.35	0.24	0.16	0.18
14	Sulphate	mg/l	2772	2700	2184	2580
15	Nitrate	mg/l	2.89	2.46	2.45	3.44
16	Nitrite	mg/l	<0.05	<0.05	<0.05	<0.05
17	Calcium	mg/l	561.12	480.96	480.96	521.04
18	Magnesium	mg/l	1725.3	1530.9	1676.7	1603.8
19	Sodium	mg/l	9038.0	8014.0	8629.0	9638.0
20	Potassium	mg/l	313.0	271.0	336.0	378.0
21	Iron	mg/l	1.42	1.30	1.32	1.10
22	Chromium	mg/l	0.12	0.11	0.13	0.12
23	Copper	mg/l	BQL	BQL	BQL	BQL
24	Arsenic	mg/l	BQL	BQL	BQL	BQL
25	Cadmium	mg/l	0.05	0.06	0.06	0.05
26	Mercury	mg/l	BQL	BQL	BQL	BQL
27	Lead	mg/l	0.07	0.09	0.06	0.08
28	Zinc	mg/l	BQL	BQL	BQL	BQL

BQL- Below Quantification Limit, (BOD-2.0 mg/l,Cu-0.1 mg/l, As-0.1mg/l, Hg-0.01 mg/l, Zinc-0.1 mg/l).

**Environmental Monitoring Report of Deendayal Port Trust,  
November-2021**

**Table 31: Marine Water Quality Monitoring Parameters for location near  
passenger Jetty One at Kandla**

Sr. No.	Parameters	Unit	Near passenger Jetty One (2)			
			23° 0'18 "N 70°13'31"E			
			Spring Tide		Neap Tide	
Tide	High Tide	Low Tide	High Tide	Low Tide		
1	pH	pH unit	7.12	7.31	7.30	7.20
2	Color	-	Colorless	Colorless	Colorless	Colorless
3	Odor	-	Odorless	Odorless	Odorless	Odorless
4	Salinity	ppt	32.0	31.6	32.2	31.4
5	Turbidity	NTU	35	28	38	42
6	Total Dissolved Solids	mg/l	39062	40035	40245.0	36627.0
7	Total Suspended Solids	mg/l	784	773	528.3	504
8	Total Solids	mg/l	39845	40808	40773.3	37131.0
9	DO	mg/l	4.3	3.9	4.2	4
10	COD	mg/l	88.0	86.0	92.0	90.0
11	BOD	mg/l	BQL	BQL	BQL	BQL
12	Silica	mg/l	0.56	0.53	0.76	0.69
13	Phosphate	mg/l	0.24	0.26	0.19	0.20
14	Sulphate	mg/l	2580	3132	2340	2700
15	Nitrate	mg/l	3.03	3.31	2.80	3.98
16	Nitrite	mg/l	BQL	BQL	BQL	BQL
17	Calcium	mg/l	601.20	681.36	561.12	601.2
18	Magnesium	mg/l	1555.2	1652.4	1676.7	1628.1
19	Sodium	mg/l	9530.0	9278.0	9116.0	9368.0
20	Potassium	mg/l	349.0	336.0	272.0	302.0
21	Iron	mg/l	1.88	1.70	1.48	1.55
22	Chromium	mg/l	0.12	0.11	0.11	0.14
23	Copper	mg/l	BQL	BQL	BQL	BQL
24	Arsenic	mg/l	BQL	BQL	BQL	BQL
25	Cadmium	mg/l	0.04	0.05	0.06	0.04
26	Mercury	mg/l	BQL	BQL	BQL	BQL
27	Lead	mg/l	0.10	0.09	0.09	0.10
28	Zinc	mg/l	BQL	BQL	BQL	BQL

BQL- Below Quantification Limit, (BOD-2.0 mg/l, Nitrite: 0.05mg/l, Cu-0.1 mg/l, As-0.1mg/l, Hg-0.01 mg/l, Zinc-0.1 mg/l).

**DCPL/DPT/20-21/19 -NOVEMBER - 2021**

**Detox Corporation Pvt. Ltd., Surat**

**Environmental Monitoring Report of Deendayal Port Trust,  
November-2021**

**Table 32: Marine Water Quality Monitoring Parameters for location Near Coal Berth**

Sr. No.	Parameters	Unit	Near Coal Berth			
			22°59'12"N 70°13'40"E			
			Spring Tide		Neap Tide	
Tide		High Tide	Low Tide	High Tide	Low Tide	
1	pH	pH unit	7.30	7.46	7.30	7.36
2	Color	-	Colorless	Colorless	Colorless	Colorless
3	Odor	-	Odorless	Odorless	Odorless	Odorless
4	Salinity	ppt	32.0	31.8	32.7	31.0
5	Turbidity	NTU	34	29	36	40
6	Total Dissolved Solids	mg/l	43205	41674	43606.0	40029.0
7	Total Suspended Solids	mg/l	590	863	500.2	604.3
8	Total Solids	mg/l	43795	42537	44106.2	40633.3
9	DO	mg/l	4	5.1	4.5	4.4
10	COD	mg/l	90.0	86.0	88.0	79.0
11	BOD	mg/l	BQL	BQL	BQL	BQL
12	Silica	mg/l	0.69	0.75	0.56	0.64
13	Phosphate	mg/l	0.28	0.34	0.17	0.20
14	Sulphate	mg/l	3240	2016	2676	2148
15	Nitrate	mg/l	3.87	4.58	2.95	2.62
16	Nitrite	mg/l	<0.05	<0.05	<0.05	<0.05
17	Calcium	mg/l	641.28	601.20	480.96	521.04
18	Magnesium	mg/l	1628.1	1749.6	1749.6	1749.6
19	Sodium	mg/l	9425.0	8408.0	9423.0	8709.0
20	Potassium	mg/l	339.0	299.0	306.0	230.0
21	Iron	mg/l	1.41	1.78	1.76	1.56
22	Chromium	mg/l	0.11	0.13	0.13	0.12
23	Copper	mg/l	BQL	BQL	BQL	BQL
24	Arsenic	mg/l	BQL	BQL	BQL	BQL
25	Cadmium	mg/l	0.06	0.05	0.05	0.07
26	Mercury	mg/l	BQL	BQL	BQL	BQL
27	Lead	mg/l	0.08	0.09	0.09	0.07
28	Zinc	mg/l	BQL	BQL	BQL	BQL

BQL- Below Quantification Limit, (BOD-2.0 mg/l,Cu-0.1 mg/l, As-0.1mg/l, Hg-0.01 mg/l,Zinc-0.1 mg/l).

**DCPL/DPT/20-21/19 -NOVEMBER - 2021**

**Detox Corporation Pvt. Ltd., Surat**

**Table 33: Marine Water Quality Monitoring Parameters for location  
Khori creek at Kandla**

Sr. No.	Parameters	Unit	KPT 4			
			Near 15/16 Berth			
			Spring Tide		Neap Tide	
			High Tide	Low Tide	High Tide	Low Tide
1	pH	pH unit	7.35	7.50	7.50	7.20
2	Color	-	Colorless	Colorless	Colorless	Colorless
3	Odor	-	Odorless	Odorless	Odorless	Odorless
4	Salinity	ppt	32.1	32.6	31.9	31.6
5	Turbidity	NTU	43	39	45	33
6	Total Dissolved Solids	mg/l	42399	39089	38986.0	39711.0
7	Total Suspended Solids	mg/l	743	577	681.8	530.3
8	Total Solids	mg/l	43142	39666	39667.8	40241.3
9	DO	mg/l	4.6	4.5	5.2	5.6
10	COD	mg/l	92.0	90.0	82.0	89.0
11	BOD	mg/l	BQL	BQL	BQL	BQL
12	Silica	mg/l	0.82	0.58	0.58	0.51
13	Phosphate	mg/l	0.28	0.25	0.22	0.20
14	Sulphate	mg/l	1620	3492	2388	2100
15	Nitrate	mg/l	1.97	3.03	2.71	2.06
16	Nitrite	mg/l	<0.05	<0.05	<0.05	<0.05
17	Calcium	mg/l	561.12	641.28	561.12	440.88
18	Magnesium	mg/l	1579.5	1603.8	1603.8	1822.5
19	Sodium	mg/l	9423.0	9014.0	9526.0	9468.0
20	Potassium	mg/l	341.0	301.0	218.0	221.0
21	Iron	mg/l	1.62	1.33	1.74	1.21
22	Chromium	mg/l	0.16	0.12	0.14	0.16
23	Copper	mg/l	BQL	BQL	BQL	BQL
24	Arsenic	mg/l	BQL	BQL	BQL	BQL
25	Cadmium	mg/l	0.04	0.05	0.06	0.04
26	Mercury	mg/l	BQL	BQL	BQL	BQL
27	Lead	mg/l	0.08	0.06	0.09	0.08
28	Zinc	mg/l	BQL	BQL	BQL	BQL

BQL- Below Quantification Limit, (BOD-2.0 mg/l,Cu-0.1 mg/l, As-0.1mg/l, Hg-0.01 mg/l, Zinc-0.1 mg/l).

**DCPL/DPT/20-21/19 -NOVEMBER - 2021**

**Detox Corporation Pvt. Ltd., Surat**



**Table 34: Marine Water Quality Monitoring Parameters for location  
Nakti Creek near Tuna Port**

Sr. No.	Parameters	Unit	Nakti Creek Near Tuna Port			
			22°57'49."N 70° 7'0.67"E			
			Spring Tide		Neap Tide	
			High Tide	Low Tide	High Tide	Low Tide
1	pH	pH unit	7.30	7.20	7.30	7.40
2	Color	-	Colorless	Colorless	Colorless	Colorless
3	Odor	-	Odorless	Odorless	Odorless	Odorless
4	Salinity	ppt	31.6	31.2	31.2	31.1
5	Turbidity	NTU	36	42	36	31
6	Total Dissolved Solids	mg/l	40770	38329	38644.0	38955.0
7	Total Suspended Solids	mg/l	766	853	494.2	474
8	Total Solids	mg/l	41536	39182	39138.2	39429.0
9	DO	mg/l	4.1	4.7	4.6	4.8
10	COD	mg/l	98.0	96.0	96.0	98.0
11	BOD	mg/l	BQL	BQL	BQL	BQL
12	Silica	mg/l	0.53	0.89	0.75	0.64
13	Phosphate	mg/l	0.24	0.24	0.21	0.18
14	Sulphate	mg/l	3456	3732	2820	2424
15	Nitrate	mg/l	2.75	3.38	2.77	4.31
16	Nitrite	mg/l	<0.05	<0.05	<0.05	<0.05
17	Calcium	mg/l	561.12	521.04	480.96	561.12
18	Magnesium	mg/l	1676.7	1725.3	1773.9	1676.7
19	Sodium	mg/l	9839.0	10125.0	10118.0	10168.0
20	Potassium	mg/l	399.0	402.0	387.0	390.0
21	Iron	mg/l	1.20	1.13	1.45	1.10
22	Chromium	mg/l	0.11	0.13	0.14	0.14
23	Copper	mg/l	BQL	BQL	BQL	BQL
24	Arsenic	mg/l	BQL	BQL	BQL	BQL
25	Cadmium	mg/l	0.05	0.07	0.07	0.08

**DCPL/DPT/20-21/19 -NOVEMBER - 2021**

**Detox Corporation Pvt. Ltd., Surat**

**Environmental Monitoring Report of Deendayal Port Trust,  
November-2021**

Sr. No.	Parameters	Unit	Nakti Creek Near Tuna Port			
			22°57'49."N 70° 7'0.67"E			
			Spring Tide		Neap Tide	
			High Tide	Low Tide	High Tide	Low Tide
26	Mercury	mg/l	BQL	BQL	BQL	BQL
27	Lead	mg/l	0.07	0.09	0.10	0.09
28	Zinc	mg/l	BQL	BQL	BQL	BQL

BQL- Below Quantification Limit, (BOD-2.0 mg/l,Cu-0.1 mg/l, As-0.1mg/l, Hg-0.01 mg/l,Zinc-0.1 mg/l).

**Table 35: Marine Water Quality Monitoring Parameters for location  
Nakti Creek Near NH-8A at Kandla**

Sr. No.	Parameters	Unit	Nakti Creek Near NH-8A			
			23° 02'01"N 70° 09'31"E			
			Spring Tide		Neap Tide	
			High Tide	Low Tide	High Tide	Low Tide
1	pH	pH unit	7.52		7.52	
2	Color	-	Colorless		Colorless	
3	Odor	-	Odorless		Odorless	
4	Salinity	ppt	31.8		32.2	
5	Turbidity	NTU	35		35	
6	Total Dissolved Solids	mg/l	41695		42152.0	
7	Total Suspended Solids	mg/l	684		452	
8	Total Solids	mg/l	42379		42604.0	
9	DO	mg/l	4.8	Sampling not possible during Low Tide	5.1	Sampling not possible during Low Tide
10	COD	mg/l	100.0		94.0	
11	BOD	mg/l	BQL		BQL	
12	Silica	mg/l	0.96		0.53	
13	Phosphate	mg/l	0.23		0.17	
14	Sulphate	mg/l	3780		2376	
15	Nitrate	mg/l	3.24		3.61	
16	Nitrite	mg/l	<0.05		<0.05	
17	Calcium	mg/l	480.96		601.2	

**DCPL/DPT/20-21/19 -NOVEMBER - 2021**

**Detox Corporation Pvt. Ltd., Surat**

**Environmental Monitoring Report of Deendayal Port Trust,  
November-2021**

Sr. No.	Parameters	Unit	Nakti Creek Near NH-8A			
			23° 02'01"N 70° 09'31"E			
			Spring Tide		Neap Tide	
			Tide		High Tide	Low Tide
18	Magnesium	mg/l	1725.3		1628.1	
19	Sodium	mg/l	10308.0		10319.0	
20	Potassium	mg/l	409.0		364.0	
21	Iron	mg/l	1.20		1.35	
22	Chromium	mg/l	0.11		0.12	
23	Copper	mg/l	BQL		BQL	
24	Arsenic	mg/l	BQL		BQL	
25	Cadmium	mg/l	0.06		0.06	
26	Mercury	mg/l	BQL		BQL	
27	Lead	mg/l	0.08		0.11	
28	Zinc	mg/l	BQL		BQL	

BQL- Below Quantification Limit, (BOD-2.0 mg/l, Cu-0.1 mg/l, As-0.1mg/l, Hg-0.01 mg/l, Zinc-0.1 mg/l).

**Table 36: Marine Water Quality Monitoring Parameters for locations Nr. Vadinar Jetty**

Sr. No.	Parameters	Unit	Nr.Vadinar Jetty			
			22°26'25.26"N 69°40'20.41"E			
			Spring Tide		Neap Tide	
			Tide		High Tide	Low Tide
1	pH	pH unit	7.60	7.70	7.41	7.52
2	Color	-	Colorless	Colorless	Colorless	Colorless
3	Odor	-	Odorless	Odorless	Odorless	Odorless
4	Salinity	ppt	31.2	32.0	31.5	31.4
5	Turbidity	NTU	45	39	42	40
6	Total Dissolved Solids	mg/l	38510	42661	40025.0	40250.0
7	Total Suspended Solids	mg/l	585	523	548.9	505
8	Total Solids	mg/l	39095	43184	40573.9	40755.0
9	DO	mg/l	4.4	4.6	4.7	4.6

**DCPL/DPT/20-21/19 -NOVEMBER - 2021**

**Detox Corporation Pvt. Ltd., Surat**

**Environmental Monitoring Report of Deendayal Port Trust,  
November-2021**

Sr. No.	Parameters	Unit	Nr.Vadinar Jetty			
			22°26'25.26"N 69°40'20.41"E			
			Spring Tide		Neap Tide	
			High Tide	Low Tide	High Tide	Low Tide
10	COD	mg/l	76.0	80.0	72.0	70.0
11	BOD	mg/l	BQL	BQL	BQL	BQL
12	Silica	mg/l	0.85	1.02	0.75	0.82
13	Phosphate	mg/l	0.22	0.25	0.18	0.17
14	Sulphate	mg/l	2580	2700	2592	2508
15	Nitrate	mg/l	2.75	3.59	3.67	3.39
16	Nitrite	mg/l	<0.05	<0.05	<0.05	<0.05
17	Calcium	mg/l	601.20	521.04	641.28	480.96
18	Magnesium	mg/l	1603.8	1676.7	1652.4	1676.7
19	Sodium	mg/l	10968.0	10848.0	11126.0	10829.0
20	Potassium	mg/l	344.0	382.0	355.0	392.0
21	Iron	mg/l	1.06	1.70	1.12	1.42
22	Chromium	mg/l	0.12	0.13	0.14	0.13
23	Copper	mg/l	BQL	BQL	BQL	BQL
24	Arsenic	mg/l	BQL	BQL	BQL	BQL
25	Cadmium	mg/l	0.05	0.04	0.08	0.07
26	Mercury	mg/l	BQL	BQL	BQL	BQL
27	Lead	mg/l	0.10	0.08	0.10	0.09
28	Zinc	mg/l	BQL	BQL	BQL	BQL

BQL- Below Quantification Limit, (BOD-2.0 mg/l,Cu-0.1 mg/l, As-0.1mg/l, Hg-0.01 mg/l, Zinc-0.1 mg/l).

**Table 36 (a): Marine Water Quality Monitoring Parameters for locations  
Nr. Vadinar SPM**

Sr. No.	Parameters	Unit	Nr.Vadinar SPM			
			22°30'56.15"N 69°42'12.07"E			
			Spring Tide		Neap Tide	
			High Tide	Low Tide	High Tide	Low Tide
1	pH	pH unit	7.40	7.60	7.45	7.26

**DCPL/DPT/20-21/19 -NOVEMBER - 2021**

**Detox Corporation Pvt. Ltd., Surat**

**Environmental Monitoring Report of Deendayal Port Trust,  
November-2021**

Sr. No.	Parameters	Unit	Nr.Vadinar SPM			
			22°30'56.15"N 69°42'12.07"E			
			Spring Tide		Neap Tide	
			Tide	High Tide	Low Tide	High Tide
2	Color	-	Colorless	Colorless	Colorless	Colorless
3	Odor	-	Odorless	Odorless	Odorless	Odorless
4	Salinity	ppt	32.2	32.1	32.0	31.8
5	Turbidity	NTU	33.0	34.0	36.0	33.0
6	Total Dissolved Solids	mg/l	41700.0	41987	40610.0	40925
7	Total Suspended Solids	mg/l	635.0	480	513.0	548
8	Total Solids	mg/l	43340.0	43924	41384.0	42000
9	DO	mg/l	4.3	4.1	4.5	4.3
10	COD	mg/l	90.0	92.0	78.0	70.0
11	BOD	mg/l	BQL	BQL	BQL	BQL
12	Silica	mg/l	0.92	0.96	0.6	0.78
13	Phosphate	mg/l	0.24	0.25	0.2	0.16
14	Sulphate	mg/l	2628.0	2364	2316.0	2556
15	Nitrate	mg/l	3.10	3.38	3.34	3.68
16	Nitrite	mg/l	<0.05	<0.05	<0.05	<0.05
17	Calcium	mg/l	481.0	561.12	521.0	561.12
18	Magnesium	mg/l	1652.4	2065.5	1676.7	1701
19	Sodium	mg/l	10318	10829	10418	10786
20	Potassium	mg/l	354	355	377	354
21	Iron	mg/l	1.60	1.80	1.27	1.90
22	Chromium	mg/l	0.14	0.14	0.13	0.12
23	Copper	mg/l	BQL	BQL	BQL	BQL
24	Arsenic	mg/l	BQL	BQL	BQL	BQL
25	Cadmium	mg/l	0.05	0.04	0.08	0.08
26	Mercury	mg/l	BQL	BQL	BQL	BQL
27	Lead	mg/l	0.09	0.08	0.11	0.09

**DCPL/DPT/20-21/19 -NOVEMBER - 2021**

**Detox Corporation Pvt. Ltd., Surat**

**Environmental Monitoring Report of Deendayal Port Trust,  
November-2021**

Sr. No.	Parameters	Unit	Nr.Vadinar SPM			
			22°30'56.15"N 69°42'12.07"E			
			Spring Tide		Neap Tide	
			High Tide	Low Tide	High Tide	Low Tide
28	Zinc	mg/l	BQL	BQL	BQL	BQL

BQL- Below Quantification Limit, (BOD-2.0 mg/l,Cu-0.1 mg/l, As-0.1mg/l, Hg-0.01 mg/l,Zinc-0.1 mg/l).

#### 5.4.1 Marine Sediments

Sediment samples were collected with Van Veen Grab from the six locations in Kandla Port Waters and two locations in Vadinar Port. Samples were collected and preserved in silver foil in ice box to prevent the contamination/decaying of the samples.

#### 5.5 Results

The Sediment Quality results are given in below from table no. 34 A & B.

**Table 34A: Results of Analysis of Sediment of Kandla & Vadinar Port  
(Spring Tide)**

Sr. No.	Parameters	Unit	KPT - 1	KPT - 2	KPT - 3	KPT - 4	KPT - 5	Jetty	SPM
1	Texture	-	Sandy Loam	Sandy Loam	Sandy Loam	Sandy Loam	Sandy Loam	Sandy Loam	Sandy Loam
2	Organic Matter	mg/kg	1.72	1.34	1.36	2.74	1.53	2.31	2.00
3	Organic Carbon	mg/kg	0.99	0.78	0.79	1.59	0.89	1.34	1.16
4	Inorganic Phosphate	mg/kg	112.0	121.0	116.0	124.0	128.0	122.0	133.0
5	Moisture	%	29.43	23.11	31.0	27.25	25.69	27.4	43.00
6	Aluminium	mg/kg	ND	ND	ND	ND	ND	ND	ND
7	Silica	mg/kg	11.2	10.5	12.3	11.4	10.5	13.2	13.00
8	Phosphate	mg/kg	2.57	2.73	9.48	6.12	11.84	5.96	6.68
9	Sulphate	mg/kg	283.0	257.0	411.0	182.0	338.0	209.0	494.7
10	Nitrite	mg/kg	0.12	0.11	0.12	0.12	0.11	0.1	0.11
11	Nitrate	mg/kg	BQL	BQL	BQL	BQL	BQL	BQL	BQL
12	Calcium	mg/kg	364.7	152.3	505.0	76.2	325.0	225.0	177.0
13	Magnesium	mg/kg	260.0	241.0	158.0	175.0	308.0	58.3	228.4

**DCPL/DPT/20-21/19 -NOVEMBER - 2021**

**Detox Corporation Pvt. Ltd., Surat**

**Environmental Monitoring Report of Deendayal Port Trust,  
November-2021**

14	Sodium	mg/kg	1819.0	2881.0	876.0	1858.0	4022.0	3159.0	8637.0
15	Potassium	mg/kg	119.0	166.0	102.0	113.0	263.0	283.0	1036.7
16	Chromium	mg/kg	60	46.6	51.2	43.2	58	46.40	65.00
17	Nickel	mg/kg	32.1	33.7	24.9	28	32.5	28.00	45.60
18	Copper	mg/kg	39.9	14.6	33	16.8	31.8	26.80	21.00
19	Zinc	mg/kg	81.90	52.70	60.80	42.70	68.70	64.30	65.90
20	Cadmium	mg/kg	2.0	BQL	BQL	BQL	BQL	BQL	BQL
21	Lead	mg/kg	18.50	5.4	9.0	5.70	11.4	10.90	5.20
22	Mercury	mg/kg	BQL	BQL	BQL	BQL	BQL	BQL	BQL
23	Arsenic	mg/kg	BQL	BQL	BQL	BQL	BQL	BQL	BQL

\*ND - Not Detected, BQL: Below Quantification Limit (NO3:10.0mg/kg, Cd: 1.0mg/kg, Hg: 1.0mg/kg, As: 1.0mg/kg)

**Table 34B: Results of Analysis of Sediment of Kandla & Vadinar Port (Neap Tide)**

Sr. No.	Parameters	Unit	KPT - 1	KPT - 2	KPT - 3	KPT - 4	KPT - 5	Jetty	SPM
1	Texture	-	Sandy Loam	Sandy Loam	Sandy Loam	Sandy Loam	Sandy Loam	Sandy Loam	Sandy Loam
2	Organic Matter	mg/kg	1.12	1.50	2.88	1.41	0.69	1.43	1.69
3	Organic Carbon	mg/kg	0.65	0.87	1.67	0.82	0.40	0.83	0.98
4	Inorganic Phosphate	mg/kg	118.0	126.0	120.0	130.0	128.0	112.0	130.0
5	Moisture	%	17.76	22.98	20.4	14.01	22.6	34.3	32.16
6	Aluminium	mg/kg	ND	ND	ND	ND	ND	ND	ND
7	Silica	mg/kg	12.0	11.3	10.5	11.3	9.8	11.4	12.60
8	Phosphate	mg/kg	17.14	2.91	7.83	0.49	2.54	15.65	2.20
9	Sulphate	mg/kg	255.0	427.0	290.0	440.0	390.0	564.0	595.0
10	Nitrite	mg/kg	0.1	0.11	0.1	0.11	0.12	0.1	0.11
11	Nitrate	mg/kg	BQL	BQL	BQL	BQL	BQL	BQL	BQL
12	Calcium	mg/kg	180.0	188.0	172.0	180.0	176.0	116.0	140.0
13	Magnesium	mg/kg	38.9	102.1	82.6	150.7	58.3	158.0	179.8

**DCPL/DPT/20-21/19 -NOVEMBER - 2021**

**Detox Corporation Pvt. Ltd., Surat**

**Environmental Monitoring Report of Deendayal Port Trust,  
November-2021**

Sr. No.	Parameters	Unit	KPT - 1	KPT - 2	KPT - 3	KPT - 4	KPT - 5	Jetty	SPM
14	Sodium	mg/kg	325.0	853.6	743.8	656.1	414.0	1895.0	1810.0
15	Potassium	mg/kg	25.7	72.3	52.3	52.3	40.0	248.0	307.0
16	Chromium	mg/kg	25.7	38.6	28.6	27.3	31.3	51.90	56.20
17	Nickel	mg/kg	18.0	29.2	20.1	18.4	17.8	32.40	72.70
18	Copper	mg/kg	12.30	20.30	6.70	9.40	7.90	22.20	41.10
19	Zinc	mg/kg	24.90	57.40	32.80	27.90	25.50	46.40	1511.00
20	Cadmium	mg/kg	BQL	BQL	BQL	BQL	BQL	BQL	BQL
21	Lead	mg/kg	4.60	6.7	7.5	3.70	3.3	24.70	29.60
22	Mercury	mg/kg	BQL	BQL	BQL	BQL	BQL	BQL	BQL
23	Arsenic	mg/kg	BQL	BQL	BQL	BQL	BQL	BQL	BQL

\*ND - Not Detected, BQL: Below Quantification Limit (NO3:10.0mg/kg,Cd: 1.0mg/kg,Hg: 1.0mg/kg, As: 1.0mg/kg)



**REPORT**  
**ON**  
**ECOLOGICAL MONITORING**  
**OF MARINE ENVIRONMENT**  
**IN**  
**DPT HARBOUR AREA, NEAR BY CREEKS**  
**AND**  
**VADINAR JETTY AND SPM**  
**FOR**  
**DEENDAYAL PORT TRUST**

November, 2021

## **INTRODUCTION:**

### **Sampling Stations:**

The monitoring of marine environment for the study of biological and ecological Parameters was carried out on 19<sup>th</sup> November, 2021 in harbour region of DPT at Kandla Creek, and on 20<sup>th</sup> November, 2021 in creeks near by the port during spring tide. The monitoring of marine environment for the study of biological and ecological parameters was repeated again on 26<sup>th</sup> November, 2021 in harbour region of DPT at Kandla Creek and on 27<sup>th</sup> November 2021 in creeks near by the port during neap tidal condition.

Plankton samples from sub surface layer was collected both during high tide period and low tide period from 3 water quality monitoring stations of KPT harbour area and one stations in Nakti creek and one station in Khori creek. Sampling at second sampling station of Nakti creek was possible only during high tide period. The same sampling schedule was repeated during consecutive Neap tide and spring tide in same month.

Plankton samples from sub surface layer were collected during high tide period and low tide period from monitoring station near Vadinar jetty at Path Finder Creek during neap tide on 11/11/2021 and spring tide period on 26/11/2021. Collected water samples were processed for estimation of Chlorophyll- a, Pheophytin- a, qualitative & quantitative evaluation of phytoplankton, qualitative & quantitative evaluation zooplanktons (density and their population).

**TABLE #1 SAMPLING LOCATIONS**

<b>monitoring requirement</b>	<b>Number of locations</b>
Kandla creek	3 in Kandla creek
Nakti creek	2 in Nakti creek
Khori Creek	1 in Khori creek
Vadinar jetty	1 near Vadinar Jetty
SPM	1 near 1 <sup>st</sup> SPM
<b>Total Number of locations</b>	<b>8</b>

### **Sampling methodology adopted:**

**A** marine sampling is an estimation of the body of information in the population. The theory of the sampling design is depending upon the underlying frequency

**DCPL/DPT/20-21/19 -NOVEMBER - 2021**

---

**Detox Corporation Pvt. Ltd., Surat**

distribution of the population of interest. The requirement for useful water sampling is to collect a representative sample of suitable volume from the specified depth and retain it free from contamination during retrieval.

50 litres of the water sample were collected from Sub surface by using bucket. From the collected water sample 1 litres of water sample were taken in an opaque plastic bottle for chlorophyll estimation, thereafter plankton samples were collected by using filtration assembly with nilyobolt cloth of 20µm mesh size.

### **Samples Processing for chlorophyll estimation:**

Samples for the chlorophyll estimation were preserved in ice box on board in darkness to avoid degradation in opaque container covered with aluminium foil. Immediately after reaching the shore after sampling, 1 litre of collected water sample was filtered through GF/F filters (pore size 0.45 µm) by using vacuum filtration assembly. After vacuum filtration the glass micro fiber filter paper was grunted in tissue grinder, macerating of glass fiber filter paper along with the filtrate was done in 90% aqueous Acetone in the glass tissue grinder with glass grinding tube. Glass fiber filter paper will assist breaking the cell during grinding and chlorophyll content was extracted with 10 ml of 90% Acetone, under cold dark conditions along with saturated magnesium carbonate solution in glass screw cap tubes. After an extraction period of 24 hours, the samples were transferred to calibrated centrifuge tubes and adjusted the volume to original volume with 90% aqueous acetone solution to make up the evaporation loss. The extract was clarified by using centrifuge in closed tubes. The clarified extracts were then decanted in clean cuvette and optical density was observed at wavelength 664, 665 nm. By using corrected optical density, Chlorophyll-a value was calculated as given in (APHA, 1998).

### **PLANKTON:**

The entire area open water in the sea is the pelagic realm. Pelagic organisms live in the open sea. In contrast to the pelagic realm, the benthic realm comprises organisms and zone of the bottom of the sea. Vertically the pelagic realm can be dividing into two zones based on light penetration; upper photic or euphotic zone and lower dark water mass, aphotic zone below the photic zone.

The term plankton is general term for organisms have such limited powers of locomotion that they are at the mercy of the prevailing water movement. Plankton is subdivided to phytoplankton and zooplankton. Phytoplankton is free floating organisms that are capable of photosynthesis and zooplankton is the various free floating animals.

**DCPL/DPT/20-21/19 -NOVEMBER - 2021**

---

***Detox Corporation Pvt. Ltd., Surat***

Pelagic zone, represents the entire ocean water column from the surface to the deepest depths, is home to a diverse community of organisms. Differences in their locomotive ability categorize the organisms in the pelagic realm into two, **plankton** and **nekton** (Lalli and Parsons, 1997). **Plankton** consists of all organisms drifting in the water and is unable to swim against water currents, whereas **Nekton** includes organisms having strong locomotive power. Ecological studies on the plankton community, which form the base of the aquatic food chain, help in the better understanding of the dynamics and functioning of the marine ecosystem. The term 'Plankton' first coined by Victor Hensen (1887), Plankton, (Greek word: *planktos* meaning "passively drifting or wandering") is defined as drifting or free-floating organisms that inhabit the pelagic zone of water. Based on their mode of nutrition planktonic organisms are categorised into phytoplankton (organisms having an autotrophic mode of nutrition) and zooplankton (organisms having a heterotrophic mode of nutrition).

#### **Phytoplankton in the marine environment:**

Phytoplankton is free floating unicellular, filamentous and colonial eutrophic organisms that grow in aquatic environments whose movement is more or less dependent upon water currents. These micro flora acts as primary producers as well as the basis of food chain, source of protein, bio purifier and bio indicators of the aquatic ecosystems of which diverse array of the life depends .They are considered as an important component of aquatic flora, play a key role in maintaining equilibrium between abiotic and biotic components of aquatic ecosystem.

The phytoplankton includes a wide range of photosynthetic and phototrophic organisms. Marine phytoplankton is mostly microscopic and unicellular floating flora, which are the primary producers that support the pelagic food-chain. The two most prominent groups of phytoplankton are diatoms (Bacillariophyceae) and dinoflagellates (Dinophyceae).The phytoplankton those normally captured in the net from the Gulf of Kutch is normally dominated by these two major groups; diatoms and dinoflagellates. Phytoplankton also include numerous and diverse collection of extremely small, motile algae which are termed micro flagellates (naked flagellates) as well as and Cyanophytes (blue-green algae).

Algae are an ecologically important group in most aquatic ecosystems and have been an important component of biological monitoring programs. Algae are ideally suited for water quality assessment because they have rapid reproduction

rates and very short life cycles, making them valuable indicators of short-term impacts.

Aquatic populations are impacted by anthropogenic stress, resulting in a variety of alterations in the biological integrity of aquatic systems. Algae can serve as an indicator of the degree of deterioration of water quality, and many algal indicators have been used to assess environmental status.

### **Zooplankton in the marine environment:**

Zooplankton includes a taxonomically and morphologically diverse community of heterotrophic organisms that drift in the waters of the world's oceans. Qualitative and quantitative studies on zooplankton community are a prerequisite to delineate the ecological processes active in the marine ecosystem. Zooplankton community plays a pivotal role in the pelagic food web as the primary consumers of phytoplankton and act as the food source for organisms in the higher trophic levels, particularly the economically essential groups such as fish larvae and fishes. They also function in the cycling of elements in the marine ecosystem. The dynamics of the zooplankton community, their reproduction, and growth and survival rate are all significant factors determining the recruitment and abundance of fish stocks as they form an essential food for larval, juvenile and adult fishes (Beaugrand et al., 2004). Zooplankton grazing in the marine environment controls the primary production and helps in determining the pelagic ecosystem (Banse, 1995). Through grazing in surface waters and following the production of sinking faecal matters and also by the active transportation of dissolved and particulate matter to deeper waters via vertical migration, they help in the transport of organic carbon to deep ocean layers and thus act as key drivers of biological pump' in the marine ecosystem. Zooplankton grazing and metabolism also, transform particulate organic matter into dissolved forms, promoting primary producer community, microbial demineralization, and particle export to the ocean's interior.

The categorisation of zooplankton into various ecological groups is based on several factors such as duration of planktonic life, size, food preferences and habitat. As they vary significantly in size from microscopic to metazoic forms, the classification of zooplankton based on size has paramount importance in the field of quantitative plankton research.

Based on the duration of planktonic life, zooplankton are categorised into Holoplankton (organisms which complete their entire lifecycle as plankton) and Meroplankton (organisms which are planktonic during the early part of their lives such as the larval stages of benthic and nektonic organisms). Tycho plankton are


organisms which live a brief planktonic life, such as the benthic crustaceans (Cumaceans, mysids, isopods) which ascend to the water column at night for feeding and certain ectoparasitic copepods, they leave the host and spend their life as plankton during their breeding cycle.

Zooplankton can be subdivided into holoplankton, i.e., permanent members of the plankton (e.g., Calanoid copepods), and meroplankton, i.e., temporary members in the plankton e.g., larvae of fish, shrimp, and crab). The meroplankton group consists of larval and young stages of animals that will adopt a different lifestyle once they mature. In contrast to phytoplankton which consist of a relatively smaller variety of organisms, Zooplankton are extremely diverse, consist of a host of larval and adult forms representing many animal phylum.

Among the zooplankton one group always dominate than others; members of sub class copepods (Phylum Athropoda) and Tintinids (Phylum Protozoa) among the net planktons. These small animals are of vital importance in marine ecosystem as one of the primary herbivores animals in the sea, and it is they provide vital link between primary producer (autotrophs) and numerous small and large marine consumers.

As their community structure and function are highly susceptible to changes in the environmental conditions regular monitoring of their distribution as well as their interactions with various physicochemical parameters is inevitable for the sustainable management of the ecosystem (Kusum et al., 2014). Of all the marine zooplankton groups, copepods mainly Calanoid copepods are the dominant groups in marine subtropical and tropical waters and exhibit considerable diversity in morphology and habitats they occupy (Madhupratap, 1991;)

It has been well established that potential of pelagic fishes viz. finfishes, crustaceans, molluscs and marine mammals either directly or indirectly depend on zooplankton. The herbivorous zooplankton is efficient grazers of the phytoplankton and is referred to as living machines transforming plant material into animal tissue. Hence they play an essential role as the intermediaries for nutrients/energy transfer between primary and tertiary trophic levels. Due to their large density, shorter lifespan, drifting nature, high group/species diversity and different tolerance to the stress, they used as the indicator organisms for the physical, chemical and biological processes in the aquatic ecosystem (Ghajbhiye, 2002).



### Spatial distribution of Plankton:

A characteristic of plankton population is that they tend to occur in patches, which are varying spatially on a scale of few meters to far as few kilometres in distance. They also vary in time scale, season as well as vertically in the water column. It is this patchiness and its constant changes in time and spot, that has made it so difficult for plankton biologist to learn about the ecology of plankton. The biological factors that causes this patchiness is due to the ability of zooplankton to migrate vertically and graze out the phytoplankton at a rapid rate that can create patchiness. Similarly the active swimming ability by certain zooplankton organisms can cause to aggregate in dense group.

At its most extreme, because the water in which plankton is suspended is constantly moving, each sample taken by the plankton biologists remain a different volume of water, so each sample is unique and replicate does not exist. Plankton may also exhibit vertical patchiness. Physical factors contribute to this type of patchiness include light intensity, nutrients and density gradients in the water column.

Phytoplankton in particular tends to be unequally distributed vertically, which leads to the existence of different concentration of a chlorophyll value between photic zone and below the photic zone.

#### **Methodology adopted for Plankton sampling:**

Mixed plankton sample were obtained from the sub surface layer at each sampling locations by towing the net horizontally with the weight .After the tow of about 15-30minutes, plankton net was pulled up and washed down to the tail and collected the plankton adhered to plankton net in the collection bucket at the bottom by springing outer and inner surface of the net with sea water, while the net was hanging with the mouth upward. For quantitative evaluation 50 L water samples were collected from subsurface layer and filtered through 20µm mesh size net by using bucket and filtration assembly.

#### **Preservation and storage:**

Both filtered plankton and those collected from the plankton net were preserved with 5% buffered formalin and stored in 1L plastic container for further processing in the laboratory.

#### **Sample concentration:**

The collected plankton samples were concentrated by using centrifuge and made up to 50 ml with 5% formalin -Glycerine mixture.

#### **Taxonomic evaluation:**

Before processing, the sample was mixed carefully and a subsample was taken with a calibrated Stempel-pipette. 1 ml of the concentrated plankton samples were transferred on a glass slide with automatic pipette. The plankton sample on the glass slides were stained by using Lugol's iodine and added glycerine to avoid drying while observation. The plankton samples were identified by using Labex triangular Research microscope with photographic attachment. Microphotographs of the plankton samples were taken for record as well as for confirming the identification. The bigger sized zooplankton was observed through dissecting stereomicroscope with magnification of 20-30 x. Plankton organisms in the whole slide were identified to the lowest axon possible. A thorough literature search was conducted for the identification of the different groups of zooplankton that were encountered

#### **Cell counts by drop count method:**

The common glass slide mounted with a 1ml of concentrated phytoplankton/zooplankton sample in glycerol and covered with cover slip 22x 60mm was placed under the compound microscope provided with a mechanical

**DCPL/DPT/20-21/19 -NOVEMBER - 2021**

---



stage. The plankton was then counted from the microscopic field of the left top corner of the slide. Then slide is moved horizontally along the right side and plankton in each microscopic field was thus counted. When first microscopic field row was finished the next consecutive row was adjusted using the mechanical device of the stage. In this way all the plankton present in entire microscopic field are counted. From this total number in 1ml of the concentrated plankton, total number of plankton in the original volume of sample filtered was calculated as units/L.

### **BENTHIC ORGANISMS:**

Benthos is those organisms that are associated with the sea bed or benthic habitats. Epi- benthic organisms live attached to a hard substratum or rooted to a shallow depth below the surface. In fauna organisms live below the sediment-water interface. Interstitial organisms live and move in pore water among sedimentary grains.

Because the benthic organisms are often collected and separated on sieves, a classification based on the overall size is used. Macro benthos include organisms whose shortest dimension is greater than or equal to 0.5 mm. Meio benthos are smaller than 0.5mm but larger than 42 $\mu$  in size.

The terms such as macro fauna and Meio fauna generally have little relevance with taxonomic classification. The terms Meio fauna and macro fauna depend on the size. Meio fauna were considered as good bioassay of community health and rather sensitive indicators of environmental changes

### **SAMPLING METHODOLOGY ADOPTED FOR SUB TIDAL REGION:**

Van veen sampler (0.09m<sup>2</sup>) was used for sampling bottom sediments. Two sets of sediments were sampled from each location, one for macro fauna and other for Meio fauna. The macro fauna in the sediments were sieved on board to separate out the organisms. The fixation of Meio fauna is normally done by bulk fixation of the sediment sample. The bulk fixation is done by using 10% formalin (Buffered with borate). The organisms were preserved with seawater as diluting agent.

### **Sample sieving:**

Sediments samples were sieved to extract the organisms. Sieving was performed carefully as possible to avoid any damage to the animals. The large portion of the sediment was split in to smaller portions and mixed with sea water in a bucket. The cohesive lumps were broken down by continuous stirring. The disaggregated sediments were then passed through the sieves.

### **Sample staining:**

**DCPL/DPT/20-21/19 -NOVEMBER - 2021**

---

Sorting of the Meio fauna from the sieve is difficult task especially in the preserved material, because organisms are not easily detectable. To facilitate the animal detection the entire sample retained on the sieve after sieving operation were stained by immersing the sieve in a flat bottom tub with 1% Rose Bengal stain; a protein stain. A staining period of 10-30 minutes is sufficient for sample detection.

### **DIVERSITY INDICES:**

On the whole, diversity indices provide more information about community composition than simply species richness (number of species present); they also, take the relative abundances of different species into account. Based on this fact, diversity indices therefore depend not only on species richness but on the evenness, or equitability, with which individuals are distributed among the different species (Magurram, A. E. (1988)

A diversity index is a measure of species diversity within a community that consists of co-occurring populations of several (two or more) different species. It includes two components: richness and evenness. Richness is the measure of the number of different species within a sample showing that more the types of species in a community, the higher is the diversity or greater is the richness. Evenness is the measure of relative abundance of the different species with in a community.

The basic idea of diversity index is to obtain a quantitative estimate of biological variability that can be used to compare biological entities composed of discrete components in space and time (Carol H.R. *eta/*. 1998). Biodiversity is commonly expressed through indices based on species richness and species abundances (Whittaker 1972, Lande 1996, Purvis and Hector 2000). Biodiversity indices are a non-parametric tool used to describe the relationship between species number and abundance. The most widely used bio diversity indices are Shannon Weiner index and Simpson's index.

A diversity Index is a single statistic that incorporates in formation on richness and evenness. The diversity measures that incorporate the two concepts may be termed heterogeneity measures (Magurran, 2004).

Any study intended to interpret causes and effect of adverse impact on Biodiversity of communities require suitable measures to evaluate specie richness and Diversity. The former is number of species in community, while latter is a function of relative frequency of different species. Species richness is the iconic measure of biological diversity (Magurran, 2004). Several indices have been created to measure the diversity of species; however, the most widely used

in the last decades are the Shannon (1948) and Simpson (1949) (Buzas and Hayek 1996; Gorelick 2006), with the components of diversity: richness (*S*) and evenness (*J*)

### **Simpson's diversity index**

Simpson's index (**D**) is a measure of diversity, which takes into account both species richness, and evenness of abundance among the species present.. The Simpson index is one of the meaningful and robust biodiversity measures available.(Magurran ,2004).

The formula for calculating D is presented as:

$$D = \frac{\sum n_i(n_i - 1)}{N(N - 1)}$$

Where  $n_i$  = the total number of organisms of each individual species

$N$  = the total number of organisms of all species

The value of D ranges from 0 to 1. With this index, 0 represents infinite diversity and, 1, no diversity. When D increases diversity decreases. Simpson's index is therefore usually expressed as 1-D or 1/D. (Magurran, 2004)

Low species diversity suggests:

- relatively few successful species in the habitat

- the environment is quite stressful with relatively few ecological niches and only a few organisms are really well adapted to that environment

- food webs which are relatively simple

- change in the environment would probably have quite serious effects

High species diversity suggests:

- a greater number of successful species and a more stable ecosystem

- more ecological niches are available and the environment is less likely to be hostile complex food webs

- environmental change is less likely to be damaging to the ecosystem as a whole

### **Species richness indices**

The species richness (**S**) is simply the number of species present in an ecosystem. Species richness Indices of species richness are widely used to quantify or monitor the effects of anthropogenic disturbance. A decline in species richness may be concomitant with severe or chronic human-induced perturbation (Fair weather 1990,) Species richness measures have traditionally been the mainstay in assessing the effects of environmental degradation on the biodiversity of natural assemblages of organisms (Clarke &Warwick, 2001)

Species richness is the iconic measure of biological diversity (Magurran, 2004). The species richness (**S**) is simply the number of species present in an ecosystem. This index makes no use of relative abundances. The term species richness was coined by McIntosh (1967) and oldest and most intuitive measure of biological diversity (Magurran, 2004).

Margalef's diversity index is a species richness index. Margalef's Species richness index (d), or indices that describe the evenness of the distribution of the numbers of individuals among species, were derived.

The value of a diversity index increases both when the number of types increases and when evenness increases. For a given number of types, the value of diversity index is maximised when all types are equally abundant (Rosenzweig, M. L. (1995).

### **Shannon-Wiener's index:**

An index of diversity commonly used in plankton community analyses is the Shannon-Wiener's index (**H**), which emphasizes not only the number of species (richness or variety), but also the apportionment of the numbers of individuals among the species (Odum 1971 and Reish 1984). Shannon-Wiener's index (**H**) reproduces community parameters to a single number by using an equation.

Shannon and Weiner index represents entropy. It is a diversity index taking into account the number of individuals as well as the number of taxa. It varies from 0 for communities with only single taxa to high values for community with many taxa each with few individuals. This index can also determine the pollution status of a water body. Normal values range from 0 to 4. This index is a combination of species present and the evenness of the species. Examining the diversity in the range of polluted and unpolluted ecosystems, Wilham and Dorris (1968) concluded that the values of the index greater than 3 indicate clean water, values in the range of 1 to 3 are characterized by moderate pollution and values less than 1 are characterized as heavily polluted

$$H' = - \sum_{j=1}^s \frac{n_j}{N} \ln \left( \frac{n_j}{N} \right)$$

**RESULTS:**

**CHLOROPHYLL-a:**

Water Samples for the chlorophyll estimation were collected from sub surface layer during high tide and low tide period of the tidal cycle for each sampling locations and analysed for Chlorophyll -a and after acidification for Pheophytin -a. Chlorophyll- a value was used as algal biomass indicator (APHA,1998) Algal biomass was estimated by converting Chlorophyll value.

In the sub surface water chlorophyll-a was varying from 0.559 -0.868 mg/m<sup>3</sup>.in harbour region of DPT in Kandla Creek during sampling done in spring tide period of November, 2021. In the nearby creeks chlorophyll-a was varying from 0.246 -0.954 mg/m<sup>3</sup>.Pheophytin -a level was below detectable limit- the all the sampling stations during springtide

In the sub surface water chlorophyll-a was varying from 0.535 -0.921mg/m<sup>3</sup>.in harbour region of DPT in Kandla Creek during sampling done in neap tide period of November , 2021 . In the nearby creeks chlorophyll-a was varying from 0.425 -1.923 mg/m<sup>3</sup>.Pheophytin -a level was below detectable limit- the all the sampling stations except KPT-4 Khori-I during low tide and high tide and KPT-5 Nakti-I during High tide period.

In the sub surface water chlorophyll-a was varying from 0.393 -0.338 mg/m<sup>3</sup>.in harbour region of DPT OOT in path finder Creek during sampling done in spring tide period of November, 2021. In the sub surface water chlorophyll-a was varying from 1.356 -0.500 mg/m<sup>3</sup>.in harbour region of DPT OOT in path finder Creek during sampling done in Neap Tide period of November, 2021

In the sub surface water chlorophyll-a was varying from 0.424 -0.290 mg/m<sup>3</sup>.in SPM region of DPT OOT in path finder Creek during sampling done in spring tide period of November, 2021. In the sub surface water chlorophyll-a was varying from 0.703 -0.409 mg/m<sup>3</sup>.in SPM region of DPT OOT in path finder Creek during sampling done in Neap Tide period of November, 2021

**TABLE #2 VARIATIONS IN CHLOROPHYLL -a PHEOPHYTIN- a AND ALGAL BIOMASS FROM SAMPLING STATIONS IN DPT HARBOUR AREA IN KANDLA CREEK ,NEAR BY CREEKS AND DPT OOT JETTY IN PATH FINDER CREEK AND SPM NEAR VADINAR DURING SPRING TIDE IN NOVEMBER,2021**

Sr. No	Station	Tide	Chlorophyll-a (mg/m <sup>3</sup> )	Pheophytin-a (mg/m <sup>3</sup> )	Algal Biomass (Chlorophyll method) mg/m <sup>3</sup>
DPT HARBOUR AREA KANDLA CREEK					
1	KPT1	High tide	0.748	BDL	50.15
		Low tide	0.559	BDL	37.45
2	KPT 2	High tide	0.677	BDL	45.36
		Low tide	0.764	BDL	51.19
3	KPT 3	High tide	0.835	BDL	55.94
		Low tide	0.868	BDL	58.16
CREEKS					
4	KPT-4 Khor-i	High tide	0.661	BDL	44.29
		Low tide	0.720	BDL	48.24
5	KPT-5 Nakti-I	High tide	0.848	BDL	56.82
		Low tide	0.954	BDL	63.92
6	KPT-5 Nakti-II	High tide	0.246	BDL	16.48
PATHFINDER CREEK VADINAR					
7	VADINAR-I jetty	Low tide	0.393	BDL	26.33
8		High tide	0.338	BDL	22.65
9	SPM	High tide	0.424	BDL	28.41
10	SPM	Low tide	0.290	BDL	19.43

BDL: Below Detectable Limit.

**TABLE #3 VARIATIONS IN CHLOROPHYLL -a PHEOPHYTIN- a AND ALGAL BIOMASS FROM SAMPLING STATIONS IN DPT HARBOUR AREA , NEAR BY CREEKS AND DPT OOT JETTY IN PATH FINDER CREEK AND SPM NEAR VADINAR DURING NEAP TIDE IN NOVEMBER,2021**

Sr. No .	Station	Tide	Chlorophyll-a (mg/m <sup>3</sup> )	Pheophytin-a (mg/m <sup>3</sup> )	Algal Biomass (Chlorophyll method) mg/m <sup>3</sup>
DPT HARBOUR AREA KANDLA CREEK					
1	KPT1	High tide	0.748	BDL	50.12
		Low tide	0.535	BDL	35.85
2	KPT 2	High tide	0.713	BDL	47.77
		Low tide	0.713	BDL	47.77
3	KPT 3	High tide	0.882	BDL	59.09
		Low tide	0.921	BDL	61.71
CREEKS					
4	KPT-4 Khori-I	High tide	1.669	0.484	111.82
		Low tide	1.178	0.380	78.93
5	KPT-5 Nakti-I	High tide	1.923	0.570	128.84
		Low tide	0.882	BDL	59.09
6	KPT-5 Nakti-II	High tide	0.425	BDL	28.47
PATHFINDER CREEK VADINAR					
7	VADINAR-I jetty	Low tide	1.356	0.415	90.85
8		High tide	0.500	BDL	33.50
9	SPM	High tide	0.703	BDL	47.10
10	SPM	Low tide	0.409	BDL	27.40

BDL: Below Detectable Limit.

**PHYTOPLANKTON POPULATION:**

For the evaluation of the Phytoplankton population in DPT harbour area and within the immediate surroundings of the port, sampling was conducted from 5 sampling locations (3 in harbour area and two in Nakti creek) during high tide period and low tide period of spring tide and neap tide.

The phytoplankton community of the sub surface water in the harbour and nearby creeks was represented by; Diatoms blue green algae and dinoflagellates during spring tide period. Diatoms were represented by 16 genera. Blue green were represented by 2 genera and dinoflagellates were represented by two genera during the sampling conducted in spring tide in November, 2021. Phytoplankton of the sampling stations at sub surface layer in the harbour area

and nearby creeks was varying from 43-198 units/ L during high tide period and 133-220 units/ L during low tide of Spring Tide.

The phytoplankton community of the sub surface water in the harbour and nearby creeks was represented by Diatoms, Blue green algae and Dinoflagellates during Neap tide period. Diatoms were represented by 20 genera Blue green algae were represented 1 genera and Dinoflagellates with two genera during the sampling conducted in Neap tide in November, 2021. Phytoplankton of the sampling stations at sub surface layer in the harbour area and nearby creeks was varying from 83-327 units/ L during high tide period and 108 -252 units/ L during low tide of Neap Tide.

For the evaluation of the Phytoplankton population in DPT OOT jetty area in Path Finder creek sampling was conducted from one sampling locations ; jetty area during high tide period and low tide of spring tide. For the evaluation of the Phytoplankton population in DPTOOT jetty area in Path Finder creek in Vadinar sampling was conducted from 2 sampling locations ; jetty area and one near SPM during high tide period and low tide of Neap tide.

The phytoplankton community of the sub surface water in the path finder creeks was represented by only Diatoms spring tide period. Diatoms were represented by 11 genera during the sampling conducted in spring tide in November, 2021. Phytoplankton of the sampling stations at sub surface layer in the harbour area was varying from 162 units/ L during high tide period and 178 units/ L during low tide of Spring Tide. Phytoplankton of the sampling stations at sub surface layer in the SPM area was varying from 154 units/ L during high tide period and 130 units/ L during low tide of Spring Tide.

The phytoplankton community of the sub surface water in the path finder creeks was represented by Diatoms, and Dinoflagellates during Neap tide period. Diatoms were represented by 15 genera and dinoflagellates by two genera during the sampling conducted in Neap tide in November, 2021. Phytoplankton of the sampling stations at sub surface path finder creek near OOT Jetty was varying from 227 units/ L during high tide period and 182 -744 units/ L during low tide of Neap Tide. Phytoplankton of the sampling stations at sub surface path finder creek near SPM area was varying from 158 units/ L during high tide period and 158 units/ L during low tide of Neap Tide.

### **Species Richness Indices and Diversity Indices:**

#### **Margalef's diversity index (Species Richness) S**



At the organismal level, the most widely used biodiversity measures are those based on the number of species present, perhaps adjusted for the number of individuals sampled, Here Margalef's Species richness index (d), or indices that describe the evenness of the distribution of the numbers of individuals among species, are derived.

Margalef's diversity index (Species Richness) S of phytoplankton communities in the Kandla creek and nearby creeks sampling stations was varying from 1.595-3.091 with an average of 2.396 during the sampling conducted in High tide period of spring tide. While Margalef's diversity index (Species Richness) S of phytoplankton communities in the Kandla creek region and nearby creeks was varying from 2.236 -2.863 with an average of 2.554 during the consecutive low tide period.

Margalef's diversity index (Species Richness) S of phytoplankton communities in the stations in Kandla creek and nearby creeks was varying from 2.339-2.984 with an average of 2.696 during the sampling conducted in High tide period of Neap tide. While Margalef's diversity index (Species Richness) S of phytoplankton communities in the Kandla creek region and nearby creeks was varying from 2.450-2.713 with an average of 2.624 during consecutive low tide.

Margalef's diversity index (Species Richness) S of phytoplankton communities in the stations was 1.769 at OOT jetty area and 1.588 at SPM area during the sampling conducted in High tide period of spring tide. While Margalef's diversity index (Species Richness) S of phytoplankton communities in the path finder creek near OOT jetty was 1.737 and 1.644 at SPM during the consecutive low tide period.

Margalef's diversity index (Species Richness) S of phytoplankton communities in the stations was 2.561 at OOT jetty area and 2.370 at SPM area during the sampling conducted in High tide period of Neap tide. While Margalef's diversity index (Species Richness) S of phytoplankton communities in the path finder creek near OOT jetty was 2.114 and SPM area was 2.195 during the consecutive low tide period.

### **Shannon-Wiener's index:**

Shannon-Wiener's Index (H) of phytoplankton communities in the sampling stations was in the range of 0.727-0.907 ( $H'(\log_{10})$ ) between selected sampling stations with an average value of 0.805 during high tide period of spring tide at Kandla creek and nearby creeks. Shannon-Wiener's Index (H) of phytoplankton communities in the sampling stations was in the range of 0.787 -0.895 ( $H'(\log_{10})$ )

**DCPL/DPT/20-21/19 -NOVEMBER - 2021**

---

between selected sampling stations with an average value of 0.853 during consecutive low tide at Kandla creek and nearby creeks. Shannon-Wiener's Index (H) of phytoplankton communities in the sampling stations was in the range of 0.695 -0.931 ( $H'(\log_{10})$ ) between selected sampling stations with an average value of 0.823 during high tide period of neap tide at Kandla creek and nearby creeks. Shannon-Wiener's Index (H) of phytoplankton communities in the sampling stations was in the range of 0.759-0.867 ( $H'(\log_{10})$ ) between selected sampling stations with an average value of 0.810 during consecutive low tide at Kandla creek and nearby creeks.

Shannon-Wiener's Index (H) of phytoplankton communities in the stations was 0.798 at OOT jetty area and 0.7551 at SPM area during the sampling conducted in High tide period of spring tide. While Shannon-Wiener's Index (H) of phytoplankton communities in the path finder creek near OOT jetty was 0.715 and 0.771 at SPM during the consecutive low tide period.

Shannon-Wiener's Index (H) of phytoplankton communities in the stations was 0.787 at OOT jetty area and 0.7330 at SPM area during the sampling conducted in High tide period of Neap tide. While Shannon-Wiener's Index (H) of phytoplankton communities in the path finder creek near OOT jetty was 0.729 and at SPM area was 0.712 during the consecutive low tide period.

Typical values are generally between 1.5 and 3.5 in most ecological studies, and the index is rarely greater than 4. The Shannon-Wiener's index increases as both the richness and the evenness of the community increase. This result indicates that diversity of phytoplankton of Kandla Harbour region and nearby creeks is less but with abundant population of few, with relatively few ecological niches and only very few opportunist organisms are really well adapted to this environment and thrive better than other species.

#### **Simpson's diversity index:**

Simpson's index (D) is a measure of diversity, which takes into account both species richness, and an evenness of abundance among the species present. The Simpson index is one of the meaningful and robust biodiversity measures available. (Magurran,2004).

Simpson diversity index (1-D) of phytoplankton communities was below 0.9 at all sampling stations in the Kandla Harbour region and nearby creeks, which was varying from 0.742- 0.830 between selected sampling stations with an average of 0.788 during high tide period of spring tide. Simpson diversity index (1-D) of phytoplankton communities was below 0.9 at all sampling stations in the Kandla Harbour region and nearby creeks, which was varying from 0.786- 0.832

between selected sampling stations with an average of 0.809 during consecutive low tide .

Simpson diversity index (1-D) of phytoplankton communities was below 0.9 at all sampling stations in Kandla Harbour region and nearby creeks, during high tide period and low tide period during neap tide also, which was varying from 0.664-0.841 with an average value of 0.774 between selected sampling stations during high tide period and varying from 0.732-0.824 with an average value of 0.771 between selected sampling stations during consecutive low tide period. Low species diversity suggests a relatively few successful species in this habitat.

Simpson diversity index (1-D) of phytoplankton communities in the stations was 0.813 at OOT jetty area and 0.779 at SPM area during the sampling conducted in High tide period of spring tide at Path finder creek . While Simpson diversity index (1-D) of phytoplankton communities in the path finder creek near OOT jetty was 0.753 and 0.794 at SPM during the consecutive low tide period in the path finder creek.

Simpson diversity index (1-D) of phytoplankton communities in the stations was 0.765 at OOT jetty area and 0.737 at SPM area during the sampling conducted in High tide period of Neap tide at Path finder Creek. While Simpson diversity index (1-D) of phytoplankton communities in the path finder creek near OOT jetty was 0.738 and at SPM area was 0.708 during the consecutive low tide period.

Low species diversity suggests a relatively few successful species in this habitat. The environment is quite stressful with relatively few ecological niches and only a few organisms are really well adapted to that environment. Any change in the environment would probably have quite serious effects.

**Table # 4 PHYTOPLANKTON VARIATIONS IN ABUNDANCE AND DIVERSITY  
IN SUB SURFACE SAMPLING STATIONS IN DPT HARBOUR AREA AT  
KANDLA CREEK AND ,NEAR BY CREEKS DURING SPRING TIDE IN  
NOVEMBER,2021**

Tide	Samplin g Station	Abundanc e In units/L	No of Species observe d /total species	% of diversit y	Margalef 's diversity index (Species Richness S)	Shanno n Weiner index H (log <sub>10</sub> )	Diversity Index (Simpson' s Index) 1-D
HIGH TIDE	1	177	17/20	85	3.091	0.9004	0.8193
	2	152	16/20	80	2.986	0.9067	0.8305
	3	192	11/20	55	1.902	0.7268	0.7421
	4	167	13/20	65	2.345	0.7525	0.7454
	5	198	14/20	70	2.458	0.7886	0.7683
	6	43	7/20	35	1.595	0.7583	0.825
LOW TIDE	1	133	15/20	75	2.863	0.8948	0.8214
	2	153	15/20	75	2.783	0.893	0.832
	3	190	14/20	70	2.478	0.8679	0.8106
	4	137	12/20	60	2.236	0.8213	0.7966
	5	220	14/20	70	2.41	0.7872	0.786

**Table # 5 PHYTOPLANKTON VARIATIONS IN ABUNDANCE AND DIVERSITY  
IN SUB SURFACE SAMPLING STATIONS IN DPT HARBOUR AREA AT  
KANDLA CREEK AND ,NEAR BY CREEKS DURING NEAP TIDE IN  
NOVEMBER,2021**

Tide	Samplin g Station	Abundan ce In units/L	No of Species observe d /total species	% of diversit y	Margalef 's diversity index (Species Richness S)	Shanno n Weiner index H (log <sub>10</sub> )	Diversity Index (Simpson' s Index) 1-D
HIGH TIDE	1	131	15/24	62.5	2.872	0.9308	0.8406
	2	120	13/24	54.16	2.507	0.8561	0.8136
	3	213	17/24	70.83	2.984	0.8315	0.7733
	4	259	14/24	58.33	2.339	0.7394	0.7223
	5	327	17/24	70.83	2.763	0.6955	0.6641
	6	83	13/24	54.16	2.716	0.8861	0.8316
LOW TIDE	1	108	13/24	54.16	2.563	0.791	0.7606
	2	134	13/24	54.16	2.45	0.8677	0.8239
	3	177	15/24	62.5	2.705	0.7892	0.7325
	4	252	16/24	66.66	2.713	0.7591	0.7444
	5	182	15/24	62.5	2.69	0.8458	0.7939

**Table # 6 ABUNDANCE OF PHYTOPLANKTON SUBSURFACE SAMPLING STATIONS IN DPT HARBOUR AREA AT KANDLA CREEK AND , NEAR BY CREEKS DURING SPRING TIDE IN NOVEMBER, 2021**

Tide	Surface	No of Sampling location	Group of phytoplankton	Phytoplankton Group range Units/L	Genera or species /total Phytoplankton	Taxon Diversity % (Group level)
HIGH TIDE	Sub surface	6	DIATOMS	41-197	16/20	80
			BLUE GREEN	0-4	2/20	10
			DINOFLAGELLATES	0-3	2/20	10
			TOTAL PHYTOPLANKTON	<b>43-198</b>	<b>20</b>	-
LOW TIDE	Sub surface	5	DIATOMS	129-216	16/20	80
			BLUE GREEN	0-4	2/20	10
			DINOFLAGELLATES	0-2	2/20	10
			TOTAL PHYTOPLANKTON	<b>133-220</b>	<b>20</b>	-

**Table # 7 ABUNDANCE OF PHYTOPLANKTON SUBSURFACE SAMPLING STATIONS IN DPT HARBOUR AREA AT KANDLA CREEK AND , NEAR BY CREEKS DURING NEAP TIDE IN NOVEMBER,2021**

Tide	Surface	No of Sampling location	Group of phytoplankton	Phytoplankton Group range Units/L	Genera or species /total Phytoplankton	Taxon Diversity % (Group level)
HIGH TIDE	Sub surface	6	DIATOMS	81-326	21/24	87.6
			BLUE GREEN	0-2	1/24	4.16
			DINOFLAGELLATES	0-2	2/24	8.33
			TOTAL PHYTOPLANKTON	<b>83-327</b>	<b>24</b>	
LOW TIDE	Sub surface	5	DIATOMS	108-251	21/24	87.6
			BLUE GREEN	0-2	1/24	4.16
			DINOFLAGELLATES	0-1	2/24	8.33
			TOTAL PHYTOPLANKTON	<b>108-252</b>	<b>24</b>	

**Table # 8 PHYTOPLANKTON VARIATIONS IN ABUNDANCE AND DIVERSITY  
IN SUB SURFACE SAMPLING STATIONS IN DPT OOTAT PATH FINDER  
CREEK , VADINAR & NEAR BY SPM, DURING SPRING TIDE IN NOVEMBER,  
2021**

Tide	Samplin g Station	Abundan ce In units/L	No of Species observe d /total species	% of diversit y	Margalef' s diversity index (Species Richness S)	Shanno n Weiner index H (log <sub>10</sub> )	Diversity Index (Simpson' s Index) 1-D
HIGH TIDE	jetty	162	10/11	90.90	1.769	0.7989	0.8132
	SPM	178	10/11	90.90	1.737	0.7149	0.7536
LOW TIDE	jetty	154	9/11	81.82	1.588	0.7441	0.7796
	SPM	130	9/11	81.82	1.644	0.7712	0.7937

**Table # 9 PHYTOPLANKTON VARIATIONS IN ABUNDANCE AND DIVERSITY  
IN SUB SURFACE SAMPLING STATIONS IN DPT OOTAT PATH FINDER  
CREEK , VADINAR & NEAR BY SPM, DURING NEAP TIDE IN NOVEMBER,  
2021**

Tide	Samplin g Station	Abundance In units/L	No of Species observe d /total species	% of diversi ty	Margalef' s diversity index (Species Richness S)	Shanno n Weiner index H (log <sub>10</sub> )	Diversity Index (Simpson' s Index) 1-D
HIGH TIDE	Jetty	227	15/17	88.24	2.581	0.7875	0.7647
	SPM	182	12/17	70.59	2.114	0.7288	0.7383
LOW TIDE	Jetty	158	13/17	76.47	2.37	0.733	0.7374
	SPM	150	12/17	70.59	2.195	0.7123	0.7087


**Table # 10 ABUNDANCE OF PHYTOPLANKTON SUBSURFACE SAMPLING STATIONS IN DPT DPTOOT AT PATH FINDER CREEK , VADINAR & NEAR BY SPM, DURING SPRING TIDE IN NOVEMBER, 2021**

Tide	Surface	No of Sampling location	Group of phytoplankton	Phytoplankton Group range Units/L	Genera or species /total Phytoplankton	Taxon Diversity % (Group level)
HIGH TIDE	Sub surface	1	DIATOMS	162-178	11/11	100
			TOTAL PHYTOPLANKTON	162-178	<b>11</b>	
LOW TIDE	Sub surface	1	DIATOMS	130-154	11/11	100
			TOTAL PHYTOPLANKTON	130-154	<b>11</b>	


**Table # 11 ABUNDANCE OF PHYTOPLANKTON SUBSURFACE SAMPLING STATIONS IN DPT DPTOOT AT PATH FINDER CREEK , VADINAR & NEAR BY SPM, DURING NEAP TIDE IN NOVEMBER, 2021**

Tide	Surface	No of Sampling location	Group of phytoplankton	Phytoplankton Group range Units/L	Genera or species /total Phytoplankton	Taxon Diversity % (Group level)
HIGH TIDE	Sub surface	2	DIATOMS	182-226	15/17	88.24
			DINOFLAGELLATES	0-1	2/17	11.76
			TOTAL PHYTOPLANKTON	<b>182-227</b>	<b>17</b>	
LOW TIDE	Sub surface	2	DIATOMS	148-157	15/17	88.24
			DINOFLAGELLATES	0-1	2/17	11.76
			TOTAL PHYTOPLANKTON	148-158	<b>17</b>	

**Taxon Diversity % of Phytoplankton during High tide and Low tide period during spring tide in Kandala creek and nearby creeks**




**Taxon Diversity % of Phytoplankton during High tide and Low tide  
period during Neap tide in Kandala creek and nearby creeks**




**Taxon Diversity % of Phytoplankton during High tide and Low tide  
period during spring tide in Path Finder Creek, Vadinar**





**Taxon Diversity % of Phytoplankton during High tide and Low tide  
period during Neap tide in Path Finder Creek, Vadinar**



**ZOOPLANKTON POPULATION:**

For the evaluation of the Zooplankton population in DPT harbour area and within the immediate surroundings of the port sampling was conducted from 6 sampling locations (3 in harbour area and two in Nakti creek and one in Khoricreek) during high tide period and low tide period of spring tide and Neap tide in November,2021. The Zooplankton community of the sub surface water in

**DCPL/DPT/20-21/19 -NOVEMBER - 2021**

the harbour and nearby creeks during spring tide was represented by mainly 4 groups, and 5 larval forms; Tintinids, Copepods, Rotifers, Urochordates and larval forms represented from the group of Crustacea, Molluscs and Polychaetes. The Zooplankton community of the sub surface water in the harbour and nearby creeks during neap tide was represented by mainly Six groups, Tintinids, Copepods, Arrow worms, Mysids, Urochordata, Ciliates and unidentified **Cnidarian member and** larval forms of Crustacea Molluscs and Echinodermata Larvae Polychaete Larvae..,

Zooplankton of the sampling stations at sub surface layer in the DPT harbour area and nearby creek was varying from 25-106x10<sup>3</sup> N/ m<sup>3</sup> during high tide and 58-85x10<sup>3</sup> N/ m<sup>3</sup> during low tide of Spring Tide period. Zooplankton of the sampling stations at sub surface layer in the DPT harbour area and nearby creek was varying from 40-143 x10<sup>3</sup> N/ m<sup>3</sup> during high tide and 83-129x10<sup>3</sup> N/ m<sup>3</sup> during low tide of Neap Tide period.

For the evaluation of the Zoo plankton population in DPT OOT jetty area in Path Finder creek and SPM in Vadinar selected 2 sampling locations (1 in jetty area and one near SPM) During spring tide sampling plankton sample were collected only at Jetty area during consecutive high tide period and low tide period. During Neap tide sampling Plankton samples were collected from jetty area and SPM during consecutive high tide period and low tide period.

The Zooplankton community of the sub surface water in the path finder creek during spring tide was represented by mainly Tintinids , Copepods and larval forms of Crustaceans, Molluscs and Polychaetes .The Zooplankton community of the sub surface water in the path Finder creeks at Jetty region and SPM during neap tide was represented by mainly three groups, Tintinids, Copepods , Urochordates and , five Larval forms were represented from the major group of Crustaceans , Molluscs , and Polychaetes..

Zooplankton of the sampling stations at sub surface layer in the DPT OOT jetty area of path finder creek was 87 x10<sup>3</sup> N/ m<sup>3</sup> during high tide and 117 x10<sup>3</sup> N/ m<sup>3</sup> during low tide of Spring Tide period. Zooplankton of the sampling stations at sub surface layer in the DPT SPM area of path finder creek was 85 x10<sup>3</sup> N/ m<sup>3</sup> during high tide and 109 x10<sup>3</sup> N/ m<sup>3</sup> during low tide of Spring Tide period. Zooplankton of the sampling stations at sub surface layer in the DPT OOT jetty area in path finder creek was recorded 54x10<sup>3</sup> N/ m<sup>3</sup> during high tide and 86x10<sup>3</sup> N/ m<sup>3</sup> during consecutive low tide period of Neap . Zooplankton of the sampling

stations at sub surface layer in the DPT SPM area in path finder creek was recorded  $72 \times 10^3$  N/  $m^3$  during high tide and  $92 \times 10^3$  N/  $m^3$  during consecutive low tide period of Neap Tide .

### **Species Richness Indices and Diversity Indices: Margalef's diversity index (Species Richness)S**

At the organismal level, the most widely used biodiversity measures are those based on the number of species present, perhaps adjusted for the number of individuals sampled, Here Margalef's Species richness index (d), or indices that describe the evenness of the distribution of the numbers of individuals among species, are derived.

Margalef's diversity index (Species Richness) S of Zooplankton communities in the stations Kandla creek region and nearby creeks was varying from 1.733-2.796 with an average of 2.196 during the sampling conducted in High tide period. Margalef's diversity index (Species Richness) S of Zooplankton communities varying from 1.871-2.217 with an average of 2.026 during the sampling conducted in low tide period during Spring tide.

Margalef's diversity index (Species Richness) S of Zooplankton communities in the Kandla creek region and nearby creeks sampling stations was varying from 2.771-3.983 with an average of 3.445 during the sampling conducted in high tide and varying from 2.635- 3.054 with an average of 3.049 during the sampling conducted in low tide during Neap tide period

Margalef's diversity index (Species Richness) S of Zooplankton communities in the sampling station near jetty at Path Finder Creek, Vadinar during the sampling conducted in consecutive High tide period and low tide of spring tide was recorded as 1.769 and 1.588 respectively..

Margalef's diversity index (Species Richness) S of Zooplankton communities near SPmat Path finder creek was varying from 2.256-2.572 during the sampling conducted in High tide period of Neap tide. While Margalef's diversity index (Species Richness) S of Zooplankton communities in the two stations at Path finder creek was varying from 2.020- 1.769 during the consecutive low tide period.

### **Shannon-Wiener's index:**

Shannon-Wiener's Index (H) of Zooplankton communities in the sampling stations in Kandla Harbour region and nearby creeks was in the range of 0.725-0.945 ( $H'(\log_{10})$ ) between selected sampling stations with an average value of 0.811 ( $H'(\log_{10})$ ) during high tide period of spring tide. Shannon-Wiener's Index (H) of Zooplankton communities in the sampling stations in Kandla Harbour

region and nearby creeks was in the range of 0.703-0.884 ( $H'(\log_{10})$ ) between selected sampling stations with an average value of 0.780 ( $H'(\log_{10})$ ) during consecutive low tide period.

Shannon-Wiener's Index (H) of Zooplankton communities in the sampling stations in Kandla Harbour region and nearby creeks was in the range of 0.782-1.119 ( $H'(\log_{10})$ ) between selected sampling stations with an average value of 1.000 ( $H'(\log_{10})$ ) during high tide period of Neap tide. Shannon-Wiener's Index (H) of Zooplankton communities in the sampling stations in Kandla Harbour region and nearby creeks was in the range of 0.855-1.059 ( $H'(\log_{10})$ ) between selected sampling stations with an average value of 0.950 ( $H'(\log_{10})$ ) during consecutive low tide period.

Shannon-Wiener's Index (H) of Zooplankton communities in the sampling station near jetty at Path Finder Creek, Vadinar during the sampling conducted in consecutive High tide period and low tide of spring tide was recorded as 0.799 and 0.7441 respectively.. Shannon-Wiener's Index (H) of Zooplankton communities in the sampling station near jetty at Path Finder Creek, Vadinar during the sampling conducted in consecutive High tide period and low tide of Neap tide was recorded as 0.742 and 0.709 respectively

Shannon-Wiener's Index (H) of Zooplankton communities in the two stations at Path finder creek was varying from 0.715-0.798 during the sampling conducted in High tide period of Spring tide. While Shannon-Wiener's Index (H) of Zooplankton communities in the two stations at Path finder creek was varying from -0.771-0.7441 during the consecutive low tide period.

Shannon-Wiener's Index (H) of Zooplankton communities in the two stations at Path finder creek was varying from 0.743-0.849 during the sampling conducted in High tide period of Neap tide. While Shannon-Wiener's Index (H) of Zooplankton communities in the two stations at Path finder creek was varying from -0.641 - 0.709 during the consecutive low tide period.

Typical values are generally between 1.5 and 3.5 in most ecological studies, and the index is rarely greater than 4. The Shannon-Wiener's index increases as both the richness and the evenness of the community increase. This result indicates that diversity of Zooplankton of Kandla Harbour region and nearby creeks stations is slightly high with very minimum diverse population but very few opportunist organisms are really well adapted to this environment and thrive better than other species.

**Simpson's diversity index:**

Simpson's index (D) is a measure of diversity, which takes into account both species richness, and an evenness of abundance among the species present. The Simpson index is one of the meaningful and robust biodiversity measures available. (Magurran, 2004).

Simpson diversity index (1-D) of Zooplankton communities was below 0.9 most of sampling stations in the Kandla Harbour region and nearby creeks during high tide and low tide of spring tide period, which was varying from 0.751-0.910 between selected sampling stations with an average of 0.804 during high tide period and was varying from 0.722- 0.854 with an average value of 0.780 between selected sampling stations during low tide

Simpson diversity index (1-D) of Zooplankton communities was below 0.9 at all sampling stations in the Kandla Harbour region and nearby creeks except few during high tide and low tide period, which was varying from 0.766 - 0.912 between selected sampling stations with an average of 0.863 during high tide period and was varying from 0.795- 0.896 with an average value of 0.843 between selected sampling stations during consecutive low tide

This low species diversity suggests a relatively few of successful species in this habitat during November, 2021 sampling.

Simpson diversity index (1-D) of Zooplankton communities in the sampling station near jetty at Path Finder Creek, Vadinar during the sampling conducted in consecutive High tide period and low tide of spring tide was recorded as 0.813 and 0.779 respectively. Simpson diversity index (1-D) of Zooplankton communities in the sampling station near SPM at Path Finder Creek, Vadinar during the sampling conducted in consecutive High tide period and low tide of spring tide was recorded as 0.753 and 0.779 respectively.

Simpson diversity index (1-D) of Zooplankton communities in the sampling station near jetty at Path Finder Creek, Vadinar during the sampling conducted in consecutive High tide period and low tide of Neap tide was recorded as 0.778 - 0.729 respectively. Simpson diversity index (1-D) of Zooplankton communities in the sampling station near SPM at Path Finder Creek, Vadinar during the sampling conducted in consecutive High tide period and low tide of spring tide was recorded as 0.817 and 0.697 respectively.

**Table # 12 ZOOPLANKTON VARIATION IN ABUNDANCE AND DIVERSITY  
IN SUB SURFACE SAMPLING STATIONS IN DPT HARBOUR AREA AT  
KANDLA CREEK AND NEAR BY CREEKS DURING SPRING TIDE IN  
NOVEMBER,2021**

Tide	Samplin g Station	Abundance In $N \times 10^3 /$ $m^3$	No of Species/gr oups observed /total species/gro up	% of diversi ty	Margalef' s diversity index (Species Richness S)	Shanno n Weiner index H ( $\log_{10}$ )	Diversity Index (Simpson' s Index) 1-D
HIGH TIDE	1	75	10/16	62.5	2.085	0.7569	0.751
	2	82	11/16	68.75	2.269	0.8385	0.8154
	3	66	10/16	62.5	2.148	0.8294	0.8224
	4	106	11/16	68.75	2.144	0.7752	0.7641
	5	101	9/16	56.25	1.733	0.7251	0.7651
	6	25	10/16	62.5	2.796	0.9451	0.91
LOW TIDE	1	69	9/16	56.25	1.889	0.8145	0.809
	2	58	10/16	62.5	2.217	0.8838	0.8542
	3	72	9/16	56.25	1.871	0.7031	0.7218
	4	69	10/16	62.5	2.126	0.7896	0.7899
	5	85	10/16	62.5	2.026	0.7112	0.7272

**Table # 13 ZOOPLANKTON VARIATION IN ABUNDANCE AND DIVERSITY  
IN SUB SURFACE SAMPLING STATIONS IN DPT HARBOUR AREA AT  
KANDLA CREEK AND NEAR BY CREEKS DURING NEAP TIDE IN NOVEMBER,**

Tide	Samplin g Station	Abundance In $No \times 10^3 /$ $m^3$	No of Species/gr oups observed /total species/gro up	% of diversi ty	Margalef' s diversity index (Species Richness S)	Shanno n Weiner index H ( $\log_{10}$ )	Diversity Index (Simpson' s Index) 1-D
HIGH TIDE	1	118	20/28	71.42	3.983	1.119	0.9122
	2	102	17/28	60.71	3.459	0.9987	0.875
	3	108	19/28	67.86	3.844	1.085	0.8974
	4	143	18/28	64.29	3.425	1.118	0.9087
	5	101	16/28	57.14	3.25	0.9028	0.8212
	6	40	11/28	39.29	2.711	0.7823	0.7667
LOW TIDE	1	83	13/28	46.43	2.716	0.8552	0.7949
	2	128	18/28	64.29	3.504	1.059	0.8958
	3	129	18/28	64.29	3.498	1.055	0.8815
	4	89	14/28	50	2.896	0.8648	0.7975
	5	95	13/28	46.43	2.635	0.9189	0.8434

**DCPL/DPT/20-21/19 -NOVEMBER - 2021**

**Detox Corporation Pvt. Ltd., Surat**

**Environmental Monitoring Report of Deendayal Port Trust,  
November-2021**

**2021**

**Table # 14 ABUNDANCE OF ZOOPLANKTON IN SUBSURFACE SAMPLING STATIONS IN DPT HARBOUR AREA AT KANDLA CREEK AND , NEAR BY CREEKS DURING SPRING TIDE IN NOVEMBER, 2021**

Tide	Surface	No of Sampling locations	Group of Zooplankton	Abundance of Zooplankton $\times 10^3$ Group Range	Genera or species /total Zooplankton	Taxon Diversity % (Group level)
HIGH TIDE	Sub surface	6	Tintinids	3-13	3/16	18.75
			Copepods	11-40	6/16	37.5
			Rotifers	0-2	1/16	6.25
			Urochordata	1-4	1/16	6.25
			Larval forms	5-52	5/16	31.25
			TOTAL ZOOPLANKTON N/ M <sup>3</sup>	<b>25-106</b>	<b>16</b>	
LOW TIDE	Sub surface	5	Tintinids	5-9	3/16	18.75
			Copepods	20-27	6/16	37.5
			Rotifers	0	1/16	6.25
			Urochordata	0-4	1/16	6.25
			Larval forms	30-53	5/16	31.25
			TOTAL ZOOPLANKTON N/M <sup>3</sup>	<b>58-85</b>	<b>16</b>	

**Table # 15 ABUNDANCE OF ZOOPLANKTON IN SUBSURFACE SAMPLING STATIONS IN DPT HARBOUR AREA IN KANDLA CREEK AND , NEAR BY CREEKS DURING NEAP TIDE IN NOVEMBER, 2021**

Tide	Surface	No of Sampling locations	Group of Zooplankton	Abundance of Zooplankton $\times 10^3$ Group Range	Genera or species /total Zooplankton	Taxon Diversity % (Group level)
HIGH TIDE	Sub surface	6	Tintinids	7-36	5/28	17.86
			Copepods	11-49	8/28	28.58
			Mysids	0-1	2/28	7.14
			Arrow worms	0-2	1/28	3.57
			Urochordata	0-2	1/28	3.57
			Ciliates	0-4	1/28	3.57
			Medusa	0-4	1/28	3.57
			Larval forms	20-58	7/28	25
			Foraminiferans	0-4	2/28	7.14
			TOTAL ZOOPLANKTON N/M <sup>3</sup>	<b>40-143</b>	<b>28</b>	
LOW TIDE	Sub surface	5	Tintinids	10-32	5/28	17.86
			Copepods	17-54	8/28	28.58
			Mysids	0-2	2/28	7.14
			Arrow worms	0-1	1/28	3.57
			Urochordata	0-2	1/28	3.57
			Ciliates	0-1	1/28	3.57
			Medusa	0-1	1/28	3.57
			Larval forms	50-62	7/28	25

**DCPL/DPT/20-21/19 -NOVEMBER - 2021**

**Detox Corporation Pvt. Ltd., Surat**



**Environmental Monitoring Report of Deendayal Port Trust,  
November-2021**

		Foraminiferans	<b>0-3</b>	2/28	7.14
		Total Zooplankton N/M3		<b>28</b>	

**Table # 16 ZOOPLANKTON VARIATION IN ABUNDANCE AND DIVERSITY  
IN SUB SURFACE SAMPLING STATIONS IN DPT OOT AREA AT PATH  
FINDER CREEK AND NEAR BY SPM DURING SPRING TIDE IN  
NOVEMBER,2021**

Tide	Samplin g Station	Abundanc e In $\times 10^3 N / m^3$	No of Species/gro ups observed /total species/gro up	% of diversit y	Margalef' s diversity index (Species Richness S)	Shannon Weiner index H (log <sub>10</sub> )	Diversity Index (Simpso n's Index) 1-D
HIGH TIDE	Jetty	87	11/13	84.62	2.239	0.6821	0.6864
	SPM	85	12/13	92.31	2.476	0.7967	0.788
LOW TIDE	Jetty	117	10/13	76.92	1.89	0.7264	0.7265
	SPM	109	10/13	76.92	1.918	0.6599	0.6624

**Table # 17 ZOOPLANKTON VARIATION IN ABUNDANCE AND DIVERSITY  
IN SUB SURFACE SAMPLING STATIONS IN DPT OOT AREA AT PATH  
FINDER CREEK AND NEAR BY SPM DURING NEAP TIDE IN  
NOVEMBER,2021**

Tide	Sampling Station	Abundanc e In $N \times 10^3 / m^3$	No of Species/gro ups observed /total species/gro up	% of diversit y	Margalef' s diversity index (Species Richness S)	Shannon Weiner index H (log <sub>10</sub> )	Diversity Index (Simpso n's Index) 1-D
HIGH TIDE	Jetty	227	15/17	88.23	2.581	0.7875	0.7647
	SPM	182	12/17	70.59	2.114	0.7288	0.7383
LOW TIDE	Jetty	158	13/17	76.47	2.37	0.733	0.7374
	SPM	150	12/17	70.59	2.195	0.7123	0.7087

**Table # 18 ABUNDANCE OF ZOOPLANKTON IN SUBSURFACE SAMPLING  
STATIONS IN DPT OOT AREA AT PATH FINDER CREEK AND NEAR BY SPM  
DURING SPRING TIDE IN NOVEMBER,2021**

Tide	Surface	No of Sampling locations	Group of Zooplankton	Abundance of Zooplankton $\times 10^3$ Group Range	Genera or species /total Zooplankto n	Taxon Diversity % (Group level)
HIGH TIDE	Sub surface	2	Tintinids	6-7	3/13	23.08
			Copepods	30-39	6/13	46.15
			Larval forms	39-51	4/13	30.77
			TOTAL ZOOPLANKTON NO/L	<b>85-87</b>	<b>13</b>	
LOW TIDE	Sub surface	2	Tintinids	15-16	3/13	23.08
			Copepods	30-35	6/13	46.15
			Larval forms	67-73	4/13	30.77

**DCPL/DPT/20-21/19 -NOVEMBER - 2021**

**Detox Corporation Pvt. Ltd., Surat**




**Environmental Monitoring Report of Deendayal Port Trust,  
November-2021**

			TOTAL ZOOPLANKTON NO/M3	<b>109-117</b>	<b>13</b>	
--	--	--	----------------------------	----------------	-----------	--


**Table # 19 ABUNDANCE OF ZOOPLANKTON IN SUBSURFACE SAMPLING STATIONS IN DPT OOT AREA AT PATH FINDER CREEK AND NEAR BY SPM DURING NEAP TIDE IN NOVEMBER, 2021**

Tide	Surface	No of Sampling locations	Group of Zooplankton	Abundance of Zooplankton x10 <sup>3</sup> Group Range	Genera or species /total Zooplankton	Taxon Diversity % (Group level)
HIGH TIDE	Sub surface	2	Tintinids	6-9	4/17	23.53
			Copepods	25-31	7/17	41.18
			Urochordata	0-1	1/17	5.88
			Larval forms	23-41	5/17	29.41
			<b>TOTAL ZOOPLANKTON</b>	<b>48-63</b>	<b>17</b>	
LOW TIDE	Sub surface	2	Tintinids	9-10	4/16	25
			Copepods	43-47	7/16	43.75
			Urochordata	0	0	0
			Larval forms	43-47	5/16	31.25
			<b>TOTAL ZOOPLANKTON NO/M3</b>	<b>77-83</b>	<b>16</b>	


**Taxon Diversity % of Zooplankton during High tide and Low tide period of Spring tide In Kandla Creek and near by Creeks**




**Taxon Diversity % of Zooplankton during High tide and Low tide period  
of Neap tide In Kandla Creek and nearby Creeks**



**Taxon Diversity % of Zooplankton during High tide and Low tide period  
of Spring tide In Path Finder Creek and near Jetty**



**Taxon Diversity % of Zooplankton during High tide and Low tide period  
of Neap tide In Path Finder Creek near jetty and nearby SPM**



**TABLE # 20 SYSTEMATIC ACCOUNT OF PHYTOPLANKTON IN THE SAMPLING LOCATIONS OF DPT HARBOUR AREA AT KANDLA CREEK AND NEARBY CREEKS DURING SPRING TIDE OF NOVEMBER 2021**

<b>GROUP</b>	<b>PHYLUM</b>	<b>CLASS</b>	<b>ORDER</b>	<b>FAMILY</b>	<b>GENUS/SPECIES</b>	<b>#</b>	<b>Relative Abundance</b>	
BLUE GREEN ALGAE	Cyanophyta	Cyanophyceae	Nostocales	Oscillatoriaceae	<i>Oscillatoria</i> sp.	B1	Rare	
					<i>Arthrospira</i> sp.	B2	Rare	
DIATOMS	Bacillariophyta	Coscinodiscophyceae	Thalassiosirales	Thalassiosiraceae	<i>Planktoniella</i> sp.	D1	Rare	
			Coscinodiscales	Coscinodiscaceae	<i>Coscinodiscus</i> sp.	D2	Abundant	
			Triceratiales	Triceratiaceae	<i>Odontella</i> sp.	D3	Occasional	
					<i>Triceratium</i> sp.	D4	Occasional	
			Biddulphiales	Biddulphiaceae	<i>Biddulphia</i> sp.	D5	Dominant	
			Hemiaulales	Bellerucheaceae	<i>Belleruche</i> sp.	D6	Rare	
			Rhizosoleniales	Rhizosoleniaceae	<i>Rhizosolenia</i> sp.	D7	Occasional	
			Chaetocerotales	Chaetocerotaceae	<i>Chaetoceros</i> sp.	D8	Occasional	
			Leptocylindrales	Leptocylindraceae	<i>Leptocylindrus</i> sp.	D9	Occasional	
			Lithodesmiales	Lithodesmiaceae	<i>Ditylum</i> sp.	D10	Frequent	
		Bacillariophyceae	Naviculales	Pleurosigmales	Pleurosigmales	<i>Pleurosigma</i> sp.	D11	Occasional
		Fragilariophyceae	Thalassionematales	Thalassionemataceae	<i>Thalassiothrix</i> sp.	D12	Frequent	
					<i>Thalassionema</i> sp.	D13	Rare	
			Fragilariales	Fragilariaceae	<i>Fragilaria</i> sp.	D14	Frequent	
					<i>Synedra</i> sp.	D15	Rare	
		Tabellariales	Tabellariaceae	<i>Tabellaria</i> sp.	D16	Rare		
DINOFLAGELLATES	Dinoflagellata / Dinzoa	Dinophyceae	Peridinales	Protoperidiniaceae	<i>Protoperidinium</i> sp.	DF1	Rare	
			Gonyaulacales	Ceratiaceae	<i>Ceratium furca</i>	DF2	Rare	

**TABLE # 21 SYSTEMATIC ACCOUNT OF PHYTOPLANKTON IN THE SAMPLING LOCATIONS IN OF DPT HARBOUR AREA AT KANDLA CREEK AND NEARBY CREEKS DURING AND NEAP TIDE OF NOVEMBER,2021**

GROUP	PHYLUM	CLASS	ORDER	FAMILY	GENUS/SPECIES	#	Relative Abundance
BLUE GREEN ALGAE	Cyanophyta	Cyanophyceae	Nostocales	Oscillatoriaceae	<i>Oscillatoria</i> sp.	B1	Rare
DIATOMS	Bacillariophyta	Coscinodiscophyceae	Thalassiosirales	Thalassiosiraceae	<i>Planktoniellasp</i>	D1	Occasional
					<i>Thalassiosirasp</i>	D2	Occasional
			Coscinodiscales	Coscinodiscaceae	<i>Coscinodiscus</i> sp.	D3	Frequent
			Triceratiales	Triceratiaceae	<i>Odontellasp</i>	D4	Rare
					<i>Triceratium</i> sp.	D5	Occasional
			Biddulphiales	Biddulphiaceae	<i>Biddulphiasp</i>	D6	Dominant
			Hemiaulales	Bellerocheaceae	<i>Bellerocheasp</i>	D7	Rare
			Rhizosoleniales	Rhizosoleniaceae	<i>Rhizosolenia</i> sp.	D8	Occasional
			Chaetocerotales	Chaetocerotaceae	<i>Chaetoceros</i> sp.	D9	Rare
			Leptocylindrales	Leptocylindraceae	<i>Leptocylindrussp</i>	D10	Rare
		Lithodesmiales	Lithodesmiaceae	<i>Ditylum</i> sp.	D11	Frequent	
		Bacillariophyceae	Bacillariales	<i>Bacillariasp.</i>	D12	Occasional	
				<i>Nitzschiasp</i>	D13	Rare	
			Naviculales	Naviculaceae	<i>Naviculasp</i>	D14	Rare
				Pleurosigmataceae	<i>Pleurosigmasp</i>	D15	Rare
			Surirellales	Entomoneidaceae	<i>Entomoneissp</i>	D16	Rare
		Fragilariophyceae	Thalassionematales	<i>Thalassiothrix</i> sp.	D17	Abundant	
				<i>Thalassionema</i> sp.	D18	Occasional	
			Fragilariales	<i>Fragilariasp</i>	D19	Frequent	
				<i>Synedrassp</i>	D20	Rare	
		Tabellariales	Tabellariaceae	<i>Tabellariasp</i>	D21	Rare	
DINO FLAGELLATES	Dinoflagellata / Dinzoa	Dinophyceae	Gonyaulacales	Ceratiaceae	<i>Ceratiumfurca</i>	DF1	Rare
					<i>Ceratiumtripos</i>	DF2	Rare

**TABLE # 22 SYSTEMATIC ACCOUNT OF PHYTOPLANKTON IN THE SAMPLING LOCATIONS IN OF DPT OOT AREA AT PATH FINDER CREEK AND NEARBY SPM AT VADINAR DURING SPRING TIDE OF NOVEMBER 2021**

<b>GROUP</b>	<b>PHYLUM</b>	<b>CLASS</b>	<b>ORDER</b>	<b>FAMILY</b>	<b>GENUS/SPECIES</b>	<b>#</b>	<b>Relative Abundance</b>
DIATOMS	Bacillariophyta	Coscinodiscophyceae	Coscinodiscales	Coscinodiscaceae	<i>Coscinodiscus sp.</i>	D1	Dominant
			Triceratiales	Triceratiaceae	<i>Triceratiumsp.</i>	D2	Rare
			Biddulphiales	Biddulphiaceae	<i>Biddulphiasp</i>	D3	Abundant
			Hemiaulales	Bellerocheaceae	<i>Bellerocheasp</i>	D4	Rare
			Rhizosoleniales	Rhizosoleniaceae	<i>Rhizosolenia sp.</i>	D5	Occasional
			Chaetocerotales	Chaetocerotaceae	<i>Chaetocerossp</i>	D6	Frequent
			Lithodesmiales	Lithodesmiaceae	<i>Ditylumsp</i>	D7	Occasional
		Bacillariophyceae	Naviculales	Pleurosigmaaceae	<i>Pleurosigmasp</i>	D8	Rare
			Bacillariales	Bacillariaceae	<i>Bacillariasp.</i>	D9	Rare
		<i>Pseudo-Nitzschiasp</i>			D10	Occasional	
		Fragilariophyceae	Thalassionematales	Thalassionemataceae	<i>Thalassiothrix sp.</i>	D11	Frequent

**TABLE # 23 SYSTEMATIC ACCOUNT OF PHYTOPLANKTON IN THE SAMPLING LOCATIONS IN OF DPT OOT AREA AT PATH FINDER CREEK AND NEARBY SPM AT VADINAR DURING AND NEAP TIDE OF NOVEMBER, 2021**

<b>GROUP</b>	<b>PHYLUM</b>	<b>CLASS</b>	<b>ORDER</b>	<b>FAMILY</b>	<b>GENUS/SPECIES</b>	<b>#</b>	<b>Relative Abundance</b>
DIATOMS	Bacillariophyta	Coscinodiscophyceae	Thalassiosirales	Thalassiosiraceae	<i>Planktoniellasp</i>	D1	Occasional
					<i>Thalassiosiras</i>	D2	Rare
			Coscinodiscales	Coscinodiscaceae	<i>Coscinodiscus sp.</i>	D3	Dominant
			Triceratiales	Triceratiaceae	<i>Triceratiumsp</i>	D4	Rare
			Biddulphiales	Biddulphiaceae	<i>Biddulphiasp</i>	D5	Abundant
			Hemiaulales	Bellerocheaceae	<i>Bellerocheasp</i>	D6	Occasional
			Rhizosoleniales	Rhizosoleniaceae	<i>Rhizosolenia sp.</i>	D7	Occasional
			Chaetocerotales	Chaetocerotaceae	<i>Chaetocerossp</i>	D8	Rare
		Lithodesmiales	Lithodesmiaceae	<i>Ditylumsp</i>	D9	Frequent	
		Bacillariophyceae	Naviculales	Pleurosigmaaceae	<i>Pleurosigmasp</i>	D10	Rare
			Bacillariales	Bacillariaceae	<i>Bacillariasp.</i>	D11	Occasional
					<i>Nitzschiasp</i>	D12	Rare
					<i>Pseudo-Nitzschiasp</i>	D13	Frequent
			Fragilariales	Fragilariaceae	<i>Synedra sp.</i>	D14	Rare
		Fragilariophyceae	Thalassionematales	Thalassionemataceae	<i>Thalassiothrix sp.</i>	D15	Occasional
DINOFLAGELLATES	Dinoflagellata / Dinozoa	Dinophyceae	Gonyaulacales	Ceratiaceae	<i>Ceratiumfusius</i>	DF1	Rare
					<i>Ceratiumfurca</i>	DF2	Rare

**TABLE #24 SYSTEMATIC ACCOUNT OF ZOOPLANKTON FROM THE SAMPLING LOCATIONS OF DPT HARBOUR AREA AT KANDLA CREEK AND NEARBY CREEKS DURING SPRING TIDE OF NOVEMBER, 2021**

<b>GROUP</b>	<b>PHYLUM</b>	<b>CLASS</b>	<b>ORDER</b>	<b>FAMILY</b>	<b>GENUS/SPECIES</b>	<b>#</b>	<b>RELATIVE ABUNDANCE</b>
TINTINIDS	PROTOZOA CILIOPHORA	Spirotrichea	Tintinnida	Tintinnidiidae	<i>Leprotintinnus</i> sp.	T1	Rare
				Codonellidae	<i>Tintinnopsis radix</i>	T2	Rare
					<i>Tintinnopsis failakkaensis</i>	T3	Occasional
COPEPODS	ARTHROPODA	Crustacea Subclass: Copepoda	Calanoida	Paracalanidae	<i>Acrocalanus</i> sp.	C1	Frequent
				Clausocalanidae	<i>Clausocalanus</i> sp.	C2	Rare
			Cyclopoida	Oithonidae	<i>Oithona</i> sp.	C3	Abundant
			Harpacticoida	Ectinosomatidae	<i>Microsetella</i> sp.	C4	Rare
				Euterpinae	<i>Euterpina</i> sp.	C5	Occasional
			Poecilostomatoida	Oncaeidae	<i>Oncaea</i> sp.	C6	Rare
ROTIFERS	ROTIFERA	Rotifera Subclass: Eurotatoria	Superorder: Monogononta Order: Ploimida	Brachionidae	<i>Brachionus plicatilis</i>	R1	Rare
UROCHORDATA	CHORDATA SUB PHYLUM UROCHORDATA	Appendicularia		Oikopleuridae	<i>Oikopleura</i> sp.	U1	Rare
CRUSTACEAN LARVAE	ARTHROPODA (CRUSTACEA)	Copepoda			Nauplius larvae of Copepods	L1	Dominant
BRACHYURA LARVAE	ARTHROPODA (CRUSTACEA)	Malacostraca Decapoda			Brachyuran Zoea larvae	L2	Occasional

**DCPL/DPT/20-21/19 -NOVEMBER - 2021**

**Detox Corporation Pvt. Ltd., Surat**



**Environmental Monitoring Report of Deendayal Port Trust, November-2021**

MOLLUSCAN LARVAE	MOLLUSCA	Gastropoda Streptoneura			Opisthobranchia larvae	L3	Rare
POLYCHAETE LARVAE	ANNELIDA	Polychaeta			Trochophore larvae	L4	Occasional
BIVALVE LARVAE	MOLLUSCA	Pelecypoda			Veliger larvae of Bivalves	L5	Rare

**TABLE # 25 SYSTEMATIC ACCOUNT OF ZOOPLANKTON FROM THE SAMPLING OF DPT HARBOUR AREA AT KANDLA CREEK AND NEARBY CREEKS DURING NEAP TIDE OF NOVEMBER, 2021**

GROUP	PHYLUM	CLASS	ORDER	FAMILY	GENUS/SPECIES	#	RELATIVE ABUNDANCE
TINTINIDS	PROTOZOA CILIOPHORA	Spirotrichea	Tintinnida	Tintinnidiidae	<i>Leprotintinnus</i> sp.	T1	Frequent
				Codonellidae	<i>Tintinnopsis gracilis</i>	T2	Occasional
					<i>Tintinnopsis radix</i>	T3	Frequent
					<i>Tintinnopsis failakkaensis</i>	T4	Occasional
				Tintinnidae	<i>Amphorides</i> sp.	T5	Rare
COPEPODS	ARTHROPODA	Crustacea Subclass: Copepoda	Calanoida	Paracalanidae	<i>Acrocalanus</i> sp.	C1	Occasional
				Eucalanidae	<i>Pareucalanus</i> sp.	C2	Rare
				Clausocalanidae	<i>Clausocalanus</i> sp.	C3	Rare
				Centropagidae	<i>Centropages</i> sp.	C4	Rare
			Cyclopoida	Oithonidae	<i>Oithona</i> sp.	C5	Abundant
			Harpacticoida	Ectinosomatidae	<i>Microsetella</i> sp.	C6	Frequent
				Euterpinae	<i>Euterpina</i> sp.	C7	Occasional
			Poecilostomatoida	Oncaeidae	<i>Oncaea</i> sp.	C8	Rare
ARROW WORMS	CHAETOGNATHA	Sagittoidea	Aphragmophora	Sagittidae	<i>Sagitta</i> sp.	A1	Rare

**DCPL/DPT/20-21/19 -NOVEMBER - 2021**

**Detox Corporation Pvt. Ltd., Surat**

**Environmental Monitoring Report of Deendayal Port Trust, November-2021**

MYSIDS	ATHROPODA CRUSTACEA	Malacostraca	Mysida, Decapoda	Solenoceridae	<i>Solenocerasp.</i>	M1	Rare
				Luciferidae	<i>Lucifer sp.</i>	M2	Rare
UROCHORDATA	CHORDATA SUB PHYLUM UROCHORDAT A	Appendicularia		Oikopleuridae	<i>Oikopleura sp.</i>	U1	Rare
CILIATES	CILIOPHORA	Oligohymenoph orea	Sessilida	Zoothamniidae	<i>Zoothamniumsp.</i>	CI1	Rare
MEDUSA	PHYLUM CNIDARIA	Hydrozoa			Unidentified medusa	ME 1	Rare
CRUSTACEAN LARVAE	ARTHROPODA (CRUSTACEA)	Copepoda			Nauplius larvae of Copepods	L1	Dominant
BRACHYURA LARVAE	ARTHROPODA (CRUSTACEA)	Malacostraca Decapoda			Brachyuran Zoea larvae	L2	Frequent
<b>GROUP</b>	<b>PHYLUM</b>	<b>CLASS</b>	<b>ORDER</b>	<b>FAMILY</b>	<b>GENUS/SPECIES</b>	<b>#</b>	<b>RELATIVE ABUNDAN CE</b>
BARNACLE LARVAE	ATHROPODA CRUSTACEA	Maxillopoda Thecostraca			Cirripede larvae	L3	Occasional
CYPHONAUTES LARVAE	BRYOZOA				Cyphonautes larvae	L4	Occasional
MOLLUSCAN LARVAE	MOLLUSCA	Gastropoda Streptoneura			Opisthobranchia larvae	L5	Rare
ECHINODERMAT A LARVAE	ECHINODERMA TA				Ophioplutes larvae/ Echinoplutes larvae	L6	Rare
POLYCHAETE LARVAE	ANNELIDA	Polychaeta			Trochophore larvae	L7	Occasional
FORAMINIFERA	FORAMINIFERA	Globothalamea	Rotalliida	Globigerinidae	<i>Globigerina sp.</i>	F1	Rare
				Rotalliidae	<i>Rotalia sp.</i>	F2	Rare

**DCPL/DPT/20-21/19 -NOVEMBER - 2021**

**Detox Corporation Pvt. Ltd., Surat**

**TABLE # 26 SYSTEMATIC ACCOUNT OF ZOOPLANKTON FROM THE SAMPLING LOCATIONS OF DPT OOT AREA AT PATH FINDER CREEK AND NEARBY SPM AT VADINAR DURING SPRING TIDE OF NOVEMBER,2021**

<b>GROUP</b>	<b>PHYLUM</b>	<b>CLASS</b>	<b>ORDER</b>	<b>FAMILY</b>	<b>GENUS/SPECIES</b>	<b>#</b>	<b>RELATIVE ABUNDANCE</b>
TINTINIDS	PROTOZOA CILIOPHORA	Spirotrichea	Tintinnida	Tintinnidiidae	<i>Leprotintinnus</i> sp.	T1	Occasional
				Codonellidae	<i>Tintinnopsis gracilis</i>	T2	Rare
					<i>Tintinnopsis radix</i>	T3	Occasional
COPEPODS	ARTHROPODA	Crustacea Subclass: Copepoda	Calanoida	Paracalanidae	<i>Acrocalanus</i> sp.	C1	Frequent
				Clausocalanidae	<i>Clausocalanus</i> sp.	C2	Rare
			Cyclopoida	Oithonidae	<i>Oithona</i> sp.	C3	Abundant
			Harpacticoida	Euterpinidae	<i>Euterpina</i> sp.	C4	Rare
			Poecilostomatoida	Oncaeidae	<i>Oncaea</i> sp.	C5	Rare
				Corycaeidae	<i>Corycaeus</i> sp.	C6	Rare
CRUSTACEAN LARVAE	ARTHROPODA (CRUSTACEA)	Copepoda			Nauplius larvae of Copepods	L1	Dominant
BIVALVE LARVAE	MOLLUSCA	Pelecypoda			Veliger larvae of Bivalves	L2	Occasional
MOLLUSCAN LARVAE	MOLLUSCA	Gastropoda Streptoneura			Opisthobranchia larvae	L3	Occasional
POLYCHAETE LARVAE	ANNELIDA	Polychaeta			Trochophore larvae	L4	Occasional

**TABLE # 27 SYSTEMATIC ACCOUNT OF ZOOPLANKTON FROM THE SAMPLING LOCATIONS OF DPT OOT AREA AT PATH FINDER CREEK AND NEARBY SPM AT VADINAR DURING NEAP TIDE OF NOVEMBER, 2021**

<b>GROUP</b>	<b>PHYLUM</b>	<b>CLASS</b>	<b>ORDER</b>	<b>FAMILY</b>	<b>GENUS/SPECIES</b>	<b>#</b>	<b>RELATIVE ABUNDANCE</b>
TINTINIDS	PROTOZOA CILIOPHORA	Spirotrichea	Tintinnida	Tintinnidiidae	<i>Leprotintinnus</i> sp.	T1	Occasional
				Codonellidae	<i>Tintinnopsis gracilis</i>	T2	Rare
					<i>Tintinnopsis radix</i>	T3	Occasional
				Codonellopsidae	<i>Codonellopsis</i> sp.	T4	Rare
COPEPODS	ARTHROPODA	Crustacea Subclass: Copepoda	Calanoida	Paracalanidae	<i>Acrocalanus</i> sp.	C1	Abundant
				Eucalanidae	<i>Subeucalanus</i> sp.	C2	Rare
				Clausocalanidae	<i>Clausocalanus</i> sp.	C3	Occasional
			Cyclopoida	Oithonidae	<i>Oithona</i> sp.	C4	Frequent
			Harpacticoida	Euterpinae	<i>Euterpina</i> sp.	C5	Rare
			Poecilostomatoida	Oncaidae	<i>Oncaea</i> sp.	C6	Rare
				Corycaidae	<i>Corycaeus</i> sp.	C7	Rare
UROCHORDATA	CHORDATA SUB PHYLUM UROCHORDATA	Appendicularia		Oikopleuridae	<i>Oikopleura</i> sp.	U1	Rare
CRUSTACEAN LARVAE	ARTHROPODA (CRUSTACEA)	Copepoda			Nauplius larvae of Copepods	L1	Dominant
BRACHYURA LARVAE	ARTHROPODA (CRUSTACEA)	Malacostraca Decapoda			Brachyuran Zoea larvae	L2	Rare
MOLLUSCAN LARVAE	MOLLUSCA	Gastropoda Streptoneura			Opisthobranchia larvae	L3	Rare
POLYCHAETE LARVAE	ANNELIDA	Polychaeta			Trochophore larvae	L4	Occasional
BIVALVE LARVAE	MOLLUSCA	Pelecypoda			Veliger larvae of Bivalves	L5	Rare

**DCPL/DPT/20-21/19 -NOVEMBER - 2021**

**Detox Corporation Pvt. Ltd., Surat**

**Environmental Monitoring Report Of Deendayal Port Trust, NOVEMBER-2021**

**BENTHIC ORGANISMS:**

Few Benthic organisms were observed in the collected sediments by using the Van-veen grabs during the sampling conducted during spring tide period and Neap tide period from DPT harbour region and nearby creek. The meiobenthic organisms during spring tide were represented by Polychaetes, and Nematodes. The polychaetes were represented by *Scyphoproctus sp.*, *Notomastus sp.*, *Dasybranchus*. The meiobenthic organisms in the collected samples were varying from 50-180 N/M<sup>2</sup> during spring tide and 60-130 N/M

**Table # 28 BENTHIC FAUNA IN THE SAMPLING LOCATIONS IN DPT HARBOUR AREA CREEKS DURING SPRING TIDE IN NOVEMBER, 2021**

Benthic fauna	ABUNDANCE IN NO/M <sup>2</sup> DIFFERENT SAMPLING STATIONS REPRESENTATION BY GROUP						
	DPT HARBOUR			CREEKS			
POLYCHAETES	KPT-1	KPT-2	KPT-3	KPT-4	KPT-5	KPT-6	
Family : Capitellidae <i>c</i>	0	40	0	20	20	NS	
Family : Capitellidae <i>Notomastus sp.</i>	40	60	40	80	30	NS	
<b>Total Polychaetes N/M<sup>2</sup></b>	40	100	40	120	50		
<b>Un identified Nematode worms</b>	10	20	40	60	20	NS	
TOTAL Benthic Fauna NUMBER/ M <sup>2</sup>	50	120	80	180	70	-	

NS : No sample

**Table # 29 BENTHIC FAUNA IN THE SAMPLING LOCATIONS IN DPT HARBOUR AREA CREEKS DURING NEAP TIDE IN NOVEMBER, 2021**

Benthic fauna	ABUNDANCE IN NO/M <sup>2</sup> DIFFERENT SAMPLING STATIONS REPRESENTATION BY GROUP						
	DPT HARBOUR			CREEKS			
POLYCHAETES	KPT-1	KPT-2	KPT-3	KPT-4	KPT-5	KPT-6	
Family : Capitellidae <i>Dasybranchus sp.</i>	10	20	10	10	20	NS	
Family : Capitellidae <i>Notomastus sp.</i>	50	60	20	40	20	NS	
Family : Glyceridae <b>Glycera</b>	10	20	10	0	0	NS	

DCPL/DPT/20-21/19 - NOVEMBER -2021

## Environmental Monitoring Report Of Deendayal Port Trust, NOVEMBER-2021

<b>Total Polychates N/M<sup>2</sup></b>	70	100	40	50	40		
<b>Un identified Nematode worms</b>	20	30	30	30	20	NS	
TOTAL Benthic Fauna NUMBER/ M <sup>2</sup>	90	130	70	80	60	-	

NS : No sample

### Meteorological Data

Automatic Weather station have been installed in Seva Sadan -3 at the Deendayal Port which records the data on Temperature (°C), Humidity (%), Wind (mph), Dew Point (°C), Wind Direction (°), Pressure, Solar radiation, heat Index and UVI.

#### Temperature

The mean day time temperature for Deendayal Port was 30.3 °C. The day-time maximum temperature was 38.6 °C. The mean night time temperature was 26.5 °C. The minimum mean night time temperature recorded was 30.6 °C.

#### Air Pressure

The mean absolute air pressure for the month of November was 1009.9 hpa, whereas the mean relative pressure was 1009.3 hpa. The maximum absolute air pressure recorded for the month of November was 1016.5 hpa.

#### Heat Index

The mean day-time heat index for the month of November was 33.8 °C. The maximum heat index recorded was 55°C.

#### Solar Radiation

The mean Solar Radiation in November was 252.2 w/m<sup>2</sup>. The maximum solar radiation recorded in the month of November was 746.6 w/m<sup>2</sup>.

#### Humidity

The mean day-time humidity was 60.0 % for the month of November and mean night time humidity was 72.4%. Maximum humidity recorded during day-time was 94.0 % and maximum humidity recorded during night-time was 93.0%.

### **Wind Velocity and Wind Direction**

The mean wind velocity for the entire month of November was 4.6 km/hour. Maximum wind velocity recorded was 29.2 Km/hr . The wind direction was mostly S to N.

### **Conclusive Summary and Remedial measures Suggested**

The AAQ monitoring at six locations of Deendayal Port indicates that the mean PM<sub>10</sub> values at four locations viz. Coal storage area, Marine Bhavan and Oil Jetty area were found above the permissible standards (100 µg/m<sup>3</sup> )andPM<sub>2.5</sub>was above permissible limits at Coal storage location(Limit 60 µg/m<sup>3</sup>).

Drinking water at all the twenty locations was found potable and was within permissible limits of BIS standards (IS 10500).

Noise quality was also within the set permissible standards of an Industrial Area. The noise level observed during day time was >75 dB (A) and at night time was >70 dB (A) during the entire monitoring period.

The sewage treated water of Deendayal Port Colony (Gopalpuri) was in line with the standards set by the Gujarat Pollution Control Board.

### **Reasons for higher Values of PM<sub>10</sub>**

Large amount of coal is handled at Berth No. 6, 7, 8 and 9. The unloading of coal directly in the truck, using grabs cause coal to spread in air as well as coal dust to fall on ground. This settled coal dust again mixes with the air while trucks travel through it.

Also, the coal laden trucks are not always covered with tarpaulin sheets and these results in spillage of coal from trucks/dumpers during its transit from vessel to yard or storage site. This also increased PM values around marine Bhavan & Coal storage area.

### **Remedial Measures**

The values of PM<sub>10</sub> during the month of November, 2021 were observed beyond the permissible limit at four locations mentioned above. Given below are the remedial measures suggest to minimize the Air pollution at Deendayal Port.

Guidelines for Coal Handling by GPCB should be strictly followed.  
(<http://gpcb.gov.in/pdf/coal-handling-guidelines.pdf>)

Except for the higher values of PM<sub>10</sub> at Coal storage site, Oil Jetty, Tuna Port and Marine Bhavan locations, the monitoring results for the present month suggest that the overall Environment Quality of Deendayal Port is satisfactory.



**SOURCE OF LITERATURE AND ADDITIONAL REFERENCE FOR  
ECOLOGICAL STUDY**

- 1) ALBERT WEST PHAL (1976) Protozoa Blackwell , London
- 2) BANERJEE R.K. (1989) Heavy metals and Benthic foraminiferal distribution along Bombay coast India. Studies in benthic foraminifera. *Tokyo University Press* Tokyo pp 151-157
- 3) Banse K (1995) Zooplankton: Pivotal role in the control of ocean production: I. Biomass and production. *ICES J Mar Sci* 52: 265-277.
- 4) Beaugrand G, and Ibanez F (2004) Monitoring marine plankton ecosystems. II: long-term changes in North Sea calanoid copepods in relation to hydroclimatic variability. *Inter Res Mar Ecol Prog Ser* 284:35-47.
- 5) DAY F. (1889) The fauna of British India Ceylon and Burma- Fishes Vol-1- Vol-2 *Taylor and Francis* London
- 6) DESIKACHARY T.V. (1989) Atlas of diatoms, Madras Science Foundation
- 7) DESIKACHARY T.V. (1959) Cyanophyta ICAP Monographs on Algae *Indian Council of Agricultural research* New Delhi
- 8) FAIZAYOUSIF AL-YAMANI & MARIA A. SABUROVA (2010) illustrative guide on the flagellates of Intertidal soft sediment *Kuwait Institute for scientific Research* Kuwait
- 9) FAIZAYOUSIF AL-YAMANI, VALERIYSKRYABIN, ALEKSANDRA GUBANOVA, SERGEY KHVOROV AND IRINA PRUSOVA (2011), Marine zooplankton Practical guide from North western Arabian gulf Vol-1 and vol-2 *Kuwait Institute for scientific Research* Kuwait
- 10) FAUVEL P. (1953), The fauna of India Annelida - Polychaeta *Indian Press* Allahabad
- 11) Gajbhiye SN, Nair VR, and Desai BN (1984). Diurnal variation of zooplankton in Malad creek, Bombay. *Indian Journal of Marine Science*. 13:75-79.
- 12) HAYWARD P.J AND RYLAND J.S. (1995) Handbook of Marine fauna of north -West Europe *oxford University Press* London
- 13) HIGGINS R.P. HAJAMARTHIEL Eds. (1998) Introduction to the study of Meio Fauna
- 14) HORACE G. BARBER AND ELIZABETH Y. HAWORTH (1981) A guide to the Morphology of DIATOMS FRUSTULES.
- 15) INGRAM HENDEY (1964) An introductory account of smaller Algae of British coastal waters part-V. Bacillariophyceae
- 16) JOHN H. WICKSTEAD (1965) an Introduction to the study of Tropical Plankton *Hutchinson Tropical Monographs*
- 17) JOYOTHIBABU, R. MADHU, N.V. MAHESHWARAN, P.A., NAIR K.K.C., VENUGOPL, P. BALASUBRAMANIAN T. (2005) Dominance of Dinoflagellates in micro zooplankton communities in the oceanic region Bay of Bengal and Andaman sea *Current science* vol.84. 10<sup>th</sup> May 2003
- 18) KASTURIRANGAN L.R. (1963) A key for the identification of the Common Planktonic Copepoda of Indian Coastal water
- 19) Kusum KK, Vineetha G, Raveendran TV, Nair VR, Muraleedharan KR, Achuthankutty CT and Joseph T (2014) Chaetognath community and their responses to varying environmental factors in the northern Indian ocean. *J Plankton Res* 36(4): 1146- 1152.

## **Environmental Monitoring Report Of Deendayal Port Trust, NOVEMBER-2021**

---

- 20) Lalli CM and Parsons TR (1997) Biological Oceanography: An Introduction. DOI <https://doi.org/10.1016/B978-0-7506-3384-0.X5056-7>.
- 21) Madhupratap M (1978) Studies on ecology of zooplankton of Cochin backwaters. *Mahasagar Bull Nat Inst Oceanogr* 11: 45-56.
- 22) Madhupratap M (1979) Distribution, community structure and species succession of copepods from Cochin Backwaters. *Indian J Ma Sci* 8: 1-8.
  
- 23) Madhupratap M (1987) Status and strategy of zooplankton of tropical Indian estuaries: A review. *Bull Plank Soc Jpn* 34: 65-81.
- 24) Madhupratap M (1999) Free living copepods of the Arabian Sea, Distribution and Research Perspectives. *I J Mar Sci* 146-149.
- 25) Madhupratap M and Haridas P (1986) Epipelagic calanoid copepods of the northern Indian Ocean. *Oceanologica Acta* 9(2):105-117.
- 26) MANAL AL-KANDARI, FAIZA Y. AL-YAMANI , KHOLOOD AL-RIFAIE ( 2009) Marine phytoplankton Atlas of Kuwait's water *Kuwait Institute for scientific Research*
- 27) MPEDA (1998) Commercial Fishes and shell fishes of India
- 28) NEWEL G.E. & NEWELL R.C. (1963) Marine plankton a Practical Guide Hutchinson Educational
- 29) NIGAM R.C. AND CHATURVEDIS.K. (2000) Foraminiferal Study from Kharo Creek , Kachchh ( Gujarat) North west coast of *India. Indian Journal of marine science* Vol.29 133-189
- 30) OLAV GIERE (1993) Meio benthology , Microscopic Fauna in Aquatic Sediments m Springer London
- 31) PERRAGALLO( 1965) Diatomees marines de france A. *Asher & Co.* Amsterdam
- 32) Robert P.. Higgins (Eds.), (1985) An introduction to the study of Meio fauna Smithsons Institution press Washington DC
- 33) STERRER W. STERRER S Eds. Marine fauna and flora of Bermuda A systematic Guide to the identification of Marine Organisms. *John Wiley and Sons* New York
- 34) Suresh Gandhi. M. (2009) Distribution of certain ecological parameters and Foraminiferal distribution in the depositional environment of Pak strait east coast of India .*Indian J. of Marine Science* Vol.33 pp 287-295
- 35) Venktaraman (1993) A systematic account of some south Indian diatoms . *Proceeding of Indian Academy of Science* Vol.X No.6 Sec.B.

\*\*\*\*\*