

DEENDAYAL PORT AUTHORITY
(Erstwhile: DEENDAYAL PORT TRUST)



www.deendayalport.gov.in

Administrative Office Building
Post Box NO. 50
GANDHIDHAM (Kutch).
Gujarat: 370 201.
Fax: (02836) 220050
Ph.: (02836) 220038

EG/WK/4751/Part (Comp.1)/ 287

Dated: 20/04/2023

The Director (Environment) & Member Secretary,
Gujarat Coastal Zone Management Authority,
Forest & Environment Department,
Govt. of Gujarat,
Block No.14, 8th floor,
Sachivalaya,
Gandhinagar – 382 010.

Sub: "Development of 7 Integrated facilities (Stage I) within the existing Kandla Port Trust limit at District Kutch (Gujarat) by M/s Kandla Port Trust Limited"– **Pointwise Compliances of the conditions stipulated in CRZ Recommendations reg.**

Ref.: 1) Letter No. ENV-I0-2014-25-E July, 1, 2015 of Director (Environment) & Member Secretary, GCZMA, Forest & Environment Department, GoG
2) Compliance Report (period up to May, 2017) submitted by KPT vide letter no. EG/WK/4751/Part (Compliance)/198 Dated: 12/6/2017.
3) Compliance Report (period up to Nov., 2017) submitted by DPT vide letter no. EG/WK/4751/Part (Compliance)/612 Dated: 15/12/2017.
4) Compliance Report (period up to May, 2018) submitted by DPT vide letter no. EG/WK/4751/Part (Compliance)/313 Dated: 14(21)/06/2018.
5) Compliance Report (period up to March, 2019) submitted by DPT vide letter no. EG/WK/4751/Part (Compliance)/114 Dated: 30(2)/03(04)/2019.
6) Compliance report (period up to October, 2019) submitted by DPT vide letter no. EG/WK/4751/Part (Comp 1)/153 Dated: 14/11/2019.
7) Compliance report (period upto November, 2020) submitted by DPT vide letter no. EG/WK/4751/Part (Comp 1) dated 29/12/2020.
8) Compliance report (period upto May, 2021) submitted by DPT vide letter no. EG/WK/4751/Part (Comp 1)/93 dated 07/10/2021.
9) Compliance report (period upto May, 2022) submitted by DPA vide letter no. EG/WK/4751/Part (Comp 1)/222 dated 30/01/2023.

Sir,

It is requested to kindly refer the above cited references for the said subject.

In this connection, it is to state that, the Gujarat Coastal Zone Management Authority vide above referred letter dated 1/7/2015 had recommended 7 project activities of Deendayal Port Authority. Subsequently, the MoEF&CC, GoI had accorded the Environmental & CRZ Clearance vide letter dated 19/12/2016 for the 7 project activities recommended by the GCZMA.

.....Cont.....

Subsequently, DPA vide above referred letters had regularly submitted compliance report of the stipulated conditions, to the Additional Secretary & Director (Env.), F & E Dept., GoG.

Now, as directed under Specific Condition No. 28 mentioned in the CRZ Clearance letter dated 1/7/2015 i.e. **A six monthly report on compliance of the conditions mentioned in this letter shall have to be furnished by the DPA on a regular basis to this Department and MoEF&CC, GoI**, please find enclosed herewith compliance report (for the period up to November, 2022) of stipulated conditions along with necessary annexures, for kind information & record please **(Annexure I)**.

Further, as per the MoEF&CC, Notification S.O.5845 (E) dated 26.11.2018, stated that **"In the said notification, in paragraph 10, in sub-paragraph (ii), for the words "hard and soft copies" the words "soft copy" shall be substituted"**. Accordingly, we are submitting herewith soft copy of the same via e-mail ID gczma.crz@gmail.com & direnv@gujarat.gov.in.

This has the approval of Chief Engineer, Deendayal Port Authority.

Yours faithfully,


Manager (Env.)

Deendayal Port Authority

Copy to:

Shri Amardeep Raju,
Scientist E, Ministry of Environment, Forest and Climate Change,
& Member Secretary (EAC-Infra.1),
Indira Paryavaran Bhawan,
3rd Floor, Vayu Wing, Jor Bagh Road, Aliganj,
New Delhi- 110 003;
E-mail: ad.raju@nic.in

Annexure -I

Annexure 1

CURRENT STATUS OF WORK (up to November, 2022)

Subject: Development of 7 integrated facilities (Stage I) within existing Deendayal Authority at Kandla.

Name of Project	Status
1. Development of oil jetty to handle liquid cargo and ship bunkering terminal at old Kandla under PP mode (jetty: 300m x 15m, approach 450 m X 10 m, back up area 5.5 HA, capacity – 3.39 MMTPA, capital dredging 1,73,660 m ³ maintenance dredging 1,56,294 m ³ (Estimated cost: 276.53 Cr.).	The Concession Agreement was executed between DPA and M/s KOTPL on 16/11/2013 to implement the project on Built, Transfer & Operate (BOT- PPP) Basis by M/s KOTPL. The award of concession was issued on 11/12/2020 to M/s KOTPL by DPA. The Project is under construction phase.
2. Multipurpose Cargo Terminal at Tekra off Tuna on BOT basis (T shape jetty 600m X 80 m Capacity 18 MMTPA, back up area 101 Ha capital dredging 1,26,57,175 m ³ maintenance dredging 18,98,576. 25 m ³ Estimated cost: 1686.66 Cr.	The Board of DPA approved the Feasibility Report in its meeting on 19.02.2021. The MoPSW,GoI vide communication dated 21/10/2022 has conveyed approval granted by the Cabinet Committee on Economic Affairs to the project. Accordingly, DPA already invited RFQ and RFP which is currently under bidding stage <u>No construction activity has started yet.</u>
3. Up gradation of Barge handling capacity at Bundar basis at Kandla capacity 3.33 MMTA back-up area 5 Ha, Estimated cost: 109.59 Cr.	The up-gradation work was completed.
4. Construction of Rail over Bridge at NH 8 A near Nakti Bridge (crossing of NH 8 A Estimated cost: 32.17 Cr.	Construction activity has not yet started.
5. Mechanization of Dry Cargo handling capacity at Kandla Port (Berth 7 and 8 capacity 7.35 MMTPA estimated cost 80.61 Cr.	Mechanization work already completed.
6. Strengthening of Oil jetty 1 (Estimated cost: 7.5 Cr.	The strengthening work completed.
7. Modification and strengthening of Cargo berth No. 6 at Kandla Port Estimated	The modification & strengthening work completed.

cost: 11.5 Cr.	
----------------	--

Out of a total of 7 project activities, construction activities of 4 projects (i.e. Sr. No. 3, 5, 6 & 7 mentioned in the EC & CRZ Clearance) have already been completed. Whereas construction activity of the project at Sr. No. 2 & 4 have not yet started.

For the current compliance period up to November, 2022, construction activity related to project No. 1 is ongoing. The compliance report submitted by the Concessionaire M/s KOTPL is attached herewith as Annexure A.

COMPLIANCE REPORT (for the period up to November, 2022)

Subject: Status of Compliance with the conditions stipulated By Gujarat State Coastal Zone Management Authority, Gandhinagar, in CRZ Recommendation Letter granted for "Development of 7 integrated facilities (Stage I) within existing Deendayal Authority at Kandla".

CRZ Recommendations: Letter No. ENV-I0-2014-25-E dated July 1, 2015, of Director (Environment) & Member Secretary, GCZMA, Forest & Environment Department, GoG.

**Note: Based on the recommendation of the GCZMA, MoEF&CC, GoI had accorded Environmental & CRZ Clearance vide letter dated 19/12/2016*

Sr. No.	Conditions in CRZ Recommendation Letter	Compliance
	Specific Conditions	
1	The provisions of the CRZ notification of 2011 and subsequent amendments issued from time to time shall be strictly adhered to by the KPT. No activity in contradiction to the Provisions of the CRZ Notification shall be carried out by the KPT.	a) For Project at Sr. No. 1 which is under construction, kindly refer compliance submitted by M/s KOTPL (concessionaire of the project) placed at Annexure A . b) Further, w.r.t. project at Sr. No. 2 & 4 (construction not yet started), it is assured that no activity in contradiction to the Provisions of the CRZ Notification shall be carried out by the DPA.
2	The KPT shall have to ensure that there shall not be any damage to the existing mangrove area.	a) For Project at Sr. No. 1 which is under construction, kindly refer compliance submitted by M/s KOTPL (concessionaire of the project) placed at Annexure A . b) Further, w.r.t. project at sr.no. 2 & 4 (construction not yet started), it is assured that due care shall be taken to protect the existing mangrove area.
3	The KPT shall prepare an emergency plan to protect existing mangroves in case of any eventuality/accident.	The final report submitted by M/s GUIDE, Bhuj (vide letter dated 21/5/2018) had already been communicated to the MoEF&CC, GoI, Bhopal & copy to the MoEF&CC, GoI, New Delhi, along with six monthly compliance report submitted vide letter dated 21/06/2018.
4	The KPT shall have to make a provision that mangrove areas get proper flushing water and free flow of water shall not be obstructed.	a) For Project at Sr. No. 1 which is under construction, kindly refer compliance submitted by M/s KOTPL (concessionaire of the project) placed at Annexure A . b) Further, w.r.t. project at sr.no. 2 & 4 (construction not yet started), it is assured that provisions shall be made that mangrove areas get proper flushing water and free flow of water shall not be obstructed.

5	The KPT shall have to abide by whatever decision is taken by the GCZMA for violations of CRZ Notification, 2011.	<p>a) For Project at Sr. No. 1 which is under construction, kindly refer compliance submitted by M/s KOTPL (concessionaire of the project) placed at Annexure A.</p> <p>b) Point noted. DPA will abide by whatever decision is taken by the GCZMA for violations of CRZ Notification, 2011.</p>
6	There shall not be violations of the order dated 9/12/2013 passed by the National Green Tribunal; and accordingly, there shall be no mangrove destruction taking place in the KPT area.	<p>a) Point Noted. It is hereby assured that due care shall be taken to protect the existing mangrove area.</p> <p>b) For Project at Sr.No. 1 which is under construction, kindly refer compliance submitted by M/s KOTPL (concessionaire of the project) placed at Annexure A.</p>
7	No dredging, reclamation or any other project-related activities shall be carried out in the CRZ area categorised as CRZ I (i), and it shall have to be ensured that the mangrove habitats and other ecologically important and significant areas, if any, in the region are not affected due to any of the project activity.	<p>a) It is hereby assured that DPA will undertake only such project activities (7 project activities) recommended by the GCZMA vide letter dated 1/7/2015 and EC & CRZ Clearance accorded by the MoEF&CC, GoI vide letter dated 19/12/2016.</p> <p>DPA issued a work order to M/s GUIDE vide its letter no. EG/WK/ 4751 /Part (Marine Ecology Monitoring) /11 dated 03/05/2021 for Regular monitoring of Marine Ecology in and around Deendayal Port Authority (Erstwhile Deendayal Port Trust) and continuous Monitoring Program covering all seasons on various aspects of the Coastal Environs for the period 2021-24. The first season report for the year 2022-2023 submitted is attached herewith as Annexure B.</p> <p>b) For Project at Sr.No. 1 which is under construction, kindly refer compliance submitted by M/s KOTPL (concessionaire of the project) placed at Annexure A.</p>
8	The KPT shall participate financially in installing and operating the Vessel Traffic Management System in the Gulf of Kachchh and shall also take the lead in preparing and operational sing and updating regularly after getting it vetted by the Indian Coast Guard.	Deendayal Port Authority had already contributed a number of Rs. 41.25 crores, i.e. 25% of the total project cost of 165 crores for installing and operating the VTMS in the Gulf of Kachchh.
9	The KPT shall strictly ensure that no creeks or rivers are blocked due to any activity at Kandla.	<p>a) For Project at Sr.No. 1 which is under construction, kindly refer compliance submitted by M/s KOTPL (concessionaire of the project) placed at Annexure A.</p> <p>b) Further, w.r.t. project at sr.no. 2 & 4</p>

		(construction not yet started), it is assured that no creeks or rivers shall be blocked due to any activity at Kandla.
10	Mangrove plantation in an area of 100 ha. Shall be carried out by the KPT within 2 years in time bound manner on Gujarat coastline either within or outside the Kandla port Trust area, and a six-monthly compliance report along with the satellite images shall be submitted to the Ministry of Environment and Forest as well as to this Department without fail.	<p>DPA had already undertaken Mangrove Plantation in an area of 1500 Ha. till date since the year 2005. A statement showing details of the mangrove plantation and the cost incurred is again placed in Annexure C.</p> <p>Further, DPA is carrying out an additional mangrove plantation of 100 ha. with the consultation of the Gujarat Ecology Commission vide Work Order No. DD/WK/3050/Pt-I/GIM/PC-44 dated 02/06/2022 (Annexure D).</p> <p>In addition to the above, DPA appointed M/s GUIDE, Bhuj, for "Regular Monitoring of Mangrove Plantation carried out by DPA" (period 15/9/2017 to 14/9/2018 vide work order dated 1/9/2017 and 24/5/2021 to 23/5/2022 vide work order dated 3/5/2021). The final report submitted by M/s GUIDE, Bhuj, for the years 2017 to 2018 as well as for the year 2021 to 2022 has been submitted in the earlier compliance report submitted.</p>
11	No activities other than those permitted by the competent authority under the CRZ Notification shall be carried out in the CRZ area.	<p>a) Point Noted. It is assured that only activities permitted by the competent authority under the CRZ Notification shall be carried out in the CRZ area.</p> <p>b) For Project at Sr. No. 1 which is under construction, kindly refer compliance submitted by M/s KOTPL (concessionaire of the project) placed at Annexure A.</p>
12	No groundwater shall be tapped for any purpose during the proposed expansion modernization activities.	<p>a) For Project at Sr.No. 1 which is under construction, kindly refer compliance submitted by M/s KOTPL (concessionaire of the project) placed at Annexure A.</p> <p>b) Further, w.r.t. Project at Sr. no.2 & 4 (construction not yet started), Water requirement will be met through procurement from GWSSB or private tankers. No ground water shall be tapped.</p>
13	All necessary permissions from different Government Departments/agencies shall be obtained by the KPT before commencing the expansion activities.	DPA had already obtained the necessary Environmental & CRZ Clearance for 7 project activities (dated 19.12.2016). Further, Consent to Establish from GPCB had already been obtained from GPCB for 7 project activities. Moreover, DPA applied for a CTE extension vide Inw Id 268644 dated 9/11/2022.
14	No effluent or sewage shall be discharged	a) For completed projects (modification/

	<p>into sea/creek or in the CRZ area and it shall be treated to conform to the norms prescribed by the GPCB and would be reused /recycled within the plant premises.</p>	<p>strengthening/ up-gradation of existing facilities), Sewage is being treated in the STP of Kandla (1.5 MLD). The treated sewages from STP of DPA are utilized for plantation / Gardening.</p> <p>In addition to the above, DPA appointed M/s Detox Corporation, Surat (NABL Accredited laboratory) for regular Monitoring of environmental parameters including effluent to confirm to the norms prescribed by the GPCB since the year 2016. The work is in progress & DPA have been monitoring data regularly to all the concerned authorities along with compliance reports submitted.</p> <p>The Environmental Monitoring Reports, as submitted by M/s Detox Corporation, is enclosed herewith as Annexure E.</p> <p>b) Further, w.r.t. Project at Sr.No.1, kindly refer to the Monitoring reports submitted by M/s KOTPL along with compliance report placed at Annexure A.</p>
15	<p>All the recommendations and suggestion given by the MANTEC Consultants Pvt. Ltd. in their Comprehensive Environment Impact Assessment report for conservation / protection and betterment of environment shall be implemented strictly by the KPT.</p>	<p>DPA has installed Mist Canon at the Port area to minimize the dust.</p> <p>Further, DPA has already installed continuous sprinkling system in coal stack yard in DPA (40 ha. area) for to prevent dust pollution. Further, to control dust pollution in other area, regular sprinkling through tankers on roads and other staking yards is being done. Regular sweeping of spilled cargo from roads is done by parties on regular basis.</p> <p>DPA appointed M/s Detox Corporation, Surat (NABL Accredited laboratory) for regular Monitoring of environmental parameters since the year 2016. DPA submitted monitoring data regularly to all the concerned authorities along with compliance reports submitted.</p> <p>The Environmental Monitoring Reports as submitted by M/s Detox Corporation is enclosed herewith as Annexure E.</p> <p>For ship waste management, DPA issued Grant of License/Permission to carry out the work of collection and disposal of "Hazardous Waste/Sludge/ Waste Oil" and "Dry Solid Waste (Non- Hazardous)" from Vessels calling at Deendayal Port" through DPA contractors. Further, it is to state that, all ships are required</p>

to follow DG Shipping circulars regarding the reception facilities at Swachh Sagar portal.

DPA assigned work to M/s GUIDE, Bhuj, for regular monitoring of Marine Ecology since the year 2017 (From 2017 – 2021), and reports of the same are being submitted regularly to the Regional Office, MoEF&CC, GoI, Gandhinagar as well as to the MoEF&CC, GoI, New Delhi along with compliance reports submitted.

The final report for the Holistic Marine Ecological Monitoring for the period up to May 2021 was submitted on 22.05.2021. Copy of the report was communicated vide earlier compliance report submitted vide letter dated 29/6/2021.

Further, it is to submit that DPA issued a work order to M/s GUIDE vide its letter no. EG/WK/4751 /Part (Marine Ecology Monitoring) /11 dated 03/05/2021 for Regular monitoring of Marine Ecology in and around Deendayal Port Authority (Erstwhile Deendayal Port Trust) and continuous Monitoring Program covering all seasons on various aspects of the Coastal Environs for the period 2021-24. The first season report for the year 2022-2023 submitted is attached herewith as **Annexure B**.

As already informed, DPA entrusted work of green belt development in and around the Port area to the Forest Department, Gujarat at Rs. 352 lakhs (Area 32 hectares). The work is completed.

Further, DPA has appointed the Gujarat Institute of Desert Ecology (GUIDE) for "Green belt development in Deendayal Port Authority and its Surrounding Areas, Charcoal site' (Phase-I)" vide Work Order No.EG/WK/4757/Part [Greenbelt GUIDE, dated 31st May 2022 (**Annexure F**).

For dredged material management, DPA assigned work to M/s GUIDE, Bhuj for analysis of dredged material since the year 2017 and the reports are being submitted from time to time along with compliance reports submitted. The second Season Report submitted by M/s GUIDE, Bhuj for the period 2021-2022 is attached herewith as **Annexure G**. Further, Dredged Material will be disposed of at

		<p>designated location as identified by the CWPRS, Pune.</p> <p>For energy conservation measures, DPA is already generating 20 MW of Wind energy. In addition to it, DPA has commissioned a 45 kWp Solar Plant at Gandhidham. Further, it is relevant to mention that, two out of four Nos. of Harbour Mobile Crane (HMC) made electric operated. Balance 02 Nos. shall be made electric operated by 2023-2024. Four Nos. of Deisel operated RTGs converted to e-RTGs. Retrofitting of hydrogen fuel cell in Tug Kalinga and Pilot Boat Niharika to be done as a pilot project under the guidance of MoPSW. Also, 14 Nos. of EV cars to be hired in this year and 03 Nos. EV Bus to be procured by the year 2023-24.</p> <p>Further, for Oil Spill Management, DPA is already having Oil Spill Contingency Plan in place and Oil Response System as per the NOS-DCP guidelines.</p>
16	The construction and operational activities shall be carried out in such a way that there is no negative impact on mangroves and other coastal /marine habitats. The construction activities and dredging shall be carried out only under the constant supervision and guidelines of the Institute of National repute like NIOT.	<p>a) For Project at Sr. No. 1 which is under construction, kindly refer compliance submitted by M/s KOTPL (concessionaire of the project) placed at Annexure A.</p> <p>b) For the remaining projects Sr. No 2 & 4 (construction not yet started), it is assured that construction activities and dredging shall be carried out only under the constant supervision and guidelines of the Institute of National repute like NIOT.</p>
17	The KPT shall contribute financially for any common study or project that may be proposed by this Department for environmental management / conservation / improvement for the Gulf of Kutch.	Point noted.
18	The construction debris and / or any other of waste shall not be disposed of into the sea, creek or the CRZ areas. The debris shall be removed from the construction site immediately after the construction is over.	<p>a) For Project at Sr.No. 1 which is under construction, kindly refer compliance submitted by M/s KOTPL (concessionaire of the project) placed at Annexure A.</p> <p>b) Further, w.r.t. project at sr.no. 2 & 4 (construction not yet started), it is assured that construction debris and/ or any other of waste shall not be disposed of into the sea, creek or the CRZ areas, and the debris shall be removed from the construction site immediately after the construction is over.</p>

19	The construction camps shall be located outside the CRZ area and the construction labour shall be provided with the necessary amenities, including sanitation, water supply and fuel and it shall be ensured that the environmental conditions are not deteriorated by the construction labours.	<p>a) No. 1 which is under construction, kindly refer compliance submitted by M/s KOTPL (concessionaire of the project) placed at Annexure A.</p> <p>b) Further, w.r.t. project at sr.no. 2 & 4 (construction not yet started), it is assured that, the construction camps shall be located outside the CRZ area, provision of the necessary amenities, including sanitation, water supply and fuel to the construction labour shall be made, and that the environmental conditions are not deteriorated by the construction labours.</p>
20	The KPT shall regularly updates its Local Oil Spill Contingency and Disaster management Plan in accordance with the National Oil Spill and Disaster Contingency Plan and shall submit the same to the MoEF, GoI and this department after having it vetted through the Indian Coast Guard.	<ul style="list-style-type: none"> ▪ Deendayal Port already has an updated Disaster Management Plan. ▪ Further, the Local Oil Spill Contingency Plan is already available with Deendayal Port Trust. ▪ DPT has also executed MOU with Oil Companies, i.e. IOCL, HPCL, BPCL etc., for combating the Oil Spill at Kandla
21	The KPT shall bear the cost of the external agency that may be appointed by this Department for supervision/monitoring of proposed activities and the environmental impacts of the proposed activities.	Point noted.
22	The KPT shall take up massive greenbelt development activities in and around Kandla and also within the KPT limits.	<p>DPA has planted about one lakhs trees in roadside dividers, colony areas at Kandla and Gopalpuri, in the green belt area of Gandhidham & Adipur Township, Sewage Treatment Plants at Gopalpuri & Kandla and some green belt development plans initiated at different locations in Township areas.</p> <p>DPA entrusted work of green belt development in and around the Port area to the Forest Department, Gujarat, at the cost of Rs. 352 lakhs (Area 32 hectares). The plantation is completed.</p> <p>Further, DPA has appointed the Gujarat Institute of Desert Ecology (GUIDE) for "Green belt development in Deendayal Port Authority and its Surrounding Areas, Charcoal site' (Phase-I)" vide Work Order No.EG/WK/4757/Part [Greenbelt GUIDE, dated 31st May 2022 (Annexure F).</p>
23	The KPT shall have to contribute financially for taking up the socio-	The details of the fund earmarked under CSR activities and CSR activities undertaken by DPA

	economic upliftment activities in this region in construction with the Forest and Environment Department and the District Collector/District Development Officer.	to date & proposed activities are placed at Annexure H .
24	A separate budget shall be earmarked for environmental management and socioeconomic activities and details thereof shall be furnished to this Department as well as the MoEF, GOI. The details with respect to the expenditure from this budget head shall also be furnished.	<p>a) The allocation made under the "Environmental Services & Clearance of other related Expenditure" scheme during BE 2021-22 is Rs. 266 Lakhs. and BE 2022-2023 is Rs. 345 Lakhs.</p> <p>b) The funds earmarked for EMP by the Concessionaire M/s KOTPL w.r.t. project at Sr.No. 1 are delineated in the compliance report submitted (Annexure A).</p>
25	A separate environmental management cell with qualified personnel shall be created for environmental monitoring and management during the construction and operational phases of the project.	<p>a) For Project at Sr.No. 1 which is under construction, kindly refer compliance submitted by M/s KOTPL (concessionaire of the project) placed at Annexure A.</p> <p>b) DPA already has an Environment Management cell. Further, DPA has also appointed an expert agency to provide Environmental Experts from time to time. Recently, DPA appointed M/s Precitech Laboratories Pvt. Ltd., Vapi, for three years vide work order dated 5/2/2021 (Annexure I).</p> <p>Further, DPA has appointed a Manager Environment on a contractual basis for 3+2 years. A copy of the office order is attached herewith as Annexure J.</p> <p>In addition to the above, DPA appointed M/s Detox Corporation, Surat (NABL Accredited laboratory) for regular Monitoring of environmental parameters including effluent to confirm to the norms prescribed by the GPCB since the year 2016. The work is in progress & DPA have been monitoring data regularly to all the concerned authorities along with compliance reports submitted.</p> <p>The Environmental Monitoring Reports, as submitted by M/s Detox Corporation, is enclosed herewith as Annexure E.</p>
26	An Environmental report indicating the changes, if any, with respect to the baseline environmental quality in the coastal and marine environment shall be submitted every year by the KPT to this Department as well as to the MoEF&CC, GOI.	DPA appointed M/s Detox Corporation, Surat (NABL Accredited laboratory) for regular Monitoring of environmental parameters including effluent to confirm to the norms prescribed by the GPCB since the year 2016. The work is in progress & DPA have been monitoring data regularly to all the concerned

		<p>authorities along with compliance reports submitted.</p> <p>The Environmental Monitoring Reports, as submitted by M/s Detox Corporation, is enclosed herewith as Annexure E.</p>
27	The KPT shall have to contribute financially to support the National Green Corps Scheme being implemented in Gujarat by the GEER Foundation, Gandhinagar, in construction with Forests and Environment Department.	Point Noted.
28	A six-monthly report on compliance of the conditions mentioned in this letter shall have to be furnished by the KPT on regular basis to this department/MoEF, GOI.	Point Noted.
29	Any other condition that may be stipulated by this department from time to time for environmental protection/management purpose shall also have to be complied with by the KPT.	Point noted.

Annexure -A



Kandla Oil Terminal Private Limited

Registered Office: "NEELADRI", 3rd Floor, No. 9, Cenotaph Road, Alwarpet, Chennai - 600 018.

Tel: +91-44-4590 2222, 4590 2299, Fax: + 91-44-4590 2200, URL : www.imc.net.in CIN: U60200TN2013PTC092551

KRO/KOTPL/17122022

December 17, 2022

The Superintending Engineer (Design)

Deendayal Port Authority

Administrative -Office

Gandhidham

Kutch 370 201

Dear Sir,

Sub. : Development of Oil Jetty to handle Liquid Bulk and Ship bunkering Terminal at Old Kandla ("Project").-
Half Yearly EC & CRZ Compliance report

The half-yearly compliance reports for the KOTPL project for the period from June 2022 to November 2022 are enclosed. (EC, CRZ & CTE)

We would appreciate your acknowledgment of receipt of these documents.

Yours sincerely,

For Kandla Oil Terminal (P) Limited

(Authorized Signatory)

17/12/2022.

CC: Independent Engineer, IITM

Encls.:

1. EC Compliance report
2. CRZ Compliance report
3. CTE Compliance report
4. Monitoring Data sheet
5. Ambient Air (Six Months)
6. Noise Monitoring (Six Months)
7. Drinking Water Report (Six Months)
8. Construction Activities Photographs

145(PL)
06/01/23

17(0)
06/01/2023

08(0)
06/01/2023

SE (Emc) &
6/1/23

Manager(Emc)/Shri Anand Dave
EMC
6/1



Monitoring Report (Up to November, 2022)
DATA SHEET

Sr. No.	Particulars	Reply
1.	Project type: River valley/ Mining/Industry/ thermal/nuclear/Other (specify)	Development of Oil Jetty to handle Liquid Bulk and Ship bunkering Terminal at Old Kandla
2.	Name of the project	Development of Oil Jetty to handle Liquid Bulk and Ship bunkering Terminal at Old Kandla
3.	Clearance Letter (s). OM no and date	MoEFCC File No. F.No.11-82/2011-IA-III Proposal No. IA/GJ/MIS/28772/2011 Dated 16 th May 2016
4.	Location a) District (s) b) State (s)	Location: a) Kutch b) Gujarat
5.	Address for Correspondence a) address of Concerned Project Chief Engineer (with pin code & telephone/telex/fax numbers b) Address of Executive project Engineer/manager/ (with pin code fax numbers)	Regional Head Kandla Regional Office Kutch 370 201 Deputy General Manager Kandla Regional Office Administrative -Office Gandhidham Kutch 370 201
6.	Salient features a) Of the Project b) Of the Environmental Management Plan	Jetty : 3.39 MMTPA Tank farm : 1,64,500 KL (As Per Concession Agreement) Approved Construction Capacity About 1,36,417 KL (As per revised statutory/OISD norms duly approved by the competent Authorities) & Allied Facilities
7.	Production Details during compliance period and (or) during the previous financial year	Project is under construction stage.
8.	Breakup of the project area a) Submergence area: forest & non-	N/A





	forest b) Others	
9.	Breakup of the project affected population with enumeration of those loing houses/dwelling units only agricultural land & landless laborer's/artisen a) SC. ST/Adivasis b) Others (please indicate whether these figures arebased on any scientific and systematic survey carried out of only provisional figures, if a survey is carried out give details and years of survey).	Not Applicable
10.	Financial details a) Project cost as originally planned and subsequent revised estimates and the year of prices reference b) Allocation made for environmental management plans with item wise and year wise break-up c) Benefit cost ratio/Internal rate of Return and the year of assessment Whether (c) includes the cost of environmental management plans so far. d) Actual expenditure incurred on the project e) Actual expenditure incurred on the environmental management plans so far.	Estimated Project cost: Rs. 233.50 Cr(Estimated by the Concessioneing Authority in RFQ. Revised project cost: Rs.343 Cr. Rs. 10 Lacs Rs.22.04 Cr. (Till November 22) Rs. 7 Lakhs
11.	Forest land requirement a) The status of approval for diversion of forest land for non-forestry use b) The status of clear felling	Nil N/A. N/A





	c) The status of compensatory a forestation, if any	N/A
	d) Comments on the viability & sustainability of compensatory a forestation programmed in the light of actual field experience so far	N/A
12.	The status of clear felling in non-forest areas (such as submergence area of reservoir, approach roads), if any with quantitative information.	N/A
13.	Status of construction a) Date of commencement (Actual and/or planned) b) Date of completion (Actual and/or planned)	Project is under construction stage. Award of concession: December, 2020. Construction schedule- 24 months. Planned date of Completion.: 4th week of July 2024
14.	Reasons for the delay if the Project is yet to start	Project is under construction stage and delayed because on Pandemic & Local Hindrances.
15.	Date of site visited a) The dates on which the project was monitored by the regional office on pervious occasion. if any b) The date site visit for this monitoring report	No
16.	Details of the correspondence with project authorities for obtaining action plans/information on status of compliance to safeguard other than the routine letters for logistic support for site visit. (The first monitoring report may contain the details of all the letters issued so far but the later reports may cover only the letters issued subsequently.)	Noted.





Subject: Point-wise Compliance Status Report for Environmental clearance for Developing integrated facility within the existing Kandla Port at Kandla, Dist: Kutch by M/s. Kandla Port Trust Limited – Reg.

Ref No: - Environmental Clearance vide Letter No- F. No. 11-82/2011-IA III dated 19.12.2016

Sr. No.	EC Conditions	Compliance Status
PART A – SPECIFIC CONDITIONS		
I	Construction activity shall be carried out strictly according to the provisions of CRZ Notification 2011 No. construction work other than those permitted in coastal Regulation Zone Notification Shall be carried out in Coastal Regulation Zone area	Noted
ii	The project proponent shall ensure that there shall be no damage to the existing mangroves patches near site and also ensure the free flow of water to avoid damage to the mangroves.	Noted
iii	The project proponent shall ensure that no creeks or rivers are blocked due to any activities at the project site, and free flow of water is maintained.	Noted
iv	The shoreline should not be disturbed due to dumping. Periodical study on shoreline changes shall be conducted, and mitigation carried out, if necessary. The details shall be submitted along with the six-monthly monitoring reports.	No shoreline is disturbed.
v	The foreshore facilities shall be set up in the stable/low or medium eroding site as demarcated in the shoreline change map by NCSCM. Further, NCSCM shall be authorized to	Noted





Sr. No.	EC Conditions	Compliance Status
	monitor the project during the construction and operation phases so as to ensure that the foreshore facilities cause minimum or no impact to the geomorphological systems.	
vi	The PP should take measures to ensure that construction materials/debris (mortar, cementing material, etc.) do not fall in the water. Construction material including labour camps should be located at adequate distance from CRZ areas.	Noted.
vii	Dredged materials should be analyzed for the presence of contaminants and also to decide the disposal options. Monitoring of dredging activities should be conducted, and the findings should be shared with the Gujarat SPCB and the Regional office of the Ministry.	No dredging activity carried out till date.
viii	PP in consultation with GCZMA should prepare a regional strategic impact assessment report with a special focus on the region where the PP started construction without permission. The cost towards this study should be borne by the PP	Noted
ix	A comprehensive and integrated conservation plan including a detailed bathymetry study and protection of creeks/mangrove area including buffer zone, mapping of coordinates, running length, HTL, and CRZ boundary should be put in the place. The plan should take note of all the conditions of approvals granted to all the project proponents in this area, and the reported cases of the disappearance of mangroves	DPA appointed Gujrat Institute of Desert Ecology, Bhuj for the said work.





Sr. No.	EC Conditions	Compliance Status
	near the project site. The preservation of the entire area to maintain the fragile ecological conditions should be a part of the plan in relation to the creek and mangrove conservation.	
x	The commitments made during the Public Hearing and recorded in the minutes shall comply with by letter and spirit. A hard copy of the action taken shall be submitted to the ministry.	Not Applicable
xi	All the conditions stipulated in the earlier clearance including the recommendations of the Environment Management Plan, and Disaster Management Plan shall be strictly complied with.	Noted
xii	Disposal sites for excavated material should be so designed that the revised land use after dumping and changes in the land use pattern does not interfere with the natural drainage.	Noted
xiii	PP shall install a continuous automatic ambient air quality monitoring system (24x7) for all relevant parameters at two locations to monitor the ambient air quality status of the project area. Data should be transferred online to CPCB and SPCB websites.	We are conducting AAQ monitoring as per CPCB guidelines and are attached as Annexure .
xiv	The groundwater shall not be tapped within the CRZ areas by the PP to meet the water requirement in any case.	Noted
xv	Necessary arrangements for the treatment of the effluents and solid wastes must be made and it must be ensured that they conform to the standards laid down by the competent authorities including the Central	The project is under the construction stage.





Sr. No.	EC Conditions	Compliance Status
	or State Pollution Control Board and under the Environment (Protection) Act, 1986.	
xvi	All the operational areas will be connected with the network of liquid waste collection corridors comprising of stormwater, oily waste and sewage collection pipelines.	The project is under the construction stage.
xvii	Automatic /online monitoring system (24x7) monitoring devices) for water pollution in respect of flow measurement and relevant pollutants in the treatment system to be installed. The data to be made available to the respective SPCB and in the Company's website.	Not Applicable
xviii	Marine ecology shall be monitored regularly also in terms of seaweeds, sea grasses, mudflats, sand dunes, fisheries, echinoderms, shrimps, turtles, corals, coastal vegetation, mangroves and other marine biodiversity components as part of the management plan. Marine ecology shall be monitored regularly also in terms of all micro, macro and mega floral and faunal components of marine biodiversity.	DPA appointed the Gujarat Institute of Desert Ecology, Bhuj for Regular Monitoring of Marine Ecology in May 2017. Present work was assigned by DPT for 2021-2024.
xix	Measures should be taken to contain, control and recover the accidental spills of fuel and cargo handle.	Noted
xx	All the mitigation measures submitted in the EIA report shall be prepared in a matrix format and the compliance for each mitigation plan shall be submitted to the RO, MoEF&CC along with half yearly compliance report.	Noted





Sr. No.	EC Conditions	Compliance Status
xxi	Ships/barges shall not be allowed to release any oily bilge waste or ballast water in the sea. Any effluents from the Jetty which have leachable characteristics shall be segregated and recycled/disposed of as per SPCB guidelines.	The project is under the construction stage.
xxii	The location of DG sets and other emission-generating equipment shall be decided keeping in view the predominant wind direction so that emissions do not affect nearby residential areas. Installation and operation of DG sets shall comply with the guidelines of CPCB.	Not Applicable
xxiii	All the mechanized handling systems and other associated equipment such as hoppers, belt conveyors, stackers cum reclaimers shall have integrated dust suppression systems. Dust suppression systems shall be provided at all transfer points.	Not applicable as this project is for the handling of liquid cargo.
xxiv	No product other than permitted under the CRZ notification, 2011 shall be stored in the CRZ area.	Noted.
xxv	It shall be ensured by the Project Proponent that the activities do not cause disturbance to the fishing activity, movements of fishing boats and destruction of mangroves during the construction and operation phase.	Noted.
xxvi	As proposed, a green belt over an area of 36.8 ha shall be developed with at least 10-meter-wide green belt on all sides along the periphery of the project area, in the downward direction, and along roadsides etc. Selection of plant species shall be as per the CPCB guidelines in	Noted.





Sr. No.	EC Conditions	Compliance Status
	consultation with the DFO.	
xxvii	Mangrove plantation in an area of 100 ha. shall be carried out by KPT within 2 years in a time bound manner. Action taken report shall be submitted to the Regional Office of MoEF &CC.	Not Applicable
xxviii	Municipal solid wastes and hazardous wastes shall be managed as per the Municipal Solid Waste Rule, 2016 and Hazardous Waste Management Rule, 2016.	Noted.
xxix	The Project Proponent shall take up and earmark adequate funds for socio-economic development and welfare measures as proposed under the CSR program. This shall be taken up on priority.	Noted.
xxx	The project proponent shall set up a separate environmental management cell for the effective implementation of the stipulated environmental safeguards under the supervision of a Senior Executive.	Noted.
xxxi	The funds earmarked for the environment management plan shall be included in the budget, and this shall not be diverted for any other purposes.	Noted.
xxxii	The proponent shall abide by all the commitments and recommendations made in the EIA/EMP report and also during their presentation to the EAC.	Noted.
xxxiii	The company shall prepare an operating manual in respect of all activities. It shall cover all safety & environmental related issues and systems. Measures to be taken for protection. One set of the environmental manual shall	Noted.





Sr. No.	EC Conditions	Compliance Status
	be made available at the project site. Awareness shall be created at each level of management. All the schedules and results of environmental monitoring shall be available at the project site office.	
xxxiv	Corporate Social Responsibility.	
a.	The Company shall have a well-laid-down Environment Policy approved by the Board of Directors.	Noted.
b.	The Environment Policy shall prescribe for standard operating processes/procedures to bring into focus any infringements/deviation/violations of the environmental or forest norms/conditions.	Noted.
c.	The hierarchical system or Administrative Order of the company to deal with environmental issues and for ensuring compliance with the environmental clearance conditions shall be furnished.	Noted.
d.	To have proper checks and balances, the company shall have a well-laid-down system of reporting of non-compliances/violations of environmental norms to the board of Directors of the company and/or shareholders or stakeholders at large.	Noted.
B. GENERAL CONDITIONS:		
(i)	The project authorities must strictly adhere to the stipulations made by the State Pollution Control Board (SPCB), State Government and any other statutory authority.	Noted.
(ii)	Full support shall be extended to the officers of this Ministry/ Regional Office at Bhopal by the	Noted.



Sr. No.	EC Conditions	Compliance Status
	project proponent during the inspection of the project for monitoring purposes by furnishing full details and an action plan including action is taken reports in respect of mitigation measures and other environmental protection activities.	
(iii)	A six-Monthly monitoring report shall need to be submitted by the project proponents to the Regional Office of this Ministry at Bhopal regarding the implementation of the stipulated conditions.	Noted.
(iv)	Ministry of Environment, Forest and Climate Change or any other competent authority may stipulate any additional conditions or modify the existing ones, if necessary, in the interest of the environment and the same shall be complied with.	Noted.
(v)	The Ministry reserves the right to revoke this clearance if any of the conditions stipulated have not complied with the satisfaction of the Ministry.	Noted.
(vi)	In the event of a change in the project profile or change in the implementation agency, a fresh reference shall be made to the Ministry of Environment, Forest and Climate Change.	Noted.
(vii)	The project proponents shall inform the Regional Office as well as the Ministry, of the date of financial closure and final approval of the project by the concerned authorities and the date of start of land development work.	Noted.
(viii)	A copy of the clearance letter	Complied



Sr. No.	EC Conditions	Compliance Status
	shall be marked to the concerned Panchayat/local NGO, if any, from whom any suggestion/representation has been made or received while processing the proposal.	
(ix)	A copy of the environmental clearance letter shall also be displayed on the website of the concerned State Pollution Control Board. The EC letter shall also be displayed at the Regional Office, District Industries centre and Collector's Office/Tehsildar's office for 30 days.	Complied.
11	These stipulations would be enforced among others under the provisions of the Water (Prevention and Control of Pollution) Act 1974, the Air (Prevention and Control of Pollution) Act 1981, the Environment (Protection) Act 1986, the Public Liability (Insurance) Act, 1991 and EIA Notification 1994, including the amendments and rules made thereafter.	Noted.
12	All other statutory clearances such as the approvals for storage of diesel from Chief Controller of Explosives, Fire Department, Civil Aviation Department, Forest Conservation Act, 1980 and Wildlife (Protection) Act, 1972 etc. shall be obtained, as applicable by project proponents from the respective competent authorities.	Till date no tank farm construction activity has been started.
13	The project proponent shall advertise in at least two local Newspapers widely circulated in the region, one of which shall be in the vernacular language	Complied



Sr. No.	EC Conditions	Compliance Status
	informing that the project has been accorded Environmental and CRZ Clearance and copies of clearance letters are available with the State Pollution Control Board and may also be seen on the website of the Ministry of Environment, Forest and Climate Change at http://www.envfor.nic.in . The advertisement should be made within Seven days from the date of receipt of the Clearance letter and a copy of the same should be forwarded to the Regional office of this Ministry at Bhopal.	
14	This Clearance is subject to a final order of the Hon'ble Supreme Court of India in the matter of Goa Foundation Vs Union of India in Writ Petition (Civil) No. 460 of 2004 as may be applicable to this product.	Noted.
15	The status of compliance with the various stipulated environmental conditions and environmental safeguards will be uploaded by the project proponent on its website.	Noted.
16	Any appeal against this Clearance shall lie with the National Green Tribunal, if preferred, within a period of 30 days as prescribed under Section 16 of the National Green Tribunal Act, 2010.	Noted.
17	A copy of the clearance letter shall be sent by the proponent to the concerned Panchayat, Zilla Parishad/Municipal Corporation, Urban Local Body and the Local NGO, if any, from whom suggestions/ representations, if any, were received while processing the proposal. The	Complied.



Sr. NO.	EC Conditions	Compliance Status
	clearance letter shall also be put on the website of the company by the proponent.	
18	The proponent shall upload the status of compliance with the stipulated EC conditions, including results of monitored data on their website and shall update the same periodically. It shall simultaneously be sent to the Regional Office of MoEFCC, the respective Zonal Office of CPCB and the SPCB.	Noted.
19	The environmental statement for each financial year ending 31 st March in Form-V as is mandated to be submitted by the project proponent to the concerned State Pollution Control Board as prescribed under the Environment (Protection) Rules, 1986, as amended subsequently, shall also be put on the website of the company along with the status of compliance of EC conditions and shall also be sent to the respective Regional Offices of MoEFCC by e-mail.	Noted.





Subject: Point-wise Compliance Status Report for CRZ clearance for Developing integrated facility within the existing Kandla Port at Kandla, Dist: Kutch by M/s. Kandla Port Trust Limited – Reg.

Ref No: - GCZMA CRZ recommendation vide Letter No- ENV-10-2014-25-E Cell dated 01.07.2015

S. No.	CRZ Conditions	Compliance Status
	SPECIFIC CONDITIONS	
1.	The provisions of the CRZ notification of 2011 shall be strictly adhered to by the KPT. No activity in contradiction to the provisions of the CRZ Notification shall be carried out by the KPT.	Noted.
2.	The KPT shall have to ensure that there shall not be any damage to the existing mangrove area.	Noted
3.	The KPT shall prepare an emergency plan to protect existing mangroves in case of any eventuality/accident	Noted
4.	The KPT shall have to make a provision that mangrove areas get proper flushing water and free flow of water shall not be obstructed.	Noted
5.	The KPT shall have to abide by whatever decision taken by the GCZMA for violations of CRZ notification 2011	Noted
6.	There shall not be violations of the order dated 9-12-2013 passed by the National Green Tribunal, and accordingly, there shall be no mangrove destruction taking place in the KPT area.	Noted
7.	No dredging, reclamation or any other project-related activities shall be carried out in the CRZ area categorized as CRZ I (i), and it shall have to be ensured that the mangrove habitats and other ecologically important and significant areas, if any, in the region are not affected due to any of the project activities.	Noted
8.	The KPT shall participate financially in installing and operating the Vessel Traffic Management System in the Gulf of Kachchh and shall also take the lead in preparing and operational sing the Regional Oil Spill Contingency plan in the Gulf of Kachchh.	Not Applicable
9.	The KPT shall strictly ensure that no creeks or rivers are blocked due to any activity at Kandla.	Noted





Kandla Oil Terminal Private Limited

S. No.	CRZ Conditions	Compliance Status
10.	Mangrove plantation in an area of 100 ha. shall be carried out by the KPT within 2 years in a time-bound manner on the Gujarat coastline either within or outside the Kandla Port Trust area, and a six-monthly compliance report along with the satellite images shall be submitted to the Ministry of Environment and Forests as well as to this Department without fail.	Not Applicable
11.	No activities other than those permitted by the competent authority under the CRZ Notification shall be carried out in the CRZ area.	Noted
12.	No groundwater shall be tapped for any purpose during the proposed expansion/modernization activities.	Noted
13.	All necessary permissions from different Government Departments/agencies shall be obtained by the KPT before commencing the expansion activities.	Noted
14.	No effluent or sewage shall be discharged into the sea/creek or in the CRZ area, and it shall be treated to conform to the norms prescribed by the Gujarat Pollution Control Board and would be reused/recycled within the plant premises.	No waste water generation during the construction phase
15.	All the recommendations and suggestions given by Mantec Consultants Pvt. Ltd. New Delhi in their Comprehensive Environment Impact Assessment report for conservation/protection and betterment of the environment shall be implemented strictly by the KPT.	Noted
16.	The construction and operational activities shall be carried out in such a way that there is no negative impact on mangroves and other coastal/marine habitats. The construction activities and dredging shall be carried out only under the constant supervision and guidelines of the Institute of National repute like NIOT.	Noted
17.	The KPT shall contribute financially to any common study or project that may be proposed by this Department for environmental management/conservation/improvement for the Gulf of Kutch.	Not applicable
18.	The construction debris and/or any other type of waste shall not be disposed of into the sea, creek or in the CRZ areas. The	Noted





Kandla Oil Terminal Private Limited

S. No.	CRZ Conditions	Compliance Status
	debris shall be removed from the construction site immediately after the construction is over.	
19.	The construction camps shall be located outside the CRZ area, and the construction labour shall be provided with the necessary amenities, including sanitation, water supply and fuel, and it shall be ensured that the environmental conditions are not deteriorated by the construction labours.	Noted.
20.	The KPT shall regularly update their Local Oil Spill Contingency and Disaster Management plan in consonance with the National Oil Spill and Disaster Contingency Plan and shall submit the same to this Department after having it vetted through the Indian Coast Guard.	Noted
21.	The KPT shall bear the cost of the external agency that may be appointed by this Department for supervision/ monitoring of proposed activities and the environmental impacts of the proposed activities.	Not applicable
22.	The KPT shall take up massive greenbelt development activities in and around Kandla and also within the KPT limits.	Not applicable
23.	The KPT shall have to contribute financially for taking up the socio-economic upliftment activities in this region in consultation with the Forests and Environment Department and the District Collector/ District Development officer.	Not applicable
24.	A separate budget shall be earmarked for environmental management and socio-economic activities, and details thereof shall be furnished to this Department as well as MoEF, GOI. The details with respect to the expenditure from this budget head shall also be furnished.	Noted
25.	A separate environmental management cell with qualified personnel shall be created for environmental monitoring and management during the construction and operational phases of the project.	Noted
26.	An environmental report indicating the changes, if any, with respect to the baseline environmental quality in the coastal and marine environment shall be submitted every year by the KPT to this Department as well as to the	Noted





S. No.	CRZ Conditions	Compliance Status
	MoEF&CC,GOI.	
27.	The KPT shall have to contribute financially to support the National Green Corps Scheme being implemented in Gujarat by the GEER Foundation, Gandhinagar, in consultation with the Forests and Environment Department	Not applicable
28.	A six-monthly report on compliance with the conditions mentioned in this letter shall have to be furnished by the KPT on a regular basis to this Department/MoEF&CC,GOI	Noted
29.	Any other conditions that may be stipulated by this Department/MoEF&CC,GOI from time to time for environmental protection/management purposes shall also have to be complied with by the KPT.	Noted.





Subject: Point-wise Compliance Status Report for Consent to Establish for Developing integrated facility within the existing Kandla Port at Kandla, Dist: Kutch by M/s. Kandla Port Trust Limited – Reg.

Ref No: - PC/CCA-KUTCH-1231/GPCB ID 44000 dated 22.12.2015 and Amendment of Consent to Establish dated 04.12.2017

Sr. No.	Condition	Compliance Status								
2.	SPECIFIC CONDITIONS:									
	1. Kandla Port Trust shall strictly adhere to all conditions of CRZ Clearance issued by the Forest & Environment Department vide order no. ENV-10-2014-25-E dated 01/07/2015.	Noted								
	2. CTE is granted conditionally that Kandla Port Trust shall not install & commission, including the construction activity of seven activities mentioned above, without obtaining Environment Clearance from MoEF&CC, New Delhi.	Noted								
	3. Kandla Port Trust shall strictly adhere to all conditions of the Terms of Reference (ToR) (vide letter no. F. No. 11-82/2011-IA.III) by MoEF&CC, New Delhi.	Noted								
3.	<u>CONDITION UNDER THE WATER ACT 1974:</u>									
	3.1 There shall be no industrial effluent generation from the loading and unloading activities at the port and other ancillary operations.	Not applicable								
	3.2 The quantity of Domestic wastewater (Sewage) shall not exceed 6.4 KL/Day.	Not applicable								
	3.3 The quality of the sewage shall conform to the following standards:	Not applicable								
	<table><tr><th>PARAMETERS</th><th>GPCB NORMS</th></tr><tr><td>BOD (5 days at 20 °C)</td><td>20 mg/L</td></tr><tr><td>Suspended solids</td><td>30 mg/L</td></tr><tr><td>Residual Chlorine</td><td>Minimum 0.5 mg/L</td></tr></table>	PARAMETERS	GPCB NORMS	BOD (5 days at 20 °C)	20 mg/L	Suspended solids	30 mg/L	Residual Chlorine	Minimum 0.5 mg/L	
PARAMETERS	GPCB NORMS									
BOD (5 days at 20 °C)	20 mg/L									
Suspended solids	30 mg/L									
Residual Chlorine	Minimum 0.5 mg/L									
	3.4 Sewage shall be disposed of through a septic tank/soak pit system.	Not applicable								
	3.5 The unit shall install meters at utilities for measuring category-wise (Category as given in Schedule II of "Water (Prevention & Control of Pollution) Cess Act-1977") consumption of water.	Not applicable								





CONDITION UNDER THE AIR ACT 1981:

4.1 There shall be no use of fuel hence there shall be no flue and process gas emission from storage handling activity and other ancillary operations.

Noted

4.2 The applicant shall provide portholes, ladder, platform etc at chimney(s) for monitoring the air emissions and the same shall be open for inspection. The chimney(s) vents attached to various sources of emission shall be designed by numbers such as S-1, S-2, etc. and these shall be painted/ displayed to facilitate identification.

Not Applicable

4.3 The concentration of the following parameters in the ambient air within the premises of the industry shall not exceed the limits specified hereunder as per National Ambient Air Quality Standards issued by MoEF&CC dated 16th November-2009.

Noted (Annexure)

Sr. No.	Pollutant	Time Weighted Average	Concentration in Ambient air in $\mu\text{g}/\text{m}^3$
1.	Sulphur Dioxide (SO_2)	Annual 24Hours	50 80
2.	Nitrogen Dioxide (NO_2)	Annual 24Hours	40 80
3.	Particulate Matter (Size less than $10\ \mu\text{m}$) OR PM10	Annual 24Hours	60 100
4.	Particulate Matter (Size less than $2.5\ \mu\text{m}$) OR PM 2.5	Annual 24Hours	40 60

4.4 The level of Noise in ambient air within the premises of the industrial unit shall not exceed the following levels:

Between 6 A.M. to 10 P.M.:75 dB(A)

Between 10 P.M. to 6 A.M.:70 dB(A)

Noted (Annexure)

5. CONDITIONS UNDER HAZARDOUS WASTE:

5.1 The applicant shall provide temporary storage facilities for each type of Hazardous Waste as per Hazardous Waste (Management, Handling & Transboundary Movement) Rules, 2008, as amended from time to time.

Noted

5.2 The applicant shall obtain membership of a common TSDF site for the disposal Hazardous. Waste as categorized in Hazardous

The project is under the construction





Waste (Management, Handling & Transboundary Movement) Rules, 2008, as amended from time to time.

GENERAL CONDITIONS

6.1 Any change in personnel, equipment or working conditions as mentioned in the consent form/order should immediately be intimated to this Board.

6.2 The waste generator shall be totally responsible for (i.e. Collection, storage, transportation and ultimate disposal) the wastes generated.

6.3 Records of waste generation, its management and annual return shall be submitted to the Gujarat Pollution Control Board in Form- 4 by 31st January of every year.

6.4 In case of any accident, details of the same shall be submitted in Form- 5 to the Gujarat Pollution Control Board.

6.5 Applicant shall comply with the relevant provision of "Public Liability Insurance Act-91".

6.6 Unit shall take all concrete measures to show tangible results in waste generation reduction, avoidance, reuse, and recycle. Action taken in this regard shall be submitted within 03 months and also along with Form 4.

6.7 Industry shall have to display online data outside the main factory gate with regard to the quantity and nature of hazardous chemicals being handled in the plant, including wastewater and air emissions and solid hazardous waste generated within the factory premises,

6.8 Adequate plantation shall be carried out all along the periphery of the industrial premises in such a way that the density of plantation is at least 1000 trees per acre of land and a green belt of 10 meters width is developed.

6.9 The applicant shall have to submit the returns in the prescribed form regarding water consumption and shall have to make payment of water cess to the Board under the Water (Prevention and Control of Pollution) Cess Act 1977.

stage

Noted

Noted

Noted

Noted

Noted

The project is under the construction stage

Noted

Noted

The project is under the construction stage





ENVIROTECH LAB PVT. LTD.



TC-10331

TEST REPORT

Customer's Name and Address:

KOTPL
OPP. SHIRVA RAILWAY CROSSING, NEAR LOC
FORESHORE TERMINALS NEW KANDLA -370210

Format No. : 7.8 F-01

Date of : 12/08/2022

Report

Report No : A2ZELPL/DW/08/2022/01

Description of Sample:

Date of Sampling	: 08/08/2022	Type of Sampling	: Drinking Water
Date of Sample Received	: 09/08/2022	Sample ID	: DW/052022/01
Sampling Location of Sampling Point	: Near office area	Sample Particular	: Drinking Water
Sample Collected / Submitted by	: A2ZELPL Team Member	Reference Method for Sampling	: IS 3025 (Part-1)
Sample Quantity /Total No.	: 2 Litre/1Nos.	Date of Analysis Start	: 09/08/2022
Details of Packing/Label/Seal	: Satisfactory	Date of Analysis Completion	: 11/08/2022
Environment condition during the test	: 25 ± 3 °C		

Test Results

Sr. No.	Parameters	Results	STANDARD Limit	Unit	Reference Method
1.	pH @°C	6.9	6.5-8.5	-	APHA, 23rd Edition 2017/4500-H* B
2.	Electrical Conductivity @°C	0.52	--	mS/cm	APHA, 23rd Edition 2017/2510-B
3.	Temperature	25	--	°C	APHA, 23rd Edition 2017/2550B
4.	Total dissolved solids	236	500	mg/L	APHA, 23rd Edition 2017/ 2540-C
5.	Total Suspended solids	21	50-150	mg/L	APHA, 23rd Edition 2017/ 2540-D
6.	Chloride as Cl	289	250	mg/L	APHA, 23rd Edition 2017/ 4500 Cl- B
7.	Total Hardness as CaCO ₃	64	200	mg/L	APHA, 23rd Edition 2017/2340-C
8.	Calcium as Ca	31	75	mg/L	APHA, 23rd Edition 2017/3500-Ca-B
9.	Magnesium as Mg.	24	30	mg/L	APHA, 23rd Edition 2017/Calculation Method 3500 -Mg B
10.	E Coil	Absent	Shall Not be detectable	Nos./100 ml	APHA 23rd Edition Part - 9000 Section : 9221D

A.A. JALOR
Tested By
(Sr. Analyst/Analyst)

Checked By/Prepared By
Dr. Jagruti P. Matang
(Technical Manager)

Akhbar Khan P. Jalori
Authorized Signatory
(Quality Manager)

This Report is issued under the following terms & Condition:

1. The results relate only to the items tested and for applicable parameter
2. This Test report shall not be reproduced in full or part for any promotional or publicity and can't be used as evidence in court of law without the written consent of A 2 Z Envirotech Lab Private Limited
3. Reanalysis of the sample will be done if requested Within 15 days from the date of reporting of sample if the samples are not consumed during analysis.

End of Report



ENVIROTECH LAB PVT. LTD.

TEST REPORT



TC-10331

Customer's Name and Address:

KOTPL
OPP. SHIRVA RAILWAY CROSSING, NEAR LOC
PORESHORE TERMINALS NEW KANDLA -370210

Format No. : 7.8 F-01
Date of : 12/08/2022
Report :
Report No : A2ZELPL/AA/08/2022/01

Description of Sample:

Date of Sampling	: 08/08/2022	Type of Sampling	:
Date of Sample Received	: 09/08/2022	Sample ID	: AA/082022/01
Sampling Location of Sampling Point	: NEAR TANK FARM AREA	Sample Particular	: AMBIENT AIR
Sample Collected / Submitted by	: A2ZELPL Team Member	Reference Method for Sampling	: -----
Sample Quantity /Total No.	: 2 Filter paper 1SO ₂ × 35 ml, 1NO ₂ × 35 ml.	Date of Analysis Start	: 09/08/2022
Details of Packing/Label/Seal	: Satisfactory	Date of Analysis Completion	: 10/08/2022
Environment condition during the test	: 25 ± 3 °C	Metrological conditions during monitoring	: Clear Sky
Instrument code	: A2ZELPL/RDS/01 A2ZELPL/FDS/01	Actual Duration of Monitoring (Hours)	: 24

Test Results

Sr. No.	Parameters	Results	Unit	NAAQS LIMITS	Reference Method
1.	Particulate Matter PM ₁₀	79	µg/m ³	100 µg/m ³	IS 5182 (Part 23)2006/ Reaffirmed 2017
2.	Particulate Matter PM _{2.5}	36	µg/m ³	60 µg/m ³	IS 5182 (Part 24)2019
3.	Sulphur Dioxide SO ₂	20	µg/m ³	80 µg/m ³	IS 5182 (Part 2)2001/ Reaffirmed 2017
4.	Nitrogen Dioxide NO ₂	38	µg/m ³	80 µg/m ³	IS 5182 (Part 6)2006/ Reaffirmed 2017

A. A. Jalori
Tested By
(Sr. Analyst/Analyst)

Checked By/Prepared By
Dr. Jagruti P. Matang
(Technical Manager)

Authorized Signatory
Akbarkhan P. Jalori
(Quality Manager)

This Report is issued under the following terms & Condition:

- The results relate only to the items tested and for applicable parameter
- This Test report shall not to be reproduced in full or part for any promotional or publicity and can't be used as evidence in court of law without the written consent of A 2 Z Envirotech Lab Private Limited
- Reanalysis of the sample will be done if requested Within 15 days from the date of reporting of sample if the samples are not consumed during analysis.

End of Report



ENVIROTECH LAB PVT. LTD.

TEST REPORT



TC-10331

Customer's Name and Address:

KOTPL
OPP. SHIRVA RAILWAY CROSSING, NEAR LOC
GORESHORE TERMINALS NEW KANDLA - 370210

Format No. : 7.8 F-01
Date of : 12/08/2022
Report :
Report No : A2ZELPL/AA/08/2022/02

Description of Sample:

Date of Sampling	: 08/08/2022	Type of Sampling	:
Date of Sample Received	: 09/08/2022	Sample ID	: AA/082022/02
Sampling Location of Sampling Point	: NEAR JEETY LAND FALL POINT	Sample Particular	: AMBIENT AIR
Sample Collected / Submitted by	: A2ZELPL Team Member	Reference Method for Sampling	: -----
Sample Quantity / Total No.	: 2 Filter paper 1SO ₂ × 35 ml, 1NO ₂ × 35 ml.	Date of Analysis Start	: 09/08/2022
Details of Packing/Label/Seal	: Satisfactory	Date of Analysis Completion	: 10/08/2022
Environment condition during the test	: 25 ± 3 °C	Metrological conditions during monitoring	: Clear Sky
Document code	: A2ZELPL/RDS/01 A2ZELPL/FDS/01	Actual Duration of Monitoring (Hours)	: 24

Test Results

Sr. No.	Parameters	Results	Unit	NAAQS LIMITS	Reference Method
5.	Particulate Matter PM ₁₀	64	µg/m ³	100 µg/m ³	IS 5182 (Part 23)2006/ Reaffirmed 2017
6.	Particulate Matter PM _{2.5}	40	µg/m ³	60 µg/m ³	IS 5182 (Part 24)2019
7.	Sulphur Dioxide SO ₂	24	µg/m ³	80 µg/m ³	IS 5182 (Part 2)2001/ Reaffirmed 2017
8.	Nitrogen Dioxide NO ₂	36	µg/m ³	80 µg/m ³	IS 5182 (Part 6)2006/ Reaffirmed 2017

A. A. Jalori
Tested By
(Sr. Analyst/Analyst)

Checked By/Prepared By
Dr. Jagruti P. Matang
(Technical Manager)

Akbar Khan P. Jalori
(Quality Manager)

This Report is issued under the following terms & Condition:

- The results relate only to the items tested and for applicable parameter
- This Test report shall not be reproduced in full or part for any promotional or publicity and can't be used as evidence in court
- of law without the written consent of A 2 Z Envirotech Lab Private Limited
- Reanalysis of the sample will be done if requested Within 15 days from the date of reporting of sample if the samples are not consumed during analysis.

End of Report

A2Z Envirotech Lab Pvt. Ltd.
Plot No. 340, Sector 1A, 2nd Floor,
Shiv House, Near Oslo Circle,
Gandhinagar - Kachchh, Gujarat - 370201

Branch:
147, Time Square Empire
Mirzapur Road, Bhuj - Kachchh
Gujarat - 370001

NABL / TC-10331
GST : 24AAU/CA0391F120
info@a2zenvirotechlab.com
www.a2zenvirotechlab.com

Akbar Khan Jalori
+91 98987 11906



ENVIROTECH LAB PVT. LTD.

TEST REPORT



TC-10331

Customer's Name and Address:

SHIRVA RAILWAY CROSSING, NEAR LOC
SHORE TERMINALS NEW KANDLA - 370210

Format No. : 7.8 F-01
Date of : 12/08/2022
Report
Report No : A2ZELPL/AN/082022/01

Description of Sample:

Date of Sampling	: 11/08/2022	Type of Sampling	:
Date of Sample Received	: 11/08/2022	Sample ID	: AN/082022/01
Sampling Location of Sampling	: -----	Sample Particular	: NOISE
Sample Collected / Submitted by	: A2ZELPL Team Member	Reference Method for Sampling	: IS 9989-1991
Instrument calibration status	: OK	Date of Analysis Start	: -----
Metreological conditions during monitoring	: CLEAR SKY	Date of Analysis Completion	: -----
Instrument code	: A2ZELPL/SLM/01	Actual Duration of Monitoring (MINUTE)	: 30

Test Results

LOCATION	Results	GPCB Limit	Unit	Reference Method
NEAR TANK FARM	63	75	dB	IS 9989-1991
NEAR JEETY LANDFALL AREA	60	75	dB	IS 9989-1991
TANK FARM RIGHT SIDE	50	75	dB	IS 9989-1991
TANK FARM LEFT SIDE	56	75	dB	IS 9989-1991

A. A. 301021
Tested By
(Sr. Analyst/Analyst)

Checked By/Prepared By
Dr. Jagruti P. Matang
(Technical Manager)

Authorized Signatory
Akbar Khan P. Jalori
(Quality Manager)

This Report is issued under the following terms & Condition:

- The results relate only to the items tested and for applicable parameter
- This Test report shall not be reproduced in full or part for any promotional or publicity and can't be used as evidence in court of law without the written consent of A 2 Z Envirotech Lab Private Limited
- Reanalysis of the sample will be done if requested Within 15 days from the date of reporting of sample if the samples are not consumed during analysis.

End of Report



ENVIROTECH LAB PVT. LTD.

TEST REPORT



TC-10331

Customer's Name and Address:

OPP. SHIRVA RAILWAY CROSSING, NEAR LOC
FORESHORE TERMINALS NEW KANDLA - 370210

Format No. : 7.8 F-01
Date of : 06/07/2022
Report :
Report No : A2ZELPL/AN/072022/01

Description of Sample:

Date of Sampling	: 05/07/2022	Type of Sampling	: -----
Date of Sample Received	: 05/07/2022	Sample ID	: AN/072022/01
Sampling Location of Sampling Point	: -----	Sample Particular	: ----
Sample Collected / Submitted by	: A2ZELPL Team Member	Reference Method for Sampling	: IS 9989-1991
Instrument calibration status	: OK	Date of Analysis Start	: ----
Metreological conditions during monitoring	: CLEAR SKY	Date of Analysis Completion	: ----
Instrument code	: A2ZELPL/SLM/01	Actual Duration of Monitoring (MINUTE)	: 30

Test Results

LOCATION	Results	CPCB Limit	Unit	Reference Method
NEAR TANK FARM	60	75	dB	IS 9989-1991
NEAR JEETY LANDFALL AREA	55	75	dB	IS 9989-1991
TANK FARM RIGHT SIDE	62	75	dB	IS 9989-1991
TANK FARM LEFT SIDE	57	75	dB	IS 9989-1991

A.A. Jagnani
Tested By
(Sr. Analyst/Analyst)

Checked By/Prepared By
Dr. Jagruti P. Matang
(Technical Manager)

Authorized Signatory
Akbarkhan P. Jalori
(Quality Manager)

This Report is issued under the following terms & Condition:

- The results relate only to the items tested and for applicable parameter
- This Test report shall not to be reproduced in full or part for any promotional or publicity and can't be used as evidence in court of law without the written consent of A 2 Z Envirotech Lab Private Limited
- Reanalysis of the sample will be done if requested Within 15 days from the date of reporting of sample if the samples are not consumed during analysis.

End of Report



ENVIROTECH LAB PVT. LTD.

TEST REPORT



TC-10331

Customer's Name and Address:

WTP,
OPP SHIRVA RAILWAY CROSSING, NEAR LOC
GORESHORE TERMINALS NEW KANDLA -370210

Format No. : 7.8 F-01
Date of : 10/07/2022
Report :
Report No : A2ZELPL/AA/07/2022/01

Description of Sample:

Date of Sampling	: 05/07/2022	Type of Sampling	:
Date of Sample Received	: 06/07/2022	Sample ID	: AA/072022/01
Sampling Location of Sampling Point	: NEAR TANK FARM AREA	Sample Particular	: AMBIENT AIR
Sample Collected / Submitted by	: A2ZELPL Team Member	Reference Method for Sampling	: -----
Sample Quantity /Total No.	: 2 Filter paper 1SO ₂ × 35 ml, 1NO ₂ × 35 ml.	Date of Analysis Start	: 06/07/2022
Details of Packing/Label/Seal	: Satisfactory	Date of Analysis Completion	: 08/07/2022
Environment condition during the test	: 25 ± 3 °C	Metrological conditions during monitoring	: Clear Sky
Instrument code	: A2ZELPL/RDS/01 A2ZELPL/FDS/01	Actual Duration of Monitoring (Hours)	: 24

Test Results

Sr. No.	Parameters	Results	Unit	NAAQS LIMITS	Reference Method
1.	Particulate Matter PM ₁₀	84	µg/m ³	100 µg/m ³	IS 5182 (Part 23)2006/ Reaffirmed 2017
2.	Particulate Matter PM _{2.5}	42	µg/m ³	60 µg/m ³	IS 5182 (Part 24)2019
3.	Sulphur Dioxide SO ₂	23	µg/m ³	80 µg/m ³	IS 5182 (Part 2)2001/ Reaffirmed 2017
4.	Nitrogen Dioxide NO ₂	46	µg/m ³	80 µg/m ³	IS 5182 (Part 6)2006/ Reaffirmed 2017

A. A. Jalori

Tested By
(Sr. Analyst/Analyst)

Checked By/Prepared By
Dr. Jagruti P. Matang
(Technical Manager)

Authorized Signatory
Akbar Khan P. Jalori
(Quality Manager)

This Report is issued under the following terms & Condition:

- The results relate only to the items tested and for applicable parameter
- This Test report shall not be reproduced in full or part for any promotional or publicity and can't be used as evidence in court of law without the written consent of A 2 Z Envirotech Lab Private Limited
- Reanalysis of the sample will be done if requested Within 15 days from the date of reporting of sample if the samples are not consumed during analysis.

End of Report

A2Z Envirotech Lab Pvt. Ltd.
Plot No. 340, Sector 1A, 2nd Floor,
Shiv House, Near Oslo Circle,
Gandhinagar - Kachchh, Gujarat - 370201

Branch:
147, Time Square Empire,
Mirjapur Road, Bhuj - Kachchh
Gujarat - 370001

NABL / TC-10331
GST : 24AAUCA0391F120
Info@azzenvirotechlab.com
www.azzenvirotechlab.com

Akbar Khan Jalori
+ 91 98987 11906



ENVIROTECH LAB PVT. LTD.

TEST REPORT



TC-10331

Customer's Name and Address:

KOTPL
OPP. SHIRVA RAILWAY CROSSING, NEAR LOC
FORESHORE TERMINALS NEW KANDLA -370210

Format No. : 7.8 F-01
Date of Report : 10/07/2022
Report No : A2ZELPL/AA/07/2022/02

Description of Sample:

Date of Sampling	: 05/07/2022	Type of Sampling	:
Date of Sample Received	: 06/07/2022	Sample ID	: AA/072022/02
Sampling Location of Sampling Point	: NEAR JEETY LAND FALL POINT	Sample Particular	: AMBIENT AIR
Sample Collected / Submitted by	: A2ZELPL Team Member	Reference Method for Sampling	: -----
Sample Quantity /Total No.	: 2 Filter paper 1SO ₂ × 35 ml, 1NO ₂ × 35 ml.	Date of Analysis Start	: 06/07/2022
Details of Packing/Label/Seal	: Satisfactory	Date of Analysis Completion	: 08/07/2022
Environment condition during the test	: 25 ± 3 °C	Metrological conditions during monitoring	: Clear Sky
Instrument code	: A2ZELPL/RDS/01 A2ZELPL/FDS/01	Actual Duration of Monitoring (Hours)	: 24

Test Results

Sr. No.	Parameters	Results	Unit	NAAQS LIMITS	Reference Method
5.	Particulate Matter PM ₁₀	80	µg/m ³	100 µg/m ³	IS 5182 (Part 23)2006/ Reaffirmed 2017
6.	Particulate Matter PM _{2.5}	46	µg/m ³	60 µg/m ³	IS 5182 (Part 24)2019
7.	Sulphur Dioxide SO ₂	20	µg/m ³	80 µg/m ³	IS 5182 (Part 2)2001/ Reaffirmed 2017
8.	Nitrogen Dioxide NO ₂	52	µg/m ³	80 µg/m ³	IS 5182 (Part 6)2006/ Reaffirmed 2017

A.A. Jalori
Tested By
(Sr. Analyst/Analyst)

Checked By/Prepared By
Dr. Jagruti P. Matang
(Technical Manager)

Authorized Signatory
Akbarkhan P. Jalori
(Quality Manager)

This Report is issued under the following terms & Condition:

- The results relate only to the items tested and for applicable parameter
- This Test report shall not be reproduced in full or part for any promotional or publicity and can't be used as evidence in court of law without the written consent of A 2 Z Envirotech Lab Private Limited
- Reanalysis of the sample will be done if requested Within 15 days from the date of reporting of sample if the samples are not consumed during analysis.

End of Report

A2Z Envirotech Lab Pvt. Ltd.
Plot No. 340, Sector 1A, 2nd Floor,
Shiv House, Near Oslo Circle,
Gandhinagar - Kachchh, Gujarat - 370201

Branch:
147, Time Square Empire
Mirzapur Road, Bhuj - Kachchh
Gujarat - 370001

NABL / TC-10331
GST: 24AAUCA0391F1Z0
info@a2zenvirotechlab.com
www.a2zenvirotechlab.com

Akbarkhan Jalori
+ 91 98987 11906



ENVIROTECH LAB PVT. LTD.



TC-10331

TEST REPORT

Customer's Name and Address:

KOTPL
OPP. SHIRVA RAILWAY CROSSING, NEAR LOC
FORESHORE TERMINALS NEW KANDLA -370210

Format No. : 7.8 F-01
Date of : 15/07/2022
Report :
Report No : A2ZELPL/DW/07/2022/01

Description of Sample:

Date of Sampling	: 12/07/2022	Type of Sampling	: Drinking Water
Date of Sample Received	: 12/07/2022	Sample ID	: DW/072022/01
Sampling Location of Sampling Point	: Near office area	Sample Particular	: Drinking Water
Sample Collected / Submitted by	: A2ZELPL Team Member	Reference Method for Sampling	: IS 3025 (Part-1)
Sample Quantity /Total No.	: 2 Litre/1Nos.	Date of Analysis Start	: 13/07/2022
Details of Packing /Label/Seal	: Satisfactory	Date of Analysis Completion	: 14/07/2022
Environment condition during the test	: 25 ± 3 °C		

Test Results

Sr. No.	Parameters	Results	STANDARD Limit	Unit	Reference Method
1	pH @°C	7.2	6.5-8.5	-	APHA, 23 rd Edition 2017/4500-H* B
2	Electrical Conductivity @°C	0.55	--	mS/cm	APHA, 23 rd Edition 2017/2510-B
3	Temperature	25	--	°C	APHA, 23 rd Edition 2017/2550B
4	Total dissolved solids	242	500	mg/L	APHA, 23 rd Edition 2017/ 2540-C
5	Total Suspended solids	22	50-150	mg/L	APHA, 23 rd Edition 2017/ 2540-D
6	Chloride as Cl	28.4	250	mg/L	APHA, 23 rd Edition 2017/ 4500 Cl- B
7	Total Hardness as CaCO ₃	62	200	mg/L	APHA, 23 rd Edition 2017/2340-C
8	Calcium as Ca	34	75	mg/L	APHA, 23 rd Edition 2017/3500-Ca-B
9	Magnesium as Mg.	23	30	mg/L	APHA, 23 rd Edition 2017/Calculation Method 3500 -Mg B
10.	E Coil	Absent	Shall Not be detectable	Nos./100 ml	APHA 23 rd Edition Part - 9000 Section : 9221D

A.A. Jalori

Tested By

(Sr. Analyst/Analyst)

Checked By/Prepared By

Authorized Signatory
Akbar Khan P. Jalori
(Quality Manager)

- This Report is issued under the following terms & Condition:
1. The results relate only to the items tested and for applicable parameter
 2. This Test report shall not be reproduced in full or part for any promotional or publicity and can't be used as evidence in court of law without the written consent of A 2 Z Envirotech Lab Private Limited
 3. Reanalysis of the sample will be done if requested Within 15 days from the date of reporting of sample if the samples are not consumed during analysis.

End of Report

Page No.: 1 of 1

A2Z Envirotech Lab Pvt. Ltd.
Plot No. 340, Sector 1A, 2nd Floor,
Shiv House, Near Oslo Circle,
Gandhidham - Kachchh, Gujarat - 370201

Branch:
147, Time Square Empire
Mirjapur Road, Bhuj - Kachchh
Gujarat - 370001

NABL / TC-10331
GST : 24AAUAC0391F120
info@a2zenvirotechlab.com
www.a2zenvirotechlab.com

Akbar Khan Jalori
+ 91 98987 11906



TC-10331

Test Results

Customer's Name and Address:

KOTPL
Opp. Shirva Railway Crossing, Near loc
Foreshore Terminals New kandla-370210

Format No. : 7.8 F-01
Date of : 29/06/2022
Report
Report No : A2ZELPL/AA/062022/05

Description of Sample:

Date of Sampling	: 26/06/2022	Sample Particular	: Ambient Air
Date of Sample Received	: 26/06/2022	Sample ID	: AA/062022/05
Sampling Location of Sampling Point	: Tank Farm Area	Instrument calibration status	: Ok
Sample Collected / Submitted by	: A2ZELPL Team Member	Reference Method for Sampling	: IS 5182
Sample Quantity /Total No.	: 2 Filter paper, 1 SO ₂ ×35 mL, 1 NO ₂ ×35 mL	Date of Analysis Start	: 27/06/2022
Details of Packing/Label/Seal	: Satisfactory	Date of Analysis Completion	: 27/06/2022
Environment condition during the test	: 25 ± 3 °C	Meteorological condition during monitoring	: Clear Sky
Instrument code	: A2ZELPL/RDS/01 A2ZELPL/FDS/01	Actual duration of Monitoring, (Hours)	: 8

Test Results

Sr.No.	Parameter	Result	Unit	Test Method
1	Particulate Matter PM ₁₀	73.81	µg/m ³	IS 5182 (Part 23) 2006/Reaffirmed 2017
2	Particulate Matter PM _{2.5}	32.45	µg/m ³	IS 5182 (Part 24) 2019
3	Sulfur Dioxide SO ₂	16.06	µg/m ³	IS 5182 (Part 2) 2001/Reaffirmed 2017
4	Nitrogen Dioxide NO ₂	22.87	µg/m ³	IS 5182 (Part 6) 2006/Reaffirmed 2017

Checked By/Prepared By
Dr. Jagruti P. Matang
(Technical Manager)

Authorized Signatory
Akbar Khan P. Jalori
(Quality Manager)

Tested By

(Sr. Analyst/Analyst)

This Report is issued under the following terms & Condition:

- The results relate only to the items tested and for applicable parameter
- This Test report shall not be reproduced in full or part for any promotional or publicity and can't be used as evidence in court of law without the written consent of A 2 Z Envirotech Lab Private Limited
- Reanalysis of the sample will be done if requested Within 15 days from the date of reporting of sample if the samples are not consumed during analysis.

End of Report

Page No.: 1 of 1



ENVIROTECH LAB PVT. LTD.



TC-10331

TEST REPORT

Customer's Name and Address

Shri. Shurva Railway Crossing, Near loc
Shore Terminals New kandla-370210

Format No. : 7.8 F-07
Date of : 28/06/2022
Report
Report No : AZZELPL/AN/062022/06

Description of Sample:

Date of Sampling	: 24/06/2022	Type of Sampling	: Ambient Noise
Date of Sample Received	: 24/06/2022	Sample ID	: AN/062022/06
Sampling Location of Sampling	:	Sample Particular	: Noise Monitoring
Sample Collected / Submitted by	: AZZELPL TEAM MEMBER	Reference Method for Sampling	: IS 9989-1991
Instrument code	: AZZELPL/SLM/01	Sampling Duration (Minute)	: 30 min
Instrument calibration status	: Ok	Date of Analysis Start	: -
Meteorological condition during monitoring	: Clear Sky	Date of Analysis Completion	: -

Test Results

Sr. No.	Location	Result	Unit	GPCB Limit	Test Method
1.	Near Tank Farm	69	dB	75	IS 9989-1991
2.	Near jeety landfall Area	59	dB	75	IS 9989-1991
3.	Tank farm right side	68	dB	75	IS 9989-1991
4.	Tank farm Left side	62	dB	75	IS 9989-1991

Authorized Signatory
Akbarkhan P. Jalori
(Quality Manager)

Checked By/Prepared By
Dr. Jagruti P. Matang
(Technical Manager)

Tested By

(Sr. Analyst/Analyst)

This Report is issued under the following terms & Condition:

- The results relate only to the items tested and for applicable parameter
- This Test report shall not to be reproduced in full or part for any promotional or publicity and can't be used as evidence in court of law without the written consent of A 2 Z Envirotech Lab Private Limited
- Reanalysis of the sample will be done if requested within 15 days from the date of reporting of sample if the samples are not consumed during analysis.

End of Report

Page No.: 1 of 1



ENVIROTECH LAB PVT. LTD.

TEST REPORT

Customer's Name and Address:

SOTPL
OPP. SHIRVA RAILWAY CROSSING, NEAR LOC
FORESHORE TERMINALS NEW KANDLA -370210

Format No. : 7.8 F-01
Date of : 28/06/2022
Report :
Report No : A2ZELPL/DW/06/2022/25

Description of Sample:

Date of Sampling	: 25/06/2022	Type of Sampling	: Drinking Water
Date of Sample Received	: 25/06/2022	Sample ID	: DW/062022/25
Sampling Location of Sampling Point	: Near office area	Sample Particular	: Drinking Water
Sample Collected / Submitted by	: A2ZELPL Team Member	Reference Method for Sampling	: IS 3025 (Part-1)
Sample Quantity /Total No.	: 2 Litre/1Nos.	Date of Analysis Start	: 26/06/2022
Details of Packing/Label/Seal	: Satisfactory	Date of Analysis Completion	: 27/06/2022
Environment condition during the test	: 25 ± 3 °C		

Test Results

Sr. No.	Parameters	Results	STANDARD Limit	Unit	Reference Method
1.	pH @°C	6.9	6.5-8.5	-	APHA, 23 rd Edition 2017/4500-H ⁺ B
2.	Electrical Conductivity @°C	0.47	--	mS/cm	APHA, 23 rd Edition 2017/2511-B
3.	Temperature	25	--	°C	APHA, 23 rd Edition 2017/2550B
4.	Total dissolved solids	263	500	mg/L	APHA, 23 rd Edition 2017/ 2540-C
5.	Total Suspended solids	23	50-150	mg/L	APHA, 23 rd Edition 2017/ 2540-D
6.	Chloride as Cl	26	250	mg/L	APHA, 23 rd Edition 2017/ 4500 Cl ⁻ B
7.	Total Hardness as CaCO ₃	61	200	mg/L	APHA, 23 rd Edition 2017/2340-C
8.	Calcium as Ca	32	75	mg/L	APHA, 23 rd Edition 2017/3500-Ca-B
9.	Magnesium as Mg,	19	30	mg/L	APHA, 23 rd Edition 2017/Calculation Method 3500 -Mg B
10.	E Coil	Absent	Shall Not be detectable	Nos./100 ml	APHA 23 rd Edition Part - 9000 Section : 9221D

A. A. Jalori

Checked By/Prepared By

Authorized Signatory
Akbar Khan P. Jalori
(Quality Manager)

Tested By

(Sr. Analyst/Analyst)

This Report is issued under the following terms & Condition:

- The results relate only to the items tested and for applicable parameter
- This Test report shall not be reproduced in full or part for any promotional or publicity and can't be used as evidence in court
- of law without the written consent of A 2 Z Envirotech Lab Private Limited
- Reanalysis of the sample will be done if requested Within 15 days from the date of reporting of sample if the samples are not consumed during analysis.

End of Report

Page No.: 1 of 1



ENVIROTECH LAB PVT. LTD.

TEST REPORT



TC-10331

Customer's Name and Address:

KOTPL
OPP. SHIRVA RAILWAY CROSSING, NEAR LOC
FORESHORE TERMINALS NEW KANDLA - 370210

Format No. : 7.8 F-01
Date of : 18/11/2022
Report :
Report No : AZZELPL/DW/11/2022/01

Description of Sample:

Date of Sampling	: 15/11/2022	Type of Sampling	: Drinking Water
Date of Sample Received	: 16/11/2022	Sample ID	: DW/112022/01
Sampling Location of Sampling Point	: Near office area	Sample Particular	: Drinking Water
Sample Collected / Submitted by	: AZZELPL Team Member	Reference Method for Sampling	: IS 3025 (Part-1)
Sample Quantity / Total No.	: 2 Litre / 1 Nos.	Date of Analysis Start	: 16/11/2022
Details of Packing/Label/Seal	: Satisfactory	Date of Analysis Completion	: 17/11/2022
Environment condition during the test	: 25 ± 3 °C		

Test Results

Sr. No.	Parameters	Results	STANDARD Limit	Unit	Reference Method
1.	pH @°C	7.3	6.5-8.5	-	APHA, 23 rd Edition 2017/4500-H-B
2.	Electrical Conductivity @°C	0.60	--	mS/cm	APHA, 23 rd Edition 2017/2511-B
3.	Temperature	25	--	°C	APHA, 23 rd Edition 2017/2550B
4.	Total dissolved solids	268	500	mg/L	APHA, 23 rd Edition 2017/ 2540-C
5.	Total Suspended solids	26	50-150	mg/L	APHA, 23 rd Edition 2017/ 2540-D
6.	Chloride as Cl	29.7	250	mg/L	APHA, 23 rd Edition 2017/ 4500 Cl-B
7.	Total Hardness as CaCO ₃	73	200	mg/L	APHA, 23 rd Edition 2017/2340-C
8.	Calcium as Ca	35	75	mg/L	APHA, 23 rd Edition 2017/3500-Ca-B
9.	Magnesium as Mg.	19	30	mg/L	APHA, 23 rd Edition 2017/Calculation Method 3500-Mg B
10.	E Coil	Absent	Shall Not be detectable	Nos./100 ml	APHA 23 rd Edition Part - 9000 Section : 9221D

A.A. Jalori

Tested By
(Sr. Analyst/Analyst)

Checked By/Prepared By
Dr. Jagruti P. Matang
(Technical Manager)

Authorized Signatory
Akbarkhan P. Jalori
(Quality Manager)

This Report is issued under the following terms & Condition:

1. The results relate only to the items tested and for applicable parameter
2. This Test report shall not be reproduced in full or part for any promotional or publicity and can't be used as evidence in court of law without the written consent of A2Z Envirotech Lab Private Limited
3. Reanalysis of the sample will be done if requested Within 15 days from the date of reporting of sample if the samples are not consumed during analysis.

Page No.: 1 of 1



ENVIROTECH LAB PVT. LTD.

TEST REPORT



TC-10331

Customer's Name and Address:

NOTPL
OPP. SHIRVA RAILWAY CROSSING, NEAR LOC
FORESHORE TERMINALS NEW KANDLA - 370210

Format No. : 7.8 F-01
Date of Report : 16/11/2022
Report No : A2ZELPL/AN/112022/01

Description of Sample:

Date of Sampling	: 15/11/2022	Type of Sampling	: -----
Date of Sample Received	: 15/11/2022	Sample ID	: AN/102022/01
Sampling Location of Sampling Point	: -----	Sample Particular	: ----
Sample Collected / Submitted by	: A2ZELPL Team Member	Reference Method for Sampling	: IS 9989-1991
Instrument calibration status	: OK	Date of Analysis Start	: ----
Metrological conditions during monitoring	: CLEAR SKY	Date of Analysis Completion	: ----
Instrument code	: A2ZELPL/SLM/01	Actual Duration of Monitoring (MINUTE)	: 30

Test Results

Sr. No.	LOCATION	Results	CPCB Limit	Unit	Reference Method
1.	NEAR TANK FARM	60	75	dB	IS 9989-1991
2.	NEAR JEETY LANDFALL AREA	45	75	dB	IS 9989-1991
3.	TANK FARM RIGHT SIDE	50	75	dB	IS 9989-1991
4.	TANK FARM LEFT SIDE	52	75	dB	IS 9989-1991

A.A. Jalori

Tested By
(Sr. Analyst/Analyst)

Checked By/Prepared By
Dr. Jagruti P. Matang
(Technical Manager)

Authorized Signatory
Akbar Khan P. Jalori
(Quality Manager)

This Report is issued under the following terms & Condition:

- The results relate only to the items tested and for applicable parameter
- This Test report shall not to be reproduced in full or part for any promotional or publicity and can't be used as evidence in court of law without the written consent of A 2 Z Envirotech Lab Private Limited
- Reanalysis of the sample will be done if requested Within 15 days from the date of reporting of sample if the samples are not consumed during analysis.

End of Report



ENVIROTECH LAB PVT. LTD.

TEST REPORT



TC-10331

Customer's Name and Address:

NDTPL
OPP. SHIRVA RAILWAY CROSSING, NEAR LOC
FORESHORE TERMINALS NEW KANDLA - 370210

Format No. : 7.8 F-01
Date of : 18/11/2022
Report :
Report No : A2ZELPL/AA/11/2022/02

Description of Sample:

Date of Sampling	: 15/11/2022	Type of Sampling	:
Date of Sample Received	: 16/11/2022	Sample ID	: AA/112022/2
Sampling Location of Sampling Point	: NEAR TANK FARM AREA	Sample Particular	: AMBIENT AIR
Sample Collected / Submitted by	: A2ZELPL Team Member	Reference Method for Sampling	: ----
Sample Quantity / Total No.	: 2 Filter paper 1SO ₂ × 35 ml, 1NO ₂ × 35 ml.	Date of Analysis Start	: 16/11/2022
Details of Packing/Label/Seal	: Satisfactory	Date of Analysis Completion	: 17/11/2022
Environment condition during the test	: 25 ± 3 °C	Metrological conditions during monitoring	: Clear Sky
Instrument code	: A2ZELPL/RDS/01 A2ZELPL/FDS/01	Actual Duration of Monitoring (Hours)	: 24

Test Results

Sr. No.	Parameters	Results	Unit	NAAQS LIMITS	Reference Method
1.	Particulate Matter PM ₁₀	72	µg/m ³	100 µg/m ³	IS 5182 (Part 23)2006/ Reaffirmed 2017
2.	Particulate Matter PM _{2.5}	32	µg/m ³	60 µg/m ³	IS 5182 (Part 24)2019
3.	Sulfur Dioxide SO ₂	28	µg/m ³	80 µg/m ³	IS 5182 (Part 2)2001/ Reaffirmed 2017
4.	Nitrogen Dioxide NO ₂	42	µg/m ³	80 µg/m ³	IS 5182 (Part 6)2006/ Reaffirmed 2017

A. A. Jalori

Tested By
(Sr. Analyst/Analyst)

Checked By/Prepared By
)

Authorized Signatory
Akbar Khan P. Jalori
(Quality Manager)

This Report is issued under the following terms & Condition:

- The results relate only to the items tested and for applicable parameter
- This Test report shall not be reproduced in full or part for any promotional or publicity and can't be used as evidence in court of law without the written consent of A 2 Z Envirotech Lab Private Limited
- Reanalysis of the sample will be done if requested Within 15 days from the date of reporting of sample if the samples are not consumed during analysis.

End of Report



ENVIROTECH LAB PVT. LTD.

TEST REPORT



TC-10331

Customer's Name and Address:

SHIRVA RAILWAY CROSSING, NEAR LOC
KORSHORE TERMINALS NEW KANDLA - 370210

Format No. : 7.8 F-01
Date of : 18/11/2022
Report :
Report No : A2ZELPL/AA/11/2022/03

Description of Sample:

Date of Sampling	: 15/11/2022	Type of Sampling	:
Date of Sample Received	: 16/11/2022	Sample ID	: AA/112022/3
Sampling Location of Sampling Point	: NEAR JEETY LAND FALL POINT	Sample Particular	: AMBIENT AIR
Sample Collected / Submitted by	: A2ZELPL Team Member	Reference Method for Sampling	: -----
Sample Quantity / Total No.	: 2 Filter paper 1SO ₂ × 35 ml, 1NO ₂ × 35 ml.	Date of Analysis Start	: 16/11/2022
Details of Packing/Label/Seal	: Satisfactory	Date of Analysis Completion	: 17/11/2022
Environment condition during the test	: 25 ± 3 °C	Metrological conditions during monitoring	: Clear Sky
Instrument code	: A2ZELPL/RDS/01 A2ZELPL/FDS/01	Actual Duration of Monitoring (Hours)	: 24

Test Results

Sr. No.	Parameters	Results	Unit	NAAQS LIMITS	Reference Method
5.	Particulate Matter PM ₁₀	72	µg/m ³	100 µg/m ³	IS 5182 (Part 23)2006/ Reaffirmed 2017
6.	Particulate Matter PM _{2.5}	36	µg/m ³	60 µg/m ³	IS 5182 (Part 24)2019
7.	Sulfur Dioxide SO ₂	26	µg/m ³	80 µg/m ³	IS 5182 (Part 2)2001/ Reaffirmed 2017
8.	Nitrogen Dioxide NO ₂	38	µg/m ³	80 µg/m ³	IS 5182 (Part 6)2006/ Reaffirmed 2017

A. A. Jalori

Tested By
(Sr. Analyst/Analyst)

Checked By/Prepared By

Authorized Signatory
Akbar Khan P. Jalori
(Quality Manager)

This Report is issued under the following terms & Condition:

- The results relate only to the items tested and for applicable parameter
- This Test report shall not be reproduced in full or part for any promotional or publicity and can't be used as evidence in court of law without the written consent of A 2 Z Envirotech Lab Private Limited
- Reanalysis of the sample will be done if requested Within 15 days from the date of reporting of sample if the samples are not consumed during analysis.

End of Report

A2Z Envirotech Lab Pvt. Ltd.
Plot No. 340, Sector 1A, 2nd Floor,
Shiv House, Near Oslo Circle,
Gandhinagar - Kachchh, Gujarat - 370201

Branch:
147, Time Square Empire
Mirzapur Road, Bhuj - Kachchh
Gujarat - 370001

NABL / TC-10331
GST: 24AAUCA0391F1Z0
info@a2zenvirotechlab.com
www.a2zenvirotechlab.com

Akbar Khan Jalori
+ 91 98987 11906



ENVIROTECH LAB PVT. LTD.

TEST REPORT



TC-10331

Customer's Name and Address:

SHIRVA RAILWAY CROSSING, NEAR LOC
SHORE TERMINALS NEW KANDLA -370210

Format No. : 7.8 F-01
Date of : 16/10/2022
Report :
Report No : AZZELPL/AA/010/2022/03

Description of Sample:

Date of Sampling	: 12/10/2022	Type of Sampling	:	
Date of Sample Received	: 13/10/2022	Sample ID	:	AA/102022/03
Sampling Location of Sampling Point	: NEAR JEETY LAND FALL POINT	Sample Particular	:	AMBIENT AIR
Sample Collected / Submitted by	: AZZELPL Team Member	Reference Method for Sampling	:	----
Sample Quantity /Total No.	: 2 Filter paper 1SO ₂ × 35 ml, 1NO ₂ × 35 ml.	Date of Analysis Start	:	13/10/2022
Details of Packing/Label/Seal	: Satisfactory	Date of Analysis Completion	:	15/10/2022
Environment condition during the test	: 25 ± 3 °C	Metrological conditions during monitoring	:	Clear Sky
Instrument code	: AZZELPL/RDS/01 AZZELPL/FDS/01	Actual Duration of Monitoring (Hours)	:	24

Test Results

Sr. No.	Parameters	Results	Unit	NAAQS LIMITS	Reference Method
5.	Particulate Matter PM ₁₀	65	µg/m ³	100 µg/m ³	IS 5182 (Part 23)2006/ Reaffirmed 2017
6.	Particulate Matter PM _{2.5}	30	µg/m ³	60 µg/m ³	IS 5182 (Part 24)2019
7.	Sulfur Dioxide SO ₂	20	µg/m ³	80 µg/m ³	IS 5182 (Part 2)2001/ Reaffirmed 2017
8.	Nitrogen Dioxide NO ₂	32	µg/m ³	80 µg/m ³	IS 5182 (Part 6)2006/ Reaffirmed 2017

Tested By
(Sr. Analyst/Analyst)

Checked By/Prepared By
Dr. Jagruti P. Matang
(Technical Manager)

Authorized Signatory
Akbar Khan P. Jalori
(Quality Manager)

This Report is issued under the following terms & Condition:

- The results relate only to the items tested and for applicable parameter
- This Test report shall not be reproduced in full or part for any promotional or publicity and can't be used as evidence in court of law without the written consent of A Z Z Envirotech Lab Private Limited
- Reanalysis of the sample will be done if requested Within 15 days from the date of reporting of sample if the samples are not consumed during analysis.

End of Report



ENVIROTECH LAB PVT. LTD.



TC-10331

TEST REPORT

Customer's Name and Address:

WTP
RPF SHIRVA RAILWAY CROSSING, NEAR LOC
FORESHORE TERMINALS NEW KANDLA -370210

Format No. : 7.R.F.01
Date of : 15/10/2022
Report
Report No. : A2ZELPL/DW/10/2022/01

Description of Sample:

Date of Sampling	12/10/2022	Type of Sampling	Drinking Water
Date of Sample Received	12/10/2022	Sample ID	DW/102022/01
Sampling Location of Sampling Point	Near office area	Sample Particular	Drinking Water
Sample Collected / Submitted by	A2ZELPL Team Member	Reference Method for Sampling	IS 3025 (Part-1)
Sample Quantity / Total No.	2 Litre / 1 Nos.	Date of Analysis Start	13/10/2022
Details of Packing / Label / Seal	Satisfactory	Date of Analysis Completion	14/10/2022
Environment condition during the test	25 ± 3 °C		

Test Results

Sr. No.	Parameters	Results	STANDARD Limit	Unit	Reference Method
1	pH @°C	6.8	6.5-8.5	-	APHA, 23rd Edition 2017/4500-H ⁺ B
2	Electrical Conductivity @°C	0.45	--	mS/cm	APHA, 23rd Edition 2017/2510-B
3	Temperature	25	--	°C	APHA, 23rd Edition 2017/2550B
4	Total dissolved solids	200	500	mg/L	APHA, 23rd Edition 2017/ 2540-C
5	Total Suspended solids	22	50-150	mg/L	APHA, 23rd Edition 2017/ 2540-D
6	Chloride as Cl	27.5	250	mg/L	APHA, 23rd Edition 2017/ 4500 Cl ⁻ B
7	Total Hardness as CaCO ₃	60	200	mg/L	APHA, 23rd Edition 2017/2340-C
8	Calcium as Ca	30	75	mg/L	APHA, 23rd Edition 2017/3500-Ca-B
9	Magnesium as Mg	22	30	mg/L	APHA, 23rd Edition 2017/Calculation Method 3500 -Mg B
10	E Coil	Absent	Shall Not be detectable	Nos./100 ml	APHA 23rd Edition Part - 9000 Section : 9210

A A - Jalori
Tested By

(Sr. Analyst/Analyst)

Checked By/Prepared By
Dr. Jagruti P. Matang
(Technical Manager)

Authorized Signatory
Akbar Khan P. Jalori
(Quality Manager)

This Report is issued under the following terms & Condition:

1. The results relate only to the items tested and for applicable parameter
2. This Test report shall not be reproduced in full or part for any promotional or publicity and can't be used as evidence in court of law without the written consent of A 2 Z Envirotech Lab Private Limited
3. Reanalysis of the sample will be done if requested Within 15 days from the date of reporting of sample if the samples are not consumed during analysis.

End of Report

Page No.: 1 of 2

A2Z Envirotech Lab Pvt. Ltd.
Plot No. 340, Sector 1A, 2nd Floor,
Shiv House, Near Oslo Circle,
Gandhinagar - Kachchh, Gujarat - 370201

Branch:
147, Time Square Empire
Mirzapur Road, Bhuj - Kachchh
Gujarat - 370001

NABL / TC-10331
GST : 24AAUCA0391F1Z0
Info@azzenvirotechlab.com
www.azzenvirotechlab.com

Akbar Khan Jalori
+ 91 98987 11906



ENVIROTECH LAB PVT. LTD.

TEST REPORT



TC-10331

Customer's Name and Address:

NOTPL
OPP. SHIRVA RAILWAY CROSSING, NEAR LOC
FORESHORE TERMINALS NEW KANDLA - 370210

Format No. : 7.8 F-01

Date of : 13/10/2022

Report

Report No. : A2ZELPL/AN/102022/01

Description of Sample:

Date of Sampling	: 12/10/2022	Type of Sampling	: -----
Date of Sample Received	: 12/10/2022	Sample ID	: AN/102022/01
Sampling Location of Sampling Point	: -----	Sample Particular	: ----
Sample Collected / Submitted by	: A2ZELPL Team Member	Reference Method for Sampling	: IS 9989-1991
Instrument calibration status	: OK	Date of Analysis Start	: -----
Metrological conditions during monitoring	: CLEAR SKY	Date of Analysis Completion	: -----
Instrument code	: A2ZELPL/SLM/01	Actual Duration of Monitoring (MINUTE)	: 30

Test Results

Sr. No.	LOCATION	Results	CPCB Limit	Unit	Reference Method
1.	NEAR TANK FARM	65	75	dB	IS 9989-1991
2.	NEAR JEETY LANDFALL AREA	55	75	dB	IS 9989-1991
3.	TANK FARM RIGHT SIDE	55	75	dB	IS 9989-1991
4.	TANK FARM LEFT SIDE	56	75	dB	IS 9989-1991

A. A. J. 2022
Tested By
(Sr. Analyst/Analyst)

Checked By/Prepared By
Dr. Jagruti P. Matang
(Technical Manager)

Authorized Signatory
Akbar Khan P. Jalori
(Quality Manager)

This Report is issued under the following terms & Condition:

- The results relate only to the items tested and for applicable parameter
- This Test report shall not to be reproduced in full or part for any promotional or publicity and can't be used as evidence in court of law without the written consent of A 2 Z Envirotech Lab Private Limited
- Reanalysis of the sample will be done if requested Within 15 days from the date of reporting of sample if the samples are not consumed during analysis.

End of Report

A2Z Envirotech Lab Pvt. Ltd.
Plot No. 340, Sector 1A, 2nd Floor,
Shiv House, Near Oslo Circle,
Gandhidham - Kachchh, Gujarat - 370201

Branch:
147, Time Square Empire
Mirjapur Road, Bhuj - Kachchh
Gujarat - 370001

NABL / TC-10331
GST : 24AAUCA0391F120
info@a2zenvirotechlab.com
www.a2zenvirotechlab.com

Akbar Khan Jalori
+ 91 98987 11906



ENVIROTECH LAB PVT. LTD.

TEST REPORT



TC-10331

Customer's Name and Address:

ADPTPL
OPP. SHIRVA RAILWAY CROSSING, NEAR LOC
FORESHORE TERMINALS NEW KANDLA -370210

Format No. : 7.8 F-01
Date of : 16/10/2022
Report :
Report No : A2ZELPL/AA/010/2022/02

Description of Sample:

Date of Sampling	: 12/10/2022	Type of Sampling	:
Date of Sample Received	: 13/10/2022	Sample ID	: AA/102022/02
Sampling Location of Sampling Point	: NEAR TANK FARM AREA	Sample Particular	: AMBIENT AIR
Sample Collected / Submitted by	: A2ZELPL Team Member	Reference Method for Sampling	: -----
Sample Quantity /Total No.	: 2 Filter paper 1SO ₂ × 35 ml, 1NO ₂ × 35 ml.	Date of Analysis Start	: 13/10/2022
Details of Packing/Label/Seal	: Satisfactory	Date of Analysis Completion	: 15/10/2022
Environment condition during the test	: 25 ± 3 °C	Metrological conditions during monitoring	: Clear Sky
Instrument code	: A2ZELPL/RDS/01 A2ZELPL/FDS/01	Actual Duration of Monitoring (Hours)	: 24

Test Results

Sr. No.	Parameters	Results	Unit	NAAQS LIMITS	Reference Method
1.	Particulate Matter PM ₁₀	69	µg/m ³	100 µg/m ³	IS 5182 (Part 23)2006/ Reaffirmed 2017
2.	Particulate Matter PM _{2.5}	32	µg/m ³	60 µg/m ³	IS 5182 (Part 24)2019
3.	Sulfur Dioxide SO ₂	22	µg/m ³	80 µg/m ³	IS 5182 (Part 2)2001/ Reaffirmed 2017
4.	Nitrogen Dioxide NO ₂	38	µg/m ³	80 µg/m ³	IS 5182 (Part 6)2006/ Reaffirmed 2017

A. A. Jalori
Tested By
(Sr. Analyst/Analyst)

Checked By/Prepared By
Dr. Jagruti P. Matang
(Technical Manager)

Authorized Signatory
Akbarkhan P. Jalori
(Quality Manager)

This Report is issued under the following terms & Condition:

- The results relate only to the items tested and for applicable parameter
- This Test report shall not be reproduced in full or part for any promotional or publicity and can't be used as evidence in court of law without the written consent of A 2 Z Envirotech Lab Private Limited
- Reanalysis of the sample will be done if requested Within 15 days from the date of reporting of sample if the samples are not consumed during analysis.

End of Report



ENVIROTECH LAB PVT. LTD.

TEST REPORT



TC-10331

Customer's Name and Address:

KOTPL
OFF SHIRVA RAILWAY CROSSING, NEAR LOC
FORESHORE TERMINALS NEW KANDLA -370210

Format No. : 7.8 F-01

Date of : 06/09/2022

Report

Report No : AZZELPL/AN/092022/01

Description of Sample:

Date of Sampling	: 05/09/2022	Type of Sampling	: -----
Date of Sample Received	: 05/09/2022	Sample ID	: AN/092022/01
Sampling Location of Sampling Point	: -----	Sample Particular	: ----
Sample Collected / Submitted by	: AZZELPL Team Member	Reference Method for Sampling	: IS 9989-1991
Instrument calibration status	: OK	Date of Analysis Start	: ----
Metrological conditions during monitoring	: CLEAR SKY	Date of Analysis Completion	: ----
Instrument code	: AZZELPL/SLM/01	Actual Duration of Monitoring (MINUTE)	: 30


Test Results

Sr. No.	LOCATION	Results	CPCB Limit	Unit	Reference Method
1.	NEAR TANK FARM	58	75	dB	IS 9989-1991
2.	NEAR JEETY LANDFALL AREA	52	75	dB	IS 9989-1991
3.	TANK FARM RIGHT SIDE	54	75	dB	IS 9989-1991
4.	TANK FARM LEFT SIDE	59	75	dB	IS 9989-1991

A.A. JALORI

Tested By
(Sr. Analyst/Analyst)

Checked By/Prepared By
Dr. Jagruti P. Matang
(Technical Manager)


Authorized Signatory
Akbarkhan P. Jalori
(Quality Manager)

This Report is issued under the following terms & Condition:

- The results relate only to the items tested and for applicable parameter
- This Test report shall not to be reproduced in full or part for any promotional or publicity and can't be used as evidence in court of law without the written consent of A 2 Z Envirotech Lab Private Limited
- Reanalysis of the sample will be done if requested Within 15 days from the date of reporting of sample if the samples are not consumed during analysis.

End of Report



ENVIROTECH LAB PVT. LTD.



TC-10331

TEST REPORT

Customer's Name and Address:

KOTPL
OPP. SHIRVA RAILWAY CROSSING, NEAR LOC
FORESHORE TERMINALS NEW KANDLA - 370210

Format No. : 7.8 F-01
Date of : 15/09/2022
Report
Report No : A2ZELPL/DW/09/2022/01

Description of Sample:

Date of Sampling	: 11/09/2022	Type of Sampling	: Drinking Water
Date of Sample Received	: 12/09/2022	Sample ID	: DW/092022/01
Sampling Location of Sampling Point	: Near office area	Sample Particular	: Drinking Water
Sample Collected / Submitted by	: A2ZELPL Team Member	Reference Method for Sampling	: IS 3025 (Part-1)
Sample Quantity / Total No.	: 2 Litre/1Nos.	Date of Analysis Start	: 12/09/2022
Details of Packing/Label/Seal	: Satisfactory	Date of Analysis Completion	: 14/09/2022
Environment condition during the test	: 25 ± 3 °C		

Test Results

Sr. No.	Parameters	Results	STANDARD Limit	Unit	Reference Method
1.	pH @°C	7.3	6.5-8.5	-	APHA, 23rd Edition 2017/4500-H° B
2.	Electrical Conductivity @°C	0.55	--	mS/cm	APHA, 23rd Edition 2017/2510-B
3.	Temperature	25	--	°C	APHA, 23rd Edition 2017/2550B
4.	Total dissolved solids	256	500	mg/L	APHA, 23rd Edition 2017/ 2540-C
5.	Total suspended solids	24	50-150	mg/L	APHA, 23rd Edition 2017/ 2540-D
6.	Chloride as Cl	29.2	250	mg/L	APHA, 23rd Edition 2017/ 4500 Cl- B
7.	Total Hardness as CaCO ₃	64	200	mg/L	APHA, 23rd Edition 2017/2340-C
8.	Calcium as Ca	32	75	mg/L	APHA, 23rd Edition 2017/3500-Ca-B
9.	Magnesium as Mg,	26	30	mg/L	APHA, 23rd Edition 2017/Calculation Method 3500 -Mg B
10.	E Coil	Absent	Shall Not be detectable	Nos./100 ml	APHA 23rd Edition Part - 9000 Section : 9221D

A.A. JALORI

Tested By

(Sr. Analyst/Analyst)

Checked By/Prepared By
Dr. Jagruti P. Matang
(Technical Manager)

Authorized Signatory
Akbar Khan P. Jalori
(Quality Manager)

This Report is issued under the following terms & Condition:

- The results relate only to the items tested and for applicable parameter
- This Test report shall not to be reproduced in full or part for any promotional or publicity and can't be used as evidence in court of law without the written consent of A 2 Z Envirotech Lab Private Limited
- Reanalysis of the sample will be done if requested Within 15 days from the date of reporting of sample if the samples are not consumed during analysis.

End of Report

Page No.: 1 of 2



ENVIROTECH LAB PVT. LTD.

TEST REPORT



TC-10331

Customer's Name and Address:

KOTPL
OPP. SHIRVA RAILWAY CROSSING, NEAR LOC
FORESHORE TERMINALS NEW KANDLA -370210

Format No. : 7.8 F-01
Date of : 15/09/2022
Report
Report No : A2ZELPL/AA/09/2022/01

Description of Sample:

Date of Sampling	: 11/09/2022	Type of Sampling	:
Date of Sample Received	: 12/09/2022	Sample ID	: AA/092022/01
Sampling Location of Sampling Point	: NEAR JEETY LAND FALL POINT	Sample Particular	: AMBIENT AIR
Sample Collected / Submitted by	: A2ZELPL Team Member	Reference Method for Sampling	: ----
Sample Quantity / Total No.	: 2 Filter paper 1SO ₂ × 35 ml, 1NO ₂ × 35 ml.	Date of Analysis Start	: 12/09/2022
Details of Packing/Label/Seal	: Satisfactory	Date of Analysis Completion	: 14/09/2022
Environment condition during the test	: 25 ± 3 °C	Metrological conditions during monitoring	: Clear Sky
Instrument code	: A2ZELPL/RDS/01 A2ZELPL/FDS/01	Actual Duration of Monitoring (Hours)	: 24

Test Results

Sr. No.	Parameters	Results	Unit	NAAQS LIMITS	Reference Method
1.	Particulate Matter PM ₁₀	72	µg/m ³	100 µg/m ³	IS 5182 (Part 23)2006/ Reaffirmed 2017
2.	Particulate Matter PM _{2.5}	30	µg/m ³	60 µg/m ³	IS 5182 (Part 24)2019
3.	Sulphur Dioxide SO ₂	22	µg/m ³	80 µg/m ³	IS 5182 (Part 2)2001/ Reaffirmed 2017
4.	Nitrogen Dioxide NO ₂	36	µg/m ³	80 µg/m ³	IS 5182 (Part 6)2006/ Reaffirmed 2017

A. A. J. J. J.
Tested By
(Sr. Analyst/Analyst)

Checked By/Prepared By
Dr. Jagruti P. Matang
(Technical Manager)

Authorized Signatory
Akbarkhan P. Jalori
(Quality Manager)

This Report is issued under the following terms & Condition:

- The results relate only to the items tested and for applicable parameter
- This Test report shall not to be reproduced in full or part for any promotional or publicity and can't be used as evidence in court of law without the written consent of A 2 Z Envirotech Lab Private Limited
- Reanalysis of the sample will be done if requested Within 15 days from the date of reporting of sample if the samples are not consumed during analysis.

End of Report

A2Z Envirotech Lab Pvt. Ltd.
Plot No. 340, Sector 1A, 2nd Floor,
Shiv House, Near Oslo Circle,
Gandhinagar - Kachchh, Gujarat - 370201

Branch:
147, Time Square Empire
Mirjapur Road, Bhuj - Kachchh
Gujarat - 370001

NABL / TC-10331
GST : 24AAUCAA0391F1Z0
Info@azzenvirotechlab.com
www.azzenvirotechlab.com

Akbarkhan Jalori
+91 98987 11906



ENVIROTECH LAB PVT. LTD.
TEST REPORT



TC-10331

Customer's Name and Address:

OPP. SHIRVA RAILWAY CROSSING, NEAR LOC
FORESHORE TERMINALS NEW KANDLA -370210

Format No. : 7.8 F-01
Date of : 15/09/2022
Report :
Report No : A2ZELPL/AA/09/2022/01

Description of Sample:

Date of Sampling	: 11/09/2022	Type of Sampling	:
Date of Sample Received	: 12/09/2022	Sample ID	: AA/092022/01
Sampling Location of Sampling Point	: NEAR TANK FARM AREA	Sample Particular	: AMBIENT AIR
Sample Collected / Submitted by	: A2ZELPL Team Member	Reference Method for Sampling	: -----
Sample Quantity /Total No.	: 2 Filter paper 1SO ₂ × 35 ml, 1NO ₂ × 35 ml.	Date of Analysis Start	: 12/09/2022
Details of Packing/Label/Seal	: Satisfactory	Date of Analysis Completion	: 14/09/2022
Environment condition during the test	: 25 ± 3 °C	Metrological conditions during monitoring	: Clear Sky
Instrument code	: A2ZELPL/RDS/01 A2ZELPL/FDS/01	Actual Duration of Monitoring (Hours)	: 24

Test Results

Sr. No.	Parameters	Results	Unit	NAAQS LIMITS	Reference Method
5.	Particulate Matter PM ₁₀	72	µg/m ³	100 µg/m ³	IS 5182 (Part 23)2006/ Reaffirmed 2017
6.	Particulate Matter PM _{2.5}	30	µg/m ³	60 µg/m ³	IS 5182 (Part 24)2019
7.	Sulphur Dioxide SO ₂	22	µg/m ³	80 µg/m ³	IS 5182 (Part 2)2001/ Reaffirmed 2017
8.	Nitrogen Dioxide NO ₂	36	µg/m ³	80 µg/m ³	IS 5182 (Part 6)2006/ Reaffirmed 2017

A.A. Jalori
Tested By
(Sr. Analyst/Analyst)

Checked By/Prepared By
Dr. Jagruti P. Matang
(Technical Manager)

A.A. Jalori
Authorized Signatory
Akbar Khan P. Jalori
(Quality Manager)

This Report is issued under the following terms & Condition:

- The results relate only to the items tested and for applicable parameter
- This Test report shall not be reproduced in full or part for any promotional or publicity and can't be used as evidence in court of law without the written consent of A 2 Z Envirotech Lab Private Limited
- Reanalysis of the sample will be done if requested Within 15 days from the date of reporting of sample if the samples are not consumed during analysis.

End of Report

Annexure -B

**Second Year Monsoon Report
(June 2022 to September 2022)**

**Regular Monitoring of Marine Ecology in and
around the Deendayal Port Authority and Continuous
Monitoring Programme**

Submitted to



DEENDAYAL PORT AUTHORITY
Administrative Office Building
Post Box No. 50, Gandhidham (Kachchh)
Gujarat-370201

Submitted by



GUJARAT INSTITUTE OF DESERT ECOLOGY
P.B. No. 83, Mundra Road, Opp. Changleshwar Temple
Bhuj-Kachchh, Gujarat-370001

October 2022

**Second Year Monsoon Report
(June 2022 to September 2022)**


**Regular Monitoring of Marine Ecology in and
around the Deendayal Port Authority and Continuous
Monitoring Programme**

Submitted to



DEENDAYAL PORT AUTHORITY
Administrative Office Building
Post Box No. 50, Gandhidham (Kachchh)
Gujarat-370201

Submitted by



GUJARAT INSTITUTE OF DESERT ECOLOGY
P.B. No. 83, Mundra Road, Opp. Changleshwar Temple
Bhuj-Kachchh, Gujarat-370001

October 2022

Project Team

Project Coordinator

Dr. V. Vijay Kumar, Director

Principal Investigator

Dr. Durga Prasad Behera	Project Scientist	Phytoplankton & Zooplankton, Physico-chemical parameters, Seaweed, Seagrass & Marine Fisheries
--------------------------------	-------------------	--

Co-Principal Investigator

Dr. R. Ravinesh	Project Scientist	Marine Biodiversity & Taxonomy
------------------------	-------------------	--------------------------------

Core Team

Dr. Jaikumar, M.	Senior Scientist	Mangrove & Mudflat
Dr. L. Prabha Devi	Advisor	Management Plan
Dr. Nikunj B. Gajera,	Scientist	Avifauna
Dr. Kapilkumar Ingle	Project Scientist	Mangrove Ecology
Dr. Dhara Dixit	Project Scientist	Halophytes

Team members

Mr. Dayesh Parmar	Project officer	GIS & Remote sensing
Mr. Sai Vineeth Perla	Senior Research Fellow	Sediment, Water & Benthic Fauna
Ms. Pallavi V. Joshi	Junior Research Fellow	Phytoplankton and Zooplankton
Miss. Bhagavati N. Kannad	Junior Research Fellow	Mangrove ecology

Monsoon (June 2022 to September 2022)

S. No	Components of the Study	Remarks
1	MoEF & CC sanction letter and details	<p>(i). EC & CRZ clearance granted by the MoEF &CC, GoI dated 19/12/16 Dev. Of 7 integrated facilities – specific condition no. xviii.</p> <p>(ii). EC & CRZ clearance granted by the MoEF &CC, GoI dated 18/2/2020 Dev. Remaining 3 integrated facilities – specific condition no. xxiii.</p> <p>(iii). EC & CRZ clearance granted by the MoEF &CC, GoI dated 19/2/2020 Dev. integrated facilities (Stage II-5 -specific condition no. xv.</p> <p>(iv). EC & CRZ clearance granted by the MoEF &CC, GoI dated 20/11/20 – Creation of waterfront facilities (OJ 8 to 11- Para VIII Marine Ecology, specific condition iv.</p>
2	Deendayal Port letter sanctioning the project	DPA work Order: WK/4751/Part/ (Marine Ecology Monitoring)/11 date 03.05.2021
3	Duration of the project	Three years-from 24.05.2021 to 23.05.2024
4	Period of the survey carried	Second Year Monsoon season (June 2022 to September 2022)
5	Survey area within the port limit	All major and minor creek systems from Tuna to Surajbari and Vira coastal area.
6	Number of sampling locations	Fifteen sampling locations in and around the DPA port jurisdiction
7	Components of the report	
7a	Mangroves	The overall average density was 4602 trees/ha of <i>A. marina</i> during monsoon 2022. The highest tree density was reported at the S-12 station in the Tuna creek area (7359 plants/ha). The lowest average tree density (2935 plants/ha) was reported in Phang creek. However, the lowest density in the individual site was recorded in site S-5 at Phang creek. The highest regeneration (140,000 plants/ha) at S-9 of Navlakhi creek and recruitment (31,500 plants/ha) class density were recorded at Kharo creek (S-7).
7b	Mudflats	The highest TOC value (0.83%) was recorded at station S-4 followed by S-2 site. The lowest TOC value was reported at S-12. It is observed that TOC values varied significantly among the sampling stations, which means that organic carbon depends on the living life forms and the type of life forms in the mudflats.

Snapshot

7c	Zooplankton	The zooplankton identified from the 15 stations falls under 10 phyla and 41 genera which are described 16 groups. The phylum Arthropoda was the predominant represented with 25 genera, including copepods, crabs, shrimps and their larvae. The highest percentage was due to the calanoid copepods (36.9%) followed by Decapoda (13.2%) and Gastropoda (8.2%).
7d	Phytoplankton	The generic number recorded during the monsoon period ranged from 24 to 33 at the sampling stations with remarkable variations concerning the composition. The maximum number (33 genera) was observed at S-11, and the minimum from S-15 represented 24 genera. The percentage composition of the various groups varied from 5 % to 47 %, of which the centrales and pennales are the dominant, constituting 47% and 27%, respectively.
7e	Intertidal Fauna	The intertidal fauna and the species diversity of the invertebrates showed the maximum for phylum Arthropoda (8 species), followed by Mollusca (6 species). The phylum Chordata was represented by two species. The overall percentage composition of the four groups of intertidal fauna at the 15 sites revealed the Arthropoda (50%), Mollusca (37%), and Chordata (13%).
7f	Sub-tidal Macrobenthos	The DPA port environment revealed that Mollusca (14 species) and Annelida (2 species) were the major constituents, followed by Arthropoda (1 species) and Cnidaria (1 species). The phylum Mollusca constituted the maximum (78%) share of the subtidal Fauna, followed by Annelida (11%), Arthropoda (5.5%) and Cnidaria (5.5%) in the total benthic samples collected.
7g	Seaweeds	No seaweed is reported in the DPA area.
7h	Seagrass	No seagrass is reported in the DPA area.
7i	Marine reptiles	One species of reptile was recorded from the DPA area.
7j	Marine mammals	One species of marine mammal was recorded from the DPA area.
7k	Halophytes	Four halophytes were recorded along the selected Deendayal Port Authority sites during the Monsoon sampling; among the halophyte species recorded, <i>Salicornia brachiata</i> alone was found in the 3 sampling locations. The percentage of <i>Salicornia brachiata</i> was found to be the highest at stations S-8 (78%) and the lowest at S-11.
7l	Avifauna	A total of 49 species belonging to 6 orders, 25 families and 38 genera were recorded from the coastal area of Deendayal Port Authority during the Monsoon season study.

Contents

S. No	Title	Page No
1	Introduction	1
1.1	Rationale of the present study	2
1.2	Scope of work	3
1.3	Study Area	4
2	Sampling of Water and sediment samples	5
2.1	Methodology	6
	Physico-chemical parameters	6
	pH and Temperature	6
	Water sample collection	6
	Salinity	6
	Total Suspended Solids (TSS)	6
	Total Dissolved Solids (TDS)	6
	Turbidity	7
	Dissolved Oxygen (DO)	7
	Phosphate	7
	Total phosphorus	7
	Nitrite	7
	Nitrate	7
	Petroleum Hydrocarbons (PHC)	7
2.2	Sediment characteristics	8
	Sediment Texture	8
	Total Organic Carbon	8
	Petroleum Hydrocarbons (PHC)	8
2.3	Biological Characteristics of water and sediment	8
	Primary productivity	8
	Phytoplankton	9
	Zooplankton	9
	Intertidal Fauna	10
	Subtidal macro benthic Fauna	10
2.4	Mudflats	12
	Sampling locations	12
	Total Organic Carbon	13
	Estimation of Bulk Density (BD)	14

2.5	Mangrove assessment	14
2.6	Halophytes	15
2.7	Marine Fishery	16
2.8	Avifauna	18
	Boat Surveys	18
2.9	Data Analysis	18
3	Results	19
3.1	Water quality assessment	19
	Temperature (°C) and pH	19
	Salinity (ppt)	19
	Dissolved oxygen (DO)	19
	Suspended Solids (TSS)	20
	Total Dissolved solids (TDS)	20
	Turbidity	20
	Water nutrients (Nitrate, Nitrite and Total Phosphorus)	20
	Petroleum Hydrocarbon (PHs)	21
3.2	Sediment	23
	Sediment texture	23
	Total Organic Carbon (TOC)	24
3.3	Biological characteristics of water and sediment	25
	Primary productivity	25
	Phytoplankton	26
3.4	Generic status	26
	Percentage composition of phytoplankton	26
	Percentage of occurrence	28
	Phytoplankton density and diversity	29
3.5	Zooplankton	33
	Phylum, group and generic status	33
	Percentage composition	33
	Percentage occurrence of zooplankton	35
	Density of zooplankton	36
3.6	Intertidal Fauna	40
	Faunal composition of Subtidal macrobenthos	40
	Cumulative Percentage composition of Fauna	42
	Intertidal Fauna density (No/m ²) variation between the stations	42
	Diversity indices	43
3.7	Subtidal Fauna (Macrobenthos)	46
	Faunal composition of Subtidal macrobenthos	46

	Subtidal Fauna density (No/m ²) variation between the stations	48
	Subtidal fauna distribution at the selected sites in the Deendayal Port area during monsoon	49
3.8	Seaweeds	52
3.9	Seagrass	52
3.10	Halophytes	53
3.11	Mangroves	55
	Tree Density	55
	Height	56
	Canopy Crown Cover	56
	Basal area	56
	Regeneration and recruitment class	58
3.12	Marine Reptiles	61
3.13	Marine Fishery	62
3.14	Marine Mammals	63
4	Mudflat	64
	Bulk density of the sediment samples	64
	Total Organic Carbon (TOC)	64
5	Avifauna	66
	Status, distribution and diversity of avifauna in different stations	66
	References	71

List of Figures

Fig No	Figure details	Page No
1	Map showing the proposed sampling locations 2021-2024	4
2	Petroleum Hydrocarbons in water ($\mu\text{g/L}$) during Monsoon 2022	21
3	Characteristics of sediment at the study stations during Monsoon 2022	23
4	Total Organic Carbon content (%) in the sediment samples in Monsoon 2022	24
5	Chlorophyll 'a' concentration at the study stations in Monsoon 2022	25
6	Number of Phytoplankton genera in Monsoon 2022	27
7	Percentage composition of phytoplankton groups in Monsoon 2022	27
8	Percentage occurrence of phytoplankton genera in Monsoon 2022	28
9	Phytoplankton density in Monsoon 2022	29
10	Different diversity indices a. Shannon b. Menhinick c. Margalef d. Simpson	30
11	Phylum and generic status of Zooplankton during Monsoon 2022	34
12	Percentage composition of zooplankton groups during Monsoon 2022	34
13	Percentage occurrence of Zooplankton groups during Monsoon 2022	35
14	Zooplankton Density in the different stations during monsoon 2022	36
15	Number of genera of intertidal fauna (Phylum) during in Monsoon 2022	41
16	Percentage composition of intertidal fauna during Monsoon 2022	41
17	Density of intertidal fauna during Monsoon 2022	42
18	Number of genera of macrobenthos during Monsoon 2022	47
19	Percentage composition of macrobenthos during Monsoon 2022	47
20	Subtidal fauna density during Monsoon 2022	48
21	Mangrove Plant density during Monsoon 2022	55
22	Plant height during Monsoon 2022	56
23	Mangrove canopy cover during Monsoon 2022	57
24	Mangrove basal area during Monsoon 2022	57
25	Bulk density of mudflat sediment during Monsoon 2022	65
26	Percentage of Total Organic Carbon in the mudflat in Monsoon 2022	65
27	Number of Avian species recorded from the Deendayal Port Area	67
28	Behavioral status of Avian species from the DPA in Monsoon 2022	67
28	Habitat wise distribution of Bird species from the DPA in Monsoon 2022	68
30	Station wise Foraging Guild status of species recorded during Monsoon 2022	69

List of Tables

S. No	Table details	Page No
1	Physico-chemical and biological parameters analysed	5
2	Physico-chemical characteristics of coastal waters during Monsoon 2022	22
3	Phytoplankton density, percentage composition and occurrence during Monsoon 2022	31
4	Diversity indices of Phytoplankton during Monsoon 2022	32
5	Zooplankton generic status during Monsoon 2022 in Deendayal Port Authority area	37
6	Diversity indices of Zooplankton along Deendayal Port Authority area during Monsoon 2022	39
7	Intertidal faunal distribution along Deendayal Port Authority area during Monsoon 2022	44
8	Diversity indices of Intertidal Fauna during Monsoon 2022	45
9	Macro-benthic faunal distribution during Monsoon 2022 in Deendayal Port Area	50
10	Diversity indices of the benthic fauna during Monsoon 2022	51
11	Percentage of Halophytes cover in the DPA during Monsoon 2022	53
12	Density of mangroves in the DPT vicinity during monsoon 2022	59
13	Regeneration and Recruitment class plants during Monsoon 2022	60

List of Photo plates

Plate No	Plate details	Page No
1	Estimation of intertidal fauna by the quadrature method	11
2	Collection of Plankton and macrobenthos in subtidal habitat	11
3	Sediment sample collection at mangrove and mudflat areas	13
4	Assessment of mangrove density, height, canopy cover & girth	15
5	Assessment and percentage cover of halophytes	16
6	Collection of fisheries information from DPA environment	17
7	Halophyte species on the intertidal zone of Deendayal Port Authority area	54
8	Mangrove species recorded along the Deendayal Port Authority area	58
9	Marine reptiles recorded along the Deendayal Port Authority area	61
10	Fish and Crab catch along the Deendayal Port Authority in monsoon 2022	62
11	Indian Ocean humpback dolphin <i>Sousa plumbea</i>	63
12	Some common birds from the Deendayal Port Authority	70

1. Introduction

Deendayal Port is located at the inner end of Gulf of Kachchh on the Kandla creek (22°59'4.93N and longitude 70°13'22.59 E) in the Kachchh district of Gujarat state, operated by Deendayal Port Authority (DPA). Being the India's busiest major port in recent years, is gearing to add substantial cargo handling capacity with private participation. Since its formation in the 1950s, the Deendayal Port provides the maritime trade requirements of states such as Rajasthan, Madhya Pradesh, Uttar Pradesh, Haryana and Gujarat. Because of its proximity to the Gulf countries, large quantities of crude petroleum are imported through this port. About 35% of the country's total export takes place through the ports of Gujarat in which the Deendayal port has a considerable contribution. Assortments of liquid and dry cargo are being handled at DPA Port. The dry cargo includes fertilizers, iron and steel, food grains, metal products, ores, cement, coal, machinery, sugar, wooden logs, etc. The liquid cargo includes edible oil, crude oil and other petroleum products. Cargo handling has increased from 117.5 MMT to 127 MMT during 2021-2022. Presently, the Port has total 1-16 dry cargo berths for handling dry cargo, 6 oil jetties, and one barge jetty at Bunder basin, dry bulk terminal at Tuna Tekra, barge jetty at Tuna and two SPMs at Vadinar for handling oil. Regular expansion or developmental activities such as the addition of jetties, allied SIPC and ship bunkering facilities are underway in order to cope with the increasing demand for cargo handling during the recent times.

A developmental initiative of this magnitude is going on since past 7 decades, which will have its own environmental repercussions. Being located at the inner end of Gulf of Kachchh, Deendayal Port Authority encompasses a number of fragile marine ecosystems that includes a vast expanse of mangroves, mudflats, creek systems and associated biota. Deendayal Port is a natural harbour located on the eastern bank of North-South trending Kandla creek at an aerial distance of 90 km from the mouth of Gulf of Kachchh. The Port's location is marked by a network of major and minor mangrove lined creek systems with a vast extent of mudflats. Coastal belt in and around the port has an irregular and dissected configuration. Due to its location at the inner end of the Gulf, the tidal amplitude is elevated, experiencing 6.66 m during mean high-water spring (MHWS) and 0.78 m during mean low water spring (MLWS) with MSL of 3.88 m. Commensurate with the increasing tidal amplitude, vast intertidal expanse is present in and around the port environment. Thus, the occurrence of mudflats on the intertidal zone enables mangrove formation to an extensive

area. Contrary to the southern coast of Gulf of Kachchh, the coral formations, seaweed and seagrass beds are absent in the northern coast due to high turbulence induced suspended sediment load in the water column, a factor again induced due to the conical Gulf geomorphology and surging tides towards its inner end.

1.1. Rationale of the present study

The ongoing developmental activities at Deendayal Port Authority has been intended for the following.

- i. The development of 3 remaining integrated facilities (Stage 1) within the existing Port at Kandla which includes development of a container terminal at Tuna off Tekra on BOT base T shaped jetty, construction of port craft jetty and shifting of SNA section of Deendayal port and railway line from NH-8A to Tuna port.
- ii. EC & CRZ clearance granted by the MoEF &CC, GoI dated 18/2/2020 Dev. Remaining 3 integrated facilities – specific condition no. xxiii.
- iii. EC & CRZ clearance granted by the MoEF &CC, GoI dated 19/2/2020 Dev. integrated facilities (Stage II-5 -specific condition no. xv.
- iv. EC & CRZ clearance granted by the MoEF &CC, GoI dated 20/11/20 – Creation of water front facilities (OJ 8 to 11- Para VIII Marine Ecology,specific condition iv).

As per the environmental clearance requirements to these developmental initiatives, by MoEF & CC, among other conditions, has specified to conduct the continuous monitoring of the coastal environment on various aspects covering the three the seasons. The regular monitoring shall include physico-chemical parameters coupled with biological indices such as mangroves, seagrasses, macrophytes and plankton on a periodic basis during the construction and operation phase of the project. Besides, the monitoring study also includes assessment of Mudflats, Fisheries, and Intertidal fauna including the macrobenthos as components of the management plan. The regular marine ecology monitoring includes Micro, Macro and Mega floral and fauna components of marine biodiversity of the major intertidal ecosystems, the water and sediment characteristics. In accord with MoEF&CC directive, DPA has consigned the project on ‘Regular Monitoring of Marine Ecology in and around the Deendayal Port Authority and Continuous Monitoring Programme” to Gujarat Institute of Desert Ecology (GUIDE), Bhuj during May, 2021. Further, Deendayal Port authorities has entrusted Gujarat Institute of Desert Ecology (GUIDE) to continue the study for another three years, i.e., 2021 – 2024. The study covers all the seasons as specified

by the specific condition of the Ministry of Environment, Forest and Climate Change (MoEF&CC). The present study is designed considering the scope of the work given in the EC conditions.

1.2. Scope of work

The scope of the present investigation includes physico-chemical and marine biological components as mentioned in the specific conditions of MoEF&CC, EC & CRZ clearance dated 19.12.2016, 18.2.2020, 19.2.2022 and 20.11.2020 with specific conditions xviii, xxiii, xv & iv respectively. A detailed holistic approach to different components of the study such as marine physico-chemical parameters of water and sediment and marine biodiversity within the Deendayal Port area will be carried out. Based on the results obtained during the project period, a detailed management plan will be drawn at the end of the project period. The biological and physico-chemical variables will be investigated during the present study on a seasonal basis i.e., monsoon, post monsoon and pre-monsoon as follows.

- ✓ Physico-chemical characteristics of water and sediment
- ✓ Detailed assessment of mangrove vegetation structure including density, diversity, height, canopy, and other vegetation characteristics.
- ✓ GIS and RS studies to assess different ecological sensitive land use and land cover categories within the Port area such as the extent of dense and sparse mangroves, mudflats, creek systems, and other land cover categories within the port limits.
- ✓ Quantitative and qualitative assessment of the intertidal fauna, composition, distribution, diversity, density, and other characteristics.
- ✓ Data collection on the species composition, distribution, diversity and density of sub-tidal benthic fauna.
- ✓ Estimation of primary productivity at the selected sampling sites located in around the DPA area.
- ✓ Investigation of the species composition, distribution, density, and diversity of phytoplankton and zooplankton.
- ✓ Recording the occurrence, diversity and distribution of halophytes, seagrasses, seaweeds and other coastal flora. Investigations on the Avifaunal density, diversity, composition, habitat, threatened and endangered species and characters. Fishery

Resources – Species composition, diversity, Catch Per Unit Effort (CPUE) and other socio-economic information.

1.2.1. Study Area

The coastal belt in and around Deendayal Port Authority jurisdiction is characterized by a network of creek systems and mudflats which are covered by sparse halophytic vegetation like scrubby to dense mangroves, creeks and salt-encrusted landmass which form the major land components. The surrounding environment in 10 km radius from the port includes built-up areas, salt pans, human habitations and port related structures on the west and north creek system, mangrove formations and mudflats in the east and south. The nearest major habitation is Gandhidham town located about 12 km away on the western part with population of 2,48,705 (as per 2011 census).




Figure 1: Map showing the sampling locations 2021-2024

2. Sampling of water and sediment samples

Sampling was carried out for the coastal water (surface) and sediment to determine physical and chemical characteristics from the prefixed sampling sites. The biological parameters (benthic and pelagic fauna, flora and productivity) were also estimated (Table.1).

Table 1: Physico-chemical and biological parameters analysed

Parameters	
Water	Mangrove & Other Flora
• pH	Mangrove
• Temperature	• Vegetation structure, density
• Salinity (ppt)	• Diversity
• Petroleum Hydrocarbons-PHC	• Height
• Dissolved oxygen	• Canopy and other vegetation characteristics
• Total Suspended Solids (TSS)	
• Total Dissolved solids (TDS)	Halophytes:
Nutrients	• Percentage of distribution
Nitrate (NO ₃)	• Diversity
• Nitrite (NO ₂)	
• Total Nitrogen	Seagrass and Seaweed
• Sediment	• Occurrence, distribution, and diversity.
• Texture	Intertidal fauna
• Total organic carbon (TOC)	• Composition, distribution, diversity, density and other characteristics.
• Biological Parameters	Avifauna
Phytoplankton- Genera, abundance, diversity and biomass	• Density, diversity, composition, habitat,
• Productivity-Chlorophyll a	• Threatened and endangered species and characters
• Zooplankton – Species, abundance, diversity	
• Macrobenthos - genera, abundance, diversity	
• Fishery Resources	
• Common fishes available	
• composition, diversity	
• Catch Per Unit Effort (CPUE)	

The water samples were collected from each pre-designated site in pre-cleaned polyethylene bottles. Prior to sampling, the bottles were rinsed with sample water to be collected and stored in an ice box for transportation to laboratory and refrigerated at 4°C till further analysis. The analysis of the water quality parameters was carried out by following standard methods (APHA, 2017). All extracting reagents were prepared using metal-free, AnalaR grade chemicals (Qualigens Fine Chemicals Division of Glaxo SmithKline Pharmaceuticals Limited, Mumbai) and double distilled water prepared from quartz double distillation assembly.

2.1. Methodology

Physico-chemical Parameters

pH and Temperature

A Thermo fisher pH / EC / Temperature meter was used for pH and temperature measurements. The instrument was calibrated with standard buffers just before use.

Salinity

A suitable volume of the sample was titrated against Silver nitrate (20 g/l) with Potassium chromate as an indicator. The chlorinity was estimated, and from that, salinity values were derived using a formula (Strickland and Parsons, 1972).

Total Suspended Solids (TSS)

About 100 ml of the water sample was filtered through pre-weighed filter paper and placed in the Hot air oven at a specified temperature as per the protocol for 1 hour. The filter paper was allowed to cool in a desiccator to obtain a constant weight by repeating the drying and desiccation steps.

Total Dissolved Solids (TDS)

The water samples were subjected for gravimetric procedure for confirmation of the readings obtained from the hand-held meter. About 100 ml of the water sample was taken in a beaker and filtered which was then dried totally in a Hot Air Oven (105°C). The TDS values were calculated using the difference in the initial and final weight of the container.

Turbidity

The sample tube (Nephelometric cuvette) was filled with distilled water and placed in the sample holder. The lid of the sample compartment was closed. By adjusting the 'SET ZERO' knob, the meter reading was adjusted to read zero. The sample tube with distilled water was removed, the 40 NTU standard solutions were filled in the tube, and the meter reading was set to read 100. Other standards were also run. The turbidity of the marine water sample was then found by filling the sample tube with the sample, and the reading was noted.

Dissolved Oxygen (DO)

DO was determined by Winkler's method (Strickland and Parsons, 1972).

Phosphate

Acidified Molybdate reagent was added to the sample to yield a phosphomolybdate complex that is reduced with Ascorbic acid to a highly coloured blue compound, which is measured at the wavelength of 690 nm in a Spectrophotometer (Shimadzu UV 5040).

Total phosphorus

Phosphorus compounds in the sample were oxidized to phosphate with alkaline Potassium persulphate at high temperature and pressure. The resulting phosphate was analyzed and described as total phosphorous.

Nitrite

Nitrite in the water sample was allowed to react with Sulphanilamide in acid solution. The resulting diazo compound was reacted with N-1-Naphthyl ethylenediamine dihydrochloride to form a highly coloured azo-dye. The light absorbance was measured at the wavelength of 543 nm in Spectrophotometer (Shimadzu UV 5040).

Nitrate

The Nitrate content was determined as nitrite (as mentioned above) after its reduction by passing the sample through a column packed with amalgamated Cadmium.

Petroleum Hydrocarbon (PHs)

The water sample (1liter) was extracted with hexane and the organic layer was separated, dried over anhydrous sulphate and reduced to 10 ml at 30°C under low pressure. Fluorescence of the extract was measured at 360 nm (excitation at 310 nm) with Saudi Arabian crude residue as a standard. The residue was obtained by evaporating lighter fractions of the crude oil at 120°C.

Sediment characteristics

Sediment samples were collected from the prefixed stations by using a Van Veen grab having a mouth area of 0.04m² or by a non-metallic plastic spatula. Sediment analysis was carried out using standard methodologies. In each location (grid), sediment samples were collected from three different spots and pooled together to make a composite sample, representative of a particular site. The collected samples were air dried and used for further analysis.

Sediment Texture

For texture analysis, specified unit of sediment sample was sieved through sieves of different mesh size as per Unified Soil Classification System (USCS). Cumulative weight retained in each sieve was calculated starting from the largest sieve size and adding subsequent sediment weights from the smaller size sieves (USDA,1951). The percentage of the various fractions was calculated from the weight retained and the total weight of the sample. The cumulative percentage was calculated by sequentially subtracting percent retained from the 100%.

Total Organic carbon

Percentage of organic carbon in the dry sediment was determined by oxidizing the organic matter in the sample by Chromic acid and estimating the excess Chromic acid by titrating against Ferrous ammonium sulphate with Ferroin as an indicator (Walkley and Black, 1934).

2.3. Biological Characteristics of water and Sediment

Primary productivity

Phytoplankton possess the plant pigment chlorophyll 'a' which is responsible for synthesizing the energy for metabolic activities of phytoplankton through the process of photosynthesis in which CO₂ is used and O₂ is released. It is an essential component to understand the consequences of pollutants on the photosynthetic efficiency of phytoplankton in the system. To estimate this, a known volume of water (500 ml) was filtered through a 0.45 µm Millipore Glass filter paper and

the pigments retained on the filter paper were extracted in 90% Acetone. For the estimation of chlorophyll 'a' and pheophytin pigments the fluorescence of the Acetone extract was measured using Fluorometer before and after treatment with dilute acid (0.1N HCL) (Strickland and Parsons,1972).

Phytoplankton

Phytoplankton samples were collected from prefixed 15 sampling sites from the coastal water in and around DPA location using standard plankton net with a mesh size of 25µm and a mouth area of 0.1256 m² (20 cm radius). The net fitted with a flow meter (Hydrobios) was towed from a motorized boat moving at a speed of 2 nautical miles/hr. Plankton adhering to the net was concentrated in the net bucket by splashing seawater transferred to a pre-cleaned and rinsed container and preserved with 5% neutralized formaldehyde and appropriately labelled indicating the details of the collection, and stored for further analysis. The Quantitative analysis of phytoplankton (cell count) was carried out using a Sedgewick-Rafter counting chamber. The density (No/l) was calculated using the formula: $N = n \times v/V$ (Where, N is the total No/liter, n is the average number of cells in 1 ml, v is the volume of concentrate; V is the total volume of water filtered. The identification was done by following the standard literature of Desikachary, (1987), Santhanam et.al. (2019) and Kamboj et.al. (2018).

Zooplankton

Zooplankton samples were collected using a standard zooplankton net made of bolting silk having 50µm with mouth area of 0.25 m² fitted with a flow meter. The net was towed from a boat for 5 minutes with a constant boat speed of 2 nautical miles/hr. The initial and final reading in the flow meter was noted down and the plankton concentrate collected in the bucket was transferred to appropriately labeled container and preserved with 5% neutralized formaldehyde. One ml of the zooplankton concentrate was added to a Sedgwick counting chamber and observed under a compound microscope and identified by following standard literature. The group/taxa were identified using standard identification keys and their number was recorded. Random cells in the counting chamber were taken for consideration and the number of zooplankton was noted down along with their binomial name. This process was repeated for five times with 1 ml sample and the average value was considered for the final calculation. For greater accuracy, the final density values were counter-checked and compared with the data collected by the settlement method.

Univariate measures such as Shannon-Wiener diversity index (H'), Margalef's species richness (d), and Pielou's evenness (J'), Simpson's dominance (D) was determined using PAST software.

Intertidal Fauna

Intertidal faunal assemblages were studied for their density, abundance and frequency of occurrence during monsoon 2022 at the pre-fixed 15 sampling locations within the DPA jurisdiction. Sample collection and assessment of intertidal communities were done in the intertidal zone during the low tide period. At each site, 1 x1 m² quadrates were placed randomly and all visible macrofaunal organisms encountered inside the quadrate were identified, counted and recorded. At each site, along the transects which run perpendicular to the waterfront, three to six replicate quadrate samples were assessed for the variability in macro-faunal population structure and the density was averaged for the entire intertidal belt. Organisms, which could not be identified in the field, were preserved in 5% formaldehyde, brought to the laboratory and identified using standard identification keys (Abott, 1954; Vine, 1986; Oliver, 1992; Rao, 2003; 2017; Psomadakis *et al.*, 2015; Apte, 2012; 2014; Naderloo 2017; Ravinesh *et al.* 2021; Edward *et al.*, 2022). Average data at each site were used to calculate the mean density (No/m²).

Subtidal macro benthic Fauna

The sampling methods and procedures were designed in such a way to obtain specimens in the best possible condition as to maximize the usefulness of the data obtained. For studying the benthic organisms, triplicate samples were collected at each station using Van Veen grab, which covered an area of 0.04m². The wet sediment was passed through a sieve of mesh size 0.5 mm for segregating the organisms. The organisms retained in the sieve were fixed in 5-7% formalin and stained further with Rose Bengal dye for ease of spotting at the time of sorting. The number of organisms in each grab sample was expressed as No. /m². All the species were sorted, enumerated and identified by following the available literature. The works of Day (1967), Hartman (1968, 1969), Rouse and Pleijel (2001), Robin *et al.*, (2003), Amr (2021), were referred for polychaetes; Crane (1975), Holthuis (1993), Naderloo (2017). Xavier *et al.*, (2020) for crustaceans; Subba Rao (1989, 2003, 2017), Apte (2012, 2014), Ramakrishna and Dey (2007), Ravinesh *et al.* (2021) and Edward *et al.*, (2022) for molluscs. Statistical analyses such as diversity indices and quadrat richness were calculated using Paleontological Statistics Software Package for Education and Data (PAST) version 3.2.1 (Hammer *et al.*, 2001).



Plate 1: Estimation of intertidal fauna by the quadrat method



Plate 2: Collection of Plankton and macrobenthos in subtidal habitat

2.4. Mudflats

Mudflats are ecologically and socio-economically vital ecosystems that bring benefits to human populations around the globe. These soft-sediment intertidal habitats, with >10% silt and clay (Dyer 1979), sustain global fisheries through the establishment of food and habitat (including important nursery habitats), support resident and migratory populations of birds, provide coastal defenses, and have aesthetic value. Mudflats are intimately linked by physical processes and dependent on coastal habitats, and they commonly appear in the natural sequence of habitats between subtidal channels and vegetated salt marshes. In some coastal areas, which may be several kilometres wide and commonly form the largest part of the intertidal area. Mudflats are characterized by high biological productivity and abundance of organisms but low in species diversity with few rare species. The mudflat biota reflects the prevailing physical conditions of the region. Intertidal mudflats can be separated into three distinct zones such as the lower tidal, middle and upper mudflats. The lower mudflats lie between mean low water neap and mean low water spring tide levels, and are often subjected to strong tidal currents. The middle mudflats are located between mean low water neaps and mean high water springs. The upper mudflats lie between the mean high-water neap and mean high water springs. The upper mudflats are the least inundated part and are only submerged at high water by spring tides (Klein, 1985). Salt marsh vegetation may colonize as far seaward as mean high water neaps. Mudflats will often continue below the level of low water spring tides and form sub-tidal mudflats (McCann, 1980). The upper parts of mudflats are generally characterized by coarse clays, the middle parts by silts, and the lower region by sandy mud (Dyer *et al.*, 2000). The intertidal mudflats are prominent sub-environments that occurred on the margin of the estuaries and low relief sheltered coastal environments. The fine-grained sediments of intertidal mudflats (70%-90%) are derived from terrestrial and marine regions (Lesuere *et.al.*, 2003). Estuarine mudflats are potential sites for deposition of organic matter derived from terrigenous, marine, atmospheric and anthropogenic sources and are mainly associated with fine grained particles (Wang *et.al.*, 2006).

Sampling locations

The Sediment samples were collected from 15 sampling locations by using sediment corer. From each site triplicate samples were collected from up to 100 cm depth with four intervals (0-25cm, 25-50cm, 50-75cm & 75-100cm) and made into composite sample for analysis. The samples were packed in zip lock bags, stored in icebox and shifted to the laboratory for subsequent analysis.



Plate 3: Sediment sample collection at mangrove and mudflat areas

Total Organic Carbon

The organic carbon content of the mudflats was estimated to assess the biological productivity of the sediment. Soil Organic Carbon (SOC) was estimated following the method of Walkley and Black (1934). In this method, organic matter (humus) in the soil gets oxidized by Chromic acid (Potassium dichromate plus concentrated H₂SO₄) by utilizing the heat evolved with the addition of H₂SO₄. The unreacted dichromate is determined by back titration with Ferrous ammonium sulphate (redox titration). Organic carbon was determined by following the below given formula:

$$\text{Oxidizable organic carbon (\%)} = \frac{10 (B - T)}{B} \times 0.003 \times \frac{100}{\text{wt. of soil}}$$

Where B = volume (mL) of Ferrous ammonium sulfate required for blank titration. T = volume of Ferrous ammonium sulfate needed for soil sample. Wt. = weight of soil (g).

Estimation of Bulk Density (BD)

The soil under field condition exists as a three-phase system viz. solid (soil particles), liquid (water) and gas (mostly air). The soil organic matter contained in a unit volume of the soil sample is called its bulk density. The amount of bulk density depends on the texture, structure and organic matter status of the soil. High organic matter content lowers the bulk density, whereas compaction increases the bulk density. To determine the bulk density of the sediment samples collected during the present study, the oven-dry weight of a known sediment volume was considered, and mass per unit volume was calculated (Maiti, 2012).

2.5. Mangrove assessment

Mangroves are widely distributed on the Deendayal Port Authority jurisdiction along the Kandla coast. The 15 mangrove sites selected at the different creeks belong to Deendayal Port Authority jurisdiction and all these stations are supposed to be sufficient to represent the mangroves status in Kandla. The mangrove stations in this study were named Tuna, Jangi, Kandla, Phan and Navlakhi based on the nearest location to the respective creek system. The Point Centered Quadrature Method (PCQM) was used for the collection of data of mangrove vegetation structure. The data included measurements of density of plants, height variations, canopy and basal area of mangrove trees as per the method of Cintron and Novelli (1984). For this method, a transect of a maximum of 200 m was applied mostly perpendicular or occasionally parallel to the creek. The sampling points considered at an interval of every 10 m and the vegetation structure of the that area were recorded. As the orientation of the transect line was already fixed, it was easy for movements within the station area for data recording. The distance between trees from the center of the sampling point for nearest 4 trees of four different directions, height of trees from the ground level, canopy length and canopy width were measured to determine the canopy cover in this study. The equipments utilized in the field were handy, and easy to use such as ranging rods, pipes and for measurement of girth at root collar above the ground (GRC), a measuring tape was used. The plants with a height <50 cm was considered as regeneration class and >50 cm but <100 cm was considered as recruitment class. Along the transects, sub-plots of 1×1 m² for regeneration and 2×2 m² were laid randomly for recruitment class of the mangrove sites.



Plate 4: Assessment of mangrove density, height, canopy cover and girth

2.6. Halophytes

To quantify and document the halophytes at Deendayal Port Authority region, quadrature method was followed. At each sampling location quadrates of various sizes have been laid during every seasonal sampling. For recording the plant density at each transect, a quadrature 1 x 1m² has been laid within the site each tree quadrates were used randomly (Misra,1968; Bonham, 1989). Four quadrates each for shrubs and herbs were laid in side each tree quadrature to assess the halophytes and the percentage cover in the study area. To enrich the species inventory, areas falling outside the quadrates were also explored and the observed species were recorded and photographed and identified using standard keys. Specimens of the various species were collected to know more information on habitat and for the preparation of herbarium.



Plate 5: Assessment and percentage cover of halophytes

2.7. Marine Fishery

Fishery resources and the diversity were assessed from the selected sampling sites. Finfish and shellfish samples were collected using a gill net with a 10 mm mesh size. The net was operated onto the water from a canoe or by a person standing in waist deep water during the high tide using a cast net. For effective sampling, points were fixed at distances within the 15 offshore sites for deploying fishing nets to calculate the Catch per Unit effort estimated per hour. The collected specimens were segregated into groups, weighed and preserved in 10% neutralized formalin solution. Finfishes were identified following Fischer and Bianchi (1984), Masuda *et al.* (1984), de Bruin *et al.* (1995) and Mohsin and Ambiak (1996). Relevant secondary information pertaining to fishery resources of Deendayal Port creek systems were gathered through technical reports, the District Fisheries department, Government gazette and other research publications.



Plate 6: Collection of fisheries information from DPA environment

2.8. Avifauna

The Avifauna population was determined along DPA mangrove strands for which the area was demarcated into fifteen major stations. In each station, creeks of varying lengths from 2 to 5 km are available. These creeks were surveyed by using boat and adopting “line transect” method. A total of fifteen boat transect (one in each site) survey was conducted in the Monsoon season (June-September, 2022). Survey was done in both terrestrial habitats like Mangrove plantations adjoining the mudflats, waste land, and aquatic habitats, like creek area, rivers and wetland.

Boat Surveys

Mangrove bird diversity was calculated by using Boat Survey method. Birds were observed from an observation post on board the boat which has given the greatest angle of clear view. Birds within a 100 meter transect on one side of the boat were counted in 10-minute blocks of time (Briggs *et al.* 1985; van Franeker, 1994). Detection of birds was done with a binocular (10 x 40) and counts were made: (1) continuously of all stationary birds (swimming, sitting on mangrove, or actively feeding) within the transect limits and (2) in a snap-shot fashion for all flying birds within the transect limits. The speed of the boat determines the forward limit of the snapshot area within a range of 100 meters. Longer or shorter forward distances were avoided by adapting the frequency of the snapshot counts. Birds that following and circling the boat were omitted from both snapshot and continuous counts. If birds arrive and then follow the boat, they were included in the count only if their first sighting falls within a normal snapshot or continuous count of the transect area. For each bird observation species, number of individuals and activity at the time of sighting, were recorded. Species richness and diversity index were calculated for different mangrove patches (i.e. fifteen station) of the study station in the Deendayal port Authority.

2.9. Data analysis

Data collected in- situ and through laboratory analysis of samples were subjected to descriptive statistical analysis (PAST and Primer 7.0) for the mean, range and distribution of different variables from the selected 15 study stations.

3. Results

Water quality assessment

The data on the mean water quality parameters measured at the time of sampling of the biological components from the 15 study sites are presented in Table1.

Temperature (°C) and pH

The water temperature at the sampling sites ranged from 23°C to 31°C. The maximum temperature of seawater was reported at S-5 and the minimum at S-6 in Kandla creek. The pH of seawater ranged from 7.1 to 8.3. The highest pH was reported at sites S-15 and S-10, however, the lowest pH 7.1 was noticed at S-14 in Kandla creek. The overall observation along the port environment revealed that the temperature fluctuation might be due to high degree of warmth in summer on the land but the pH range did not show major fluctuations among the sampling locations.

Salinity (ppt)

Salinity of the water strongly influences the abundance and distribution of marine biota in coastal and marine environments. The salinity ranged from 28 ppt to 40 ppt with the average value of 37 ppt. Minimum salinity was observed S-7 and maximum at S-9, S-13 & S-15. The poor rainfall induced aridity in the Gulf of Kachchh (GoK) region renders Gulf waters hypersaline round the year. In addition, GoK is known to be a negative water body where evaporation exceeds precipitation.

Dissolved oxygen (DO)

Dissolved oxygen is the amount of oxygen dissolved in water and is a fundamental requirement of all biota and chemical processes in the aquatic environment. The concentration varies mainly due to photosynthesis and respiration by plants and animals in water. Generally, the coastal waters are having high level of dissolved oxygen due to the dissolution from the atmosphere through diffusion process on the surface layer (CCME,1999). The dissolved oxygen in the coastal waters of Deendayal port authority area ranged from 4.5 mg/L to 6.9 mg. The highest DO concentration was observed at station S-7 and lowest was observed at station-15.

Suspended Solids (TSS)

The total suspended solids (TSS) concentration at the 15 sampling sites ranged from 127 mg/L to 403 mg/L with the average of 255 mg/L. The highest TSS values was reported at S-15 in the Phang creek followed by 354 mg/L in S-3 oil jetty. The minimum TSS value was recorded at S-7 which was 127 mg/L.

Total Dissolved solids (TDS)

The total dissolved solids (TDS) in the water consist of inorganic salts and dissolved materials which mostly comprises of anions and cations. The TDS of the samples varied from 1967 mg/L to 11,288 mg/L with an average of 5,703 mg/L. The maximum value was reported at S-6.

Turbidity

The turbidity of the water samples from the study sites ranged between 44 NTU and 147 NTU with the average of 76 NTU. The lowest value was reported at S-3 and a highest value at S-6 followed by S-6 (170 NTU).

Water nutrients (Nitrate, Nitrite and Total Phosphorus)

The nutrients influence growth, metabolic activities and reproduction of biotic components in the aquatic environment. The distribution of nutrients mainly depends upon tidal conditions, season and fresh water influx from land. The nitrate concentration ranged from 0.01 mg/L to 0.02 mg/L with an average of 0.01 mg/L. The highest nitrate concentration was observed at station S-7 and the lowest at station S-11. There was no remarkable variation in concentration of nitrate among the study station. Similarly, nitrite values varied between 0.05 mg/L to 0.94 mg/L. The highest concentration was observed at station S-13 and lowest concentration was observed at station S-2. The highest concentration might be due to influx effluents from industries producing metals, dyes and celluloid in the periphery of port authority. The Total phosphorus values among the study station ranged from 0.02 mg/L to 0.96 mg/L with an average of 0.47 mg/L. The highest phosphorus concentration was observed at station S-13 near veera of Kandla creek and lowest concentration was observed at station S-11 in Jhangi creek. Highest concentration might be due to leaching of phosphatic fertilizer while handling of cargo port area.

Petroleum Hydrocarbons (PHs)

Due to urbanization and modernization, petrochemical products are in heavy demand. Petroleum hydrocarbons (PHs) represent short-chain hydrocarbons like aromatic, paraffin, alicyclic complexes, and non-hydrocarbon mixtures such as thiol, and asphaltene, naphthenic acid, phenol, thiol, heterocyclic nitrogen, sulfuric amalgams and metalloporphyrin. Due to the hydrophobic nature of the PHs, they possess low solubility in water and a high persistence level in soil, water as well as sediments (Babu *et al.*, 2019). PHs are significant toxic compounds representing one of the major wide-scale environmental threats caused due to the coastal oil refining, production, leaks or accidental spilling, transport, shipping activities, offshore oil production and other anthropogenic activities. The release of such compounds into the environment irrespective of it being accidental or due to any anthropogenic activities leads to soil as well as water pollution. This in turn poses catastrophic health effects either directly or indirectly on all the forms of life thereby deteriorating the overall ecosystem. In the current study, the presence of PHs in water samples collected along all the 15 sampling sites were detected and estimated. The PHs ranged from 2.2 µg/L to 9.9 µg/L. The PHs detected from the individual sites have been represented in (Fig 2). The highest concentration of the PHs was detected at S-1 site (Tuna creek) while the lowest was noted for S-13 (Veera). A high level of the PHs content was noted down at site S-1 too followed by the rest of the sites.




Figure 2: Petroleum hydrocarbons in water (µg/L) during Monsoon 2022

Table 2: Physico-chemical characteristics of coastal waters during Monsoon 2022

Parameters	S-1	S-2	S-3	S-4	S-5	S-6	S-7	S-8	S-9	S-10	S-11	S-12	S-13	S-14	S-15
Temp (°C) (Air)	27	28	31	33	36	26	31	29	30	32	26	34	29	29	34
Temp (°C) (Water)	25	25	29	30	31	23	29	26	27	30	23	30	27	26	29
pH	8	8.09	7.9	7.5	7.8	7.8	7.7	7.6	8.2	7.9	7.9	8.06	8.2	7.1	8.3
Salinity (ppt)	34.7	36.7	39.2	38.7	36.5	36.2	28.3	35.8	39.9	38.8	36.4	39	40.2	38.2	40.1
Dissolved oxygen (mg/L)	4.86	4.66	6.69	5.27	5.87	4.66	6.89	6.28	5.06	5.87	4.66	6.48	5.27	5.47	4.45
Total Suspended Solids (TSS) (mg/L)	200	236	354	132	347	234	127	172	342	232	334	190	272	252	403
Total Dissolved solids (TDS) (mg/L)	3970	4676	2985	3851	7885	1967	5988	4320	7549	11288	8983	3886	5676	4792	7733
Turbidity (NTU)	48	58	147	95	93	44	45	93	119	108	57	58	58	52	63
Nitrate (NO ₃) (mg/L)	0.01	0.01	0.01	0.01	0.02	0.02	0.01	0.01	0.02	0.01	0.01	0.01	0.07	0.02	0.01
Nitrite (NO ₂) (mg/L)	0.39	0.05	0.36	0.39	0.41	0.74	0.38	0.53	0.58	0.27	0.73	0.39	0.94	0.63	0.55
Total Phosphorus (mg/L)	0.35	0.64	0.46	0.41	0.39	0.90	0.76	0.30	0.04	0.06	0.02	0.35	0.96	0.85	0.63
PHs (µg/L)	9.85	4.8	8.8	3.75	3.35	4	3.4	3.55	2.8	2.9	4.75	2.4	2.15	2.45	3.2
Chlorophyll a (mg/L)	0.19	0.20	0.21	0.18	0.13	0.15	0.19	0.15	0.16	0.14	0.19	0.21	0.15	0.16	0.22

3.2. Sediment

Sediment texture

The percentage composition of the soil particles in the sediment analyzed from the 15 sampling sites are presented in Fig.3. There were noticeable variations in the soil fractions, sand, silt and clay, among the stations. In the present study the highest percentage of clay was reported at S-7 followed by S-9. The highest percentage of sand was observed at S-1 followed by S-14 station. As per the observations, the percentage of silt content was less compared to clay and sand in many sampling sites except S-1 and S-14. The nature of soil texture was characterized by the proportion of clay, sand and silt fractions. The Soil texture revealed the dominance of silty-clay type in all the stations with less variations among them. This consistently high clay-loam value may be attributed to the winnowing activity of sediment transport system. The absence of perennial flow of freshwater into the coast along with lack of wave induced sand transport from open sea are the possible reasons for this uniform pattern of soil texture.




Figure 3: Characteristics of sediment at the study stations in Monsoon 2022

Total Organic Carbon (TOC)

In the present study, the total organic carbon content varied from 0.63% to 0.84% (Fig.4). The highest values of TOC were reported at S-11 followed by S-15. The lowest TOC value was recorded at the S-7. The distribution of total organic carbon closely followed the distribution of sediment type i.e., sediment low in clay content contained relatively low organic carbon.




Figure 4: Total Organic Carbon content (%) in the sediment during Monsoon 2022

3.3. Biological characteristics of water and sediment

Primary productivity

Chlorophyll 'a' the photosynthetic pigment which can be used as a proxy for phytoplankton productivity and thus is an essential water quality parameter. Generally, the primary production of the water column is assessed from Chlorophyll 'a' concentration. It is well known that half of the global primary production being mediated by the activity of microscopic phytoplankton.

In the present study, Chlorophyll 'a' concentration ranged from 0.13 mg/L to 0.22 mg/L. The highest concentration 0.22 mg/L was reported at S-15 (Fig.5) followed by S-12 (0.21) and S-3 (0.20mg/L). The photosynthetic pigment chlorophyll a which is a measure of the population density of phytoplankton during the monsoon period showed narrow range of variations among the sites. The Chlorophyll 'a' content was very low at S-5.




Figure 5: Chlorophyll 'a' concentration at the study stations in Monsoon 2022

3.4. Phytoplankton

Phytoplankton are free-floating, photosynthetic, aquatic microorganisms, which are distributed either actively by their locomotory organs (flagella) or passively by water currents. Most of the phytoplankton survive on the open surface waters of lakes, rivers and oceans. The phytoplankton community is mainly represented by algal representatives including both prokaryotes and eukaryotic genera. Plankton populations are mostly represented by members of Cyanobacteria, Chlorophyta, Dinophyta, Euglenophyta, Haptophyta, Chrysophyta, Cryptophyta, and Bacillariophyta. Planktonic representative taxa are absent in other algal divisions like Phaeophyta and Rhodophyta.

Generic Status

There were four groups of phytoplankton occurred during monsoon along the DPA, Kandla coast and its peripheral creek system which include Diatom (Pennales, Centrales), Dinophyceae and Cyanophyceae. The number of genera recorded during the monsoon period was 24 to 33 at the sampling stations with remarkable variations with respect to the composition. The maximum number (33) genera were observed at S-11 and the minimum from S-15 representing 24 genera. As far as generic status is concerned the centrales diatom contributed a greater number of genera (16) followed by Pennales (10) (Fig.6 & Table 3). Among the 4 groups of phytoplankton, the genera *Pseudonitzschia*, *Rhizosolenia*, *Coscinodiscus*, *Eucampia*, *Melosira* and *Planktoniella* occurred at all the sites.

Percentage composition of phytoplankton

The cumulative percentage composition of the five groups of phytoplankton from all the study sites is presented in Fig.7. The percentage composition varied from 5 % to 47 % of which the centrales and pennales are the dominant constituting 47% and 27% respectively. The diatoms pennales and centrales together formed 74% of the phytoplankton population by number of genera as well as number of individuals while the rest is constituted by Dinophyceae (10%) and Cyanophyceae (12%) and Chlorophyceae (4%) during the monsoon 2022.




Figure 6: Number of Phytoplankton genera in Monsoon 2022




Figure 7: Percentage composition of phytoplankton groups in Monsoon 2022

Percentage of occurrence

The percentage occurrence denotes the number of representations by a genus among the sites sampled. The percentage occurrence of different phytoplankton genera varied from 27% to 100% with an average of 78%. Seven phytoplankton genera have the highest percentage of occurrence (100%) (fig 8) followed by *Pleurosigma*, *Gyrosigma*, *Thalassionema* and *Aphanizomenon* (93%) occurrence during the monsoon season




Figure 8: Percentage occurrence of phytoplankton genera in Monsoon 2022

Phytoplankton density and diversity

The density signifies the abundance of plankton which is measured as cell/ individual/L. The phytoplankton density varied from 1,760 No/L to 16,960 No/L with the average 13,483 No/L. The highest phytoplankton density was observed at station S-4 (16,960 No/L) followed by S-12 (16,480 No/L), whereas the lowest 1,760 No/L at S-1(fig.9). Diversity indices have become part of standard methodology in the ecological studies particularly, impact analysis and biodiversity monitoring of the environments (PEET,1974). Biodiversity indices reflects the biological variability which can be used for comparison with space and time. Various species diversity indices respond differently to different environmental and behavioral factors of biotic communities. Among the different stations, the phytoplankton taxa varied from 24 to 33 (Table-4). During monsoon the Margalef and Menhinik richness indices were maximum at stations S-11 (4.28& 0.79). The Shannon diversity index was maximum 3.31 (S-11) and minimum 2.93 at S-15. The Simpson index clearly reflexes the species dominance (genera) at S-11 (0.96) and the low value (0.94) was noticed at S-12.




Figure 9: Phytoplankton density in Monsoon 2022

As per Shannon Wiener's rules for the aquatic environment i.e., both soil and water are classified as very good when H' value is greater than four (>4), whereas the good quality represents the H' value with a range of 4-3, similarly moderate-quality (H' value 3-2), poor quality (H' value 2-1) and very poor-quality H' value significantly less than one (<1). Presently Deendayal Port Authority and its periphery environment has been influenced by contaminants deposited from industries and the cargo movements. Accordingly, species diversity decreases at sites with poor water quality. As deduced from the Shannon diversity index values between 2.93 to 3.31 representing the moderate quality of environmental status dominated by the few genera such as *Pleurosigma*, *Gyrosigma*, *Thalassionema* and *Aphanizomenon*. A community dominated by relatively few species indicates environmental stress (Plafkin *et al.*, 1989). According to Staub *et al.* (1970) species diversity index value between 3.0 to 4.5 represents slightly polluted and the lightly polluted environment, the index value characterizes 2.0-3.0, similarly, moderately polluted environment shows index value of 1.0-2.0 and finally, the heavily polluted environment index value is 0.0-1.0. While considering the overall index values it is inferred that the study sites can be included under the category of lightly polluted.




Figure 10: Different diversity indices a. Shannon Index b. Menhinick Index c. Margalef Index d. Simpson Index

Table 3: Phytoplankton density, percentage composition and occurrence during Monsoon 2022

Group	Genera	S-1	S-2	S-3	S-4	S-5	S-6	S-7	S-8	S-9	S-10	S-11	S-12	S-13	S-14	S-15	PO	PC
Pennales	<i>Amphora</i>	0	0	0	0	160	0	160	0	0	0	20	0	160	0	160	33	0.3
	<i>Bacillaria</i>	0	0	160	0	160	0	0	0	0	0	20	0	160	0	0	27	0.2
	<i>Ditylum</i>	160	0	0	960	480	0	0	640	0	320	100	640	480	0	0	53	1.9
	<i>Pseudonitzschia</i>	1760	320	480	640	960	640	1280	800	320	480	100	1600	960	640	1280	100	6.1
	<i>Pleurosigma</i>	160	320	640	1120	320	960	640	320	0	320	20	480	320	960	640	93	3.6
	<i>Rhizosolenia</i>	160	160	320	480	160	800	960	320	1120	160	40	640	160	800	960	100	3.6
	<i>Synedra</i>	320	0	320	160	0	0	0	160	0	0	20	320	0	0	0	40	0.6
	<i>Fragilaria</i>	480	320	480	0	0	160	0	1600	800	320	80	160	0	160	0	67	2.3
	<i>Gyrosigma</i>	160	320	160	640	800	320	480	1120	320	0	20	480	800	320	480	93	3.2
	<i>Thalassionema</i>	320	480	960	1600	1280	1120	800	480	800	640	60	0	1280	1120	800	93	5.8
Centrales	<i>Bellerochea</i>	800	480	1120	960	800	640	1760	960	1280	640	40	160	800	640	1760	100	6.3
	<i>Biddulphia</i>	160	0	320	160	160	480	0	160	0	640	40	320	160	480	0	73	1.5
	<i>Cheatoceros</i>	160	0	0	160	160	320	160	480	320	640	40	160	160	320	160	87	1.6
	<i>Coscinodiscus</i>	1440	640	320	480	640	800	160	320	1120	960	60	640	640	800	160	100	4.5
	<i>Cyclotella</i>	320	160	160	0	160	0	640	0	320	0	60	0	160	0	640	60	1.3
	<i>Eucampia</i>	800	960	320	1120	160	320	640	640	160	160	40	480	160	320	640	100	3.4
	<i>Hemidiscus</i>	0	0	160	0	0	160	320	0	320	0	40	160	0	160	320	53	0.8
	<i>Lauderia</i>	160	0	1600	800	320	640	160	320	480	320	60	0	320	640	160	87	3.0
	<i>Leptocylindricus</i>	320	480	1120	320	0	160	480	0	160	320	20	640	0	160	480	80	2.3
	<i>Lampriscus</i>	1120	800	480	800	640	480	0	160	320	480	120	1600	640	480	0	87	4.0
	<i>Melosira</i>	640	1760	960	1280	640	320	160	480	800	480	140	960	640	320	160	100	4.8
	<i>Navicula</i>	480	0	160	0	640	320	320	320	160	0	40	160	640	320	320	80	1.9
	<i>Odontella</i>	320	160	480	320	640	320	160	320	160	0	0	160	640	320	160	87	2.1
	<i>Planktoniella</i>	800	160	320	1120	960	480	640	800	1440	640	40	480	960	480	640	100	4.9
	<i>Phaeodactylum</i>	0	640	0	320	0	480	0	160	320	160	20	0	0	480	0	53	1.3
	<i>Triceratium</i>	160	160	160	1120	480	0	0	160	800	960	40	1120	480	0	0	73	2.8

Dinophyceae	<i>Ceratium</i>	160	0	160	160	160	160	1120	480	0	0	20	800	960	320	1120	73	2.8
	<i>Prorocentrum</i>	160	800	480	160	480	480	480	0	0	160	40	0	320	0	320	73	1.9
	<i>Photoperidinium</i>	640	1280	0	640	0	0	960	320	480	160	100	960	320	1120	160	80	3.5
	<i>Noctiluca</i>	160	800	160	480	160	160	160	320	160	0	0	0	160	0	0	67	1.3
Cyanophyceae	<i>Aphanizomenon</i>	160	160	160	160	160	160	320	160	640	800	40	480	1120	320	0	93	2.4
	<i>Cosomarium</i>	0	640	640	480	640	640	0	960	1600	1280	140	800	480	800	640	87	4.8
	<i>Trichodesmium</i>	160	160	0	320	0	0	1120	1120	960	800	80	1760	960	1280	640	80	4.6
Chlorophyceae	<i>Chlorella</i>	800	320	160	0	160	160	960	0	0	160	40	320	640	160	320	80	2.1
	<i>Scenedesmus</i>	1280	480	160	0	160	160	160	800	320	640	20	0	160	480	0	80	2.4
Total genera		30	25	29	27	28	27	26	28	26	25	33	26	30	27	24		
Density No/L		14720	12960	13120	16960	12640	11840	15200	14880	15680	12640	1760	16480	15840	14400	13120		

Table 4: Diversity indices of Phytoplankton during Monsoon 2022

Diversity Indices	S-1	S-2	S-3	S-4	S-5	S-6	S-7	S-8	S-9	S-10	S-11	S-12	S-13	S-14	S-15
Shannon_H	3.07	2.98	3.09	3.09	3.10	3.12	3.01	3.12	3.03	3.05	3.31	3.02	3.20	3.13	2.93
Simpson_1-D	0.94	0.94	0.94	0.95	0.95	0.95	0.94	0.95	0.94	0.95	0.96	0.94	0.95	0.95	0.94
Margalef	3.02	2.53	2.95	2.67	2.86	2.77	2.60	2.81	2.59	2.54	4.28	2.58	3.00	2.72	2.43
Menhinick	0.25	0.22	0.25	0.21	0.25	0.25	0.21	0.23	0.21	0.22	0.79	0.20	0.24	0.23	0.21
Dominance_D	0.06	0.06	0.06	0.05	0.05	0.05	0.06	0.05	0.06	0.05	0.04	0.06	0.05	0.05	0.06

3.5. Zooplankton

Zooplankton are highly sensitive to changes caused by physical and chemical factors in aquatic ecosystems and their distribution deliver information regarding the productivity and pollution of the particular area (Gajbhiye and Desai, 1981). Zooplankton are distributed in a wide range of habitats extending from the neuston to benthos and play vital roles influencing fisheries, oceanography and climate (Terdalkar and Pai, 2001). It has various significant roles in the estuarine ecosystem and connecting link between nutrient cycling and phytoplankton, primary production and many commercial fisheries in estuaries and coastal waters and form a chief food for a variety of pelagic consumers including coelenterates, ctenophores, fish larva forage fish and some benthic organisms such as sponges and molluscs (Day *et al.*, 1989).

Phylum, group and generic status

The zooplankton identified from the 15 stations falls under 10 phyla and 41 genera belonging to the 16 groups (Table 5). The phylum Arthropoda was the predominant, represented with 25 genera including copepods, crabs, shrimps and their larva. The phylum Arthropoda dominated in the samples with major groups Calanoida, Harpacticoida, Cyclopoida, (Copepoda) Decapoda, and the larval forms of crustaceans. There were 14 genera of copepods in the samples. Among copepods, the Calanoida ranked first in terms of generic representation particularly *Acartia* sp, *Acrocalanus* sp, *Aetideus* sp. and *Calanus* sp. (figure-11).

Percentage composition

The overall percentage of the various groups of zooplankton varied from 0.3% to 36.9%. The highest percentage was due to the calanoid copepods (36.9%) followed by Decapoda (13.2%) and Gastropoda (8.2%). The group which contributed the least was *Chaetognatha* (0.3%) followed by Nematoda (0.4%) (Fig.12). Among the zooplankton groups calanoid group was observed predominantly at all sites.




Figure 11: Phylum and generic status of zooplankton during Monsoon 2022




Figure 12: Percentage composition of zooplankton groups during Monsoon 2022

Percentage occurrence of zooplankton

The percentage occurrence of zooplankton communities varied from 33% to 100 %. There were 9 zooplankton genera that exhibited 100% of occurrence (Fig.12) followed by the copepods *Microsetella*, *Aerocalanus*, *Copelata*, *Eucalanus* and the Cyphonautes larva (93%) from the study sites (Table5).




Figure 13: Percentage occurrence of Zooplankton groups during Monsoon 2022

Density of zooplankton

Zooplankton population density values during the Monsoon 2022 at the 15 sampling sites ranged from 12,640 No/L to 21,120 No/L with an overall average of 16,789 No/L (Table 5). Station-wise, the highest density of 21,120 No/L was recorded in S-7 followed by S-2 (18,880 No/L) and lowest density was reported at S-5 (12,640 No/L) (Figure 14).

Diversity Index

The Shannon diversity index of the zooplankton ranged between 3.05 to 3.41. Similarly, Margalef and Menhinick species richness index also varied from 2.75 to 3.70, and 0.22 to 0.29 respectively representing the moderate quality of the environment. (Table 6).




Figure 14: Zooplankton Density in the different stations during Monsoon 2022

Table 5: Zooplankton generic status during Monsoon 2022 in Deendayal Port Authority area

Phylum	Group	Genera	S-1	S-2	S-3	S-4	S-5	S-6	S-7	S-8	S-9	S-10	S-11	S-12	S-13	S-14	S-15	PO	PC
Protozoa	Foraminifera	<i>Globigerina</i>	160	160	0	0	0	160	320	0	0	0	320	160	0	480	320	53	0.8
Ciliophora	Tintinnida	<i>Tintinnopsis</i>	480	320	320	160	480	160	320	640	320	480	640	320	160	320	320	100	2.2
Chaetognath		<i>Sagitta</i>	160	160	0	0	0	0	0	160	0	160	0	0	0	160	0	33	0.3
Nematoda		<i>Nemadodes</i>	320	160	0	0	0	160	0	160	0	0	0	0	0	160	0	33	0.4
Annelida		<i>Polychaete larva</i>	1120	480	320	160	0	0	960	480	0	0	640	0	320	800	640	67	2.4
Arthropoda	Calanoida	<i>Acartia</i>	480	1280	800	1760	320	480	640	960	640	1280	800	320	480	800	1600	100	5.0
		<i>Acrocalanus</i>	640	320	480	160	320	640	1120	320	960	640	320	0	320	160	480	93	2.7
		<i>Aetideus</i>	320	800	640	160	160	320	480	160	800	960	320	1120	160	320	640	100	2.9
		<i>Calanus</i>	480	320	0	320	0	320	160	0	0	0	160	0	0	160	320	53	0.9
		<i>Calanopia</i>	1120	800	320	480	320	480	0	0	160	0	1600	800	320	640	160	80	2.9
		<i>Centropages</i>	320	480	0	160	320	160	640	800	320	480	1120	320	0	160	480	87	2.3
		<i>Eucalanus</i>	640	480	160	320	480	960	1600	1280	1120	800	480	800	640	480	0	93	4.1
		<i>Labidocera</i>	320	160	480	800	480	1120	960	800	640	1760	960	1280	640	320	160	100	4.3
		<i>Nannocalanus</i>	160	320	320	160	0	320	160	160	480	0	160	0	640	320	320	80	1.4
		<i>Paracalanus</i>	320	160	320	160	0	0	160	160	320	160	480	320	640	320	160	87	1.5
		<i>Pontella</i>	1760	480	800	1440	640	320	480	640	800	160	320	1120	960	480	640	100	4.4
		<i>Pseudodiaptomus</i>	0	0	160	320	160	160	0	160	0	640	0	320	0	480	0	53	1.0
		<i>Rhincalanus</i>	320	480	160	800	960	320	1120	160	320	640	640	160	160	320	480	100	2.8
		<i>Temora</i>	320	160	0	0	0	160	0	0	160	320	0	320	0	320	160	53	0.8
	Harpacticoida	<i>Corycaeus</i>	480	0	0	160	0	1600	800	320	640	160	320	480	320	480	0	73	2.3
		<i>Euterpina</i>	160	640	800	320	480	1120	320	0	160	480	0	160	320	160	640	87	2.3

Deendayal Port Authority 2nd Year Monsoon (June-September 2022)

		<i>Microsetella</i>	960	160 0	128 0	112 0	800	480	800	640	480	0	160	320	480	960	1600	93	4.6
	Cyclopoida	<i>Oithona</i>	112 0	960	800	640	176 0	960	128 0	640	320	160	480	800	480	112 0	960	100	5.0
		<i>Oncaea</i>	320	160	160	480	0	160	0	640	320	320	320	160	0	320	160	80	1.4
	Decapoda	<i>Caridean larva</i>	0	160	160	320	160	480	320	640	320	160	320	160	0	0	160	80	1.3
		<i>Euphausia</i>	320	480	640	800	160	320	112 0	960	480	640	800	144 0	640	320	480	100	3.8
		<i>Nauplius larva</i>	160	0	160	0	640	0	320	0	480	0	160	320	160	160	0	60	1.0
		<i>Mysis</i>	160	0	160	160	160	160	112 0	480	0	0	160	800	960	320	1120	80	2.3
		<i>Phyllosoma</i>	160	800	480	160	480	480	480	0	0	160	320	0	320	0	320	73	1.7
		<i>Lucifer</i>	640	128 0	0	640	0	0	960	320	480	160	800	960	320	112 0	160	80	3.1
	Crustacean larva	<i>Barnacle nauplius</i>	160	800	160	480	160	160	160	320	160	0	0	0	160	0	0	67	1.1
	Malacostraca	<i>Brachyuran larva</i>	320	160	320	960	320	320	480	480	0	0	160	0	160 0	800	320	80	2.5
Bryozoan		<i>Cyphonautes larva</i>	160	160	160	160	160	160	320	160	640	800	320	480	112 0	320	0	93	2.0
Mollusca	Gastropod	<i>Creseis sp</i>	0	640	640	480	640	640	0	960	160 0	128 0	112 0	800	480	800	640	87	4.3
		<i>Gastropod larva</i>	160	160	0	320	0	0	112 0	112 0	960	800	640	176 0	960	128 0	640	73	3.9
	Bivalve	<i>Veliger larva</i>	0	320	0	0	0	0	480	112 0	160	320	640	0	160	320	0	53	1.4
Echinoder mata		<i>Bipinnaria larva</i>	800	320	160	0	160	160	960	0	0	160	320	320	640	160	320	80	1.8
Hemichord ata		<i>Tornaria larva</i>	128 0	480	160	0	160	160	160	800	320	640	160	0	160	480	0	80	2.0
Chordata	Appendicula ria	<i>Oikopleura</i>	800	160	0	800	0	0	480	320	0	160	480	640	480	0	160	67	1.8
		<i>Copelata</i>	160	960	160 0	128 0	112 0	800	320	800	640	480	0	640	320	160	480	93	3.9
	Fish	<i>Fish larva</i>	160	112 0	960	800	640	176 0	0	128 0	640	320	160	640	320	320	320	100	3.7
		Total genera	37	37	30	33	27	33	33	33	30	30	34	30	32	37	31		
		Density No/L	179 20	188 80	140 80	174 40	126 40	161 60	211 20	190 40	158 40	156 80	168 00	182 40	158 40	168 00	1536 0		

Table 6. Diversity indices of Zooplankton along Deendayal Port Authority area during Monsoon 2022

	S-1	S-2	S-3	S-4	S-5	S-6	S-7	S-8	S-9	S-10	S-11	S-12	S-13	S-14	S-15
Taxa_S	37	37	30	33	27	33	33	33	30	30	34	30	32	37	31
Shannon_H	3.34	3.36	3.14	3.22	3.05	3.20	3.31	3.31	3.23	3.15	3.32	3.19	3.27	3.41	3.20
Simpson_1-D	0.96	0.96	0.95	0.95	0.94	0.95	0.96	0.96	0.95	0.95	0.96	0.95	0.96	0.96	0.95
Margalef	3.68	3.66	3.04	3.28	2.75	3.30	3.21	3.25	3.00	3.00	3.39	2.96	3.21	3.70	3.11
Menhinick	0.28	0.27	0.25	0.25	0.24	0.26	0.23	0.24	0.24	0.24	0.26	0.22	0.25	0.29	0.25



3.6. Intertidal Fauna

The intertidal zone is the area above the water level at low tide and submerged at high tide. Intertidal habitats are found along the margins of the sea and include rocky shores, mudflats, salt marshes, and estuaries. The intertidal diversity was documented during monsoon at the prefixed 15 sampling locations within the DPA jurisdiction. All the macroinvertebrates and vertebrate samples were collected from the sampling stations during the low tide. At each site, 1x 1 m² quadrat was placed randomly, and all visible macro-faunal organisms encountered inside the quadrat were identified, counted and recorded. At each site along the transects that run perpendicular to the waterfront, three to six replicate quadrat samples were assessed for the variability in macro-faunal population structure (Davidson *et al.*, 2004; Ravinesh and Biju Kumar, 2013). The density of the different faunal groups was averaged for the entire intertidal belt. Organisms, which could not be identified in the field, were preserved in 5% formaldehyde, brought to the laboratory and identified using standard identification keys (Abott, 1954; Vine, 1986; Oliver, 1992; Rao, 2003; 2017; Psomadakis *et al.*, 2015; Apte, 2012; 2014; Naderloo 2017; Ravinesh *et al.*, 2021; Edward *et al.*, 2022). The invertebrates' taxonomic composition, relative abundance, species richness and diversity were determined (Zar, 1984) to describe the mangrove environment's overall biodiversity at DPA premises. Statistical analyses such as diversity indices and richness were calculated using Paleontological Statistics Software Package for Education and Data (PAST) version 3.2.1 (Hammer *et al.*, 2001).

Faunal composition of intertidal macrobenthos

The intertidal ecological survey has been conducted at the prefixed 15 locations within the vicinity of the Deendayal port Authority. The species diversity of the invertebrate phyla showed the maximum for phylum Arthropoda (8 species), which is followed by Mollusca (6 species). The phylum Chordata was represented by two species (Table 7 & Fig.15).




Figure 15: Number of genera of intertidal fauna (Phylum) during in Monsoon 2022




Figure 16: Percentage composition of intertidal fauna during Monsoon 2022

Cumulative percentage composition of Fauna

The overall percentage composition of the three groups of intertidal fauna at the 15 sites was followed, ie Arthropoda (50%), Mollusca (37%), and Chordata (13%), as shown in figure 16.

Intertidal Fauna density (No/m²) variation between the stations

The number of individuals of the Fauna collected from the intertidal zone of the mangroves are presented in Fig 17. It was observed that the faunal density was the highest in stations S- 3 and S-4 while the least from S-10.




Figure 17: Density of intertidal fauna during Monsoon 2022

The Intertidal faunal diversity documented during the monsoon period of 2022 has shown that the highest number of animals were collected from S-3, and the lowest was from S-10. The most common species were the crustaceans such as *Parasesarma plicatum* and *Austruca iranica*. The lowest density noticed was that of *Littoraria pallescens* (Table.7)

Diversity indices

Table.8 presents the various diversity indices calculated for the different fauna recorded from the 15 sites adjoining the DPA port area, Kandla. Diversity indices were calculated for the subtidal fauna in which the Dominance diversity (D) values varied from 0.12 (S-5, S-15) to 0.27 (S -3). Shannon diversity (H') values varied from 1.50 (S-10) to 2.31 (S-5). The Simpson_1-D varied from 0.73 (S -3) to 0.88 (S-5, S-15). The Evenness values varied from 0.42 to 0.83, with the maximum in S-3 and the minimum at S-12. The Margalef index ranged from 1.04 to 2.15, the maximum at S-13 and the minimum at S-3.



Table 7: Intertidal faunal distribution along Deendayal Port Authority area during Monsoon 2022

Intertidal Fauna	S-1	S-2	S-3	S-4	S-5	S-6	S-7	S-8	S-9	S-10	S-11	S-12	S-13	S-14	S-15
Arthropoda															
<i>Scylla serrata</i>	0	3	0	0	5	0	6	0	0	0	5	0	0	3	4
<i>Austruca sindensis</i>	0	17	6	8	11	0	18	23	12	15	19	17	0	4	9
<i>Austruca iranica</i>	12	19	16	31	21	24	28	26	31	39	41	52	11	26	19
<i>Parasesarma plicatum</i>	56	72	32	52	23	42	26	53	85	19	36	42	38	52	28
<i>Dotilla blanfordi</i>	0	1	2	0	2	0	1	0	0	0	0	0	0	2	3
<i>Eurycarcinus orientalis</i>	2	0	0	1	2	5	1	0	0	0	0	0	0	0	2
<i>Amphibalanus amphitrite</i>	0	23	0	56	11	0	0	38	0	0	0	21	0	0	14
<i>Tubuca dussumieri</i>	3	2	1	6	9	1	2	1	8	2	1	6	0	0	5
Mollusca															
<i>Pirenella cingulata</i>	2	8	123	19	0	11	35	0	12	0	8	0	31	6	0
<i>Telescopium telescopium</i>	0	0	2	3	0	0	6	0	2	0	5	0	2	0	1
<i>Bakawan rotundata</i>	8	0	5	0	2	0	15	0	0	0	12	0	0	2	8
<i>Littoraria pallescens</i>	0	1	2	0	2	0	0	0	0	0	0	0	0	0	0
<i>Platevindex martensi</i>	0	0	1	0	2	0	0	0	0	0	5	0	0	2	1
<i>Optedicerus breviculum</i>	35	42	52	12	7	42	0	0	34	0	15	25	0	0	19
Chordata															
<i>Periophthalmus waltoni</i>	25	11	15	21	12	7	8	9	11	4	2	9	11	8	26
<i>Scartelaos histophorus</i>	26	12	11	13	25	31	32	19	12	21	23	19	27	31	18
Total	169	211	268	222	134	163	178	169	207	100	172	191	120	136	157

Table 8: Diversity indices of Intertidal Fauna during Monsoon 2022

Indices	S-1	S-2	S-3	S-4	S-5	S-6	S-7	S-8	S-9	S-10	S-11	S-12	S-13	S-14	S-15
Dominance_D	0.21	0.19	0.27	0.16	0.12	0.20	0.14	0.21	0.23	0.26	0.15	0.17	0.23	0.24	0.12
Shannon_H	1.77	1.95	1.70	2.02	2.31	1.75	2.12	1.69	1.77	1.50	2.11	1.89	1.56	1.70	2.30
Simpson_1-D	0.79	0.81	0.73	0.84	0.88	0.80	0.86	0.79	0.77	0.74	0.85	0.83	0.77	0.76	0.88
Evenness_e^H/S	0.65	0.58	0.42	0.69	0.72	0.72	0.69	0.77	0.65	0.75	0.69	0.83	0.79	0.55	0.71
Margalef	1.56	2.06	2.15	1.85	2.65	1.37	2.12	1.17	1.50	1.09	2.14	1.33	1.04	1.83	2.57



3.7. Subtidal Fauna (Macrobenthos)

Subtidal ecosystems are permanently submerged due to tidal influence, whereas intertidal ecosystems are found between the high tide and low tide, experiencing fluctuating influences of land and sea. Macrobenthos are an important component of estuarine and marine ecosystems. At large scales, food may be the prime limiting factor for benthic biomass. Depending on the system's characteristics, grazing by benthic suspension feeders may be the most important factor determining system dynamics. The sampling methods and procedures were designed in such a way as to obtain specimens in the best possible condition to maximize the usefulness of the data obtained. For studying the benthic organisms, triplicate samples were collected at each station using Van Veen grab, which covered an area of 0.04m². The wet sediment was passed through a sieve of mesh size 0.5 mm for segregating the organisms. The organisms retained in the sieve were fixed in 5-7% formalin and stained further with Rose Bengal dye for the ease of spotting at the time of sorting (Ravinesh and Biju Kumar, 2022). The number of organisms in each grab sample was expressed as No /m². All the species were sorted, enumerated and identified by following available literature. The works of Day (1967), Hartman (1968, 1969), Rouse and Pleijel (2001), Robin et al., (2003), Amr (2021), were referred for polychaetes; Crane (1975), Holthuis (1993), Naderloo (2017). Xavier *et al.*, (2020) for crustaceans; Subba Rao (1989, 2003, 2017), Apte (2012,2014), Ramakrishna and Dey (2007), Ravinesh *et al.* (2021) and Edward *et al.*, (2022). for molluscs. Statistical analyses such as diversity indices and quadrat richness were calculated using Paleontological Statistics Software Package for Education and Data (PAST) version 3.2.1 (Hammer *et al.*, 2001).

Faunal composition of subtidal macrobenthos

The number of macrobenthic species of the various groups recorded (Fig.18) from the DPA port environment revealed that Mollusca (14 species) and Annelida (2 species) were the major constituents, while the Arthropoda (1 species) and Cnidaria (1 species) were comparatively low in the species composition.

The percentage composition of the four phyla that occurred during the monsoon is shown in (Fig 19) The phylum Mollusca is represented by maximum (78%) share of the subtidal Fauna, followed by Annelida (11%), Arthropoda (5.5%) and Cnidaria (5.5%) in the total benthic samples collected.




Figure 18. Number of genera of macrobenthos during Monsoon 2022




Figure 19: Percentage composition of macrobenthos during Monsoon 2022

Subtidal Faunal density (No/m²) variation between the stations

The number of individuals of the animals collected from the different sites are shown in Fig 20. The density of the Fauna was high at S-7 (24No/m²), and the lowest number (6/m²) was noticed at S-13 during the monsoon season 2022.




Figure 20: Subtidal fauna density during Monsoon 2022

Subtidal fauna distribution at the selected sites in the Deendayal Port area during monsoon

The table.9 depicts the subtidal microbenthic faunal diversity documented in the monsoon 2022. The highest diversity was documented from stations S-7, S-14, S-4 and S-1 and the lowest from stations S-9,10 and S- 6. The most common species are *Optedicerus breviculum*, *Glaucanome angulata* and *Pirenella cingulata*. The least diversity was documented for *Turritella* sp, *Stephensonactis* sp and *Natica* sp were found significantly less diversity. The Table.10 presents the various diversity indices calculated for the different Fauna recorded from the 15 sites adjoining

the DPA port area, Kandla. Diversity indices were calculated for the subtidal fauna in which the Dominance diversity (D) values varied from 0.12 (S- 4) to 0.24 (S -9). Shannon diversity (H') values varied from 1.52 (S-9) to 2.27 (S-4). The Simpson_1-D varied from 0.76 (S -9) 0.87 (S-3, S-15). The Evenness values varied from 0.72 to 0.96, with the maximum in S-3 and the minimum at S-14. The Margalef index ranged from 1.67 to 3.03, the maximum at S-3 and the minimum at S-15.



Table 9: Macro-benthic faunal distribution during Monsoon 2022 in Deendayal Port Area

	S-1	S-2	S-3	S-4	S-5	S-6	S-7	S-8	S-9	S-10	S-11	S-12	S-13	S-14	S-15	% of Occurrence
Cnidaria																
<i>Stephensonactis</i> sp.	0	0	0	0	0	0	0	2	0	0	0	0	0	0	0	0.9
Annelida																
<i>Lumbrineria</i> sp.	0	2	0	0	0	2	0	0	2	0	0	0	0	0	0	2.6
<i>Nereis</i> sp.	0	0	3	0	0	0	0	0	0	1	0	0	1	0	1	2.6
Arthropoda																
<i>Ampithoe</i> sp.	1	0	0	0	0	0	0	0	0	0	0	0	0	0	2	1.3
Mollusca																
<i>Umbonium vestiarius</i>	0	0	0	3	0	1	2	1	0	0	0	0	0	1	0	3.5
<i>Mitrella blanda</i>	0	0	0	2	0	1	0	2	2	0	3	0	0	5	0	6.5
<i>Clypeomorus bifasciata</i>	1	0	2	0	1	0	0	1	0	0	0	3	0	0	2	4.3
<i>Natica</i> sp.	0	0	0	1	0	0	1	0	0	0	0	0	0	0	0	0.9
<i>Optedicerus breviculum</i>	5	1	2	1	2	2	4	5	4	1	1	3	1	2	1	15.2
<i>Pirenella cingulata</i>	5	2	3	1	2	1	1	2	2	1	1	2	1	1	2	11.7
<i>Turritella</i> sp.	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0.4
<i>Mactra</i> sp.	0	1	0	3	0	0	2	1	0	0	0	2	0	2	3	6.1
<i>Glaucanome angulata</i>	4	1	2	1	2	3	5	0	0	2	3	2	0	1	0	11.3
<i>Pelecypora</i> sp.	0	0	1	2	0	1	3	0	0	2	1	1	0	0	1	5.2
<i>Gafrarium divaricatum</i>	2	0	0	2	1	0	0	0	0	0	0	2	1	1	0	3.9
<i>Meretrix</i> sp.	0	2	0	4	3	0	1	2	0	3	0	0	0	0	1	6.9
<i>Solen</i> sp.	1	0	2	0	0	0	4	0	1	0	2	0	0	7	1	7.8
<i>Protapes cor</i>	1	3	2	1	1	2	1	2	0	1	3	1	2	1	0	9.1
Total	20	12	17	21	12	13	24	18	11	11	14	16	6	22	14	100
Total No/m ²	500	300	425	525	300	325	350	450	275	275	350	400	150	550	350	

Table10: Diversity indices of the benthic fauna during Monsoon 2022

Indices	St_1	St_2	St_3	St_4	St_5	St_6	St_7	St_8	St_9	St_10	St_11	St_12	St_13	St_14	St_15
Dominance_D	0.19	0.17	0.13	0.12	0.17	0.15	0.14	0.15	0.24	0.17	0.17	0.14	0.22	0.18	0.13
Shannon_H	1.84	1.86	2.04	2.27	1.86	1.99	2.13	2.06	1.52	1.85	1.83	2.01	1.56	1.98	2.11
Simpson_1-D	0.82	0.83	0.87	0.88	0.83	0.85	0.86	0.85	0.76	0.83	0.83	0.86	0.78	0.82	0.87
Evenness_e^H/S	0.79	0.92	0.96	0.88	0.92	0.92	0.84	0.87	0.91	0.91	0.89	0.94	0.95	0.72	0.91
Margalef	2.34	2.42	2.47	3.29	2.42	2.73	2.83	2.77	1.67	2.50	2.27	2.53	2.23	2.91	3.03



3.8. Seaweeds

Along the Gujarat coast which is represented by 1600 km coastline, harbors 198 species of which 109 species from 62 genera belonging to Rhodophyta, 54 species of 23 genera to Chlorophyta, and 35 species from 16 genera to Ochrophyta (Jha *et.al.*, 2009). According to Mantri *et.al.* (2020) there are 13 potential sites for the occurrence of seaweed density and diversity. The survey conducted by CSIR-CSMCRI (Jha *et.al.*, 2009) confirmed the presence of industrially important taxa, namely, *Gelidiella acerosa*, *Gelidium micropterum*, *G. pusillum*, *Ahnfeltia plicata*, *Gracilaria dura*, *G. debilis*, *Gracilariopsis longissima* (formerly *G. verrucosa*), *Hypnea musciformis*, *Meristotheca papulosa*, *Porphyra sp*, *Asparagopsis taxiformis* (Rhodophyta), *Sargassum tenerrimum*, *S. plagiophyllum*, *S. swartzii*, *Turbinaria ornata* (Ochrophyta), *Ulva prolifera* (formerly *Enteromorpha prolifera*), *Ulva compressa* (formerly *Enteromorpha compressa*), and *Ulva flexuosa* (formerly *Enteromorpha tubulosa*) (Chlorophyta) from the coastal waters of Gujarat. In the present study, an attempt was made to describe the occurrence, diversity and other ecological features of seaweeds within Deendayal Port jurisdiction. It was found that except for some drifted species *Enteromorpha* and *Chaetomorpha* at S-13 and S-14 of Vira coast (Plate-6) no natural seaweed beds are seen in the different locations within DPA environment.

Seaweeds grow in the rocky intertidal and sub tidal habitats that offer a hard substratum for attachment. Low turbidity level in the water column with high nutrient content is a major habitat requirement that enables photosynthesis. Total dissolved solids (TDS) load in the Deendayal Port area creek waters ranged from 32088 to 42086 mg/L and suspended solids value between 88-223 mg/L restricts the photosynthetic activity of seaweeds which are highly sensitive to light. Hence, seaweed formations are absent in the creek systems of the Deendayal Port coastal environment.

3.9. Seagrass

Similar to seaweeds, sea grasses were also absent in the creek systems of Deendayal Port area and in the adjacent coastal stretches of Kachchh due to inherent habitat conditions. Sea grasses generally thrive in shallow coastal waters and are adapted to live in submerged conditions from mid intertidal to depth as much as 50 m when light penetration is sufficient; conditions contrary to the one prevailing in Deendayal Port and the nearby creek systems explain the total absence of sea grasses.

3.10. Halophytes

The halophytes are the plants that are adopted in coastal estuaries and salt marshes. It is common in arid and desert milieu which often have substantial salt accumulation. Technically it is the plant which has tolerance to moderate to high salt concentration in its growth substrate. Halophytes, that survive to reproduce in environments where the salt concentrations around 200 mM NaCl or more, constitute about 1% of the world's flora. (Timothy *et al.*, 2008). Halophytes are classified based on their growth conditions as obligate halophytes, facultative halophytes, and habitat-indifferent halophytes. In the present study, four major halophytes recorded along the selected Deendayal Port Authority sites during the Monsoon sampling, were *Salicornia brachiata*, *Aeluropus lagopoides*, *Salvadora persica* and *Sesuvium portulacastrum*. Among the halophyte species recorded, *Salicornia brachiata* alone was found in the 3 sampling locations. (Table-11 and Plate-12). The percentage of *Salicornia brachiata* was found to be the highest at station S-8 (78%) and the lowest in S-11.

Table 11: Percentage of Halophytes cover in the DPA during Monsoon 2022

	S-1	S-2	S-3	S-4	S-5	S-6	S-7	S-8	S-9	S-10	S-11	S-12	S-13	S-14	S-15
<i>Aeluropus lagopoides</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Salicornia brachiata</i>	0	0	0	0	0	0	0	78%	63%	0	57%	0	0	0	0
<i>Salvadora persica</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Sesuvium portulacastrum</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0





a. *Salicornia brachiata* b. *Aeluropus lagopoides* c. *Salvadora persica* d. *Sesuvium portulacastrum*

Plate 7: Halophyte species on the intertidal zone of Deendayal Port Authority area

3.11. Mangroves

In India, the second largest mangrove cover is located in the Gujarat state which accounts for 1175 km² (23.66%) cover of mangroves. However, it is also the fact that, this mangrove cover is predominance of *Avicennia marina*. In Gujarat, the Gulf of Kachchh shows major part of mangrove abundance, particularly of *A. marina*. The arid and hot environment of this area make it mono-species formation of *A. marina* within DPA area of Kandla.

Tree Density

In this study, totally 13 sites were surveyed for recoding the mangrove growth parameters and the density of plants. The overall average density of mangrove was 4602 plants per hectare. Among all sampling stations, the mean plant density was maximum at Tuna creek (6199/ha), followed by Kandla creek (5205/ha). Considering the sampling sites individually the highest tree density was reported at S-12 station in the Tuna creek area (7359/ha). The lowest average tree density (2935 trees/ha) was reported in Phang creek, however, the lowest density (individual site) was recorded in the site S-5 at Phang creek. From this study, it is clear that geomorphology and environmental characteristics of the Kandla coastal regions play an important role in the formation of variability in mangrove (Fig.21 & Table 12).




Figure 21. Mangrove Plant density during Monsoon 2022

Height

The overall mean height of the mangroves from the study sites along the DPA port environment was 105 cm. The highest average tree height was found at Phang creek area (167 cm) followed by Navlakhi creek (160 cm). The highest tree height was recorded in station S-9 of Navlakhi creek, followed by S-4 of Kandla creek (Fig. 22).




Figure 22. Plant height during Monsoon 2022

Canopy Crown Cover

The canopy cover of sampling stations exhibited wide variation and the average was 2.54 m². The sites S-5, S-9 and S-10 showed relatively large canopy cover. However, the lowest canopy cover was reported at S-2 and S-7 stations located at Tuna creek and Kharo creek respectively (Fig.23).

Basal area

The overall average basal area (GBH) of the mangroves of the DPA environment was 14.64 cm. Station wise the maximum mean basal area (21 cm) was at S-4 located in the Kandla creek followed by S-5 and S-11 in Phang creek and Jangi creek respectively. The minimum basal area reported to all sites was 7 cm (Fig.24). The highest value of DBH indicates the mangrove plants have multiple stems or main branches arising close to the ground from a single buttress or base. This type of growth pattern is characteristics of mangroves particularly *Avicennia marina* and *Aegiceros corniculatum*




Figure 23. Mangrove canopy cover during Monsoon 2022




Figure 24. Mangrove basal area during Monsoon 2022

Regeneration and Recruitment class

During the monsoon, generally higher values of regeneration class of mangroves is expected, but the average density was 60167 plants/ha and that of recruitment class 15434 plants/ha. The highest regeneration (140000 plants/ha) at S-9 of Navlakhi creek and recruitment (31500 plants/ha) class density were recorded at Kharo creek (S-7). The lowest regeneration class and recruitment plant density were found at S-14 station of Vira coast site. The highest density of recruitment class after the S-7 site was observed at S-8 and S-9 sites of Navlakhi creek.



Plate 8: Mangrove species recorded along the Deendayal Port area

a. Avicennia marina b. Aegiceras corniculatum c. Ceriops tagal d. Rhizophora mucronata

Table 12: Density of mangroves in the DPA vicinity during monsoon 2022

Sampling stations	Density (Tree/Ha)	Tree height (m)			Canopy cover (m)			Basal Area (cm)		
		Min	Max	Avg.	Min	Max	Avg.	Min	Max	Avg.
Tuna creek										
S-2	5038	110.00	230.00	153.00	0.24	6.48	1.00	7.00	36.00	13.00
S-12	7359	100.00	300.00	158.00	0.42	11.55	2.00	7.00	43.00	15.00
Mean	6198.64	105.00	265.00	155.50	0.33	9.02	1.50	7.00	39.50	14.00
Phang creek										
S-5	2311	110.00	220.00	149.00	0.88	11.20	5.00	7.00	50.00	19.00
S-10	3558	100.00	310.00	185.00	0.63	10.50	4.00	9.00	43.00	18.00
Mean	2934.70	105.00	265.00	167.00	0.76	10.85	4.50	8.00	46.50	18.50
Kandla creek										
S-3	3669	100.00	160.00	130.00	0.05	5.04	2.00	7.00	32.00	14.00
S-4	6400	110.00	310.00	189.00	0.16	6.48	2.00	8.00	50.00	21.00
S-15	5545	110.00	220.00	149.00	0.77	7.20	3.00	7.00	30.00	16.00
Mean	5204.96	106.67	230.00	156.00	0.33	6.24	2.33	7.33	37.33	17.00
Kharo creek										
S-7	5144	100.00	300.00	133.00	0.30	6.25	1.00	7.00	43.00	10.00
Jangi creek										
S-6	3483	100.00	190.00	132.00	0.17	3.99	2.00	8.00	14.00	11.00
S-11	3906	110.00	185.00	139.00	2.24	3.42	2.90	9.00	30.00	19.00
Mean	3694.59	105.00	187.50	135.50	1.21	3.71	2.45	8.50	22.00	15.00
Navlakhi creek										
S-8	5045	100.00	210.00	125.00	0.35	8.00	2.00	7.00	25.00	10.00
S-9	3290	110.00	420.00	196.00	0.30	42.25	4.00	7.00	85.00	16.00
Mean	4167.65	105.00	315.00	160.50	0.33	25.13	3.00	7.00	55.00	13.00
Vira coast										
S-14	4867.50	110.00	210.00	132.00	0.48	8.00	3.00	7.00	35.00	15.00
Overall average	4601.71	105.24	253.21	148.50	0.53	9.88	2.54	7.40	39.76	14.64

Table 13: Regeneration and Recruitment class plants during Monsoon 2022

Station	Tree density- No/ha (1)	Regeneration density- No/ha (2)	Recruitment density- No/ha (3)	Ratio of 1:3	Ratio of 2:3
Tuna creek				1 to	to 1
S-2	5038	68000	13250	2.63	5.13
S-12	7359	70000	16500	2.24	4.24
Mean	6198.64	69000	14875	2.40	4.64
Phang creek					
S-5	2311	24000	3750	1.62	6.40
S-10	3558	75000	17500	4.92	4.29
Mean	2934.70	49500	10625	3.62	4.66
Kandla creek					
S-3	3669	79000	17000	4.63	4.65
S-4	6400	56000	8250	1.29	6.79
S-15	5545	23000	3750	0.68	6.13
Mean	5204.96	52667	9667	1.86	5.45
Kharo creek					
S-7	5144	77000	31500	6.12	2.44
Jangi creek					
S-6	3483	49000	13250	3.80	3.70
S-11	3906	79000	18000	4.61	4.39
Mean	3694.59	64000	15625	4.23	4.10
Navlakhi creek					
S-8	5045	52000	26500	5.25	1.96
S-9	3290	140000	19500	5.93	7.18
Mean	4167.65	96000	23000	5.52	4.17
Vira coast					
S-14	4867.50	13000	2750	0.56	4.73
Overall average	4601.71	60166.67	15434.52	3.35	3.90

3.12. Marine Reptiles

During the field surveys, one reptilian species, the saw-scaled viper *Echis carinatus sochureki* was recorded at site S-3 located in the northern part of Sat Saida bet opposite to oil jetty during monsoon season. This species was spotted on the ground among the mangrove trees. The literature describes the species as aggressive and strikes at a lightning speed, the observed specimen was active. In monsoon, the maximum number of this snake was recorded in S-10 located on the northern part of Sat Saida bet.



Plate 9: Marine reptiles recorded along the Deendayal Port Authority area

3.13. Marine Fishery

Marine fish production of India during the financial year 2019-2020 was 37.27 lakhs tons (Fisheries statistics 2021). The production varied from 0.2 to 7.01 lakh tons and Gujarat state contributed the highest production (Fisheries statistics 2021). The Ichthyofauna diversity of the Gulf of Kachchh includes a total of 20 orders, 47 families and 96 species (Katira & Kardani 2017). Along the Sikka coast of Jamnagar where 112 ichthyofauna species belonging to 50 families, 12 orders, and 84 genera has been reported. Similarly, the locality of Jamnagar Marine National Park, Gulf of Kachchh reported 109 ichthyofauna species belonging to 58 families, 19 orders, and 93 genera (Brahmane et al. 2014). Apart from this, a recent study conducted by Sidat *et al.*, (2021) reported 96 species which include 20 order and 47 families. During the field observation, in the gill net catches *Mugil cephalus*, *Planiliza klunzingeri*, *Planiliza planiceps*, *Planiliza macrolepis* (Plate 9) were observed of which *Mugil cephalus* catch was the maximum during monsoon season of (20 kg) followed by mud crab (30 kg).



Plate 10: Fish and Crab catch along the Deendayal Port Authority in monsoon 2022

3.14. Marine Mammals

Sousa plumbea (Cuvier, 1829) is commonly referred to as the Indian Ocean humpback dolphin. During the field surveys, the Indian Ocean humpback dolphin (*Sousa plumbea*) was recorded at the site between the S-3 and S-4 opposite the oil jetty during monsoon season. The length of the humpback dolphin is approximately 1.7 to 2m. Humpback dolphins feed mostly on small fishes, sometimes shrimps; occur mostly in small groups (mostly 12 or less); have limited nearshore movements and in most parts of their range, exhibit a fission/fusion type of social organization. The evaluation of the conservation status of a species and its subsequent listing as a Threatened species is a function of its risk of extinction, which is influenced primarily by population dynamics (population size and trends, population structure) and the key biological and environmental factors influencing those dynamics (distribution, behaviour, life history, habitat use and the effects of human activities).



Plate 11. Indian Ocean humpback dolphin *Sousa plumbea*

4. Mud flat

Mudflats and mangroves establish a major ecosystem of the DPA coastal region and the significance of ecosystem services rendered by mudflat is endorsed in Coastal Regulation Zone (CRZ, 2011) as it accords special status to highly productive zone. Mudflat has an assemblage of plant-animal-geomorphological entities. DPA has been surrounded by two major ecosystems such as mangroves and mudflats which support a number of ecosystem services like nursery grounds for fish and shellfishes and breeding/feeding grounds for the birds (Spencer and Harvey, 2012). The TOC concentration is direct indicator of mudflat productivity and blue carbon sequestration.

Bulk density of the sediment samples

The data on the bulk density of the sediment samples are presented in (Fig.25). The bulk density of mangrove soil at Deendayal Port Authority coastal region ranged from 1.26 g/cm³ to 1.34 g/cm³. The highest bulk density was recorded at S-4 and S-12 sites followed by S-15. The lowest bulk density was recorded at site S-8 located at Tuna creek and S-1.

Total Organic Carbon (TOC)

The highest TOC value (0.83%) was recorded at station S-4 followed by S-2 site. Lowest TOC value was reported at site S-12 (Fig.26). It is observed that TOC values varied significantly among the sampling stations which means that organic carbon is dependent on the living life forms and variations in the life forms in the mudflats. The TOC concentration is a direct indicator of mudflat productivity and blue carbon sequestration. The data on monsoon samplings revealed that the different sampling sites of Deendayal Port Authority jurisdiction have considerable variations with respect to organic carbon.




Figure 25: Bulk density of mudflat sediment during Monsoon 2022




Figure 26: Percentage of Total Organic Carbon in the mudflat in Monsoon 2022

5. Avifauna

A large amount of research on bird diversity emphasizes the general negative effects of land conversion to human dominated habitats (Brooks *et al.* 1997; Castelletta *et al.* 2000). But human dominated and coastal habitats vary a lot and therefore the effect on birds can be very different. Birds depend on the habitats where they occurred, so the response of the species in particular habitat may always differ according to the habitat changes (Tworek, 2002, Winter & Faaborg, 1999; Cornelius *et al.* 2000; Zhanette 2000; Zhanette *et al.* 2000; Johnson & Igl, 2001; Beier *et al.* 2002; Herkert *et al.* 2003; Kurosawa & Askins, 2003). A total of 49 species belonging to six orders, 25 families and 38 genera were recorded from the coastal area of Deendayal Port during this study (Annexure 1). Among these, 26 species were aquatic and 23 species were terrestrial, which included three species listed as Near Threatened in the IUCN (2022), Red List.

Order Charadriiformes i.e. aquatic birds (including raptors and most water birds) constituted the predominant groups representing 58% of all species recorded from the study area followed by order Passeriformes (31%), i.e., perching birds (including babblers, drongos, mynas, sunbirds, doves, warblers, larks, chats, wagtails, robins). The families with a greater number of species were Ardeidae (eight spp.), Scolopacidae (seven spp.), Charadriidae (three spp.), Columbidae (three spp.), Laridae (two spp.), and Passeridae (one spp.). Among the recorded species, four were migrants, 10 were local migrants or resident migrants, 35 were breeding resident. During the present investigation, birds with diverse food habits viz., Aquatic (20 spp.), Insectivores (12 spp.), Granivores (eight spp.), Piscivores (six spp.), Omnivores (one spp.) Frugivores (one spp.), and Nectarivores (one spp.) were observed. The overall Shannon diversity (H') was 3.6 with species richness index for study area 1.2. The overall species evenness index value for study area was 0.77 and Equitability 0.93 (Table 13).

Status, distribution and diversity of avifauna in different stations:

A Total of fifteen sites were surveyed, of which the maximum number of species was found in Site 1 & 2 (33 spp.) followed by Site 9 (27 spp.) and Site 10 & 15 (26 spp.). Site 5 recorded the least richness (16 spp.) (Fig. 27).




Figure 27. Number of Avian species recorded from the Deendayal Port Area




Figure 28. Behavioral status of Avian species from the DPA in Monsoon 2022

Site wise migratory status showed that maximum migratory species were found in S- 2, S-9,S-13,S-14 & S-15 (three spp.) followed by S- 1,S-4,S-5,S-8 & S-11 (two spp.) (Fig. 28). From the study area all the species were categorized into two habitats i.e. terrestrial and aquatic. Survey for terrestrial and aquatic avifauna showed that maximum terrestrial avifaunal richness was recorded from S-2 (17 spp.) followed by site S-1 (15 spp.), S-11 (13 spp.) and site S-9 (12 spp.); while aquatic avifaunal species richness was more in site S-1 (18 spp.) followed by S- 15 (17 spp.), S-2 (16 spp.) and S- 8 (15 spp.) (Fig. 29).




Figure 29. Habitat wise distribution of Bird species from the DPA in Monsoon 2022

During the present investigation birds with diverse food habits were observed, viz., Aquatic, Insectivores, Granivores, Piscivores, Frugivores, Omnivores and Nectarivores. All the sites have found more number of aquatic birds species (maximum 16 species recorded from S- 15) followed by Insectivores (Maximum 8 species recorded from Site 1&2), granivore (maximum 8 species recorded from S-2) and piscivores (maximum 4 species recoded from S-3,S-6,S-8&S-11) and least species found of frugivores, omnivores and nectarivores (Fig.30)




Figure 30. Station wise Foraging Guild status of species recorded during Monsoon 2022

Data collected from point counts allows us to calculate species diversity, richness and species composition. The results showed that the maximum diversity was found from the S-1 ($H' 3.3$) followed by S-2 ($H' 3.2$) and the minimum diversity recorded from site 12 ($H' 2.6$) and S-5 ($H' 2.5$). The maximum species richness was recorded from Site 1 (2.9 spp.) and the minimum from Site 12 (2.1 spp.). These changes in individual species abundance, whether they occur independently of one another (Wiens, 1989) or are influenced by interactions with other bird species are governed by the degree of anthropogenic pressure including disturbance to habitat of species (Block & Brennan, 1993). The distribution and abundance of many bird species are mainly determined by the configuration and composition of the vegetation that comprises a major element of their habitat (Cody, 1985; Block & Brennan, 1993). As vegetation changes along complex geographical and environmental gradients, particular bird species may appear, increase in abundance, decrease, and disappear, when habitat becomes more or less suitable for its persistence. Totally 16% species were found rarely distributed in the study area while 36% species were very common. Aquatic and Insectivores species form the major groups while each of the frugivores, omnivores and nectarivores constitute about 2% of all species. Although more than 67% of the birds in the study area were Aquatic and insectivores, food competition was reduced by the

utilization of different habitat types and distinct feeding behaviour. Largely insectivorous birds like babblers (Sylviidae) and drongos (Corvidae) feed on fruits and seeds of plants particularly during winter season due to the shortage of insect food. Wetland birds were dominated largely by the aquatics followed by insectivore and grainivore species (Annexure 1). The present season study shows 49 different types of birds belonging to six orders and 25 families from the coastal area of Deendayal Port. The richness of avifauna is little low, indicator of ecological health of the coastal area of Deendayal Port. Proper and in-depth study, awareness, regarding the importance of birds and their role in ecosystem, to the local peoples through different massive programs will ultimately help the protection of birds of this region



Great Egret *Ardea alba*



Grey Heron *Ardea cinerea*



Great Cormorant *Phalacrocorax carbo*



Western Reef Egret *Egretta gularis*

Plate 12: Some common Birds from the Deendayal Port Authority

References

1. Abott, R.T. (1954). *American Sea shells*. Dvan Nostrand Company Inc, Newyork, pp 541.
2. Amr, Z.S. (2021) *The state of biodiversity in Kuwait*. Gland, Switzerland: IUCN; The State of Kuwait, Kuwait: Environmental Public Authority.
3. APHA (2017). *Standard Methods for the Examination of Water and Wastewater*, 23rd edition. American Public Health Association, 1546.
4. Apte DA (2012) *Field Guide to the Marine Life of India* - 1st Edition. Stusa Mudra Private Limited Mumbai, 502.
5. Apte, D.A. (2014) *Sea Shells of India. An Illustrated Guide to Common Gastropods*. Bombay Natural History Society & Oxford University Press, Mumbai, 197
6. Barnes RD (1980). *Invertebrate Zoology*, Saunders College, Philadelphia 108pp.
7. Brahmane, V.T., Temkar, G.S., Metar, S.Y., Sikotaria, K.M. & Desai, A.Y. (2014). Ichthyofaunal diversity in the vicinity of marine protected areas, Jamnagar, Gulf of Kachchh, India. *Asian Journal of Advanced Basic Science*, 3: 78–88.
8. Briggs, K.T., Tyler, W.B. & Lewis, D.B. (1985). Comparison of ship and aerial surveys of birds at sea. *Journal of Wildlife Management*, 49:405-411.
9. Chapgar, B.F. (1957). *Marine crabs of Bombay state*, Taraporevala Marine Biological Station, Bombay pp 88.
10. Crane, J. 1975. *Fiddler crabs of the world*. Ocypodidae: Genus Uca. Princeton University Press, Princeton, New Jersey.
11. Davidson, I.C., Crook, A.C., & Barnes, D.K.A. (2004). Quantifying Spatial Patterns of Intertidal Biodiversity: Is Movement Important?. *Marine Ecology*, 25 (1), 15–34.
12. Day, J.H. (1967). *A Monograph on the Polychaeta of Southern Africa part I Errantia*. Trustees of the British Museum (Natural History) London, 458pp.
13. Day, J.H. (1967). *A monograph on the Polychaeta of Southern Africa*. British Museum (Natural History). London. vol 1 & vol 2, 878pp.
14. Day, J.W., Hall. C.A.S., Kemp W.M. and Araneibia Y.A.C. (1989) *Estuarine Ecology*. John Wiley Sons, Inc.
15. De Bruin, G.H.P., Russell, B.C. & Bogush, A. (1995). *FAO species identification field guide for fishery purposes The Marine Fishery Resources of Sri Lanka*, Food and Agricul-

- tural Organization of the United Nations, Rome 110pp.
16. Desikachary, T.V. (1987). *Atlas of diatoms*, 3 and 25. Madras Science Foundation Madras: plates, 22-4000
 17. Dyer, K.R., Christie, M.C. & Wright, E.W. (2000). The classification of intertidal mudflats. *Continental Shelf Research*, 20(10-11): 1039-1060.
 18. Edward, J.K.P., Ravinesh, R. & Biju Kumar, A. 2022. *Molluscs of the Gulf of Mannar, India and Adjacent Waters: A Fully Illustrated Guide*, (Dekker, H. & Oliver, P.G. Eds.). Suganthi Devadason Marine Research Institute, Tuticorin & Department of Aquatic Biology & Fisheries, University of Kerala, India, 524pp.
 19. Fauvel, P. (1953). *The Fauna of India including Pakistan, Ceylon, Burma and Malaya. Annelida: Polychaeta*, Allahabad. 507pp.
 20. Fischer, W. & Bianchi, G. (1984). *FAO species identification sheets for fishery purposes Western Indian Ocean*, Fishing area 51 Prepared and prints with the support of the Danish International Development Agency DANIDA Rome, Food and Agricultural Organization of the United Nations, I-IV 20-55
 21. Gajbhiye, S. N. & Desai. B.N. (1981). Zooplankton variability in polluted and unpolluted waters off Bombay. *Mahasagar.*, 14(3): 173-182.
 22. Hammer, Ø., Harper, D.A.T. & Ryan, P.D. (2001). PAST: Paleontological statistics software package foreducation and data analysis. version 3.2.1.
 23. Hartman, O. (1969). *Atlas of the sedentariate polychaetous annelids from California*. Allan Hancock Foundation, University of Southern California. Los Angeles, 812.
 24. Hartman, O. (1968). *Atlas of the errantiate polychaetous annelids from California*. Allan Hancock Foundation, University of Southern California. Los Angeles, 828.
 25. Holthuis, L.B. (1993). *The Recent genera of the caridean and stenopodidean shrimps (Crustacea, Decapoda): With an appendix on the order Amphionidacea*. Nationaal Natuurhistorisch Museum Leiden. 328.
 26. Jha, B., C.R.K. Reddy, M.C. Thakur and M.U. Rao. (2009). *Seaweeds of India: the diversity and distribution of seaweeds of the Gujarat coast*. Springer, Dordrecht. 198
 27. Joshi, A., Parmar, E.A.R., Temkar, G.S., Desai, A.Y. & Bhatt, A.J. (2018). Ichthyofaunal biodiversity of Kharakuva Fish Market, Veraval, Gujarat, India. *International Journal of Bio-resource and Stress Management* 9: 596-605.

28. Kamboj, R.D., Salvi, H., Patel, R. & Bhagat, R. (2018) *Monograph on Phytolankton of Gulf of Kachchh*. Gujarat Ecological aeduction and Research (GEER) Foundation . 182
29. Klein, G.D. (1985). *Intertidal Flats and Intertidal Sand Bodies*, pp187-224. In: Davis, R.A. (eds) *Coastal Sedimentary Environments*. Springer, New York, NY.
30. Lyla, P.S., Velvizhi, S. & Khan, A.S. (1999). A Monograph on the amphipods of Parangipettai coast, Annamalai University, India pp78.
31. Mantri, V.A., Kavale, M.G & Mudassar A.K (2020) Seaweed Biodiversity of India: Reviewing Current Knowledge to Identify Gaps, Challenges, and Opportunities. Diversity. 1-22.
32. Masuda, H., Amaoka, K., Araka, C., Vyeno, T. & Yoshino T (1984). *The Fishes of Japanese Archipelago*. Tokai University Press, Japan 437.
33. McCann, S.B. (1980). *Classification of tidal environments*, In, McCann, SB Ed, *Sedimentary Processes and Animal Sediment Relationships in Tidal Environments*, Short Course Notes, Geological Association Canada, St Johns, Newfoundland, 1: 1-24.
34. Mohsin, A.K.M. & Ambiak, M.A. (1996). *Marine Fishes and Fisheries of Malaysia and Neighboring Countries*, University Pertanian Malaysia Press, Serdang 743.
35. Mohsin, A.K.M. & Ambiak, M.A. (1996). *Marine Fishes and Fisheries of Malaysia and Neighboring Countries*. University Pertanian Malaysia Press, Serdang 743.
36. Naderloo, R. (2017). *Atlas of Crabs of the Persian Gulf*. Springer International Publishing AG, Switzerland, 445pp.
37. Oliver, P.G. (1992) *Bivalved Seashells of the Red Sea*. National Museum of Wales, Cardiff, 330.
38. Oza, R.M., Krishnakumar, G.R., Mairh, O.P. & Zaidi, S.H. (2001) Cultivation of *Ulva fasciata* Delili on the coast of Diu, west coast of India. *Seaweed Resarch utilisation*, 23, 5–12
39. Psomadakis, P.N., Osmany, H.B. & Moazzam, M. (2015). *Field identification guide to the living marine resources of Pakistan*. FAO Species Identification Guide for Fishery Purposes, Rome, FAO. 386.
40. Ramakrishna and Dey (2007) *Hand book on Indian Freshwater Molluscs*. Published by the Director, Zoological Survey of India, Kolkata, 399pp.
41. Rao N.V.S. (1989) *Handbook of freshwater molluscs of India*. Zoological Survey of India,

Calcutta

42. Rao, N.V.S. (2003). *Indian Sea Shells (Part I). Polyplacophora and Gastropoda*. Zoological Survey of India, Kolkata, 416.
43. Rao, N.V.S. (2017) *Indian Seashells, Part B Bivalvia*. Zoological Survey of India, Kolkata, 676.
44. Ravinesh, R. and Biju Kumar, A. (2022) *Collection, preservation, and documentation of estuarine and marine benthic invertebrates*. pp 33-82. In: Prince S.G., Salom, G.T.V. and Krishnakumar, S. (Eds) *Ecology and Biodiversity of Benthos*, Elsevier Radarweg 29, PO Box 211, 1000 AE Amsterdam, Netherlands.
45. Ravinesh, R., and Biju Kumar A. (2013) Comparison of intertidal biodiversity associated with natural rocky shore and sea wall: A case study from the Kerala coast, India. *Indian Journal of Geo-Marine Sciences*, 42(2): 223-235.
46. Ravinesh, R., Biju Kumar, A. and Anjana, V.L (2021) Diversity and distribution of molluscan fauna of Asthamudi estuary, Kerala, India, *Wetlands Ecology and Management*. 29 (5), 745-765.
47. Robin S.W., Pat, H.A. & Glasby, C.J. (2003). *Polychaetes: An Interactive Identification Guide*. CSIRO Publishing, Melbourne.
48. Rouse, G.W. & Pleijel, F. (2001). *Polychaetes*. Oxford University Press: Oxford, UK, 354 pp.
49. Santhanam, P., Pachiappan, P., and Begum, A. (2019). *Methods of Collection, Preservation and Taxonomic Identification of Marine Phytoplankton*. pp25-61. In: Santhanam, P., Begum, A., Pachiappan, P. (eds) *Basic and Applied Phytoplankton Biology*. Springer, Singapore.
50. Sidat Azaz., Mukherji P., Trivedi T. & Mankodi P.C. (2021) Ichthyofauna species diversity of Gulf of Kachchh, Gujarat, India Case study: Jakhau and Mandvi coast. *Iranian Journal of Ichthyology*. 8(2): 134-150
51. Subba Rao N.V., Surya Rao, K.V. & Maitra, S. (1991). *Marine molluscs*. State Fauna Series 1, *Fauna of Orissa (Part 3)*. Zoological Survey of India, Calcutta, 1-175.
52. Terdalkar, S. & Pai. I.K. (2001) Statistical approaches for computing diversity of zooplankton in the Andaman Sea. *Tropical Ecology*, 42, 243-250.

53. van Franeker, J.A. (1994). A comparison of methods for counting seabirds at sea in the Southern Ocean. *Journal of Field Ornithology*, 65:96-108
54. Vine, P. (1986). *Red Sea Invertebrates*. Immel Publishing, London. 224 pp.
55. Walkley, A, & Black, I.A. (1934). An examination of the Degljareff method for determining soil organic matter and a proposed modification of the chromic acid titration method. *Soil Science*, 37: 29-38.
56. Xavier, J.C., Cherel, Y., Boxshall, G., Brandt, A., Coffey, T., Forman, J., Havermans, C., Jażdżewska, A.M., Kouwenberg, K., Schiaparelli, S., Schnabel, K., Siegel, V., Tarling, G.A., Thatje, S., Ward, P. & Gutt, J. (2020) *Crustacean guide for predator studies in the Southern Ocean*. Scientific Committee on Antarctic Research, Cambridge, UK. 253 .
57. Zar, J.H. (1984) *Biostatistical Analysis*. 2nd Edition, Prentice-Hall, Inc., Englewood Cliffs, 718.



Annexure -C

DEENDAYAL PORT TRUST

DETAILS OF MANGROVE PLANTATION ALREDY CARRIED OUT & Proposed To be Carried Out :

Sr. No	Name of the Organization	Total Mangrove Plantation carried out in Hectares till date and place of plantation and agency	Cost incurred
(A) <u>MANGROVE PLANTATION ALREDY CARRIED OUT</u>			
1	DEENDAYAL PORT TRUST (CRZ Recommendation 13 th to 16 th CB issued by the GCZMA) (Total 1000 ha.)	20 Hectares – 2005-06 Satsida Bet,Kandla, by GUIDE,Bhuj 50 Hectares – 2008-09 Nakti Creek,Kandla by Patel Construction 100 Hectares – 2010-11 Nakti Creek ,Kandla by GEC. (Board 29/1/2010) 200 Hectares – 2011-12 by Forest Department, GoG at Satsaida Bet 300 Hectares – 2012-13 by Forest Department, GoG at Satsaida Bet 330 Hectares – 2013-14 by Forest Department, GoG at Satsaida Bet TOTAL 1000 HA.	Rs. 8.8 lakhs Rs. 27.4 lakhs Rs.24.5 lakhs Rs. 66.5 lakhs Rs. 157.5 lakhs (total 630 hectares)
2	Creation of Berthing & allied Facilities off- tekra near Tuna (Outside Kandla Creek) – EC & CRZ Clearance. (Total 500 ha. – 250Ha. by DPT & 250 ha by Adani (concessionaire) MOU signed with GEC during Vibrant Gujarat Summit 2015 for 300 Ha.	300 Hectares – 2015-17 by GEC at Kantiyajal, Bharuch District	Rs. 90.0 lakhs
3.	EC & CRZ Clearance dated 19/12/2016 for Developing 7 integrated facilities (Condition 100 Ha)	100 Ha. –2018- 20 by GEC	Rs. 45 lakhs
TOTAL MANGROVE Plantation till date by DPT 1400 Ha. – Total 419.7 lakhs			

(B) Proposed Mangrove Plantation

1.	Development of Integrated facilities (Stage-II) within the existing Deendayal Port Trust (Erstwhile Kandla Port Trust) at District Kutch, Gujarat. (1. Setting up of Oil Jetty No.7 ; 2. Setting up of Barge jetty at Jafarwadi ; 3. Setting up of Barge port at Veera; 4. Administrative office building at Tuna Tekra; 5. Road connecting from Veera barge jetty to Tuna gate by M/s Deendayal Port Trust (Erstwhile : Kandla Port Trust) - <u>Environmental & CRZ Clearance accorded by the MoEF&CC,Gol dated 19/12/2020.</u>	<u>50 Ha. as per CRZ Recommendation issued by the GCZMA dated 29/6/2016.</u>	Rs. 45 lakhs
2.	Development of 3 Remaining Integrated Facilities (stage I) within the existing Deendayal Port Trust (Erstwhile : Kandla Port Trust) at Gandhidham, Kutch, Gujarat - <u>Environmental & CRZ Clearance accorded by the MoEF&CC,Gol dated 18/2/2020.</u>	<u>50 Ha. as per CRZ Recommendation issued by the GCZMA dated 29/6/2016. .</u>	

Annexure -D



DEENDAYAL PORT AUTHORITY

(Erstwhile Deendayal Port Trust)

Ministry of Ports, Shipping & Waterways, Govt. of India

Mech. Engg. Deptt.



Tel: (02836)220636 / 270184
FAX: (02836) 270184 / 270475
Email :- cmepdpt@gmail.com
cmepdpt@deendayalport.gov.in

Office of the Chief Mechanical Engineer,
Port & Customs Building,
New Kandla (Kutch), Gujarat-370210

No. DD/WK/3050/Pt-I/ Gm/PC-44

Date: 02.06.2022

Sir,

To,
Gujarat Ecology Commission
Forest & Environment Department
Block No. 18, First Floor, Udhog Bhavan,
Gandhinagar, Gujarat

Sub: Work Order to carry out Mangrove Plantation-reg.

The Competent Authority, Deendayal Port Authority has been pleased to approve:

1. To carry out mangrove plantation in 100 Ha. area with consultation of concern Gujarat Ecology Commission and at tentative estimated cost amounting to Rs. 50,00,000/- (excluding GST) for the said mangrove Plantation to be carried out in an area of 100 Ha. as per the stages mentioned by them in the MoU as follows:

Sr. No.	Terms and Condition	Rs. (in lakhs)
1	50% of the project cost of 100 Ha. Mangrove Plantation after signing the MoU.	Rs. 25.00
2	40% of the project cost of 100 Ha Mangrove Plantation after nursery preparation.	Rs. 20.00
3	10% of the project cost of 100 Ha Mangrove Plantation after plantation and submission of First year progress report.	Rs. 5.00
Total		50.00

2. To sign MoU with the Gujarat Ecology Commission, Government of Gujarat during the ensuing Vibrant Gujarat Summit 2022, regarding proposed Mangrove Plantation to be carried out in an area of 100 Hectares through the Gujarat Ecology Commission.
3. To authorize Dy. CME & CME (I/c) to sign MoU with the Gujarat Ecology Commission, Government of Gujarat during upcoming Vibrant Gujarat Summit 2022 for proposed Mangrove Plantation in an area of 100 Hectares through GEC.

The Expenditure shall be chargeable under Code 841/587/9744 WC-13001

Authority: Approved by Board vide Resolution No. 30 in the board meeting held on 27.05.2022

Chief Mechanical Engineer(I/c)
Deendayal Port Authority

Copy to:

- 1) SE(M)
- 2) A.O. (Works Audit)

Annexure -E

ENVIRONMENTAL MONITORING REPORT FOR DEENDAYAL PORT AUTHORITY



REPORT : DCPL/DPA/21-22/31

Mont : November 01

Issue : 00

Revision : 00

Prepare : DETOX CORPORATION PVT. LTD.,

Index		
Sr. No.	Name of Chapters	Page No.
	Executive Summary	3
A	Ambient Air	3
B	Weather	4
C	Marine Ecology (Flora and Fauna)	4
D	Drinking Water Quality	4
E	Monitoring Performance of Sewage Treatment Plant	5
F	Noise	5
1	Introduction- Deendayal Port Authority	7
2	Ambient Air Quality Monitoring	9
2.1	Ambient Air Quality Monitoring	11
2.2	Results	12
2.3	Observations and Conclusion	31
3	Meteorological Observations	32
3.1	Meteorological Data	33
4	Drinking Water Quality Monitoring	36
4.1	Drinking Water Monitoring Methodology	37
4.2	Results	37
4.3	Results & Discussion	45
4.4	Conclusions	48
5	Noise Monitoring	50
5.1	Method of Monitoring	50
5.2	Results & Discussion	50
5.3	Conclusions	51
6	Soil Monitoring	53
6.1	Methodology	53
6.2	Results	54
6.3	Discussion	55
6.4	Conclusion	55
7	Sewage Treatment Plant Monitoring	57
7.1	Methodology for STP Monitoring	57
7.2	Results	58
7.3	Conclusions	65
8	Marine Water Monitoring	67
8.1	Marine Water Quality and Results	68
8.2	Results and Discussion of Marine water samples	77
8.3	Conclusions	79
9	Marine Sediment Monitoring	81
9.1	Results	82
9.2	Discussion of Marine Sediment samples	84
9.3	Conclusions	84
10	Marine Ecological Monitoring	85
10.1	Introduction	86
10.2	Results	95
11	Conclusive Summary & Remedial Measures	136
12	References	142



EXECUTIVE SUMMARY

ENVIRONMENTAL MONITORING PLAN FOR DEENDAYAL PORT ENVIRONMENTAL MONITORING REPORT- NOVEMBER, 2022

1. EXECUTIVE SUMMARY

Monitoring of various environmental aspects of the Deendayal port by M/s Detox Corporation Pvt. Ltd. has been carried out through collection of samples, analysis of the same, comparing results with respect to the national standards and any other relevant standards by GBCB/CPCB/MoEF & CC to understand status of various parameters in the Environment of the Deendayal Port. The results shall address the identified impacts and suggest measures to minimize the environmental impact due to various operations at Deendayal Port.

A) Ambient Air

The monitoring of Ambient Air quality at 6-locations at Deendayal Port Authority Kandla and 2- location at Vadinar Port on 24 hourly basis for TSPM, PM₁₀, PM_{2.5}, SO₂, NO₂, NH₃, CO₂, CO, C₆H₆ and NMHC in twice a week 24 hourly at uniform intervals (as per NAAQS) at Gopalpuri, Tuna Port, Marine Bhavan Building, Coal storage area, Estate building, Oil jetty and at Vadinar port, Vadinar Jetty and Vadinar colony area using respirable dust sampler, Fine particulate sampler and gaseous sampler.

The Maximum TSPM values in month of November 2022 were found 846 µg/m³ at Coal Storage area on 25.11.2022 and minimum 107 µg/m³ at Gopalpuri Hospital on 01.11.2022. The Maximum PM₁₀ values were 654 µg/m³ at Coal Storage area on 25.11.2022 and minimum was 67 µg/m³ at Gopalpuri Hospital 01.11.2022. Maximum PM_{2.5} values were 187 µg/m³ at Coal Storage area on 25.11.2022 and minimum was 34 µg/m³ at Gopalpuri on 01.11. 2022. The PM₁₀ and PM_{2.5} values were found for all monitoring locations (Marine Bhavan Building, Oil Jetty, Estate Office, Gopalpuri, Coal Storage Area and Tuna Port) to exceed the Standard limit (NAAQS).

At Gopalpuri location the mean concentration of PM₁₀ was 127 µg/m³ & PM_{2.5} was 66 µg/m³ which are slightly exceed the Standard limit (NAAQS).

The AAQ monitoring for Vadinar at Admin building the mean TSPM, PM₁₀ and PM_{2.5} were 237 µg/m³, 138 µg/m³ and 97 µg/m³ respectively which was exceed the Standard limit (NAAQS) the while at Signal Building the mean TSPM, PM₁₀ and PM_{2.5} were 113 µg/m³, 74 µg/m³ and 38 µg/m³ respectively slightly exceed the Standard limit (NAAQS).

The overall values of November for Gaseous SO₂, NO₂, NH₃, CO₂, CO, C₆H₆ concentration were within the permissible limit at all location and NMHC were found BQL (Below Quantification Limit).

B) Weather

The mean day time temperature at Deendayal Port was 27.92 °C. The day-time maximum temperature was 32.9°C and minimum was 21.1 °C. The mean night time temperature recorded was 25.47 °C. The night-time maximum temperature was 29.7°C and minimum was 20.0 °C. The mean Solar Radiation in November month was 167.27 w/m². The maximum solar radiation was recorded 759 w/m² in 4th November, 2022 and the minimum solar radiation was recorded 1.80 w/m² in 30th November, 2022. The mean Relative humidity was 69.00 % for the month of November. Maximum Relative humidity was recorded 99.0 % and minimum Relative humidity was recorded 34.0 %. The average wind velocity for the entire month of November was 1.21 m/s. Maximum wind velocity was recorded 10.19 m/s. The wind direction was mostly West-South.

C) Marine Ecology (Flora and Fauna) / Marine Water / Sediments:

The results obtained from the study for the month of November 2022 for biological and ecological parameters in marine water for Arabian Sea at surrounding area of Deendayal Port Authority (DPA) Kandla and Vadinar were not affected by Port activities.

D) Drinking Water Quality

The drinking water being supplied to Deendayal Port Authority was safe for drinking purpose. At all drinking water monitoring stations around port area were in line with the standard limit as per the drinking water specifications given in IS 10500:2012 as per tested parameters only.

The average results for 20 locations were as: pH were found Min 7.24 and maximum 7.52, TDS were found min 300.0 mg/l and Max found 1060.0 mg/l, Chloride were found Min 140.31 mg/l and Max 576.28 mg/l, Total Hardness were found Min 270.0 mg/l and Max 380.0 mg/l and Calcium were found Min 34.47 mg/l and Max 43.29 mg/l, color were colorless and odor were odorless. In all water samples BOD, Heavy metal like manganese, Hexavalent chromium, Copper, Cadmium, Arsenic, Mercury, Lead, zinc all are found BQL (Below Quantification Limit). The bacterial count (E-coli & Coliform) is absent in all drinking water samples.

E) Monitoring Performance of Sewage Treatment Plant

It was seen that the performance of STP at Deendayal Township Gopalpuri, DPA STP Plant Kandla and Vadinar STP plant was satisfactory by overall. The treatment plant was well maintained during [November 2022] with considerable removal efficiency achieving the standards prescribed for final disposal. At Gopalpuri STP, the pollutant removal efficiency for TSS, BOD and COD was ranged from 49.66-81.04%, 58.97-68.42% and 45.45-73.33% respectively. At Kandla STP, removal efficiency for TSS, BOD and COD was ranged from 53.47-73.49%, 46.15-76.74% and 50.00-82.35% respectively & at Vadinar STP removal efficiency for TSS, BOD and COD was ranged from 42.09-56.69%, 50.00-78.12% and 60.00-84.61% respectively. At all STP location treated waste water the pH were ranged from 7.21-7.42, Total Suspended Solids were found 16.9-67.9 mg/l, Residual Chlorine were below Detection Limit (< 0.5), COD were found 20-60 mg/l and 3day BOD @ 27 °C were found 7.0-16.0 mg/l.

F) Noise

Noise sources in port operations include cargo handling, vehicular traffic, and loading / unloading containers and ships. The Day Time Noise Level (SPL) in all 10 locations at Deendayal Port Authority ranged from 53.2 dB(A) to 70.4 dB(A) while at Vadinar port 3 location ranged from 52.5 dB(A) to 60.6 dB(A) which was within the permissible limits of 75 dB(A) for the industrial area for the daytime. The Night Time Average Noise Level (SPL) in all locations of Deendayal Port Authority ranged from 45.4 dB to 61.7 dB(A) while at Vadinar port ranged from 52.5 dB (A) to 60.6 dB(A) which was within the permissible limits of 70 dB(A) for the industrial area for the night time.



CHAPTER-1

INTRODUCTION

DEENDAYAL PORT AUTHORITY

1.0 Introduction

About Deendayal Port

The Deendayal Port is situated in the Kandla Creek and is 90 Kms. From the mouth of Gulf of Kachchh. Latitude: 23° 01" N Longitude: 70° 13"E. Deendayal Port's journey began in 1931 with construction of RCC Jetty by Maharao Khengarji. After partition, Deendayal Port's success story has continued and it rise to the No. 1 Port in India in the year 2007-08 and since then retained the position for the 15 consecutive year. On 31.03.2016, Deendayal Port created history by handling 100 MMT cargoes in a year, the first Major Port to achieve the milestone. Kandla, also known as the Deendayal Port Authority is a seaport in Kutch District of Gujarat state in western India, near the city of Gandhidham. Located on the Gulf of Kutch, it is one of major ports on west coast. Kandla was constructed in the 1950s as the chief seaport serving western India, after the partition of India from Pakistan left the port of Karachi in Pakistan. The Port of Deendayal is located on the Gulf of Kutch on the northwestern coast of India some 256 nautical miles North West of the Port of Karachi in Pakistan and over 430 nautical miles north-northwest of the Port of Mumbai (Bombay). It is the largest port of India by volume of cargo handled. Kandla history Deendayal Port Authority, India's busiest major port in recent years, is gearing to add substantial cargo handling capacity with private sector participation. Deendayal port Authority creates a new record by handling 127.10 million metric tons of cargo during the FY 2021-22, as against 117.566 million metric tons in FY 2020-21. Showing a growth of 8.11 %. Incidentally, DPA is the only major Indian port of handle more than 127 MMT cargo throughout and it has also registered the highest cargo throughput in its history. While the port has flagged off several projects related to infrastructure creation, DPA has successfully awarded the work of augmentation of liquid cargo handling capacity by revamping the existing pipeline network at the oil jetty area in Sept. 2021. Even as much of this growth has come from handling of crude oil imports, mainly for Essar Oil's Vadinar refinery in Gujarat, the port is also taking measures to boost non-POL cargo. Last fiscal, POL traffic accounted for 63 per cent of the total cargo handled at Deendayal Port, as against 59% in 2007-08. The Deendayal Port Authority had commissioned the Off-shore Oil Terminal facilities at Vadinar in the year 1978, for which M/s. Indian Oil Corporation Limited (IOCL) provided Single Bouy Mooring (SBM) system, having a capacity of 54 MMTPA, which was first of its kind in India. Further, significant. Quantum of infrastructural up-gradation has been affected & excellent maritime infrastructure been created at Vadinar for the 32 MMTPA Essar Oil Refinery in Jamnagar District. Monitoring of various environmental aspects of the Deendayal port by M/s Detox Corporation Pvt. Ltd. has been carried out through collection of samples, analysis of the same, comparing results with respect to the prescribed standards by GPCB/CPCB/MoEF& CC. The results shall address the identified impacts and suggest measures to minimize the environmental impact due to various operations at Deendayal Port. The environmental monitoring is carried out as per the Environment Management and Monitoring Plan submitted by Detox Corporation Pvt. Ltd.

CHAPTER-2

AMBIENT AIR QUALITY MONITORING

2. Introduction

Air pollutants are added in the atmosphere from variety of sources that change the composition of atmosphere and affect the biotic environment. The concentration of air pollutants depend not only on the quantities that are emitted from air pollution sources but also on the ability of the atmosphere to either absorb or disperse these emissions. The air pollution concentration vary spatially and temporarily causing the air pollution pattern to change with different locations and time due to changes in meteorological and topographical condition. Air pollution occurs when harmful substances including particulates and biological molecules are introduced into earth's atmosphere. It may cause diseases, allergies or death of humans; it may also cause harm to other living organisms such as animals and food crops, and may damage the natural or built environment. Human activity and natural processes can both generate air pollution. A physical, biological or chemical alteration to the air in the atmosphere can be termed as pollution. It occurs when any harmful gases, dust, smoke enters into the atmosphere and makes it difficult for plants, animals and humans to survive as the air becomes dirty. The consequences of industrialization and the demand for improved quality of life has been increased exposure to air pollution (Vallero, 2014). An air pollutant is a substance in the air that can have adverse effects on humans and the ecosystem. The substance can be solid particles, liquid droplets, or gases. A pollutant can be of natural origin or man-made. Pollutants are classified as primary or secondary. Any gas could qualify as pollution if it reached a high enough concentration to do harm. Theoretically, that means there are dozens of different pollution gases. In practice, about ten different substances cause most concern. Heavy metals represent a class of omnipresent pollutants, with toxic potential, in some cases even at low exposure levels. They concentrate in each tropic level because of their weak mobility, so the concentration in plants is higher than in soil, in herbivore animals higher than in plants, in carnivores' tissues higher than in herbivore, the highest concentration being reached at the end of the tropic chain, at big predacious and human bodies.

Globally, one of the main contributors to emissions of atmospheric pollutants and a significant user of energy is the industrial sector (Conti et al. 2015).

The concentration of air pollutants depends not only on the quantities that are emitted from the polluting sources, but also on the ability of the atmosphere to either absorb or disperse such emissions (USEPA, 2008).

Nowadays, the shipping sector provides low-cost and reliable delivery services in the economic field (Arunachalam et al. 2015). Nevertheless, shipping-related activities have a considerable impact on air pollution, especially in coastal areas but also globally (Buccolieri et al. 2016). The primary air pollutants are PM, VOCs, NO_x, O₃, SO₂, and CO (Bailey and Solomon 2004). As a consequence, a wide range of options toward “greener” seaports is needed (Bailey and Solomon 2004). Some of these measures are easy to adopt such as the regulation of fuel quality (by using low-sulfur alternative fuels), the speed reduction (Lack et al. 2011), and the use of alternative transportation equipment (Lai et al. 2011).

Clean air is the basic requirement of all living organisms. In recent times, due to population growth, urban sprawl, industrial development, and vehicular boom, the quality of air is deteriorating and being polluted. Pollutants of major public health concerns include particulate matter, carbon monoxide, ozone, nitrogen dioxide, and sulfur dioxide, which pose serious threats to human health and hygiene. In the present study, prime particulate pollutants (PM₁₀, PM_{2.5}), and gaseous pollutants (SO₂, and NO₂) were estimated at seven stations in and around Dahej Port, Gujarat, India (Soni and Jagruti Patel, 2017).

Among particulate pollutants, particulate matter (PM) is a ubiquitous entity, and is especially a grave problem due to its higher suspension rate into the atmosphere, and adverse health effects on plants, animals, humans, and materials in the form of visibility reduction, soiling of buildings, etc. (Horaginamani and Ravichandran, 2010; Chaurasia *et al.*, 2013).

The sources of air pollutants include vehicles, industries, domestic sources and natural sources. Because of the presence of high amount of air pollutants in the ambient air, the health of the population and property is getting adversely affected. In order to arrest the deterioration in air quality, Govt. of India has enacted Air (Prevention and Control of Pollution) Act in 1981. The responsibility has been further emphasized under Environment (Protection) Act, 1986. It is necessary to assess the present and anticipated air pollution through continuous air quality survey/monitoring programs. Therefore, Central Pollution Control Board had started National Ambient Air Quality Monitoring (NAAQM) Network during 1984 - 85 at national level. The programme was later renamed as National Air Quality Monitoring Programme (NAMP).

2.1 Ambient Air Quality Monitoring

As per the Environmental Monitoring Plan of Deendayal Port Authority, Air monitoring was carried out at six identified locations at Deendayal Port and two locations at Vadinar Port.

Table: 1. Ambient Air Sampling Location

Sr. No.	Name of Location	Location Code	Latitude	Longitude	Remarks
1.	Marine Bhavan	AL-1	23° 0' 26.524"N	70° 13' 22.414"E	DPA-Kandla
2.	Oil Jetty	AL-2	23° 1' 45.613"N	70° 13' 11.052"E	
3.	Estate Office	AL-3	23° 1' 11.273"N	70° 12' 48.657"E	
4.	Gopalpuri Hospital	AL-4	23° 4' 53.551"N	70° 8' 7.047"E	
5.	Coal Storage Area	AL-5	22° 59' 31.812"N	70° 13' 9.979"E	
6.	Tuna Port	AL-6	22° 59' 15.291"N	70° 58' 57.018"E	
7.	Signal Building	AL-7	22° 26' 26.750"N	69° 40' 22.127"E	DPA-Vadinar
8.	Admin Building	AL-8	22° 26' 25.223"N	69° 40' 19.358"E	

● Air Quality Monitoring Methodology

Air quality is measured in all the stations, for 24 hour for Total Suspended Particulate Matter (TSPM), PM₁₀, PM_{2.5}, SO₂, NO₂, NH₃ & Benzene and Grab-sampling for CO & CO₂ measurements. The Air samplers are operated for a period of 24 hours and after a continuous operation of 8 hours for gaseous parameters. The absorbing reagents for SO₂:- Absorbing Reagent TCM (Potassium Tetrachloromercurate 0.04M): Mercuric Chloride, Potassium Chloride and EDTA used. For NO₂:- Absorbing Reagent Sodium Hydroxide (NaOH): Sodium Hydroxide and Sodium Arsenite used. For NH₃ need Conc. Sulphuric Acid and Distilled water was used. By replacing 3 times the reagents per day for each parameter namely, SO₂, NO₂, NH₃. The GFA filter paper and PTFE Membrane bound filter paper are used for a period of 24 hours to obtain one sample each of TSPM, PM₁₀ & PM_{2.5}. The AAQ samples are collected two consecutive days a week as per CPCB guidelines, from all the eight locations as mentioned in the EMP.

2.2 Results

The ambient air quality monitoring data for six stations, viz. Marine Bhavan, Oil Jetty, Port Colony, Gopalpuri Hospital, Tuna Port and Nr. Coal Storage Area for the month of November 2022 are given in Tables 2 to 7. The ambient air quality monitoring data for two stations at Vadinar (Nr. Admin Building & Nr. Signal Building) are given in Tables 8 to 9.

The Movement of heavy transport with uncovered coal transportation, raw road around ambient location may be causes fugitive dust emission from dry conditions. Particulate Matter then enters the atmosphere through the action of wind, vehicular movement, or other activities. The dust produces tends to float in air and spread all around the vicinity. Direction and speed of wind affect the dispersion of the dust particulate matter. Humidity of air also has strong effect on the spreading of particulate matter. With increasing humidity, moisture particles eventually grow in size to a point where 'dry deposition' occurs, reducing PM₁₀ concentrations in the atmosphere.

Location 1: Marine Bhavan (AL1)

Table 2 : Results of Air Pollutant Concentration at Marine Bhavan

	Date	TSPM [µg/m3]	PM10 [µg/m3]	PM2.5 [µg/m3]	SO2 [µg/m3]		NOx [µg/m3]		NH3 [µg/m3]	
Sampling Period		24hr	24hr	24hr	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
NAAQMS Limit			100 µg/m3	60 µg/m3		80 µg/m3		80 µg/m3		400 µg/m3
AL1 – 1	01.11.2022	435	302	121	3.93	3.93	5.19	14.43	2.07	4.11
					6.04		23.66		6.33	
					1.81		14.43		3.91	
AL1 – 2	04.11.2022	344	228	106	3.32	2.52	17.31	12.70	2.42	3.72
					2.72		8.66		5.18	
					1.51		12.12		3.57	
AL1 – 3	08.11.2022	398	281	116	2.31	3.84	25.39	17.31	4.72	3.57
					6.34		17.89		2.42	
					2.88		8.66		3.57	
AL1 – 4	11.11.2022	445	315	124	3.63	6.35	17.89	13.08	4.03	3.61
					9.07		12.70		4.72	
					6.35		8.66		2.07	
AL1 – 5	15.11.2022	364	253	110	4.53	4.53	11.54	13.85	4.60	3.07
					6.35		19.62		2.88	
					2.72		10.39		1.73	
AL1 - 6	18.11.2022	442	315	121	8.46	4.84	23.08	16.54	3.22	4.37
					3.32		8.66		5.87	
					2.72		17.89		4.03	
AL1 - 7	22.11.2022	375	266	106	3.32	4.43	17.89	18.47	4.83	4.45
					7.55		25.97		5.87	
					2.42		11.54		2.65	
AL1 – 8	25.11.2022	483	350	129	4.53	4.63	23.66	21.55	3.22	3.68
					6.95		28.86		5.29	
					2.42		12.12		2.53	
AL1 – 9	29.11.2022	534	383	142	6.35	5.84	17.89	19.04	3.57	3.57
					8.46		25.97		4.95	
					2.72		13.27		2.19	
Monthly Average		424	299	119		4.55		16.33		3.79
Standard Deviation		61	48	12		1.12		3.03		0.44

Table 2 : Results of Air Pollutant Concentration at Marine Bhavan

	Date	C6H6 [µg/m ³]	HC	CO [mg/m ³]	CO ₂ [ppm]
Sampling Period		8 hr		Grab Sampling	Grab Sampling
NAAQMS limit		5.0 µg/m ³	ppm	4.0 mg/m ³	-
AL1 – 1	01.11.2022	1.09	BQL	1.44	444
AL1 – 2	04.11.2022	1.2	BQL	1.54	374
AL1 – 3	08.11.2022	1.17	BQL	1.08	538
AL1 – 4	11.11.2022	1.1	BQL	1.14	470
AL1 – 5	15.11.2022	1.11	BQL	1.26	481
AL1 - 6	18.11.2022	1.1	BQL	1.64	500
AL1 - 7	22.11.2022	1.12	BQL	1.35	620
AL1 - 8	25.11.2022	1.16	BQL	1.69	511
AL1 - 9	29.11.2022	1.21	BQL	1.16	522
Monthly Average		1.14	-	1.37	495.56
Standard Deviation		0.05	-	0.22	67.59

* NMHC- Non- Methane Hydrocarbons

BQL- Below Quantification Limit (Quantification Limit – NMHC: 0.5 ppm)

At Marine Bhavan, the overall values of TSPM, PM₁₀, PM_{2.5}, SO₂, NO₂ and NH₃ is attributed mainly by motor vehicle emission produced from various types of automobiles (both diesel and petrol driven). Moreover, the loading and unloading of Food Grains and Timber at Jetty no. 1 and 2 also contributes to the high levels of TSPM and PM₁₀. The mean TSPM value at Marine Bhavan was 424 µg/m³, the mean PM₁₀ value was 299 µg/m³, and PM_{2.5} value was 119 µg/m³ which is above the permissible limit prescribed by NAAQS. The average values of SO₂, NO₂ and NH₃ were 4.55 µg/m³, 16.33 µg/m³ & 3.79 µg/m³ respectively; these values were within the standard limit prescribed by NAAQS.

The levels of Benzene, Hydrocarbons (HC) and CO were within the permissible limit at Marine Bhavan. The mean Benzene concentration was 1.14 µg/m³, well below the permissible limit of 5.0 µg/m³. NMHC's were below the detectable limit and Carbon Monoxide concentration was 1.37 mg/m³, well below the permissible limit of 4.0 mg/m³ prescribed by NAAQS.

Location 3: Oil Jetty (AL2)

Table 2 : Results of Air Pollutant Concentration at Oil Jetty

Table 2 : Results of Air Pollutant Concentration at Oil Jetty										
	Date	TSPM [µg/m3]	PM10 [µg/m3]	PM2.5 [µg/m3]	SO2 [µg/m3]		NOx [µg/m3]		NH3 [µg/m3]	
Sampling Period		24hr	24hr	24hr	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
NAAQMS Limit			100 µg/m3	60 µg/m3		80 µg/m3		80 µg/m3		400 µg/m3
AL2 -1	01.11.2022	150	99	50	2.42	3.22	6.35	13.66	2.88	4.53
					4.53		13.27		6.79	
					2.72		21.35		3.91	
AL2 -2	04.11.2022	253	180	70	2.72	3.53	5.77	11.73	0.81	3.18
					3.32		17.89		4.03	
					4.53		11.54		4.72	
AL2 -3	08.11.2022	235	166	67	2.59	2.50	5.19	14.04	2.19	2.80
					3.46		13.27		2.65	
					1.44		23.66		3.57	
AL2 -4	11.11.2022	275	194	76	6.35	4.53	10.39	14.24	2.42	2.42
					4.53		20.20		3.80	
					2.72		12.12		1.04	
AL2 – 5	15.11.2022	245	169	71	3.02	4.53	8.66	14.04	3.57	2.38
					6.65		16.16		2.30	
					3.93		17.31		1.27	
AL2 – 6	18.11.2022	185	119	53	5.74	4.94	14.43	13.47	4.95	3.84
					2.72		17.31		3.57	
					6.35		8.66		2.99	
AL2 – 7	22.11.2022	373	252	109	3.02	4.03	20.20	14.24	3.80	3.80
					6.35		12.12		5.53	
					2.72		10.39		2.07	
AL2 -8	25.11.2022	292	199	86	1.81	3.83	14.43	14.43	3.57	4.76
					6.35		19.62		4.72	
					3.32		9.23		5.99	
AL1 – 9	29.11.2022	299	194	97	3.63	4.63	5.19	13.47	2.88	3.49
					7.55		23.66		4.95	
					2.72		11.54		2.65	
Monthly Average		256	175	75		3.97		13.70		3.47
Standard Deviation		65	45	19		0.79		0.81		0.85

Table 3 : Results of Air Pollutant Concentration at Oil Jetty					
	Date	C₆H₆ [µg/m³]	*NMHC	CO [mg/m³]	CO₂ [ppm]
Sampling Period		8 hr		Grab Sampling	Grab Sampling
NAAQMS limit		5.0 µg/m³		4.0 mg/m³	-
AL2-1	01.11.2022	1.17	BQL	1.22	467
AL2-2	04.11.2022	1.01	BQL	1.53	451
AL2-3	08.11.2022	1.1	BQL	1.65	502
AL2-4	11.11.2022	1.19	BQL	1.04	447
AL2 -5	15.11.2022	1.24	BQL	1.27	634
AL2 -6	18.11.2022	1.16	BQL	1.22	531
AL2-7	22.11.2022	1.2	BQL	1.28	800
AL2-8	25.11.2022	1.06	BQL	1.89	1023
AL2-9	29.11.2022	1.22	BQL	1.46	576
Monthly Average		1.15	-	1.40	603.44
Standard Deviation		0.08	-	0.26	193.07

* NMHC- Non- Methane Hydrocarbons

BQL- Below Quantification Limit (Quantification Limit – NMHC: 0.5 ppm)

Oil Jetty Area, the overall values of TSPM, PM₁₀, PM_{2.5}, SO₂, NO₂ and NH₃ was mainly by motor vehicle emission produced from various types of vehicles at Oil Jetty Area. The mean TSPM value at Oil Jetty was 256 µg/m³. The mean PM₁₀ value was 175 µg/m³ and mean PM_{2.5} value was 75 µg/m³ which was above the permissible limit. The average values of SO₂, NO₂ and NH₃ were within the permissible limit prescribed by NAAQS. The mean concentration of SO₂, NO₂ and NH₃ were 3.97 µg/m³, 13.70 µg/m³ and 3.47 µg/m³ respectively.

The levels of Benzene, Hydrocarbons (HC) and CO were within the permissible limit at Oil Jetty. The mean Benzene concentration was 1.15 µg/m³ which was well below the permissible limit of 5.0 µg/m³. NMHC's were below the detectable limit and Carbon Monoxide concentration was 1.40 mg/m³, well below the permissible limit of 4.0 mg/m³.

Location 3: Kandla Colony – Estate Office (AL-3)

Table 4 : Results of Air Pollutant Concentration at Estate Office

Table 4 : Results of Air Pollutant Concentration at Estate Office										
	Date	TSPM [µg/m3]	PM10 [µg/m3]	PM2.5 [µg/m3]	SO2 [µg/m3]		NOx [µg/m3]		NH3 [µg/m3]	
Sampling Period		24hr	24hr	24hr	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
NAAQMS Limit			100 µg/m3	60 µg/m3		80 µg/m3		80 µg/m3		400 µg/m3
AL3 – 1	01.11.2022	245	172	69	1.51	2.32	10.39	9.62	3.68	5.10
					3.32		13.27		7.02	
					2.12		5.19		4.60	
AL3 – 2	04.11.2022	577	445	130	4.53	2.32	5.19	10.39	3.57	2.49
					1.51		17.31		2.88	
					0.91		8.66		1.04	
AL3 – 3	08.11.2022	440	321	109	6.05	3.94	19.04	12.31	4.72	3.64
					2.59		12.12		2.42	
					3.17		5.77		3.80	
AL3 – 4	11.11.2022	518	403	111	3.32	4.23	18.47	10.58	1.38	2.42
					2.72		8.66		3.57	
					6.65		4.62		2.30	
AL3 – 5	15.11.2022	451	340	107	1.81	3.73	23.08	15.97	3.22	2.42
					6.04		14.43		2.30	
					3.32		10.39		1.73	
AL3 – 6	18.11.2022	459	346	112	4.53	4.43	16.16	15.97	5.76	4.14
					2.72		8.66		4.72	
					6.04		23.08		1.96	
AL3 – 7	22.11.2022	453	325	116	2.42	4.33	19.62	17.31	3.91	3.84
					4.23		23.66		5.18	
					6.35		8.66		2.42	
AL3 – 8	25.11.2022	337	252	83	6.04	3.93	15.00	15.58	3.80	3.91
					3.32		23.08		5.76	
					2.42		8.66		2.19	
AL1 – 9	29.11.2022	491	359	129	4.84	4.63	17.89	16.16	3.57	3.57
					6.95		24.24		5.18	
					2.12		6.35		1.96	
Monthly Average		441	329	107		3.76		13.77		3.50
Standard Deviation		98	80	20		0.87		3.00		0.91

Table 4 : Results of Air Pollutant Concentration at Estate Office

Sampling Period	Date	C ₆ H ₆ [µg/m ³]	*NMHC	CO [mg/m ³]	CO ₂ [ppm]
		8 hr		Grab Sampling	Grab Sampling
		5.0 µg/m ³		4.0 mg/m ³	-
NAAQMS limit					
AL3 -1	01.11.2022	1.06	BQL	1.27	508
AL3 -2	04.11.2022	1.1	BQL	1.19	508
AL3 -3	08.11.2022	1.1	BQL	1.65	502
AL3 -4	11.11.2022	1.09	BQL	1.83	429
AL3 - 5	15.11.2022	1.09	BQL	1.76	813
AL3 - 6	18.11.2022	1.2	BQL	1.14	559
AL3 - 7	22.11.2022	1.19	BQL	2.18	1022
AL3 - 8	25.11.2022	1.11	BQL	2	1026
	29.11.2022	1.06	BQL	1.22	537
Monthly Average		1.11	-	1.58	656.00
Standard Deviation		0.05	-	0.39	234.02

* NMHC- Non- Methane Hydrocarbons

BQL- Below Quantification Limit (Quantification Limit – NMHC: 0.5 ppm)

The overall values of TSPM, PM₁₀, PM_{2.5}, SO₂, NO₂ and NH₃ at Kandla Port Colony (Estate Office) was attributed by vehicle emission produced from trucks and heavy duty vehicles that pass through the road outside Kandla Port Colony. The mean TSPM values at Estate Office were 441 µg/m³, the mean PM₁₀ value was 329 µg/m³, and PM_{2.5} value was 107 µg/m³ which was above the permissible limit prescribed by NAAQS. The average values of SO₂, NO₂ and NH₃ were 3.76 µg/m³, 13.77 µg/m³ and 3.50 µg/m³ respectively and were all within the permissible limit.

The levels of Benzene, Hydrocarbons (HC) and CO were within the permissible limit at Kandla Port Colony. The mean Benzene concentration was 1.11 µg/m³, well below the permissible limit of 5.0 µg/m³. NMHC's were below the detectable limit and Carbon Monoxide was 1.58 mg/m³, well below the permissible limit of 4.0 mg/m³.

Location 4: Gopalpuri Hospital (AL-4)

Table 5 : Results of Air Pollutant Concentration at Gopalpuri Hospital

Table 5 : Results of Air Pollutant Concentration at Gopalpuri Hospital										
	Date	TSPM [µg/m3]	PM10 [µg/m3]	PM2.5 [µg/m3]	SO2 [µg/m3]		NOx [µg/m3]		NH3 [µg/m3]	
Sampling Period		24hr	24hr	24hr	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
NAAQMS Limit			100 µg/m3	60 µg/m3		80 µg/m3		80 µg/m3		400 µg/m3
AL4 -1	01.11.2022	107	67	34	1.21	2.22	5.77	6.93	2.42	2.53
					3.02		10.39		4.14	
					2.42		4.62		1.04	
AL4 -2	04.11.2022	177	117	54	0.91	2.22	5.19	10.00	1.61	2.49
					4.53		8.66		2.42	
					1.21		16.16		3.45	
AL4 -3	08.11.2022	148	101	44	1.15	2.21	6.93	9.81	1.73	1.69
					2.88		17.31		2.42	
					2.59		5.19		0.92	
AL4 -4	11.11.2022	184	111	68	1.51	2.62	6.93	12.89	1.04	2.30
					3.63		14.43		2.42	
					2.72		17.31		3.45	
AL4 – 5	15.11.2022	202	125	72	2.12	2.42	12.12	12.70	2.42	2.49
					3.63		8.66		3.45	
					1.51		17.31		1.61	
AL4 – 6	18.11.2022	233	153	78	1.21	2.92	8.66	12.89	2.42	2.49
					4.84		17.89		1.61	
					2.72		12.12		3.45	
AL4 – 7	22.11.2022	268	168	94	0.60	2.22	5.77	12.70	1.73	2.88
					3.32		14.43		3.68	
					2.72		17.89		3.22	
AL4 – 8	25.11.2022	202	142	56	2.12	3.42	14.43	12.50	2.07	2.99
					5.14		17.89		4.03	
					3.02		5.19		2.88	
AL1 – 9	29.11.2022	249	157	91	3.02	4.03	8.66	11.54	1.38	2.49
					6.35		20.20		3.80	
					2.72		5.77		2.30	
Monthly Average		197	127	66		2.70		11.33		2.49
Standard Deviation		50	32	20		0.65		2.05		0.37

Table 5 : Results of Air Pollutant Concentration at Gopalpuri Hospital					
Sampling Period	Date	C₆H₆ [µg/m³]	*NMHC	CO [mg/m³]	CO₂ [ppm]
		8 hr		Grab Sampling	Grab Sampling
NAAQMS limit		5.0 µg/m³		4.0 mg/m³	-
AL4 -1	01.11.2022	1.14	BQL	1.26	503
AL4 -2	04.11.2022	1.15	BQL	1.26	450
AL4 -3	08.11.2022	1.03	BQL	1.73	506
AL4 -4	11.11.2022	1.02	BQL	1.82	462
AL4 – 5	15.11.2022	1.09	BQL	1.04	1048
AL4 – 6	18.11.2022	1.14	BQL	1.32	543
AL4 – 7	22.11.2022	1.16	BQL	1.83	758
AL4 – 8	25.11.2022	1.22	BQL	1.8	816
AL4 – 9	29.11.2022	1.16	BQL	1.36	665
Monthly Average		1.12	-	1.49	639.00
Standard Deviation		0.07	-	0.30	201.83

* NMHC- Non- Methane Hydrocarbons

BQL- Below Quantification Limit (Quantification Limit – NMHC: 0.5 ppm)

The overall values of TSPM, PM₁₀, PM_{2.5}, SO₂, NO₂ and NH₃ at Gopalpuri Hospital was attributed by vehicle emission produced from light motor vehicles of the colony residents. The mean TSPM values at Gopalpuri Hospital were 197 µg/m³, the mean PM₁₀ value was 127 µg/m³ and PM_{2.5} was 66 µg/m³ which was exceed the standard limit. The average values of SO₂, NO₂ and NH₃ were 2.70 µg/m³, 11.33 µg/m³ and 2.49 µg/m³ respectively and were all within the permissible limit.

The levels of Benzene, Hydrocarbons (HC) and CO were within the permissible limit at Gopalpuri Hospital. The mean Benzene concentration was 1.12 µg/m³, well below the permissible limit of 5.0 µg/m³. NMHC's were below the detectable limit and Carbon monoxide concentration was 1.49 mg/m³ which is well below the permissible limit of 4.0 mg/m³.

Location 5: Coal Storage Area (AL-5)

Table 6 : Results of Air Pollutant Concentration at Coal Storage Area

Table 6 : Results of Air Pollutant Concentration at Coal Storage Area										
	Date	TSPM [µg/m3]	PM10 [µg/m3]	PM2.5 [µg/m3]	SO2 [µg/m3]		NOx [µg/m3]		NH3 [µg/m3]	
Sampling Period		24hr	24hr	24hr	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
NAAQMS Limit			100 µg/m3	60 µg/m3		80 µg/m3		80 µg/m3		400 µg/m3
AL6 – 1	01.11.2022	779	598	175	2.72	4.33	6.35	16.54	3.68	5.06
					6.65		25.97		8.17	
					3.63		17.31		3.34	
AL6 – 2	04.11.2022	635	492	137	2.12	3.53	23.08	17.70	6.79	6.60
					5.44		12.12		8.17	
					3.02		17.89		4.83	
AL6 – 3	08.11.2022	538	412	125	8.94	5.00	23.66	21.74	2.53	3.88
					3.46		12.12		2.07	
					2.59		29.43		7.02	
AL6 – 4	11.11.2022	815	635	178	4.53	4.73	18.47	17.70	5.87	4.41
					2.72		8.66		2.65	
					6.95		25.97		4.72	
AL6 – 5	15.11.2022	792	614	176	6.35	6.65	18.47	13.66	4.72	3.88
					9.07		10.39		3.68	
					4.53		12.12		3.22	
AL6 – 6	18.11.2022	771	595	171	9.37	7.15	20.20	17.12	4.83	4.37
					5.74		8.08		2.53	
					6.35		23.08		5.76	
AL6 – 7	22.11.2022	706	543	156	4.84	4.53	10.39	18.47	4.83	5.03
					6.04		23.66		5.99	
					2.72		21.35		4.26	
AL6 – 8	25.11.2022	846	654	187	3.32	5.24	17.31	19.81	3.91	4.95
					7.86		25.97		6.91	
					4.53		16.16		4.03	
AL1 – 9	29.11.2022	801	621	172	5.14	5.64	16.16	18.28	3.57	4.30
					9.07		28.86		6.22	
					2.72		9.81		3.11	
Monthly Average		743	574	164		5.20		17.89		4.72
Standard Deviation		99	78	21		1.14		2.22		0.84

Table 6 : Results of Air Pollutant Concentration at Coal Storage Area

Sampling Period	Date	C ₆ H ₆ [µg/m ³]	*NMHC	CO [mg/m ³]	CO ₂ [ppm]
		8 hr		Grab Sampling	Grab Sampling
NAAQMS limit		5.0 µg/m ³		4.0 mg/m ³	-
AL5 – 1	01.11.2022	1.1	BQL	1.12	483
AL5 – 2	04.11.2022	1.06	BQL	1.48	475
AL5 – 3	08.11.2022	1.08	BQL	1.66	421
AL5 – 4	11.11.2022	1.06	BQL	1.69	492
AL5 – 5	15.11.2022	1.06	BQL	1.06	702
AL5 – 6	18.11.2022	1.22	BQL	1.18	483
AL5 – 7	22.11.2022	1.11	BQL	1.86	564
AL5 – 8	25.11.2022	1.2	BQL	1.54	777
AL5 – 9	29.11.2022	1.22	BQL	1.89	895
Monthly Average		1.12	-	1.50	588.00
Standard Deviation		0.07	-	0.31	164.11

* NMHC- Non- Methane Hydrocarbons

BQL- Below Quantification Limit (Quantification Limit – NMHC: 0.5 ppm)

The overall values of TSPM, PM₁₀, PM_{2.5}, SO₂, NO₂ and NH₃ at Coal Storage Area was comparatively highest among all the locations of Air Quality monitoring in Kandla Port. High values of TSPM, PM₁₀, PM_{2.5}, SO₂, NO₂ at this location was due to lifting of coal with grab and other coal handling processes near Berth no. 6 & 7. Moreover, the traffic was also heavy around this place for transport of coal thus emissions produced from heavy vehicles. The mean TSPM values at Coal storage were 743 µg/m³, the mean PM₁₀ value was 574 µg/m³, and the PM_{2.5} value was 164 µg/m³ which was above the permissible limit prescribed by NAAQS. The average values of SO₂, NO₂ and NH₃ were 5.20 µg/m³, 17.89 µg/m³ and 4.72 µg/m³ respectively and were all within the permissible limit.

The levels of Benzene, Hydrocarbons (HC) and CO were within the permissible limit at Coal Storage Area. The mean Benzene concentration was 1.12 µg/m³, well below the permissible limit of 5.0 µg/m³. NMHC's were below the detectable limit and Carbon Monoxide concentration was 1.50 mg/m³, well below the permissible limit of 4.0 mg/m³.

Location 6: Tuna Port (AL-6)

Table 7 : Results of Air Pollutant Concentration at Tuna Port

Table 7 : Results of Air Pollutant Concentration at Tuna Port										
	Date	TSPM [µg/m3]	PM10 [µg/m3]	PM2.5 [µg/m3]	SO2 [µg/m3]		NOx [µg/m3]		NH3 [µg/m3]	
Sampling Period		24hr	24hr	24hr	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
NAAQMS Limit			100 µg/m3	60 µg/m3		80 µg/m3		80 µg/m3		400 µg/m3
AL5 -1	01.11.2022	141	88	47	0.91	1.61	2.89	6.16	2.07	2.84
					2.72		12.12		4.03	
					1.21		3.46		2.42	
AL5 – 2	04.11.2022	232	166	64	1.51	2.22	6.35	7.89	1.38	2.76
					3.02		5.19		4.49	
					2.12		12.12		2.42	
AL5 – 3	08.11.2022	184	120	55	1.44	2.40	10.39	13.08	1.73	2.61
					3.46		11.54		2.65	
					2.31		17.31		3.45	
AL5 – 4	11.11.2022	233	153	78	2.12	2.32	11.54	11.54	1.27	1.57
					3.93		17.89		1.04	
					0.91		5.19		2.42	
AL5 – 5	15.11.2022	221	145	74	1.21	2.32	6.35	12.12	3.57	2.49
					3.32		12.12		2.30	
					2.42		17.89		1.61	
AL5 – 6	18.11.2022	248	162	83	1.81	2.01	17.31	17.12	2.30	10.21
					1.21		23.66		15.57	
					3.02		10.39		12.76	
AL5 – 7	22.11.2022	214	139	74	1.51	2.52	8.66	8.46	3.57	2.84
					2.72		12.70		2.88	
					3.32		4.04		2.07	
AL5 – 8	25.11.2022	255	175	77	2.72	3.02	8.66	8.08	3.45	3.30
					4.84		11.54		4.72	
					1.51		4.04		1.73	
AL1 – 9	29.11.2022	245	155	87	1.51	3.63	12.70	11.73	1.04	2.88
					6.04		17.31		5.18	
					3.32		5.19		2.42	
Monthly Average		219	145	71		2.45		10.69		3.50
Standard Deviation		36	27	13		0.58		3.37		2.56

Table 7 : Results of Air Pollutant Concentration at Tuna Port

		C₆H₆ [µg/m³]		CO [mg/m³]	CO₂ [ppm]
Sampling Period	Date	8 hr	*NMHC	Grab Sampling	Grab Sampling
NAAQMS limit		5.0 µg/m³		4.0 mg/m³	-
AL6 -1	01.11.2022	1.12	BQL	1.43	543
AL6 – 2	04.11.2022	1.17	BQL	1.41	463
AL6 – 3	08.11.2022	1.13	BQL	1.39	410
AL6 – 4	11.11.2022	1.13	BQL	1.74	509
AL6 – 5	15.11.2022	1.17	BQL	1.08	911
AL6 – 6	18.11.2022	1.17	BQL	1.1	528
AL6 – 7	22.11.2022	1.06	BQL	1.88	565
AL6 – 8	25.11.2022	1.1	BQL	1.89	999
	29.11.2022	1.22	BQL	1.89	895
Monthly Average		1.14	-	1.53	647.00
Standard Deviation		0.05	-	0.33	222.45

* NMHC- Non- Methane Hydrocarbons

BQL- Below Quantification Limit (Quantification Limit – NMHC: 0.5 ppm)

The mean TSPM values at Tuna Port was 219 µg/m³, the mean PM₁₀ value was 145 µg/m³ and the mean PM_{2.5} value was 71 µg/m³ which was exceed the standard limit prescribed by NAAQS. The average values of SO₂, NO₂ and NH₃ were 2.45 µg/m³, 10.69 µg/m³ and 3.50 µg/m³ respectively and were all within the standard limit prescribed by NAAQS.

The levels of Benzene, Hydrocarbons (HC) and CO were within the permissible limit at Tuna Port. The mean Benzene concentration was 1.14 µg/m³, well below the permissible limit of 5.0 µg/m³. NMHC's were below the detectable limit and Carbon Monoxide concentration was 1.53 mg/m³, well below the permissible limit of 4.0 mg/m³.

Location 7: Admin Building (Vadinar) (AL-7)

Table 8 : Results of Air Pollutant Concentration at Admin Building										
	Date	TSPM [µg/m3]	PM10 [µg/m3]	PM2.5 [µg/m3]	SO2 [µg/m3]		NOx [µg/m3]		NH3 [µg/m3]	
Sampling Period		24hr	24hr	24hr	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
NAAQMS Limit			100 µg/m3	60 µg/m3		80 µg/m3		80 µg/m3		400 µg/m3
AL7 -1	01.11.2022	150	98	51	2.20	3.52	9.53	10.59	5.36	5.28
					4.84		16.51		2.81	
					3.52		5.72		7.66	
AL7 -2	04.11.2022	177	115	61	3.08	4.69	17.78	21.81	2.81	6.13
					7.03		21.60		8.93	
					3.96		26.04		6.64	
AL7 -3	08.11.2022	193	113	73	6.15	6.30	6.99	11.43	3.83	7.49
					8.79		20.96		10.47	
					3.96		6.35		8.17	
AL7 -4	11.11.2022	200	121	78	3.96	6.01	17.78	15.24	10.47	6.81
					5.28		22.23		5.87	
					8.79		5.72		4.08	
AL7 -5	15.11.2022	179	108	69	1.76	5.28	7.62	18.00	3.06	5.62
					5.71		26.04		5.87	
					8.35		20.33		7.91	
AL7 -6	18.11.2022	223	121	96	2.64	4.54	8.89	15.03	5.62	5.70
					4.40		16.51		8.17	
					6.59		19.69		3.32	
AL1 -7	22.11.2022	162	104	57	4.84	5.28	14.61	14.61	13.02	9.10
					7.03		5.72		8.68	
					3.96		23.50		5.62	
AL1-8	25.11.2022	237	138	97	6.59	4.40	9.53	15.24	7.91	8.00
					3.96		14.61		5.62	
					2.64		21.60		10.47	
AL1-9	28.11.2022	203	112	87	3.96	3.66	6.99	13.76	5.62	6.04
					2.20		14.61		7.91	
					4.84		19.69		4.60	
Monthly Average		191	114	74		4.85		15.08		6.68
Standard Deviation		28	12	17		0.96		3.34		1.28

Table 8 : Results of Air Pollutant Concentration at Admin Building Vadinar

Sampling Period	Date	C ₆ H ₆ [µg/m ³]	*NMHC	CO [mg/m ³]	CO ₂ [ppm]
		8 hr		Grab Sampling	Grab Sampling
		5.0 µg/m ³		4.0 mg/m ³	-
AL7 -1	01.11.2022	1.08	BQL	1.43	225
AL7 -2	04.11.2022	1.13	BQL	1.54	236
AL7 -3	08.11.2022	1.17	1.81	1.53	455
AL7 -4	11.10.2022	1.14	BQL	1.61	443
AL7 -5	15.10.2022	1.03	BQL	1.1	347
AL7 -6	18.10.2022	1.06	BQL	1.57	416
AL7 -7	22.10.2022	1.10	BQL	1.05	372
AL7 -8	25.10.2022	1.20	BQL	1.79	464
AL7 -9	28.10.2022	1.13	BQL	1.42	487
Monthly Average		1.12	-	1.46	388
Standard Deviation		0.06	-	0.25	75

*NMHC- Non- Methane Hydrocarbons

BQL- Below Quantification Limit (Quantification Limit – NMHC: 0.5 ppm)

At Admin Building, Vadinar the mean TSPM value was 191 µg/m³, the mean PM₁₀ value was 114 µg/m³ and the mean PM_{2.5} value was 74 µg/m³ which was slightly exceed the standard limit. The average values of SO₂, NO₂ and NH₃ concentrations were 4.85 µg/m³, 15.08 µg/m³ and 6.68 µg/m³ respectively and were all within the permissible limit.

The levels of Benzene, Hydrocarbons (HC) and CO were within the permissible limit at Vadinar Port. The mean Benzene concentration was 1.12 µg/m³, well below the permissible limit of 5.0 µg/m³. NMHC's were below the detectable limit and Carbon Monoxide concentration was 1.46 mg/m³, well below the permissible limit of 4.0 mg/m³.

Location 8: Signal Building (Vadinar) (AL-8)

Table 9 : Results of Air Pollutant Concentration at Signal Building, Vadinar										
	Date	TSPM [µg/m3]	PM10 [µg/m3]	PM2.5 [µg/m3]	SO2 [µg/m3]		NOx [µg/m3]		NH3 [µg/m3]	
Sampling Period		24hr	24hr	24hr	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)	8 hr	24hr (Avg.)
NAAQMS Limit			100 µg/m3	60 µg/m3		80 µg/m3		80 µg/m3		400 µg/m3
AL8 -1	01.11.2022	113	74	38	3.96	4.40	6.99	13.34	2.30	7.15
					6.59		19.05		8.68	
					2.64		13.97		10.47	
AL8 -2	04.11.2022	146	93	49	2.64	4.40	14.61	15.88	5.36	6.13
					4.84		22.23		8.42	
					5.71		10.80		4.60	
AL8 -3	08.11.2022	124	82	42	3.08	3.52	14.61	16.73	5.62	5.62
					5.28		26.04		7.91	
					2.20		9.53		3.32	
AL8 -4	11.11.2022	175	105	67	2.20	4.40	8.26	13.76	8.93	9.02
					7.03		19.05		12.76	
					3.96		13.97		5.36	
AL8 -5	15.11.2022	152	97	52	3.52	4.98	5.72	13.13	6.89	7.57
					4.84		13.34		10.98	
					6.59		20.33		4.85	
AL8 -6	18.11.2022	176	111	61	3.08	3.81	15.24	17.57	7.15	8.42
					3.96		26.04		7.91	
					4.40		11.43		10.21	
AL8 -7	22.11.2022	214	118	93	3.52	5.71	5.72	12.91	7.91	8.25
					5.28		13.34		6.38	
					8.35		19.69		10.47	
AL8-8	25.11.2022	219	125	92	3.08	4.54	9.53	11.01	5.36	6.04
					4.84		17.78		8.17	
					5.71		5.72		4.60	
AL8-9	28.11.2022	154	97	57	5.71	3.81	10.80	16.94	7.15	8.76
					3.96		22.23		8.93	
					1.76		17.78		10.21	
Monthly Average		164	100	61		4.40		14.59		7.44
Standard Deviation		36	16	20		0.67		2.25		1.27

Table 9 : Results of Air Pollutant Concentration at Signal Building Vadinar

		C₆H₆ [µg/m³]		CO [mg/m³]	CO₂ [ppm]
Sampling Period	Date	8 hr	*NMHC	Grab Sampling	Grab Sampling
NAAQMS limit		5.0 µg/m³		4.0 mg/m³	-
AL8 -1	01.11.2022	1.06	BQL	1.5	467
AL8 -2	04.11.2022	1.05	BQL	1.46	501
AL8 -3	08.11.2022	1.14	1.81	1.31	489
AL8 -4	11.11.2022	1.16	BQL	1.38	439
AL8 -5	15.11.2022	1.17	BQL	1.29	231
AL8 -6	18.11.2022	1.10	BQL	1.31	244
AL8 -7	22.11.2022	1.00	BQL	1.34	227
AL8 -8	25.11.2022	1.05	BQL	1.37	261
AL8 -9	28.11.2022	1.02	BQL	1.29	234
Monthly Average		1.16	-	1.46	442
Standard Deviation		0.05	-	0.27	63

* NMHC- Non- Methane Hydrocarbon

BQL- Below Quantification Limit (Quantification Limit – NMHC: 0.5 ppm)

At Signal Building, Vadinar the mean TSPM value was 164 µg/m³, the mean PM₁₀ value was 100 µg/m³ which was boundary line of the permissible limit, the mean PM_{2.5} value was 61 µg/m³ which was within the permissible limit. The average values of SO₂, NO₂ and NH₃ concentrations were 4.40 µg/m³, 14.59 µg/m³ and 7.44 µg/m³ respectively and were all within the standard limit.

The levels of Benzene, Hydrocarbons (HC) and CO were within the permissible limit at Vadinar Port. The mean Benzene concentration was 1.16 µg/m³, well below the standard limit of 5.0 µg/m³. NMHC's were below the detectable limit and Carbon Monoxide concentration was 1.46 mg/m³, well below the standard limit of 4.0 mg/m³.

Fig. No:-1 Average ambient air quality (PM) month of November-2022 at DPA and Vadinar Sampling Station




Fig. No:-2. Average ambient air quality (Gaseous) month of November-2022 at DPA and Vadinar sampling location




Fig. No:-3. Average ambient air quality (Gaseous) month of November-2022 at DPA and Vadinar sampling location





Fig. No:-4. Average ambient air quality (Gaseous) month of November-2022 at DPA and Vadinar sampling location



2.3 Observations and Conclusion

During the monitoring period, the overall Ambient Air Quality of the port area was found within permissible levels for various gaseous pollutants. However, Total Suspended Particulate matter as TSPM, Particulate matter as PM₁₀ and PM_{2.5} was found to exceed the limits at locations at all ambient air sampling location.

The concentration of PM₁₀ and PM_{2.5} were slightly exceeded at Gopalpuri and Tuna Port.

The mean concentration of PM₁₀ and PM_{2.5} were slightly exceeded at Admin building Vadinar & at Signal building Vadinar was very close to the standard limit.

CHAPTER-3

METEOROLOGICAL OBSERVATIONS

4.1 Meteorological Data

Automatic Weather station (ID KAZPHOEN424) have been installed in Seva Sadan-3 at the Deendayal Port which records the data on Temperature (°C), Relative Humidity (%), Wind speed (m/s), Wind Direction (°), Solar radiation (w/m²) and Rainfall mm.

Meteorological factors play an important role in environmental pollution studies particularly in pollutant transport irrespective of their entry into the environment. The wind speed and direction play a major role in dispersion of environment pollutants. Effects of pollution on receptors animate and inanimate depends on atmospheric condition.

Temperature

At Deendayal Port, the day time temperature was found range 21.1-32.9⁰C. The average day time temperature was 27.92°C. The night time temperature was range from 20.0-29.7⁰C. The mean night time temperature recorded was 25.47 °C.

Solar Radiation

The mean Solar Radiation in November month was 167.27 w/m². The maximum solar radiation was recorded 759.0 w/m² in 4th November, 2022 and the minimum solar radiation was recorded 1.80 w/m² in 30th November, 2022.

Rainfall


Rain fall of November month was recorded 0.00 mm.

Relative Humidity

The mean Relative humidity was 69.00 % for the month of November. Maximum Relative humidity was recorded 99.0 % and minimum Relative humidity was recorded 34.0 %.

Wind Velocity and Wind Direction

Velocity and direction of wind have a significant role in the dispersion of air borne materials and therefore determines the air quality of the area. The average wind velocity for the entire month of November was 1.21 m/s. Maximum wind velocity was recorded 10.19 m/s. The wind direction was mostly North-East.



CHAPTER-4

DRINKING WATER QUALITY MONITORING

4.0 Drinking Water Quality Monitoring

Drinking Water Quality Monitoring was carried out at twenty stations at Kandla, Vadinar & Township Area of Deendayal Port.

Table No:-10. Drinking Water Sampling Location

Sr. No.	Name of Location	Location Code	Latitude	Longitude
1.	Nirman Building	DL-1	23° 0' 27"N	70° 13' 21"E
2.	P & C Building	DL-2	23° 0' 33"N	70° 13' 20"E
3.	North Gate	DL-3	23° 0' 26.97"N	70° 13' 21.87"E
4.	KPT-Canteen	DL-4	23° 2' 17.2674"N	70° 13' 18.2814"E
5.	West Gate	DL-5	23° 59' 40.48"N	70° 12' 50.96"E
6.	Wharf Area	DL-6	22° 59' 52.2"N	70° 13' 22.95"E
7.	Sevasadan-3	DL-7	23° 0' 22.55"N	70° 13' 15.34"E
8.	Workshop	DL-8	23° 0' 33.74"N	70° 13' 20.05"E
9.	Custom Building	DL-9	23° 1' 8.70"N	70° 12' 52.0"E
10.	Kandla Colony	DL-10	23° 11' 14.9"N	70° 12' 48.4"E
11.	KPT Hospital	DL-11	23° 1' 5.02"N	70° 12' 44.38"E
12.	A.O. Building	DL-12	23° 3' 42.89"N	70° 8' 41.5"E
13.	Gopalpuri School	DL-13	23° 5' 1.03"N	70° 7' 55.42"E
14.	Gopalpuri Guest House	DL-14	23° 4' 43.14"N	70° 7' 51.92"E
15.	E-Type Quarters	DL-15	23° 4' 59.90"N	70° 7' 56.72"E
16.	F-Type Quarters	DL-16	23° 4' 38.45"N	70° 8' 8.63"E
17.	Gopalpuri Hospital	DL-17	23° 4' 54.09"N	70° 8' 7.5"E
18.	Tuna Port	DL-18	23° 58' 23.06"N	70° 5' 35.6"E
19.	Vadinar Jetty	DL-19	22° 25' 51.73"N	69° 41' 36.62"E
20.	Vadinar Colony	DL-20	22° 30' 26.25"N	69° 39' 45.03"E

4.1 Drinking Water Monitoring Methodology

Samples for physico-chemical analysis were collected in 2 Carboys and samples for microbiological parameters were collected in sterilized bottles. These samples were then analyzed in laboratory for various drinking water parameters at Kandla Lab/Surat.

The Sampling was done as per IS: 3025 Part-1, analysis was done as per IS: 3025/APHA standard methods and, the analysis results compare with IS 10500:2012. The water samples were analyzed for various parameters, viz. Color , Odor, Turbidity , Conductivity , pH , Chlorides , TDS, Total Hardness, Iron , Sulphate, Salinity , DO, BOD, Na, K, Ca, Mg, F, NO₃, NO₂, Mn, Cr-6, Cu, Cd, As, Hg, Pb, Zn, Bacterial Count (CFU) .

4.2 Results

The Drinking Water Quality monitoring data for 20 stations are given in below from table No. 11 to Table No. 17

Table 11: Drinking Water Quality Monitoring Parameters for Nirman Building, P & C Building and Main Gate (North) at Kandla.

Sr. No.	Parameter	Unit	Nirman Building 1	P & C Building	Main Gate North	Acceptable Limits as per IS 10500 :2012 2012	Permissible Limits in the absence of Alternate Source as per IS 10500 : 2012
1	pH	-	7.35	7.33	7.41	7.35	6.5 to 8.5
2	Total Dissolved Solids	mg/l	690	670	670	690	2000
3	Turbidity	NTU	0	1	1	0	5
4	Odor	-	Odorless	Odorless	Odorless	Agreeable	Agreeable
5	Color	-	Colorless	Colorless	Colorless	5	15
6	Conductivity	µs/cm	1229	1194	1211	NS*	NS*
7	Biochemical Oxygen	mg/l	BQL	BQL	BQL	NS*	NS*
8	Chloride as Cl	mg/l	576.28	355.79	340.76	250	1000
9	Ca as Ca	mg/l	43.29	41.68	39.28	75	200
10	Mg as Mg	mg/l	58.8060	57.3480	56.3760	30	100
11	Total Hardness	mg/l	350	340	330	200	600
12	Iron as Fe	mg/l	BQL	BQL	BQL	0.3	No Relaxation
13	Fluorides as F	mg/l	0.35	0.37	0.31	1	1.5
14	Sulphate as SO ₄	mg/l	35.80	30.20	28.30	200	400
15	Nitrite as NO ₂	mg/l	BQL	BQL	BQL	NS*	NS*
16	Nitrate as NO ₃	mg/l	12.70	16.70	15.50	45	No Relaxation
17	Salinity	‰	1.04	0.64	0.62	NS*	NS*
18	Sodium as Na	mg/l	204.00	180.00	192.00	NS*	NS*
19	Potassium as K	mg/l	3.22	3.15	3.18	NS*	NS*
20	Manganese	mg/l	BQL	BQL	BQL	0.1	0.3
21	Hexavalent Chromium	mg/l	BQL	BQL	BQL	NS*	NS*
22	Copper	mg/l	BQL	BQL	BQL	0.05	1.5
23	Cadmium	mg/l	BQL	BQL	BQL	0.003	NS*
24	Arsenic	mg/l	BQL	BQL	BQL	0.01	0.05
25	Mercury	mg/l	BQL	BQL	BQL	0.001	NS*
26	Lead	mg/l	BQL	BQL	BQL	0.01	NS*
27	Zinc	mg/l	BQL	BQL	BQL	5	15
28	Bacterial Count	CFU/10 0ml	Absent	Absent	Absent	Absent	Absent

*NS: Not Specified

BQL- Below Quantification Limit, (BOD-2.0 mg/l, Fe- 0.009 mg/l, Mn- 0.01 mg/l, Cr⁺⁶- 0.03 mg/l, Cu- 0.004 mg/l, Cd- 0.003 mg/l, As- 0.003mg/l, Hg- 0.001 mg/l, Pb- 0.006mg/l, Zinc- 0.021 mg/l).

Table 12: Drinking Water Quality Monitoring Parameters for Canteen, West Gate – I & Wharf Area at Kandla

Sr. No.	Parameter	Unit	Canteen	West Gate – I	Wharf Area	Acceptable Limits as per IS 10500 :	Permissible Limits in the absence of Alternate Source as per IS 10500 : 2012
1	pH	-	7.48	7.52	7.36	7.48	6.5 to 8.5
2	Total Dissolved Solids	mg/l	640	650	680	640	2000
3	Turbidity	NTU	0	1	0	0	5
4	Odor	-	Odorless	Odorless	Odorless	Agreeable	Agreeable
5	Color	-	Colorless	Colorless	Colorless	5	15
6	Conductivity	µs/cm	1166	1152	1196	NS*	NS*
7	Biochemical Oxygen Demand	mg/l	BQL	BQL	BQL	NS*	NS*
8	Chloride as Cl	mg/l	335.75	360.80	350.78	250	1000
9	Ca as Ca	mg/l	40.88	38.48	40.08	75	200
10	Mg as Mg	mg/l	62.6940	66.5820	53.4600	30	100
11	Total Hardness	mg/l	360	370	320	200	600
12	Iron as Fe	mg/l	BQL	BQL	BQL	0.3	No Relaxation
13	Fluorides as F	mg/l	0.32	0.30	0.35	1	1.5
14	Sulphate as SO ₄	mg/l	31.20	28.30	26.00	200	400
15	Nitrite as NO ₂	mg/l	BQL	BQL	BQL	NS*	NS*
16	Nitrate as NO ₃	mg/l	6.60	11.40	5.80	45	No Relaxation
17	Salinity	‰	0.61	0.65	0.63	NS*	NS*
18	Sodium as Na	mg/l	202.00	200.00	-	NS*	NS*
19	Potassium as K	mg/l	3.38	3.48	3.16	NS*	NS*
20	Manganese	mg/l	BQL	BQL	BQL	0.1	0.3
21	Hexavalent Chromium	mg/l	BQL	BQL	BQL	NS*	NS*
22	Copper	mg/l	BQL	BQL	BQL	0.05	1.5
23	Cadmium	mg/l	BQL	BQL	BQL	0.003	NS*
24	Arsenic	mg/l	BQL	BQL	BQL	0.01	0.05
25	Mercury	mg/l	BQL	BQL	BQL	0.001	NS*
26	Lead	mg/l	BQL	BQL	BQL	0.01	NS*
27	Zinc	mg/l	BQL	BQL	BQL	5	15
28	Bacterial Count	CFU/100ml	Absent	Absent	Absent	Absent	Absent

*NS: Not Specified,

BQL- Below Quantification Limit, (Nitrite - 0.05 mg/l, BOD-2.0 mg/l, Fe-0.009 mg/l, Mn- 0.01 mg/l, Cr+6- 0.03 mg/l, Cu-0.004 mg/l, Cd-0.003 mg/l, As-0.003mg/l, Hg-0.001 mg/l, Pb-0.006mg/l, Zinc-0.021 mg/l).

DCPL/DPA/21-22/31– November-2022

Table 13: Drinking Water Quality Monitoring Parameters for Sewa sadan-3, Workshop I and Custom Building at Kandla

Sr. No.	Parameter	Unit	Sewa Sadan – 3	Workshop	Custom Building	Acceptable Limits as per IS 10500 : 2012	Permissible Limits in the absence of Alternate Source as per IS 10500 : 2012
1	pH	-	7.45	7.38	7.29	6.5 to 8.5	6.5 to 8.5
2	Total Dissolved Solids	mg/l	700	670	910	500	2000
3	Turbidity	NTU	0	1	1	1	5
4	Odor	-	Odorless	Odorless	Odorless	Agreeable	Agreeable
5	Color	-	Colorless	Colorless	Colorless	5	15
6	Conductivity	µs/cm	1213	1164	1564	NS*	NS*
7	Biochemical	mg/l	BQL	BQL	BQL	NS*	NS*
8	Chloride as Cl	mg/l	365.81	370.82	340.76	250	1000
9	Ca as Ca	mg/l	42.48	37.68	39.28	75	200
10	Mg as Mg	mg/l	59.2920	59.7780	53.9460	30	100
11	Total Hardness	mg/l	350	340	320	200	600
12	Iron as Fe	mg/l	BQL	BQL	BQL	0.3	No Relaxation
13	Fluorides as F	mg/l	0.41	0.30	0.35	1	1.5
14	Sulphate as SO ₄	mg/l	24.90	34.20	27.2	200	400
15	Nitrite as NO ₂	mg/l	BQL	BQL	BQL	NS*	NS*
16	Nitrate as NO ₃	mg/l	6.90	3.90	11.00	45	No Relaxation
17	Salinity	‰	0.66	0.67	0.62	NS*	NS*
18	Sodium as Na	mg/l	-	-	-	NS*	NS*
19	Potassium as K	mg/l	3.26	4.03	3.29	NS*	NS*
20	Manganese	mg/l	BQL	BQL	BQL	0.1	0.3
21	Hexavalent Chromium	mg/l	BQL	BQL	BQL	NS*	NS*
22	Copper	mg/l	BQL	BQL	BQL	0.05	1.5
23	Cadmium	mg/l	BQL	BQL	BQL	0.003	NS*
24	Arsenic	mg/l	BQL	BQL	BQL	0.01	0.05
25	Mercury	mg/l	BQL	BQL	BQL	0.001	NS*
26	Lead	mg/l	BQL	BQL	BQL	0.01	NS*
27	Zinc	mg/l	BQL	BQL	BQL	5	15
28	Bacterial Count	CFU/100ml	Absent	Absent	Absent	Absent	Absent

*NS: Not Specified,

BQL- Below Quantification Limit, (Nitrite - 0.05 mg/l, BOD-2.0 mg/l, Fe-0.009 mg/l, Mn- 0.01 mg/l, Cr+6- 0.03 mg/l, Cu-0.004 mg/l, Cd-0.003 mg/l, As-0.003mg/l, Hg-0.001 mg/l, Pb-0.006mg/l, Zinc-0.021 mg/l).

DCPL/DPA/21-22/31– November-2022

Table 14: Drinking Water Quality Monitoring Parameters for Port Colony Kandla, Hospital Kandla and A.O. Building at Gandhidham.

Sr. No.	Parameter	Unit	Port Colony Kandla	Hospital Kandla	A.O. Building	Acceptable Limits as per IS 10500 : 2012	Permissible Limits in the absence of Alternate Source as per IS 10500 :
1	pH	-	7.39	7.31	7.24	6.5 to 8.5	6.5 to 8.5
2	Total Dissolved Solids	mg/l	760	710	1060	500	2000
3	Turbidity	NTU	1	0	0	1	5
4	Odor	-	Odorless	Odorless	Odorless	Agreeable	Agreeable
5	Color	-	Colorless	Colorless	Colorless	5	15
6	Conductivity	µs/cm	1328	1251	1821	NS*	NS*
7	Biochemical	mg/l	BQL	BQL	BQL	NS*	NS*
8	Chloride as Cl	mg/l	335.75	345.77	365.81	250	1000
9	Ca as Ca	mg/l	41.68	42.48	40.88	75	200
10	Mg as Mg	mg/l	50.0580	54.4320	62.6940	30	100
11	Total Hardness	mg/l	310	330	360	200	600
12	Iron as Fe	mg/l	BQL	BQL	BQL	0.3	No Relaxation
13	Fluorides as F	mg/l	0.35	0.32	0.46	1	1.5
14	Sulphate as SO ₄	mg/l	28.10	24.50	24.50	200	400
15	Nitrite as NO ₂	mg/l	BQL	BQL	BQL	NS*	NS*
16	Nitrate as NO ₃	mg/l	20.20	7.40	15.60	45	No Relaxation
17	Salinity	‰	0.61	0.62	0.66	NS*	NS*
18	Sodium as Na	mg/l	192.80	193.60	194.50	NS*	NS*
19	Potassium as K	mg/l	4.13	4.18	3.26	NS*	NS*
20	Manganese	mg/l	BQL	BQL	BQL	0.1	0.3
21	Hexavalent Chromium	mg/l	BQL	BQL	BQL	NS*	NS*
22	Copper	mg/l	BQL	BQL	BQL	0.05	1.5
23	Cadmium	mg/l	BQL	BQL	BQL	0.003	NS*
24	Arsenic	mg/l	BQL	BQL	BQL	0.01	0.05
25	Mercury	mg/l	BQL	BQL	BQL	0.001	NS*
26	Lead	mg/l	BQL	BQL	BQL	0.01	NS*
27	Zinc	mg/l	BQL	BQL	BQL	5	15
28	Bacterial Count	CFU/100ml	Absent	Absent	Absent	Absent	Absent

*NS: Not Specified,

BQL- Below Quantification Limit, (Nitrite - 0.05 mg/l, BOD-2.0 mg/l, Fe-0.009 mg/l, Mn- 0.01 mg/l, Cr+6- 0.03 mg/l, Cu-0.004 mg/l, Cd-0.003 mg/l, As-0.003mg/l, Hg-0.001 mg/l, Pb-0.006mg/l, Zinc-0.021 mg/l).

Table 15: Drinking Water Quality Monitoring Parameters for School Gopalpuri, Guest House) and E - Type Quarter at Gopalpuri, Gandhidham

Sr. No.	Parameter	Unit	Gopalpuri School	Guest House	E - Type Quarter	Acceptable Limits as per IS 10500 : 2012	Permissible Limits in the absence of Alternate Source as per IS 10500 : 2012
1	pH	-	7.3	7.24	7.26	6.5 to 8.5	6.5 to 8.5
2	Total Dissolved Solids	mg/l	830	950	1030	500	2000
3	Turbidity	NTU	1	1	0	1	5
4	Odor	-	Odorless	Odorless	Odorless	Agreeable	Agreeable
5	Color	-	Colorless	Colorless	Colorless	5	15
6	Conductivity	µs/cm	1435	1638	1769	NS*	NS*
7	Biochemical Oxygen Demand	mg/l	BQL	BQL	BQL	NS*	NS*
8	Chloride as Cl	mg/l	355.79	350.78	340.76	250	1000
9	Ca as Ca	mg/l	39.28	43.29	39.28	75	200
10	Mg as Mg	mg/l	61.2360	61.2360	51.5160	30	100
11	Total Hardness	mg/l	350	360	310	200	600
12	Iron as Fe	mg/l	BQL	BQL	BQL	0.3	No Relaxation
13	Fluorides as F	mg/l	0.45	0.42	0.47	1	1.5
14	Sulphate as SO ₄	mg/l	24.90	26.00	30.20	200	400
15	Nitrite as NO ₂	mg/l	BQL	BQL	BQL	NS*	NS*
16	Nitrate as NO ₃	mg/l	7.10	8.30	12.60	45	No Relaxation
17	Salinity	‰	0.64	0.63	0.62	NS*	NS*
18	Sodium as Na	mg/l	199.00	193.80	193.00	NS*	NS*
19	Potassium as K	mg/l	3.90	3.26	3.18	NS*	NS*
20	Manganese	mg/l	BQL	BQL	BQL	0.1	0.3
21	Hexavalent Chromium	mg/l	BQL	BQL	BQL	NS*	NS*
22	Copper	mg/l	BQL	BQL	BQL	0.05	1.5
23	Cadmium	mg/l	BQL	BQL	BQL	0.003	NS*
24	Arsenic	mg/l	BQL	BQL	BQL	0.01	0.05
25	Mercury	mg/l	BQL	BQL	BQL	0.001	NS*
26	Lead	mg/l	BQL	BQL	BQL	0.01	NS*
27	Zinc	mg/l	BQL	BQL	BQL	5	15
28	Bacterial Count	CFU/100 ml	Absent	Absent	Absent	Absent	Absent

*NS: Not Specified,

BQL- Below Quantification Limit, (Nitrite - 0.05 mg/l, BOD-2.0 mg/l, Fe-0.009 mg/l, Mn- 0.01 mg/l, Cr+6- 0.03 mg/l, Cu-0.004 mg/l, Cd-0.003 mg/l, As-0.003mg/l, Hg-0.001 mg/l, Pb-0.006mg/l, Zinc-0.021 mg/l).

Table 16: Drinking Water Quality Monitoring Parameters for F-Type Quarter, Hospital Gopalpuri and Tuna Port.

Sr. No.	Parameter	Unit	F - Type Quarter	Hospital Gopalpuri	Tuna Port	Acceptable Limits as per IS 10500 : 2012	Permissible Limits in the absence of Alternate Source as per IS 10500 : 2012
1	pH	-	7.28	7.42	7.51	6.5 to 8.5	6.5 to 8.5
2	Total Dissolved Solids	mg/l	1050	990	600	500	2000
3	Turbidity	NTU	1	1	-	1	5
4	Odor	-	Odorless	Odorless	Odorless	Agreeable	Agreeable
5	Color	-	Colorless	Colorless	Colorless	5	15
6	Conductivity	µs/cm	1796	1700	1044	NS*	NS*
7	Biochemical Oxygen Demand	mg/l	BQL	BQL	BQL	NS*	NS*
8	Chloride as Cl	mg/l	345.77	360.80	380.85	250	1000
9	Ca as Ca	mg/l	38.48	40.88	32.87	75	200
10	Mg as Mg	mg/l	61.7220	62.6940	72.41	30	100
11	Total Hardness	mg/l	350	360	380	200	600
12	Iron as Fe	mg/l	BQL	BQL	BQL	0.3	No Relaxation
13	Fluorides as F	mg/l	0.42	0.45	0.43	1	1.5
14	Sulphate as SO ₄	mg/l	26.00	26.10	24.50	200	400
15	Nitrite as NO ₂	mg/l	BQL	BQL	BQL	NS*	NS*
16	Nitrate as NO ₃	mg/l	10.30	6.80	3.00	45	No Relaxation
17	Salinity	‰	0.62	0.65	0.69	NS*	NS*
18	Sodium as Na	mg/l	201.00	201.00	193.60	NS*	NS*
19	Potassium as K	mg/l	3.15	3.16	3.21	NS*	NS*
20	Manganese	mg/l	BQL	BQL	BQL	0.1	0.3
21	Hexavalent Chromium	mg/l	BQL	BQL	BQL	NS*	NS*
22	Copper	mg/l	BQL	BQL	BQL	0.05	1.5
23	Cadmium	mg/l	BQL	BQL	BQL	0.003	NS*
24	Arsenic	mg/l	BQL	BQL	BQL	0.01	0.05
25	Mercury	mg/l	BQL	BQL	BQL	0.001	NS*
26	Lead	mg/l	BQL	BQL	BQL	0.01	NS*
27	Zinc	mg/l	BQL	BQL	BQL	5	15
28	Bacterial Count	CFU/100ml	Absent	Absent	Absent	Absent	Absent

*NS: Not Specified, BQL- Below Quantification Limit, (Nitrite - 0.05 mg/l, BOD-2.0 mg/l, Fe-0.009 mg/l, Mn- 0.01 mg/l, Cr+6- 0.03 mg/l, Cu-0.004 mg/l, Cd-0.003 mg/l, As-0.003mg/l, Hg-0.001 mg/l, Pb-0.006mg/l, Zinc-0.021 mg/l).

Table 17: Drinking Water Quality Monitoring Parameters for Vadinar Jetty and Port Colony at Vadinar.

Sr. No.	Parameter	Unit	Vadinar Jetty	Port Colony Vadinar	Acceptable Limits as per IS 10500 : 2012	Permissible Limits in the absence of Alternate Source as per IS 10500 : 2012
1	pH	-	7.4	7.43	6.5 to 8.5	6.5 to 8.5
2	Total Dissolved Solids	mg/l	320	300	500	2000
3	Turbidity	NTU	0.00	1.00	1	5
4	Odor	-	Odorless	Odorless	Agreeable	Agreeable
5	Color	-	Colorless	Colorless	5	15
6	Conductivity	µs/cm	570	300	NS*	NS*
7	Biochemical Oxygen Demand	mg/l	BQL	BQL	NS*	NS*
8	Chloride as Cl	mg/l	160.36	140.31	250	1000
9	Ca as Ca	mg/l	36.87	34.47	75	200
10	Mg as Mg	mg/l	43.25	52.00	30	100
11	Total Hardness	mg/l	270	300	200	600
12	Iron as Fe	mg/l	BQL	BQL	0.3	No Relaxation
13	Fluorides as F	mg/l	0.25	0.22	1	1.5
14	Sulphate as SO ₄	mg/l	0.75	0.24	200	400
15	Nitrite as NO ₂	mg/l	BQL	BQL	NS*	NS*
16	Nitrate as NO ₃	mg/l	15.60	12.70	45	No Relaxation
17	Salinity	‰	0.29	0.25	NS*	NS*
18	Sodium as Na	mg/l	191.6	192.0	NS*	NS*
19	Potassium as K	mg/l	BQL	BQL	NS*	NS*
20	Manganese	mg/l	BQL	BQL	0.1	0.3
21	Hexavalent Chromium	mg/l	BQL	BQL	NS*	NS*
22	Copper	mg/l	BQL	BQL	0.05	1.5
23	Cadmium	mg/l	BQL	BQL	0.003	NS*
24	Arsenic	mg/l	BQL	BQL	0.01	0.05
25	Mercury	mg/l	BQL	BQL	0.001	NS*
26	Lead	mg/l	BQL	BQL	0.01	NS*
27	Zinc	mg/l	BQL	BQL	5	15
28	Bacterial Count	CFU/100ml	Absent	Absent	Absent	Absent

*NS: Not Specified,

BQL- Below Quantification Limit, (Nitrite - 0.05 mg/l, BOD-2.0 mg/l, Fe-0.009 mg/l, Mn- 0.01 mg/l, Cr+6- 0.03 mg/l, Cu-0.004 mg/l, Cd-0.003 mg/l, As-0.003mg/l, Hg-0.001 mg/l, Pb-0.006mg/l, Zinc-0.021 mg/l).

4.3 Results & Discussion

The colour of all drinking water samples was found Colourless and odour of the samples also agreeable. All parameters were found within the specified limit as per the Drinking water Standard.

pH

The pH is measure of the intensity of acidity or alkalinity and the concentration of hydrogen ion in water. At DPA Site the pH values for drinking water samples ranged from 7.24-7.52 and mean value was 7.36 while at Vadinar pH ranged from 7.40-7.43 and mean value was 7.42. All the sampling points showed pH values within the prescribed limit by Indian Standards.

Turbidity

The selected drinking water sample location turbidity range from 0-1NTU at all location of DPA and Vadinar in month of November. The Turbidity values were within the permissible limit at all sampling location prescribed limit by Indian standards.

Total Dissolved Solids (TDS)

Water has the ability to dissolve a wide range of inorganic and some organic minerals or salts such as potassium, calcium, sodium, bicarbonates, chlorides, magnesium, sulfates etc.

TDS values at DPA varied between 600-1060 mg/l. The average TDS value was found 792 mg/l. The minimum value for TDS was 600 mg/l at Hospital Gopalpuri and maximum was 980 mg/l at Tuna Port while at Vadinar TDS ranged from 280-300 mg/l and mean was 290.0 mg/l. The TDS values were within the permissible limit at all sampling location prescribed limit by Indian standards.

Conductivity

Electrical Conductivity is the ability of a solution to transfer (conduct) electric current. Conductivity is used to measure the concentration of dissolved solids which have been ionized in a polar solution such as water. The conductivity in the samples collected during the month of November DPA ranged from 1044.0 $\mu\text{S}/\text{cm}$ at Tuna Port to 1821.0 $\mu\text{S}/\text{cm}$ at A.O. Building and mean value was 1381.72 $\mu\text{S}/\text{cm}$ while at Vadinar ranged from 300-570 $\mu\text{S}/\text{cm}$ and mean was 435 $\mu\text{S}/\text{cm}$.

BOD

BOD value in the studied area of DPA and Vadinar was found Below Quantification Limit (<2.0 mg/l). IS 10500:2012 does not show any standard values for BOD in drinking water.

Chlorides

Excessive chloride concentration increase rates of corrosion of metals in the distribution system. This can lead to increased concentration of metals in the supply. The Chloride value in the studied area of DPA ranged from 335.75-576.28 mg/l. The mean value was 365.53 mg/l. The minimum chloride was 335.75 mg/l at Port colony and maximum was 576.28 mg/l at Nirmal Building while at Vadinar location chloride ranged from 140.31-160.36 mg/l and mean was 150.33 mg/l. The Chloride was found within the Permissible limit of the Drinking Water Standard.

Calcium

Calcium is most abundant element on the earth crust and is very important for human cell physiology and bones. About 95% calcium in human body stored in bones and teeth. The high deficiency of calcium in humans may caused rickets, poor blood clotting, bones fracture etc. and the exceeding limit of calcium produced cardiovascular diseases.

The Calcium value in the studied area of DPA ranged from 32.87-43.29 mg/l. The mean value was 40.12 mg/l. The minimum calcium was 32.87 mg/l at Tuna Port and maximum was 43.29 mg/l at Gopalpuri Hospital while at Vadinar location Calcium ranged from 34.47-36.87 and mean was 35.67 mg/l. All the locations had calcium within the prescribed limits of 75-200 mg/L.

Magnesium

The magnesium value in the studied area of DPA ranged from 50.06-72.41 mg/l. The mean value was 59.24 mg/l. The minimum magnesium was 50.06 mg/l at Port Colony and maximum was 74.41 mg/l at Tuna Port while at Vadinar location magnesium ranged from 43.25-52.00 and mean was 47.61 mg/l. All the locations had magnesium within the prescribed limits of 30-100 mg/L.

Total Hardness

Total Hardness value in the studied area of DPA ranged from 310.0 mg/l at Port Colony to 380.0 mg/l at Tuna Port and mean value was 343.89 mg/l while at Vadinar location total hardness ranged from 270.0-300.00 mg/l and mean was 285.0 mg/l. The values of total

hardness were found within the Permissible limit of the Drinking Water Standard (200-600 mg/L). These results clear, that hardness of water is according to the IS standards and it is not harmful for local inhabitants.

Iron

Iron values in the studied area of DPA & Vadinar were Below Quantification Limit (0.009 mg/l) and hence well below the permissible limit as per Indian Standards are 0.3 mg/L.

Fluoride

Fluoride value in the studied area of DPA varied between 0.3-0.47 mg/l and mean was 0.38 mg/l. The minimum value was 0.3 mg/ at West gate workshop and maximum was 0.47 mg/l at E-Type and mean was 0.38 mg/l while at Vadinar location fluoride ranged from 0.22-0.25 mg/l and mean was 0.24 mg/l. The Fluoride values were well below the permissible limit as per Indian Standards is 1.0-1.5 mg/L. Moderate amounts lead to dental effects, but long-term ingestion of large amounts can lead to potentially severe skeletal problems.

Sulphate

Sulphate value in the studied area of DPA varied between 24.5–35.8 mg/l and mean was 27.83 mg/l. The minimum value was 24.5 mg/ at A.O. Building, Hospital Kandla and Tuna Port and maximum was 35.8 mg/l at Nirmal Building while at Vadinar location Sulphate ranged from 0.24-0.75 mg/l and mean was 0.50 mg/l. All the sampling points showed Sulphate values within the prescribed limits by Indian Standards (200-400 mg/L). Sulphate content in drinking water exceeding the 400 mg/L imparts bitter taste.

Nitrites (NO₂) and Nitrates (NO₃)

The all values of Nitrite were found BQL (<0.05 mg/l) and Nitrate were well within the permissible limit of the Drinking water Standard.

Salinity

Salinity in drinking water in the present samples collected at DPA ranged from 0.61 ‰ at Canteen to 1.04 ‰ at Nirmal Building and average salinity was 0.66 ‰ while at Vadinar sampling location salinity ranged from 0.25-0.29 ‰. There are no prescribed Indian standards for salinity in Drinking water.

Sodium and Potassium Salts

Sodium values in the samples collected at DPA ranged from 180 - 204 mg/l and average was 195.74 mg/l while at Vadinar sodium ranged from 191.6- 192.0 mg/l and average was 191.8 mg/l . Potassium salts ranged at DPA ranged from 3.15 to 4.18 mg/l while average was 3.42 mg/l while at Vadinar sampling locations potassium were BQL (<2.0 mg/l). There are no prescribed limits of Sodium and Potassium in Indian standards for Drinking water.

Heavy Metals in Drinking Water

In the present study period drinking water samples were analyzed for Mn, Cr, Cu, Cd, As, Hg, Pb and Zn. All these heavy metals were well Below the Quantification limits prescribed by the Indian Standards.

Bacteriological Study

Analysis of the bacteriological parameter (E-coli and total coliform) at all location shows that Bacteria were not detectable. This shows that drinking water samples were safe for human consumption as per tested parameters.

4.4 Conclusions

These results were compared with permissible limits as prescribed in IS 10500:2012 – Drinking Water Specification. It was seen from the analysis data that during the study period at selected sampling location the water was safe for human consumption as per analyzed parameters at all drinking water monitoring stations.

CHAPTER-5

NOISE MONITORING

5.0 Noise Level Monitoring

Noise sources in port operations include cargo handling, vehicular traffic, and loading / unloading containers and ships. Noise Monitoring was done at 13 stations at Kandla, Vadinar and Township area.

5.1 Method of Monitoring

Sampling was done at all stations for 24 hour period. Data was recorded using automated sound level meter. The intensity of sound was measured in sound pressure level (SPL) and common unit of measurement is decibel (dB).

5.2 Results

Table 18: Noise Monitoring data for ten locations of Deendayal Port and three locations of Vadinar Port

Sr. No.	Location	Day Time Average Noise Level (SPL) in dB(A)	Night Time Average Noise Level (SPL) in dB(A)
	Sampling Time	6:00 am to 10:00 PM	10:00PM to 6:00 AM
1	Marine Bhavan	60.8	51.9
2	Nirman Building 1	69.9	52.0
3	Tuna Port	53.2	45.4
4	Main Gate North	63.3	51.9
5	West Gate I	67.7	58.1
6	Canteen Area	68.2	51.2
7	Main Road	66.3	52.2
8	ATM Building	69.1	51.1
9	Wharf Area /Jetty Area	70.4	61.7
10	Port & Custom Office	54.7	50.2
Vadinar Port			
11	Entrance Gate of Vadinar Port	55.0	53.5
12	Nr. Port Colony, Vadinar	60.6	57.6
13	Nr. Vadinar Jetty	52.5	51.0

5.3 Conclusions

Transportation systems are the main source of noise pollution in urban areas. Construction of buildings, highways, and roads cause a lot of noise, due to the usage of air compressors, bulldozers, loaders, dump trucks, and pavement breakers. Noise sources in port operations include cargo handling, vehicular traffic, and loading / unloading containers and ships.

Noise sources in port operations include cargo handling, vehicular traffic, and loading / unloading containers and ships. The Day Time Noise Level (SPL) in all 10 locations at Deendayal Port Authority ranged from 53.2 dB(A) to 70.4 dB(A) while at Vadinar port 3 location ranged from 52.5 dB(A) to 60.6 dB(A) which was within the permissible limits of 75 dB(A) for the industrial area for the daytime. The Night Time Average Noise Level (SPL) in all locations of Deendayal Port Authority ranged from 45.4 dB to 61.7 dB(A) while at Vadinar port ranged from 52.5 dB (A) to 60.6 dB(A) which was within the permissible limits of 70 dB(A) for the industrial area for the night time.

CHAPTER-6

SOIL MONITORING

6.0 Soil Monitoring

Sampling and analysis of soil samples were undertaken at six locations within the study area (Deendayal Port and Vadinar Port) as a part of EMP. The soil sampling locations are initially decided based on the locations as provided in the tender document of the Deendayal Port.

Table No.:-19. Soil Sampling Location

Sr. No.	Name of Location	Location Code	Latitude	Longitude	Remarks
1.	Tuna Port	SL-1	22° 58' 10.18"N	70° 6' 3.7"E	Near main gate of Port
2.	IFFCO Plant	SL-2	23° 26' 8.37"N	70° 13' 4.4"E	10 m away from main gate
3.	Khori creek	SL-3	22° 58' 10.18"N	70° 6' 3.7"E	Sand from creek after tide
4.	Nakti Creek	SL-4	23° 2' 1.10"N	70° 9' 33.6"E	
5.	DPA admin site	SL-5	22° 26' 30.9"N	69° 40' 37.03"E	Vadinar
6.	DPA colony	SL-6	22° 23' 57.09"N	69° 42' 49.42"E	

6.1 Methodology

The soil samples were collected in the month of November 2022. The samples collected from the all locations are homogeneous representative of each location. At random locations were identified at each location and soil was dug from 30 cm below the surface. It was uniformly mixed before homogenizing the soil samples. The samples were filled in polythene bags, labeled in the field with number and site name and sent to laboratory for analysis.

6.2 Results

Table-20: Chemical Characteristics of Soil in the Study Area for Tuna port, IFFCO, Khori Creek, Nakti Creek, DPA admin site, DPA colony.

Sr. No.	Parameter	Unit	Station Name					
			SL1	SL2	SL3	SL4	SL5	SL6
			Tuna Port	IFFCO Plant	Khori Creek	Nakti Creek	DPA Admin Site	DPA Colony
			Near main gate of Port	10 m away from main	Sand from creek after tide		Vadinar	
1	Texture		Sandy Loam	Sandy Loam	Sandy Loam	Sandy Loam	Sandy Loam	Sandy Loam
2	pH	-	7.79	7.80	7.54	7.58	8.14	7.54
3	Electrical Conductivity	µs/cm	35000.0	36100.0	26,820.00	12,700.0	155.0	594.0
4	Phosphorus	mg/kg	10.3	10.5	9.19	8.49	6.00	4.80
5	Moisture	%	15.9	20.3	20.90	3.50	7.20	10.10
6	Total Organic	%	4.04	1.7	3.64	7.80	2.30	2.00
7	Alkalinity	mg/kg	900.0	1000.0	800.0	500.0	800.0	600.0
8	Total Nitrogen	%	BQL	BQL	BQL	BQL	BQL	BQL
9	Sulphate	mg/kg	820.00	982.00	1,080.00	810.00	30.0	70.0
10	Chloride	mg/kg	15598.0	14275.0	12,600.00	2,950.00	140.00	525.00
11	Calcium	mg/kg	2,605.00	2,505.00	31,600.00	3,086.00	1,729.00	1,849.00
12	Sodium	mg/kg	5657	7136.0	7,649.00	4,675.00	33.02	116.90
13	Potassium	mg/kg	552	694	708.00	437.00	44.60	44.52
14	Copper as Cu	mg/kg	27.4	15.5	30.50	14.50	54.10	31.60
15	Lead as Pb	mg/kg	7.4	7.4	9.50	6.30	74.10	75.30
16	Nickel as Ni	mg/kg	39.40	32.70	44.40	27.20	30.30	32.00
17	Zinc as Zn	mg/kg	62.4	77.40	79.20	56.50	50.60	86.00
18	Cadmium as Cd	mg/kg	BQL	BQL	BQL	BQL	BQL	BQL

BQL- Below Quantification Limit, (TN: 0.001%, Cd: 1.0mg/kg)

6.3 Discussion

- DPA Kandla soil sampling data shows that value of pH ranges from 7.54 at Khori Creek to 7.80 at IFFCO Plant while the average value was 7.68. At Vadinar sampling location pH were 7.54 at DPA colony and 8.14 at DPA Admin Site.
- The Electrical Conductivity of DPA Kandla soil sample ranged from 12700.0 $\mu\text{S}/\text{cm}$ at Nakti Creek (Sand from creek after tide) to 36100 $\mu\text{S}/\text{cm}$ at IFFCO Plant and mean was 27655 $\mu\text{S}/\text{cm}$ while Vadinar soil sampling location conductivity were 155 $\mu\text{S}/\text{cm}$ at DPA Admin Site and 594 $\mu\text{S}/\text{cm}$ at DPA Colony site.
- Total organic Carbon of DPA Kandla soil sample ranged from 1.7 % at IFFCO Plant to 7.80 % at Nakti Creek (Sand from creek after tide) and mean was 4.30 % while Vadinar soil sample were 2.0 % at DPA Colony and 2.30 % at DPA admin Site.
- The concentration of Phosphorus in the soil samples of DPA Kandla varies from 8.49 mg/kg at Nakti Creek (Sand from creek after tide) and 10.5 mg/kg at IFFCO Plant and mean was 9.62 mg/kg while the Vadinar soil sample for Phosphorus were 4.80 mg/kg at DPA Colony and 6.00 mg/kg at DPA Admin Site.
- Chloride in soil sample of DPA ranged from 2950.00 mg/kg at Nakti Creek (Sand from creek after tide) to 15598 mg/kg at Tuna Port and mean was 11356 mg/kg while Vadinar soil sample were 140 mg/kg at DPA admin and 525 mg/kg at DPA Colony.
- The Concentration of Potassium in the soil samples of DPA Kandla ranged from 437 mg/kg at Nakti creek and 708 mg/kg at Khori Creek and mean was 597.75 mg/kg while the Vadinar soil sample for Potassium were 44.52 mg/kg at DPA Colony Site and 44.60 mg/kg at DPA Admin Site.
- The concentration of Sodium in the soil samples of DPA Kandla ranged from 4675.0 mg/kg at Nakti creek and 7649.0 mg/kg at Khori Creek and mean was 6279 mg/kg while the Vadinar soil sample for Sodium were 33.00 mg/kg at DPA Admin Site and 117 mg/kg at DPA Colony.

These differences in NPK in soil at different locations are due to the dissimilar nature of soil at each of the locations. Samples SL3 & SL4 (Khori Creek & Nakti Creek) were coastal soil; where as other locations are inland locations and have different chemical properties.

Heavy Metals in the Soil

Traces of Copper, Lead, Nickel and Zinc were observed in the soil samples collected from all the four locations of Deendayal Port Authority Kandla and two locations of Vadinar Port. Cadmium metal was below detection limit in the Soil.

6.4 Conclusion

The soils of Deendayal Port Authority Kandla and Vadinar Port appears to be neutral to basic with varying levels of Chloride, Sulphate, NPK and Calcium. As the nature of soil at different locations are different with respect to its proximity to the sea, the samples showed high degree of variations in their chemical properties.

CHAPTER-7

SEWAGE TREATMENT PLANT MONITORING

7.0 Sewage Treatment Plant Monitoring

This involves safe collection of waste water (spent/used water) from wash areas, bathroom, industrial units, etc., waste from toilets of various buildings and its conveyance to the treatment plant and final disposal in conformity with the requirement and guidelines of State Pollution Control Board and other statutory bodies.

7.1 Methodology for STP Monitoring

To monitor the working efficiency of Sewage Treatment Plant (STP), STP Inlet and Outlet Samples were collected once a week. Locations selected are namely Gopalpuri Township, Deendayal Port and Vadinar. Samples were collected in 1 lit. Carboys and were analyzed in laboratory for various parameters.

A new STP with an improved capacity of 1 MLD is being constructed at Gopalpuri Colony.

Table No. 21. Sewage Treatment Plant

Sr. No.	Location of STP	Types of Treatment	STP Capacity	Treated water Utilization
1.	Gopalpuri Township	MBBR	450 KLD	Plantation and Gardening
2.	Deendayal Port, Kandla	MBBR	600 KLD	Discharge to marine through pipeline, Plantation, Gardening
3.	Vadinar Port Colony	MBBR	1.5 MLD	Plantation and Gardening

7.2 Results

Table 22: Sewage Water Monitoring at Kandla STP (1st Week)

Date of Sampling	03.11.2022
-------------------------	------------

Sr. No.	Parameters	Unit	Results		GPCB Prescribed Limit
			DPA STP I/L	DPA STP O/L	
1	pH	-	7.55	7.42	6.5 - 8.5
2	Total Suspended Solids	mg/l	100.6	46.8	100
3	Residual Chlorine	mg/l	-	<0.5	-
4	COD	mg/l	80.8	30.3	100
5	BOD @ 27 °C	mg/l	22	11	30
Aeration Tank					
6	MLSS	mg/l	14.0		
7	MLVSS	%	99.73		

Table 23: Sewage Water Monitoring at Kandla STP (2nd Week)

Date of Sampling	10.11.2022
-------------------------	------------

Sr. No.	Parameters	Unit	Results		GPCB Prescribed Limit
			DPA STP I/L	DPA STP O/L	
1	pH	-	7.41	7.36	6.5 - 8.5
2	Total Suspended Solids	mg/l	127	52.6	100
3	Residual Chlorine	mg/l	-	<0.5	-
4	COD	mg/l	90.9	40.4	100
5	BOD @ 27 °C	mg/l	23	11	30
Aeration Tank					
6	MLSS	mg/l	18.0		
7	MLVSS	%	85.00		

Table 24: Sewage Water Monitoring at Kandla STP (3rd Week)

Date of Sampling	17.11.2022
-------------------------	------------

Sr. No.	Parameters	Unit	Results		CPCB Prescribed Limit
			DPA STP I/L	DPA STP O/L	
1	pH	-	7.48	7.29	6.5 - 8.5
2	Total Suspended Solids	mg/l	86.4	22.9	100
3	Residual Chlorine	mg/l	-	<0.5	-
4	COD	mg/l	101	50.5	100
5	BOD @ 27 °C	mg/l	26	14	30
Aeration Tank					
6	MLSS	mg/l	20.0		
7	MLVSS	%	98.0		

Table 25: Sewage Water Monitoring at Kandla STP (4th Week)

Date of Sampling	24.10.2022
-------------------------	------------

Sr. No.	Parameters	Unit	Results		GPCB Prescribed Limit
			DPA STP I/L	DPA STP O/L	
1	pH	-	7.41	7.29	6.5 - 8.5
2	Total Suspended Solids	mg/l	164.2	58.7	100
3	Residual Chlorine	mg/l	-	<0.5	-
4	COD	mg/l	171.7	30.3	100
5	BOD @ 27 °C	mg/l	43	10	30
Aeration Tank					
6	MLSS	mg/l	20.0		
7	MLVSS	%	89.0		

Table 26: Sewage Water Monitoring at Gopalpuri STP (1st Week)

Date of Sampling	03.11.2022
-------------------------	------------

Sr. No.	Parameters	Unit	Results		GPCB Prescribed Limit
			DPA STP I/L	DPA STP O/L	
1	pH	-	7.47	7.31	6.5 - 8.5
2	Total Suspended Solids	mg/l	121.2	61	100
3	Residual Chlorine	mg/l	-	<0.5	-
4	COD	mg/l	111.1	60.6	100
5	BOD @ 27 °C	mg/l	32	13	30
Aeration Tank					
6	MLSS	mg/l	22.0		
7	MLVSS	%	97.16		

Table 27: Sewage Water Monitoring at Gopalpuri STP (2nd Week)

Date of Sampling	10.11.2022
-------------------------	------------

Sr. No.	Parameters	Unit	Results		GPCB Prescribed Limit
			DPA STP I/L	DPA STP O/L	
1	pH	-	7.35	7.27	6.5 - 8.5
2	Total Suspended Solids	mg/l	189	67.9	100
3	Residual Chlorine	mg/l			-
4	COD	mg/l	141.4	60.6	100
5	BOD @ 27 °C	mg/l	37	15	30
Aeration Tank					
6	MLSS	mg/l	16.0		
7	MLVSS	%	89.6		

Table 28: Sewage Water Monitoring at Gopalpuri STP (3rd Week)

Date of Sampling	17.11.2022
-------------------------	------------

Sr. No.	Parameters	Unit	Results		GPCB Prescribed Limit
			Gopalpuri STP I/L	Gopalpuri STP O/L	
1	pH	-	7.41	7.36	6.5 - 8.5
2	Total Suspended Solids	mg/l	127	52.6	100
3	Residual Chlorine	mg/l			-
4	COD	mg/l	90.9	40.4	100
5	BOD @ 27 °C	mg/l	23	11	30
Aeration Tank					
6	MLSS	mg/l	08.0		
7	MLVSS	%	98.0		

Table 29: Sewage Water Monitoring at Gopalpuri STP (4th Week)

Date of Sampling	24.11.2022
-------------------------	------------

Sr. No.	Parameters	Unit	Results		GPCB Prescribed Limit
			Gopalpuri STP I/L	Gopalpuri STP O/L	
1	pH	-	7.48	7.28	6.5 - 8.5
2	Total Suspended Solids	mg/l	110.2	42.1	100
3	Residual Chlorine	mg/l	-	<0.5	-
4	COD	mg/l	78	40	100
5	BOD @ 27 °C	mg/l	24.0	12.0	30
Aeration Tank					
6	MLSS	mg/l	18.0		
7	MLVSS	%	90.0		

Table 30: Sewage Water Monitoring at Vadinar STP (1st Week)

Date of Sampling	03.11.2022
-------------------------	------------

Sr. No.	Parameters	Unit	Results		GPCB Prescribed Limit
			Vadinar STP I/L	Vadinar STP O/L	
1	pH	-	7.35	7.25	6.5 - 8.5
2	Total Suspended Solids	mg/l	74.9	39.5	100
3	Residual Chlorine	mg/	-	<0.5	-
4	COD	mg/l	101	40.4	100
5	BOD @ 27 °C	mg/l	26.0	10.0	30

Table 31: Sewage Water Monitoring at Vadinar STP (2nd Week)

Date of Sampling	10.11.2022
-------------------------	------------

Sr. No.	Parameters	Unit	Results		GPCB Prescribed Limit
			Vadinar STP I/L	Vadinar STP O/L	
1	pH	-	7.38	7.21	6.5 - 8.5
2	Total Suspended Solids	mg/l	69.6	40.3	100
3	Residual Chlorine	mg/l	-	<0.5	-
4	COD	mg/l	131.3	50.5	100
5	BOD @ 27 °C	mg/l	32.0	7.0	30

Table 32: Sewage Water Monitoring at Vadinar STP (3rd Week)

Date of Sampling	17.11.2022
-------------------------	------------

Sr. No.	Parameters	Unit	Results		GPCB Prescribed Limit
			Vadinar STP I/L	Vadinar O/L	
1	pH	-	7.51	7.42	6.5 - 8.5
2	Total Suspended Solids	mg/l	38.6	16.9	100
3	Residual Chlorine	mg/l	-	<0.5	-
4	COD	mg/l	80.8	20.2	100
5	BOD @ 27 °C	mg/l	24.0	12.0	30

Table 33: Sewage Water Monitoring at Vadinar STP (4th Week)

Date of Sampling	24.11.2022
-------------------------	------------

Sr. No.	Parameters	Unit	Results		GPCB Prescribed Limit
			Vadinar STP I/L	Vadinar STP O/L	
1	pH	-	7.61	7.42	6.5 - 8.5
2	Total Suspended Solids	mg/l	76.9	33.3	100
3	Residual Chlorine	mg/l	-	<0.5	-
4	COD	mg/l	131.3	20.2	100
5	BOD @ 27 °C	mg/l	20.0	8.0	30

Table No. 34. General Standards for discharge of Environmental Pollutant Part-A

Sr. No.	Parameter	Inland Surface Water	Land Irrigation	Marine Coastal Areas
1.	pH	5.5-9.0	5.5-9.0	5.5-9.0
2.	Total Suspended Solids (mg/l)	100	200	100
3.	Residual Chlorine (mg/l)	1.0	-	1.0
4.	BOD (mg/l)	30	100	100
5.	COD (mg/l)	250	-	250

Sources:-CPCB**7.3 Results & Discussion**

The STP Sample carried out to evaluate the efficiency and performance of the wastewater treatment plant at Gopalpuri, Kandla and Vadinar STP. The performance of these plants is an essential parameter to monitor because the treated sewage water is discharged for irrigation purposes and discharge into marine. Wastewater samples were collected from different unit operations of the plant i.e, the inlet, aeration tank and the final treated outlet. These samples were analyzed for various physico-chemical characteristics such as pH, TSS, Residual Chlorine, COD, BOD, MLSS and MLVS.

The final treated outlet observed pH values were within the allowed range at STP Gopalpuri, STP Kandla & STP Vadinar ranged from 7.22 -7.35, 7.29-7.42 & 7.21-7.42 respectively. The wastewater treatment makes it suitable for irrigation. These values are below the allowed limit of the GPCB.

- The final treated outlet observed Total suspended solid values at Gopalpuri, DPA Kandla & Vadinar ranged from 27.10-67.90 mg/l, 22.90-58.70 mg/l & 16.60-40.30 mg/l respectively. These values are below the allowed limit of the GPCB.
- The final treated outlet observed Residual Chlorine values were <0.5 at Gopalpuri, DPA Kandla & Vadinar. These values are below the allowed limit of the CPCB.
- The final treated outlet observed COD values were at Gopalpuri, DPA Kandla & Vadinar ranged from 40.40-60.60 mg/l, 30.30-50.50 mg/l & 20.20-50.50 mg/l respectively. These values are below the allowed limit of the CPCB.

- The main focus of wastewater treatment plants is supposed to reduce the BOD in the effluent discharged to natural waters. Wastewater treatment plants are designed to function as bacteria farms, where bacteria are fed oxygen and organic waste. The final treated outlet observed BOD values were at Gopalpuri, DPA Kandla & Vadinar ranged from 12.0-16.0 mg/l, 10.0-14.0 mg/l & 7.0-12.0 mg/l respectively. These values are below the allowed limit of the GPCB.

7.4 Conclusions:

All parameters for STP outlet are within limit prescribed by CPCB. After the final treatment, it is found that the treated water is satisfactory.

CHAPTER-8

MARINE WATER MONITORING

8.0 Marine Water Monitoring

Marine Water Quality

The Forty Second Amendment to the Constitution in 1976 underscored the importance of ‘green thinking’. Article 48A enjoins the state to protect and improve the environment and safeguard the forests and wildlife in the country. Further, Article 51A (g) states that the “fundamental duty of every citizen is to protect and improve the natural environment including forests, lakes, rivers and wildlife and to have compassion for living creatures”.

Policy Statement for Abatement of Pollution (1992) has suggested developing relevant legislation and regulation, fiscal incentives, voluntary agreements and educational programs and information campaigns. It emphasizes the need for integration by incorporating environmental considerations into decision making at all levels by adopting frameworks namely, pollution prevention at source, application of best practicable solution, ensure polluter pays for control of pollution, focus on heavily polluted areas and river stretches and involve public in decision-making. The National Conservation Strategy and Policy Statement on Environment and Development, (1992) aimed at “integrating environmental concerns with developmental imperatives to meet the challenges by redirecting the thrust of our developmental process so that the basic needs of our people could be fulfilled by making judicious and sustainable use of natural resources.” The priorities mentioned in this policy document include the sustainable use of land and water resources, prevention and control of pollution and preservation of biodiversity.

The National Water Policy, (2002) contains provisions for developing, conserving, sustainable utilizing and managing this important water resources and need to be governed by national perspectives.

Sampling Stations

The monitoring of marine environment for the study of biological and ecological parameters was carried out on 01st & 02nd November-2022 in harbor regions of DPA & Vadinar during Neap tide period of New moon phase of Lunar Cycle. The monitoring of marine environment for the study of biological and ecological parameters was repeated again on 8th & 9th November-2022 in harbor regions of DPA & Vadinar during Spring tide period first quarter of Lunar Cycle.

Plankton samples from sub surface layer was collected both during high tide period and low tide period from 3 water quality monitoring stations of DPA harbor area and two stations in Nakti creek and one station in Khorī creek. The same sampling schedule was repeated during consecutive spring tide and neap tide in same month. Plankton samples from sub surface layer was collected both during high tide period and low tide period from 1 water quality monitoring stations near Vadinar jetty area during spring tide and neap tide in this month. Collected water samples were processed for estimation

of Chlorophyll- a, Pheophytin- a, qualitative & quantitative evaluation of phytoplankton, qualitative & quantitative evaluation zooplanktons (density and their population).

Sampling Locations

Offshore monitoring requirement	Number of locations
Offshore Installations	3 in Kandla creek 2 in Nakti creek 1 in Khorī creek 1 near Vadinar Jetty 1 near 1 st SBM
Total Number of locations	8

8.1 Marine Water Quality and Results

Marine water quality of marine waters of Deendayal Port Harbor waters, Khorī & Nakti Creeks and two locations of Vadinar are monitored for various physico-chemical parameters during spring and neap tide of each month. The results of marine water quality from table no 35 to 42. During low tide DPA-6 Nakti-II location monitoring was not possible due to non-availability of marine water.

Table 35: Marine Water Quality Monitoring Parameters for Location Near DPA Colony

Sr. No.	Parameters	Unit	Kandla Creek Near DPA Colony (1)			
			23°0'58"N 70°13'22."E			
			Spring Tide		Neap Tide	
	Tide		High Tide	Low Tide	High Tide	Low Tide
1	pH	-	7.61	7.58	7.55	7.46
2	Color	-	Agreeable	Agreeable	Agreeable	Agreeable
3	Odor	-	Agreeable	Agreeable	Agreeable	Agreeable
4	Salinity	‰	19.0	19.9	20.4	19.0
5	Turbidity	NTU	38	35	42	35
6	Total Dissolved Solids	mg/l	34152.0	30868.0	30941.0	31974.0
7	Total Suspended Solids	mg/l	639.6	600.6	646.4	595.6
8	Total Solids	mg/l	34791.6	31468.6	31587.4	32569.6
9	DO	mg/l	5.8	5.6	5.7	5.5
10	COD	mg/l	88.0	79.0	82.0	86.0
11	BOD	mg/l	BQL	BQL	BQL	BQL
12	Silica	mg/l	1.06	0.82	0.99	0.91
13	Phosphate	mg/l	0.48	0.31	0.09	0.04
14	Sulphate	mg/l	3580	3407	3708.0	3658
15	Nitrate	mg/l	4.70	0.50	0.75	0.42
16	Nitrite	mg/l	<0.05	<0.05	BQL	BQL
17	Calcium	mg/l	521.04	440.88	561.12	480.96
18	Magnesium	mg/l	1773.9	1749.6	1701	1773.9
19	Sodium	mg/l	8011.0	8399.0	8396.0	8699.0
20	Potassium	mg/l	299.0	385.0	391.0	395.0
21	Iron	mg/l	BQL	BQL	0.88	0.57
22	Chromium	mg/l	BQL	BQL	BQL	BQL
23	Copper	mg/l	BQL	BQL	BQL	BQL
24	Arsenic	mg/l	BQL	BQL	BQL	BQL
25	Cadmium	mg/l	BQL	BQL	BQL	BQL
26	Mercury	mg/l	BQL	BQL	BQL	BQL
27	Lead	mg/l	BQL	BQL	BQL	BQL
28	Zinc	mg/l	BQL	BQL	BQL	BQL

BQL- Below Quantification Limit, (Nitrite - 0.05 mg/l, BOD-2.0 mg/l, Cu-0.1 mg/l, As-0.1mg/l, Hg-0.01 mg/l, Zinc-0.1 mg/l).

Table 36: Marine Water Quality Monitoring Parameters for Location Near Passenger Jetty One at Kandla

Sr. No.	Parameters	Unit	Near passenger Jetty One (2)			
			23° 0'18 "N 70°13'31"E			
			Spring Tide		Neap Tide	
			High Tide	Low Tide	High Tide	Low Tide
1	pH	-	7.43	7.28	7.33	7.41
2	Color	-	Agreeable	Agreeable	Agreeable	Agreeable
3	Odor	-	Agreeable	Agreeable	Agreeable	Agreeable
4	Salinity	‰	20.8	20.4	19.9	18.6
5	Turbidity	NTU	43	48	36	41
6	Total Dissolved Solids	mg/l	35468.0	37102.0	34662.0	33398.0
7	Total Suspended Solids	mg/l	679.7	665.5	703.7	663.8
8	Total Solids	mg/l	36147.7	37767.5	35365.7	34061.8
9	DO	mg/l	5.9	6.2	5.6	5.2
10	COD	mg/l	86.0	94.0	90.0	92.0
11	BOD	mg/l	BQL	BQL	BQL	BQL
12	Silica	mg/l	1.26	0.86	1.33	0.85
13	Phosphate	mg/l	0.29	0.13	0.33	0.19
14	Sulphate	mg/l	3571	3470	4072	3407
15	Nitrate	mg/l	3.40	2.70	1.17	4.36
16	Nitrite	mg/l	<0.05	<0.05	BQL	BQL
17	Calcium	mg/l	561.12	601.20	601.2	521.04
18	Magnesium	mg/l	1701	1603.8	1749.6	1701
19	Sodium	mg/l	9142.0	9345.0	9247.0	9219.0
20	Potassium	mg/l	370.0	385.0	370.0	380.0
21	Iron	mg/l	0.47	BQL	1.76	0.30
22	Chromium	mg/l	BQL	BQL	BQL	BQL
23	Copper	mg/l	BQL	BQL	BQL	BQL
24	Arsenic	mg/l	BQL	BQL	BQL	BQL
25	Cadmium	mg/l	BQL	BQL	BQL	BQL
26	Mercury	mg/l	BQL	BQL	BQL	BQL
27	Lead	mg/l	BQL	BQL	BQL	BQL
28	Zinc	mg/l	BQL	BQL	BQL	BQL

BQL- Below Quantification Limit, (Nitrite - 0.05 mg/l, BOD-2.0 mg/l, Nitrite: 0.05mg/l Cu-0.1 mg/l, As-0.1mg/l, Hg-0.01 mg/l, Zinc-0.1 mg/l).

Table 37: Marine Water Quality Monitoring Parameters for location Near Coal Berth

Sr. No.	Parameters	Unit	Near Coal Berth			
			22°59'12"N 70°13'40"E			
			Spring Tide		Neap Tide	
			High Tide	Low Tide	High Tide	Low Tide
1	pH	-	7.37	7.51	7.53	7.25
2	Color	-	Agreeable	Agreeable	Agreeable	Agreeable
3	Odor	-	Agreeable	Agreeable	Agreeable	Agreeable
4	Salinity	‰	18.6	18.1	19.5	20.8
5	Turbidity	NTU	33	42	38	45
6	Total Dissolved Solids	mg/l	39222.0	37586.0	37123.0	36668.0
7	Total Suspended Solids	mg/l	540.2	638.4	620.6	580.2
8	Total Solids	mg/l	39762.2	38224.4	37743.6	37248.2
9	DO	mg/l	7.3	6.4	7.1	6.5
10	COD	mg/l	81.0	874.0	88.0	84.0
11	BOD	mg/l	BQL	BQL	BQL	BQL
12	Silica	mg/l	0.56	0.98	0.69	1.76
13	Phosphate	mg/l	0.06	0.56	0.12	0.61
14	Sulphate	mg/l	4222	3458	2981	3758
15	Nitrate	mg/l	2.20	4.60	2.68	4.70
16	Nitrite	mg/l	<0.05	<0.05	BQL	BQL
17	Calcium	mg/l	480.96	641.28	641.28	721.44
18	Magnesium	mg/l	1628.1	1628.1	1676.7	1603.8
19	Sodium	mg/l	8346.0	9380.0	9245.0	9814.0
20	Potassium	mg/l	391.0	300.0	392.0	384.0
21	Iron	mg/l	BQL	BQL	BQL	1.34
22	Chromium	mg/l	BQL	BQL	BQL	BQL
23	Copper	mg/l	BQL	BQL	BQL	BQL
24	Arsenic	mg/l	BQL	BQL	BQL	BQL
25	Cadmium	mg/l	BQL	BQL	BQL	BQL
26	Mercury	mg/l	BQL	BQL	BQL	BQL
27	Lead	mg/l	BQL	BQL	BQL	BQL
28	Zinc	mg/l	BQL	BQL	BQL	BQL

BQL- Below Quantification Limit, (Nitrite - 0.05 mg/l, BOD-2.0 mg/l, Cu-0.1 mg/l, As-0.1mg/l, Hg-0.01 mg/l, Zinc-0.1 mg/l).

Table 38: Marine Water Quality Monitoring Parameters for location Khori creek at Kandla

Sr. No.	Parameters	Unit	Khori creek			
			Near 15/16 Berth			
			Spring Tide		Neap Tide	
			High Tide	Low Tide	High Tide	Low Tide
1	pH	-	7.48	7.27	7.34	7.21
2	Color	-	Agreeable	Agreeable	Agreeable	Agreeable
3	Odor	-	Agreeable	Agreeable	Agreeable	Agreeable
4	Salinity	‰	20.4	19.5	18.6	17.7
5	Turbidity	NTU	35	31	43	39
6	Total Dissolved Solids	mg/l	32557.0	34294.0	30473.0	33329.0
7	Total Suspended Solids	mg/l	641.2	616.3	594.7	731.2
8	Total Solids	mg/l	33198.2	34910.3	31067.7	34060.2
9	DO	mg/l	7.6	6.3	7.3	6.8
10	COD	mg/l	85.0	96.0	92.0	96.0
11	BOD	mg/l	BQL	BQL	BQL	BQL
12	Silica	mg/l	0.78	1.04	1.39	1.18
13	Phosphate	mg/l	0.44	0.67	0.35	0.42
14	Sulphate	mg/l	4047	3646	3157	3170
15	Nitrate	mg/l	3.70	1.10	1.34	5.20
16	Nitrite	mg/l	<0.05	<0.05	BQL	BQL
17	Calcium	mg/l	561.12	480.96	480.96	561.12
18	Magnesium	mg/l	1725.3	1676.7	1701	1628.1
19	Sodium	mg/l	9112.0	8436.0	7966.0	8696.0
20	Potassium	mg/l	299.0	385.0	382.0	377.0
21	Iron	mg/l	0.44	BQL	0.17	0.31
22	Chromium	mg/l	BQL	BQL	BQL	BQL
23	Copper	mg/l	BQL	BQL	BQL	0.02
24	Arsenic	mg/l	BQL	BQL	BQL	BQL
25	Cadmium	mg/l	BQL	BQL	BQL	BQL
26	Mercury	mg/l	BQL	BQL	BQL	BQL
27	Lead	mg/l	BQL	BQL	BQL	BQL
28	Zinc	mg/l	BQL	BQL	BQL	BQL

BQL- Below Quantification Limit, (Nitrite - 0.05 mg/l, BOD-2.0 mg/l, Cu-0.1 mg/l, As-0.1mg/l, Hg-0.01 mg/l, Zinc-0.1 mg/l).

Table 39: Marine Water Quality Monitoring Parameters for location Nakti Creek near Tuna Port

Sr. No.	Parameters	Unit	Nakti Creek Near Tuna Port			
			22°57'49."N 70° 7'0.67"E			
			Spring Tide		Neap Tide	
	Tide		High Tide	Low Tide	High Tide	Low Tide
1	pH	-	7.41	7.36	7.48	7.23
2	Color	-	Agreeable	Agreeable	Agreeable	Agreeable
3	Odor	-	Agreeable	Agreeable	Agreeable	Agreeable
4	Salinity	‰	19.0	18.6	19.0	19.5
5	Turbidity	NTU	45	36	40	42
6	Total Dissolved Solids	mg/l	30214.0	28996.0	31047.0	31957.0
7	Total Suspended Solids	mg/l	642.7	526.2	682.5	606.8
8	Total Solids	mg/l	30856.7	29522.2	31729.5	32563.8
9	DO	mg/l	8.1	7.5	6.4	7.2
10	COD	mg/l	94.0	112.0	98.0	100.0
11	BOD	mg/l	BQL	BQL	BQL	BQL
12	Silica	mg/l	1.12	1.20	1.42	1.22
13	Phosphate	mg/l	0.71	0.37	0.46	0.12
14	Sulphate	mg/l	4172	3846	3445	3433
15	Nitrate	mg/l	1.50	1.70	5.12	1.69
16	Nitrite	mg/l	<0.05	<0.05	BQL	BQL
17	Calcium	mg/l	440.88	641.28	601.2	521.04
18	Magnesium	mg/l	1725.3	1555.2	1701	1773.9
19	Sodium	mg/l	8639.0	9143.0	8655.0	7939.0
20	Potassium	mg/l	395.0	386.0	384.0	386.0
21	Iron	mg/l	BQL	0.33	0.34	0.18
22	Chromium	mg/l	BQL	BQL	BQL	BQL
23	Copper	mg/l	BQL	BQL	BQL	BQL
24	Arsenic	mg/l	BQL	BQL	BQL	BQL
25	Cadmium	mg/l	BQL	BQL	BQL	BQL
26	Mercury	mg/l	BQL	BQL	BQL	BQL
27	Lead	mg/l	BQL	BQL	BQL	BQL
28	Zinc	mg/l	BQL	BQL	BQL	BQL

BQL- Below Quantification Limit, (Nitrite - 0.05 mg/l,BOD-2.0 mg/l,Cu-0.1 mg/l, As-0.1mg/l, Hg-0.01 mg/l,Zinc-0.1 mg/l).

Table 40: Marine Water Quality Monitoring Parameters for location Nakti Creek Near NH-8A at Kandla

Sr. No.	Parameters	Unit	Nakti Creek Near NH-8A			
			23° 02'01"N 70° 09'31"E			
			Spring Tide		Neap Tide	
			High Tide	Low Tide	High Tide	Low Tide
1	pH	-	7.45	Sampling not possible during Low Tide	7.45	Sampling not possible during Low Tide
2	Color	-	Agreeable		Agreeable	
3	Odor	-	Agreeable		Agreeable	
4	Salinity	‰	19.9		20.8	
5	Turbidity	NTU	45		44	
6	Total Dissolved Solids	mg/l	30288.0		32796.0	
7	Total Suspended Solids	mg/l	529.6		595.7	
8	Total Solids	mg/l	30817.6		33391.7	
9	DO	mg/l	7.4		6.9	
10	COD	mg/l	118.0		110.0	
11	BOD	mg/l	BQL		BQL	
12	Silica	mg/l	1.02		0.16	
13	Phosphate	mg/l	0.75		0.46	
14	Sulphate	mg/l	4109		4961	
15	Nitrate	mg/l	2.70		3.52	
16	Nitrite	mg/l	<0.05		BQL	
17	Calcium	mg/l	681.36		641.28	
18	Magnesium	mg/l	1506.6		1628.1	
19	Sodium	mg/l	9280.0		8528.0	
20	Potassium	mg/l	427.0		427.0	
21	Iron	mg/l	BQL		0.54	
22	Chromium	mg/l	BQL		BQL	
23	Copper	mg/l	BQL		BQL	
24	Arsenic	mg/l	BQL		BQL	
25	Cadmium	mg/l	BQL		0.01	
26	Mercury	mg/l	BQL		BQL	
27	Lead	mg/l	BQL		BQL	
28	Zinc	mg/l	BQL		BQL	

BQL- Below Quantification Limit, (Nitrite - 0.05 mg/l, BOD-2.0 mg/l, Cu-0.1 mg/l, As-0.1 mg/l, Hg-0.01 mg/l, Zinc-0.1 mg/l).

Table 41: Marine Water Quality Monitoring Parameters for locations Nr. Vadinar Jetty

Sr. No.	Parameters	Unit	Nr.Vadinar Jetty			
			22°26'25.26"N 69°40'20.41"E			
			Spring Tide		Neap Tide	
			High Tide	Low Tide	High Tide	Low Tide
1	pH	-	7.43	7.26	7.36	7.29
2	Color	-	Agreeable	Agreeable	Agreeable	Agreeable
3	Odor	-	Agreeable	Agreeable	Agreeable	Agreeable
4	Salinity	‰	20.4	20.8	19.0	19.9
5	Turbidity	NTU	39	42	38	42
6	Total Dissolved Solids	mg/l	35265.0	37685.0	36325.0	36681.0
7	Total Suspended Solids	mg/l	585.3	590.8	681.4	657.6
8	Total Solids	mg/l	35850.3	38275.8	37006.4	37338.6
9	DO	mg/l	5.7	5.4	6.3	5.8
10	COD	mg/l	87.0	89.0	96.0	92.0
11	BOD	mg/l	BQL	BQL	BQL	BQL
12	Silica	mg/l	0.55	0.45	0.36	0.28
13	Phosphate	mg/l	0.18	0.42	0.33	0.19
14	Sulphate	mg/l	3608	3558	3683	3645
15	Nitrate	mg/l	2.35	1.09	1.00	2.43
16	Nitrite	mg/l	<0.05	<0.05	BQL	BQL
17	Calcium	mg/l	480.96	601.20	521.04	480.96
18	Magnesium	mg/l	1603.8	1652.4	1676.7	1749.6
19	Sodium	mg/l	9448.0	7368.0	7810.0	8912.0
20	Potassium	mg/l	371.0	354.0	452.0	456.0
21	Iron	mg/l	BQL	BQL	0.31	BQL
22	Chromium	mg/l	BQL	BQL	BQL	BQL
23	Copper	mg/l	BQL	BQL	BQL	BQL
24	Arsenic	mg/l	BQL	BQL	BQL	BQL
25	Cadmium	mg/l	BQL	BQL	BQL	BQL
26	Mercury	mg/l	BQL	BQL	BQL	BQL
27	Lead	mg/l	BQL	BQL	BQL	BQL
28	Zinc	mg/l	0.29	BQL	0.77	0.35

BQL- Below Quantification Limit, (Nitrite - 0.05 mg/l, BOD-2.0 mg/l, Cu-0.1 mg/l, As-0.1mg/l, Hg-0.01 mg/l, Zinc-0.1 mg/l).

Table 42: Marine Water Quality Monitoring Parameters for locations Nr. Vadinar SPM

Sr. No.	Parameters	Unit	Nr. Vadinar SPM			
			22°30'56.15"N 69°42'12.07"E			
			Spring Tide		Neap Tide	
			High Tide	Low Tide	High Tide	Low Tide
1	pH	-	7.37	7.22	7.41	7.35
2	Color	-	Agreeable	Agreeable	Agreeable	Agreeable
3	Odor	-	Agreeable	Agreeable	Agreeable	Agreeable
4	Salinity	‰	19.0	17.7	19.5	18.6
5	Turbidity	NTU	37	40	37	39
6	Total Dissolved Solids	mg/l	39961.0	39198.0	42642.0	40730.0
7	Total Suspended Solids	mg/l	545.5	493.6	714.3	657.9
8	Total Solids	mg/l	40506.5	39691.6	43356.3	41387.9
9	DO	mg/l	6.1	5.5	5.6	6.1
10	COD	mg/l	95.0	98.0	96.0	94.0
11	BOD	mg/l	BQL	BQL	BQL	BQL
12	Silica	mg/l	0.47	0.37	0.34	0.30
13	Phosphate	mg/l	1.08	0.19	0.46	0.28
14	Sulphate	mg/l	3495	3796	3745	4008
15	Nitrate	mg/l	3.86	2.18	4.95	2.10
16	Nitrite	mg/l	<0.05	<0.05	BQL	BQL
17	Calcium	mg/l	561.12	400.80	681.36	641.28
18	Magnesium	mg/l	1628.1	1676.7	1555.2	1628.1
19	Sodium	mg/l	8473.0	10386.0	9131.0	8526.0
20	Potassium	mg/l	452.0	406.0	413.0	441.0
21	Iron	mg/l	BQL	BQL	0.24	BQL
22	Chromium	mg/l	BQL	BQL	BQL	BQL
23	Copper	mg/l	BQL	BQL	BQL	BQL
24	Arsenic	mg/l	BQL	BQL	BQL	BQL
25	Cadmium	mg/l	BQL	BQL	BQL	BQL
26	Mercury	mg/l	BQL	BQL	BQL	BQL
27	Lead	mg/l	BQL	BQL	BQL	BQL
28	Zinc	mg/l	0.28	BQL	0.40	BQL

BQL- Below Quantification Limit, (Nitrite - 0.05 mg/l, BOD-2.0 mg/l, Cu-0.1 mg/l, As-0.1mg/l, Hg-0.01 mg/l, Zinc-0.1 mg/l)

8.2 Results & Discussion for Marine water samples

Marine water quality of Deendayal Port Harbor waters, Khorī and Nakti Creeks and two locations of Vadinar are monitored for various physico-chemical parameters during spring and neap tide of each month. The Heavy metal analyzed and mostly found below quantification limit.

pH

During spring tide the pH values was ranged from 7.27-7.61 at DPA Kandla and 7.22-7.43 at Vadinar while during Neap Tide pH values was ranged from 7.21-7.55 at DPA Kandla and 7.29-7.41 at Vadinar.

Color and Odor

All marine samples for Odor and Color were found agreeable at all sampling locations.

Turbidity

During spring tide the Turbidity values was ranged from 31-48 NTU at DPA Kandla and 37-42 NTU at Vadinar while during Neap Tide Turbidity values was ranged from 35-45 NTU at DPA Kandla and 37-42 NTU at Vadinar. Turbidity is the amount of particulate matter that is suspended in water. Turbidity measures the scattering effect that suspended solids have on light: the higher the intensity of scattered light, the higher the turbidity (Yap et al, 2011). Materials that cause water to be turbid include clay, silt, finely divided organic and inorganic matter, soluble colored organic compounds, plankton and microscopic organisms (Lawler, 2004). The turbidity affects the amount of light penetrating to the plants for photosynthesis.

Total Dissolved Solids (TDS)

TDS values in the studied area during Spring Tide varied between 28966- 39222 mg/l at DPA Kandla and 35265-39961 mg/l at Vadinar while during Neap Tide TDS values was varied 30473-37123 mg/l at DPA Kandla and 36325-42642 mg/l at Near Vadinar.

Calcium

Calcium value in the studied area during Spring Tide varied between 440.9-681.4 mg/l at DPA Kandla and 400.8-601.2 mg/l at Vadinar while during Neap Tide calcium values between 481.0-721.4 mg/l at DPA Kandla and 481.0-681.4 mg/l at Vadinar.

Magnesium

Magnesium value in the studied area during Spring Tide varied between 1506.6-1773.9 mg/l at DPA Kandla and 1603.8-1676.7 mg/l at Vadinar while during Neap Tide magnesium values between 1603.80-173.9 mg/l at DPA Kandla and 1555.2 -1749.60 at Vadinar. Calcium and magnesium both play an important role in antagonizing the toxic effects of various ions and neutralizing the excess acid produced (Narayan R. et. al., 2007)

Nitrate

Nitrate value in the studied area during Spring Tide varied between 0.5-4.7 mg/l at DPA Kandla and 1.09-3.86 mg/l at Vadinar while during Neap Tide Nitrate values between 0.42-5.2 mg/l at DPA Kandla and 1.0-4.95 at Vadinar.

The variations were observed due to variation in phytoplankton excretion, oxidation of ammonia, reduction of nitrate and by recycling of nitrogen and bacterial decomposition of planktonic detritus (Asha and Diwakar, 2007).

Iron

Iron values in the studied area during Spring Tide ranged from 0.33-0.47 mg/l at DPA Kandla and at Vadinar were BQL (<0.10) while during Neap Tide Iron values ranged from 0.17-1.76 mg/l at DPA Kandla and 0.24-0.31 mg/l at Vadinar.

Sulphates

Sulphate values in the studied area during Spring Tide ranged from 3407-4222 mg/l at DPA Kandla and 3495-3796 mg/l at Vadinar while during Neap Tide the Sulphate values was varied 2981-4961 mg/l at DPA Kandla and 3645-4008mg/l at Vadinar.

Salinity

Salinity values in the studied area during Spring Tide varied ranged 18.11 to 20.82 ‰ at DPA Kandla and 17.65 to 20.82 ‰ at Vadinar while during Neap Tide the Salinity values was varied 17.65 to 20.82 ‰ at DPA Kandla and 18.55 to 19.92 ‰ at Vadinar.

Sodium and Potassium Salts

During Spring Tide the Sodium values ranged from 8011-9380 mg/l at DPA Kandla & 7368-10386 mg/l at Vadinar and Potassium salts ranged from 299-427 mg/l at DPA Kandla & 354-452 mg/l at Vadinar while during Neap Tide the Sodium values was ranges from 7939-

9814 mg/l at DPA Kandla & 7810-9131 mg/l at Vadinar and Potassium salts ranged from 370-427 mg/l at DPA Kandla & 413-456 mg/l at Vadinar.

DO

The DO refers to the amount of oxygen dissolved in the water and it is particularly important in limnology {(aquatic ecology) (Weiss 1970)}. The fate and behavior of DO is of critical importance to marine organisms in determining the severity of adverse impacts (Best et al. 2007). The major factor controlling dissolved oxygen concentration is biological activity: photosynthesis producing oxygen while respiration and nitrification consume oxygen (Best et al. 2007). From the studied samples, DO in marine water during Spring Tide was found in ranges from 5.6-8.1 mg/l at DPA Kandla and 5.4-6.1 mg/l at Vadinar while during Neap Tide 5.2-7.3 mg/l at DPA Kandla and 5.6-6.3 mg/l at Vadinar.

BOD

BOD in marine water at all sampling location in the studied samples were found BQL (<2.0 mg/l).

Heavy Metals in Marine Water

In the present study period marine water samples were analyzed for Cr, Cu, Cd, As, Hg, Pb and Zn. Maximum heavy metals parameters were well Below the Quantification limits.

9.3 Conclusion

In the present study period marine water samples were analyzed and found inline as per Primary Water Quality criteria for class-IV WATERS (For Harbour Waters).

CHAPTER-9

MARINE SEDIMENT MONITORING

9.0 Marine Sediments

The deep-sea ocean floor is made up of sediment. This sediment is composed of tiny particles such as fine sand, silt, clay, or animal skeletons that have settled on the ocean bottom. Over long periods of time, some of these particles become compressed and form stratified layers. Scientists that study these layers look at particle size, particle composition, and origin to help them create historical records of the deep ocean floor. This process is called weathering. Weathering can be either mechanical or chemical. Mechanical weathering can occur as ice, wind, or water wears away the rock's surface. Chemical weathering can occur as rocks are dissolved by a chemical such as acid rain. The particles created as a result of weathering are called terrigenous sediments. These particles are transported to the ocean by wind and by rivers and streams. Once the particles enter the ocean, they are dispersed by waves, currents, and tides. The heaviest and largest particles that reach the oceans, such as sand, settle very quickly to the bottom as a result of gravity. Sand is deposited near the coast whereas the smaller silt and clay particles are transported farther distances offshore before they settle to the bottom. Sediments are an important component of aquatic ecosystems because they provide nutrients and habitat for aquatic organisms (Benhamed et al. 2016). However, human activities result in accumulation of toxic substances such as heavy metals in marine sediments. Heavy metals are well-known environmental pollutants due to their toxicity, persistence in the environment, and bioaccumulation. Metals affect the ecosystem because they are not removed from water by self-purification, but accumulate in sediments and enter the food chain (Astakhov et al. 2015).

Sediment samples were collected with Van Veen Grab from the six locations in Kandla Port Waters and two locations in Vadinar Port. Benthic surface grab samplers look like giant metal jaws. They dig into the bottom and take a bite of the sediment. These samplers are good for collecting softer, sandy or silty sediments that do not contain rocks. A box corer is a cross between a surface sampler and a sediment corer. It is a special device that is used to collect an undisturbed sample of the very top surface layers and the sediment underneath. Samples were collected and preserved in silver foil in ice box to prevent the contamination/decaying of the samples.

10.1 Results

The Sediment Quality results are given in below from table no. 43 & 44.

Table 43: Results of Analysis of Sediment of Kandla & Vadinar Port (Neap Tide)

Sr. No.	Parameters	Unit	DPA – 1	DPA - 2	DPA - 3	DPA - 4	DPA - 5	Jetty	SPM
1	Texture	-	Sandy Loam	Sandy Loam	Sandy Loam	Sandy Loam	Sandy Loam	Sandy Loam	Sandy Loam
2	Organic Matter	mg/kg	1.32	0.6	0.1	0.1	0.16	1.14	1.59
3	Organic Carbon	mg/kg	0.76	0.35	0.07	0.06	0.09	0.66	0.91
4	Inorganic Phosphate	mg/kg	89.00	90.00	101.00	92.00	100.00	90.00	100.00
5	Moisture	%	3.90	2.37	4.12	3.00	4.10	3.40	4.00
6	Aluminum	mg/kg	ND	ND	ND	ND	ND	ND	ND
7	Silica	mg/kg	7.30	7.68	8.90	9.30	9.10	8.90	9.60
8	Phosphate	mg/kg	5.20	4.99	4.09	5.25	9.00	3.28	10.40
9	Sulphate	mg/kg	759.00	849.00	555.00	496.00	768.00	732.00	496.00
10	Nitrite	mg/kg	0.11	0.11	0.10	0.10	0.12	0.10	0.11
11	Nitrate	mg/kg	BQL	BQL	BQL	BQL	BQL	BQL	BQL
12	Calcium	mg/kg	2765.00	1523.00	861.00	961.00	981.00	1162.00	2485.00
13	Magnesium	mg/kg	1372.00	1300.00	1020.00	1263.00	1032.00	1089.00	2065.00
14	Sodium	mg/kg	2410.0	2760.0	2644.0	2940.0	2722.0	1394.00	1082.00
15	Potassium	mg/kg	404.00	459.00	390.00	510.00	447.00	811.0	560.0
16	Chromium	mg/kg	61.30	71.90	66.00	53.30	56.40	42.80	49.70
17	Nickel	mg/kg	26.80	31.70	29.00	23.00	24.10	13.80	29.20
18	Copper	mg/kg	17.40	19.40	17.80	15.50	15.80	13.80	47.10
19	Zinc	mg/kg	43.40	55.80	49.80	41.80	46.00	32.00	64.30
20	Cadmium	mg/kg	BQL	BQL	BQL	BQL	BQL	BQL	BQL
21	Lead	mg/kg	5.20	6.20	5.70	9.80	8.40	12.00	BQL
22	Mercury	mg/kg	BQL	BQL	BQL	BQL	BQL	BQL	BQL
23	Arsenic	mg/kg	BQL	BQL	BQL	BQL	BQL	BQL	BQL

*ND - Not Detected, BQL: Below Quantification Limit (NO₃:10.0mg/kg, Cd: 1.0mg/kg, Hg: 1.0mg/kg, As: 1.0mg/kg).

Table 44 : Results of Analysis of Sediment of Kandla & Vadinar Port (Spring Tide)

Sr. No.	Parameters	Unit	DPA – 1	DPA - 2	DPA - 3	DPA - 4	DPA - 5	Jetty	SPM
1	Texture	-	Sandy Loam	Sandy Loam	Sandy Loam	Sandy Loam	Sandy Loam	Sandy Loam	Sandy Loam
2	Organic Matter	mg/kg	0.91	0.50	1.52	0.37	0.27	1.45	1.68
3	Organic Carbon	mg/kg	0.52	0.29	0.87	0.21	0.15	0.83	0.97
4	Inorganic Phosphate	mg/kg	98.00	90.00	80.00	78.00	100.00	88.00	90.00
5	Moisture	%	17.00	8.70	15.00	6.60	4.80	14.24	13.14
6	Aluminum	mg/kg	ND	ND	ND	ND	ND	ND	ND
7	Silica	mg/kg	7.20	8.26	9.02	5.50	7.80	9.20	10.02
8	Phosphate	mg/kg	7.87	9.29	6.16	5.75	9.49	11.61	10.80
9	Sulphate	mg/kg	745.00	862.00	585.00	490.00	510.00	590.00	396.00
10	Nitrite	mg/kg	0.11	0.12	0.12	0.11	0.10	0.10	0.11
11	Nitrate	mg/kg	BQL	BQL	12.00	16.6	26.2	BQL	BQL
12	Calcium	mg/kg	1723.00	1057.00	1320.00	1220.00	1390.00	1907.00	1643.00
13	Magnesium	mg/kg	1044.00	716.00	1090.00	690.00	896.00	1563.00	2320.00
14	Sodium	mg/kg	2733.00	2720.00	2578.00	2107.00	1558.00	1042.00	952.00
15	Potassium	mg/kg	302.00	332.00	378.0	357.0	87.8	384.00	325.00
16	Chromium	mg/kg	38.00	24.40	51.70	16.10	60.00	48.90	69.20
17	Nickel	mg/kg	15.60	9.50	21.70	6.00	24.70	19.70	28.30
18	Copper	mg/kg	7.80	BQL	11.30	31.40	16.40	12.10	19.90
19	Zinc	mg/kg	30.10	21.90	35.70	13.70	44.90	31.50	51.90
20	Cadmium	mg/kg	BQL	BQL	BQL	BQL	BQL	BQL	BQL
21	Lead	mg/kg	BQL	BQL	BQL	BQL	BQL	BQL	BQL
22	Mercury	mg/kg	BQL	BQL	BQL	BQL	BQL	BQL	BQL
23	Arsenic	mg/kg	BQL	BQL	BQL	BQL	BQL	BQL	BQL

*ND - Not Detected, BQL: Below Quantification Limit (NO₃:10.0 mg/kg, Cd: 1.0 mg/kg, Hg: 1.0mg/kg, As: 1.0mg/kg)

9.2 Discussion of Marine Sediment samples

Marine Sediments of Deendayal Port Harbor waters, Khorī and Nakti Creeks and two locations of Vadinar are monitored for various physico-chemical parameters during spring and neap tide of each month. The Heavy metal analyzed and found below quantification limit.

9.3 Conclusion

The sediment types are majority Sandy loamy. Also maximum heavy metals parameters found below Quantification limit wise, Pb, Cd, Hg, As, Al was not Detected and Nitrate for some locations.

CHAPTER-11

MARINE ECOLOGICAL MONITORING

10.0 INTRODUCTION:

10.1 Sampling Stations:

The monitoring of marine environment for the study of biological and ecological Parameters was carried out on 01st November 2022 in harbour region of DPA at Kandla Creek, and on 02nd November 2022 in creeks near by the port during Neap tide. The monitoring of marine environment for the study of biological and ecological parameters was repeated again on 08th November, 2022 in harbour region of DPA at Kandla Creek and on 09th November, 2022 in creeks near by the port during spring tidal condition.

Plankton samples from sub surface layer was collected both during high tide period and low tide period from 3 water quality monitoring stations of DPA harbour area and two stations in Nakti creek and one station in Khori creek. Sampling at second sampling station of Nakti creek was possible only during high tide period.

Plankton samples from sub surface layer were collected during high tide period and low tide period from monitoring station near Vadinar Jetty at Path Finder Creek during Neap tide on 01/11/2022 and Spring tide period on 08/11/2022. Collected water samples were processed for estimation of Chlorophyll- a, Pheophytin- a, qualitative and quantitative evaluation of phytoplankton, qualitative and quantitative evaluation of zoo plankton density and their population.

TABLE 43. SAMPLING LOCATIONS

monitoring requirement	Number of locations
Kandla creek	3 in Kandla creek
Nakti creek	2 in Nakti creek
Khori Creek	1 in Khori creek
Vadinar jetty	1 near Vadinar Jetty
SPM	1 near I st SPM
Total Number of locations	8

Sampling methodology adopted:

A marine sampling is an estimation of the body of information in the population. The theory of the sampling design is depending upon the underlying frequency distribution of the population of interest. The requirement for useful water sampling is to collect a representative sample of suitable volume from the specified depth and retain it free from contamination during retrieval.

50 litres of the water sample were collected from Sub surface by using bucket. From the collected water sample 1 litres of water sample was taken in an opaque plastic bottle for chlorophyll estimation, thereafter plankton samples were collected by using filtration assembly with Nylobolt cloth of 20µm mesh size. . During low tide DPA-6 Nakti-II location monitoring was not possible due to non-availability of marine water.

Samples Processing for chlorophyll estimation:

Samples for chlorophyll estimation were preserved in ice box on board in darkness to avoid degradation in opaque container covered with aluminium foil. Immediately after reaching the shore after sampling, 1 litre of collected water sample was filtered through GF/F filters (pore size 0.45 µm) by using vacuum filtration assembly. After vacuum filtration the glass micro fiber filter paper was grunted in tissue grinder, macerating of glass fiber filter paper along with the filtrate was done in 90% aqueous Acetone in the glass tissue grinder with glass grinding tube. Glass fiber filter paper will assist breaking the cell during grinding and chlorophyll content was extracted with 10 ml of 90% Acetone, under cold dark conditions along with saturated magnesium carbonate solution in glass screw cap tubes. After an extraction period of 24 hours, the samples were transferred to calibrated centrifuge tubes and adjusted the volume to original volume with 90% aqueous acetone solution to make up the evaporation loss. The extract was clarified by using centrifuge in closed tubes. The clarified extracts were then decanted in clean cuvette and optical density was observed at wavelength 664, 665 nm. By using corrected optical density, Chlorophyll-a value was calculated as given in (APHA, 2017).

PLANKTON:

The entire area open water in the sea is the pelagic realm. Pelagic organisms live in the open sea. In contrast to the pelagic realm, the benthic realm comprises organisms and zone of the bottom of the sea. Vertically the pelagic realm can be dividing into two zones based on light penetration; upper photic or euphotic zone and lower dark water mass, aphotic zone below the photic zone.

The term plankton is a general term for organisms which have such limited powers of locomotion that they are at the mercy of the prevailing water movement. Plankton is subdivided to phytoplankton and zooplankton. Phytoplanktons are free floating organisms that are capable of photosynthesis and zooplankton is the various free-floating animals.

Pelagic zone, represents the entire ocean water column from the surface to the deepest depths, is home to a diverse community of organisms. Differences in their locomotive ability categorize the organisms in the pelagic realm into two, *plankton* and *nekton* (Lalli and Parsons, 1997). *Plankton* consists of all organisms drifting in the water and is unable to swim against water currents, whereas *Nekton* includes organisms having strong locomotive power. Ecological studies on the plankton community, which form the base of the aquatic food chain, help in the better understanding of the dynamics and

functioning of the marine ecosystem. The term 'Plankton' first coined by Victor Hensen (1887), Plankton, (Greek word: *planktos* meaning "passively drifting or wandering") is defined as drifting or free-floating organisms that inhabit the pelagic zone of water. Based on their mode of nutrition planktonic organisms are categorised into phytoplankton (organisms having an autotrophic mode of nutrition) and zooplankton (organisms having a heterotrophic mode of nutrition).

Phytoplankton in the marine environment:

Phytoplanktons are free floating unicellular, filamentous and colonial eutrophic organisms that grow in aquatic environments whose movement is more or less dependent upon water currents. These micro flora acts as primary producers as well as the basis of food chain, source of protein, bio-purifier and bio-indicators of the aquatic ecosystems of which diverse array of the life depends. They are considered as an important component of aquatic flora, play a key role in maintaining equilibrium between abiotic and biotic components of aquatic ecosystem.

The phytoplankton includes a wide range of photosynthetic and phototrophic organisms. Marine phytoplankton is mostly microscopic and unicellular floating flora, which are the primary producers that support the pelagic food-chain. The two most prominent groups of phytoplankton are Diatoms (Bacillariophyceae) and Dinoflagellates (Dinophyceae). The phytoplankton those normally captured in the net from the Gulf of Kutch is normally dominated by these two major groups; Diatoms and Dinoflagellates. Phytoplankton also include numerous and diverse collection of extremely small, motile algae which are termed micro flagellates (naked flagellates) as well as and Cyanophytes (Blue-green algae).

Algae are an ecologically important group in most aquatic ecosystems and have been an important component of biological monitoring programs. Algae are ideally suited for water quality assessment because they have rapid reproduction rates and very short life cycles, making them valuable indicators of short-term impacts.

Aquatic populations are impacted by anthropogenic stress, resulting in a variety of alterations in the biological integrity of aquatic systems. Algae can serve as an indicator of the degree of deterioration of water quality, and many algal indicators have been used to assess environmental status.

Zooplankton in the marine environment:

Zooplankton includes a taxonomically and morphologically diverse community of heterotrophic organisms that drift in the waters of the world's oceans. Qualitative and quantitative studies on zooplankton community are a prerequisite to delineate the ecological processes active in the marine ecosystem. Zooplankton community plays a pivotal role in the pelagic food web as the primary consumers of phytoplankton and act as the food source for organisms in the higher trophic levels, particularly the economically essential groups such as fish larvae and fishes. They also function in the cycling of elements in the marine ecosystem. The dynamics of the zooplankton community, their reproduction, and growth and survival rate are all significant factors determining the recruitment and

abundance of fish stocks as they form an essential food for larval, juvenile and adult fishes (Beaugrand et al., 2004). Zooplankton grazing in the marine environment controls the primary Production and helps in determining the pelagic ecosystem (Banse, 1995). Through grazing in surface waters and following the production of sinking faecal matters and also by the active transportation of dissolved and particulate matter to deeper waters via vertical migration, they help in the transport of organic carbon to deep ocean layers and thus act as key drivers of 'biological pump' in the marine ecosystem. Zooplankton grazing and metabolism also, transform particulate organic matter into dissolved forms, promoting primary producer community, microbial demineralization, and particle export to the ocean's interior.

The categorisation of zooplankton into various ecological groups is based on several factors such as duration of planktonic life, size, food preferences and habitat. As they vary significantly in size from microscopic to metazoic forms, the classification of zooplankton based on size has paramount importance in the field of quantitative plankton research.

Based on the duration of planktonic life, zooplankton are categorised into Holoplankton (organisms which complete their entire lifecycle as plankton) and Meroplankton (organisms which are planktonic during the early part of their lives such as the larval stages of benthic and nektonic organisms). Tychoplankton are organisms which live a brief planktonic life, such as the benthic crustaceans (Cumaceans, mysids, isopods) which ascend to the water column at night for feeding and certain ectoparasitic copepods, they leave the host and spend their life as plankton during their breeding cycle.


Zooplankton can be subdivided into holoplankton, i.e., permanent members of the plankton (e.g., Calanoid copepods), and meroplankton, i.e., temporary members in the plankton e.g., larvae of fish, shrimp, and crab). The meroplankton group consists of larval and young stages of animals that will adopt a different lifestyle once they mature. In contrast to phytoplankton which consist of a relatively smaller variety of organisms, Zooplankton are extremely divers, consist of a host of larval and adult forms representing many animal phylum.

Among the zooplankton one group always dominate than others; members of sub class copepods (Phylum Athropoda) and Tintinids (Phylum Protozoa) among the net planktons. These small animals are of vital importance in marine ecosystem as one of the primary herbivores animals in the sea, and it is they provide vital link between primary producer (autotrophs) and numerous small and large marine consumers.

As their community structure and function are highly susceptible to changes in the environmental conditions regular monitoring of their distribution as well as their interactions with various physicochemical parameters is inevitable for the sustainable management of the ecosystem (Kusum et al., 2014). Of all the marine zooplankton groups, copepods mainly Calanoid copepods are the

dominant groups in marine subtropical and tropical waters and exhibit considerable diversity in morphology and habitats they occupy (Madhupratap, 1991 ;)

It has been well established that potential of pelagic fishes viz. finfishes, crustaceans, molluscs and marine mammals either directly or indirectly depend on zooplankton. The herbivorous zooplanktons are efficient grazers of the phytoplankton and are referred to as living machines transforming plant material into animal tissue. Hence they play an essential role as the intermediaries for nutrients/energy transfer between primary and tertiary trophic levels. Due to their large density, shorter lifespan, drifting nature, high group/species diversity and different tolerance to the stress, they used as the indicator organisms for the physical, chemical and biological processes in the aquatic ecosystem (Ghajibhiye, 2002).



Spatial distribution of Plankton:

A characteristic of plankton population is that they tend to occur in patches, which are varying spatially on a scale of few meters to far as few kilo metres in distance. They also vary in time scale, season as well as vertically in the water column. It is this patchiness and its constant changes in time and spot, that has made it so difficult for plankton biologist to learn about the ecology of plankton. The biological factors that causes this patchiness is due to the ability of zooplankton to migrate vertically and graze out the phytoplankton at a rapid rate that can create patchiness. Similarly the active swimming ability by certain zooplankton organisms can cause to aggregate in dense group.

DCPL/DPA/21-22/31– November-2022

At its most extreme, because the water in which plankton is suspended is constantly moving, each sample taken by the plankton biologists remain a different volume of water, so each sample is unique and replicate does not exist.

Plankton in the month of November also exhibit vertical patchiness. Physical factors contribute to this type of patchiness include light intensity, nutrients and density gradients in the water column.

Phytoplankton in particular tends to be unequally distributed vertically, which leads to the existence of different concentration of a chlorophyll value between photic zone and below the photic zone.

Methodology adopted for Plankton sampling:

Preservation and storage:

Both filtered plankton and those collected from the plankton net were preserved with 5% buffered formalin and stored in 1L plastic container for further processing in the laboratory.

Sample concentration:

The collected plankton samples were concentrated by using centrifuge and made up to 50 ml with 5% formalin -Glycerine mixture.

Taxonomic evaluation:

Before processing, the sample was mixed carefully and a subsample was taken with a calibrated Stempel-pipette. 1 ml of the concentrated plankton samples were transferred on a glass slide with automatic pipette. The plankton sample on the glass slides were stained by using Lugol's iodine and added glycerin to avoid drying while observation. The plankton samples were identified by using Labex triangular Research microscope with photographic attachment. Microphotographs of the plankton samples were taken for record as well as for confirming the identification. The bigger sized zooplankton was observed through dissecting stereomicroscope with magnification of 20-30 x. Plankton organisms in the whole slide were identified to the lowest taxon possible. A thorough literature search was conducted for the identification of the different groups of phytoplankton and zooplankton that were encountered

Cell counts by drop count method:

The common glass slide mounted with a 1ml of concentrated phytoplankton/zooplankton sample in glycerol and covered with cover slip 22 mm x 60 mm was placed under the compound microscope provided with a mechanical stage. The plankton was then counted from the microscopic field of the left top corner of the slide. Then slide is moved horizontally along the right side and plankton in each microscopic field was thus counted. When first microscopic field row was finished the next consecutive row was adjusted using the mechanical device of the stage. In this way all the plankton present in entire microscopic field are counted. From this total number in 1ml of the concentrated plankton, total amount of phytoplankton in the original volume of sample filtered was calculated as units/L and Zooplankton as N/m^3 .

BENTHIC ORGANISMS:

Benthos is those organisms that are associated with the sea bed or benthic habitats. Epi- benthic organisms live attached to a hard substratum or rooted to a shallow depth below the surface. In fauna organisms live below the sediment–water interface. Interstitial organisms live and move in pore water among sedimentary grains.

Because the benthic organisms are often collected and separated on sieves, a classification based on the overall size is used. Macro benthos include organisms whose shortest dimension is greater than or equal to 0.5 mm. Meio benthos are smaller than 0.5mm but larger than 42 μ in size.

The terms such as macro fauna and Meio fauna generally have little relevance with taxonomic classification. The terms Meio fauna and macro fauna depend on the size. Meio fauna were considered as good bioassay of community health and rather sensitive indicators of environmental changes

SAMPLING METHODOLOGY ADOPTED FOR SUB TIDAL REGION:

Van veen sampler (0.09m²) was used for sampling bottom sediments. Two sets of sediments were sampled from each location, one for macro fauna and other for Meio fauna. The macro fauna in the sediments were sieved on board to separate out the organisms. The fixation of Meio fauna is normally done by bulk fixation of the sediment sample. The bulk fixation is done by using 10% formalin (Buffered with borate). The organisms were preserved with seawater as diluting agent.

Sample sieving:

Sediments samples were sieved to extract the organisms. Sieving was performed carefully as possible to avoid any damage to the animals. The large portion of the sediment was split in to smaller portions and mixed with sea water in a bucket. The cohesive lumps were broken down by continuous stirring. The disaggregated sediments were then passed through the sieves.

Sample staining:

Sorting of the Meio fauna from the sieve is difficult task especially in the preserved material, because organisms are not easily detectable. To facilitate the animal detection the entire sample retained on the sieve after sieving operation were stained by immersing the sieve in a flat bottom tub with 1% Rose Bengal stain; a protein stain. A staining period of 10-30 minutes is sufficient for sample detection.

DIVERSITY INDICES:

On the whole, diversity indices provide more information about community composition than simply species richness (number of species present); they also, take the relative abundances of different species into account. Based on this fact, diversity indices therefore depend not only on species richness but on the evenness, or equitability, with which individuals are distributed among the different species (Magurram, A. E. (1988)

A diversity index is a measure of species diversity within a community that consists of co-occurring populations of several (two or more) different species. It includes two components: richness and evenness. Richness is the measure of the number of different species within a sample showing that more the types of species in a community, the higher is the diversity or greater is the richness. Evenness is the measure of relative abundance of the different species within a community.

The basic idea of diversity index is to obtain a quantitative estimate of biological variability that can be used to compare biological entities composed of discrete components in space and time (Carol H. R. *et al.* 1998). Biodiversity is commonly expressed through indices based on species richness and species abundances (Whittaker 1972, Lande 1996, Purvis and Hector 2000). Biodiversity indices are a non-parametric tool used to describe the relationship between species number and abundance. The most widely used bio diversity indices are Shannon Weiner index and Simpson's index.

A diversity Index is a single statistic that incorporates information on richness and evenness. Any study intended to interpret causes and effect of adverse impact on Biodiversity of communities require suitable measures to evaluate specie richness and Diversity. The former is number of species in community, while latter is a function of relative frequency of different species. Species richness is the iconic measure of biological diversity (Magurran, 2004). Several indices have been created to measure the diversity of species; however, the most widely used in the last decades are the Shannon (1948) and Simpson (1949) (Buzas and Hayek 1996; Gorelick 2006), with the components of diversity: richness (S) and evenness (J)

Simpson's diversity index

Simpson's index (D) is a measure of diversity, which takes into account both species richness, and evenness of abundance among the species present. The Simpson index is one of the meaningful and robust biodiversity measures available. (Magurran, 2004).

The formula for calculating D is presented as:

$$D = \frac{\sum n_i(n_i - 1)}{N(N - 1)}$$

Where n_i = the total number of organisms of each individual species

N = the total number of organisms of all species

The value of D ranges from 0 to 1. With this index, 0 represents infinite diversity and, 1, no diversity. When D increases diversity decreases. Simpson's index is therefore usually expressed as $1-D$ or $1/D$. (Magurran, 2004)

Low species diversity suggests:

- relatively few successful species in the habitat
- the environment is quite stressful with relatively few ecological niches and only a few organisms are really well adapted to that environment

DCPL/DPA/21-22/31– November-2022

- food webs which are relatively simple
- change in the environment would probably have quite serious effects

High species diversity suggests:

- a greater number of successful species and a more stable ecosystem
- more ecological niches are available and the environment is less likely to be hostile complex food webs
- environmental change is less likely to be damaging to the ecosystem as a whole

Species richness indices

The species richness(S) is simply the number of species present in an ecosystem. Species richness Indices of species richness are widely used to quantify or monitor the effects of anthropogenic disturbance. A decline in species richness in may be concomitant with severe or chronic human-induced perturbation (Fair Fair weather 1990) Species richness measures have traditionally been the mainstay in assessing the effects of environmental degradation on the biodiversity of natural assemblages of organisms (Clarke & Warwick, 2001)

Species richness is the iconic measure of biological diversity (Magurran, 2004). The species richness(S) is simply the number of species present in an ecosystem. This index makes no use of relative abundances. The term species richness was coined by Mc Intosh (1967) and oldest and most intuitive measure of biological diversity (Magurran, 2004).

Margalef's diversity index is a species richness index. Margalef's Species richness index (d), or indices that describe the evenness of the distribution of the numbers of individuals among species, were derived.

The value of a diversity index increases both when the number of types increases and when evenness increases. For a given number of types, the value of diversity index is maximised when all types are equally abundant [Rosenzweig, M. L. (1995)]

Shannon-Wiener's index:

An index of diversity commonly used in plankton community analyses is the Shannon-Wiener's index (H), which emphasizes not only the number of species (richness or variety), but also the apportionment of the numbers of individuals among the species (Odum 1971 and Reish 1984). Shannon-Wiener's index (H) reproduces community parameters to a single number by using an equation.

Shannon and Weiner index represents entropy. It is a diversity index taking into account the number of individuals as well as the number of taxon. It varies from 0 for communities with only single taxa to high values for community with many taxon each with few individuals. This index can also determine the pollution status of a water body. Normal values range from 0 to 4. This index is a combination of species present and the evenness of the species. Examining the diversity in the range

of polluted and unpolluted ecosystems, Wilham and Dorris (1968) concluded that the values of the index greater than

3 indicate clean water, values in the range of 1 to 3 are characterized by moderate pollution and values less than 1 are characterized as heavily polluted

10.2:- RESULTS:

CHLOROPHYLL-a:

$$H' = - \sum_{j=1}^s \frac{n_j}{N} \ln \left(\frac{n_j}{N} \right)$$

In the sub surface water chlorophyll-a was varying from 0.472-0.969 mg/m³ with an average value 0.645 mg/m³ in harbour region of DPA in Kandla Creek during sampling done in spring tide period of November 2022. In the nearby creeks chlorophyll-a was varying from 0.359-0.717 mg/m³ with an average value 0.552 mg/m³ Pheophytin –a level was below detectable limit- the all the sampling stations during springtide. Even though the plankton diversity and abundance were more during the spring tide sampling, the chlorophyll-content was detected lesser than expected because, the phytoplankton communities were mainly represented by diatoms *Skeletonema* sp. *Coscinodiscus* sp. and *Chaetoceros* sp.

In the sub surface water chlorophyll-a was varying from 0.338-0.547 mg/m³ with an average value 0.437 mg/m³ in harbour region of DPA in Kandla Creek during sampling done in Neap tide period of November 2022. In the nearby creeks chlorophyll-a was varying from 0.205- 0.440mg/m³ with an average value 0.370 mg/m³. Pheophytin–a level was below detectable limit- the all the sampling stations. During neap tide sampling phytoplankton communities were mainly represented by *Coscinodiscus* sp. and *Ditylum* sp.

In the sub surface water chlorophyll-a was varying from 0.598-0.968 mg/m³ in harbour region of DPA OOT in path finder Creek during sampling done in spring tide period of November 2022. In the sub surface water chlorophyll-a was varying from 0.709 - 0.987mg/m³ in harbour region of DPA OOT in path finder Creek during sampling done in Neap Tide period of November 2022

TABLE:-45 VARIATIONS IN CHLOROPHYLL-a PHEOPHYTIN-a AND ALGAL BIOMASS FROM SAMPLING STATIONS IN DPA HARBOUR AREA IN KANDLA CREEK ,NEAR BY CREEKS AND DPA OOT JETTY IN PATH FINDER CREEK AND SPM NEAR VADINAR DURING SPRING TIDE IN NOVEMBER 2022

Sr. No.	Station	Tide	Chlorophyll-a (mg/m ³)	Pheophytin- a (mg/m ³)	Algal Biomass (Chlorophyll method) mg/m ³
DPA HARBOUR AREA KANDLA CREEK					
1	KPT1	High tide	0.969	BDL	64.92
		Low tide	0.647	BDL	43.35
2	KPT 2	High tide	0.511	BDL	34.24
		Low tide	0.521	BDL	34.91
3	KPT 3	High tide	0.749	BDL	50.18
		Low tide	0.472	BDL	31.62
CREEKS					
4	KPT-4 Khor-I	High tide	0.638	BDL	42.75
		Low tide	0.359	BDL	24.05
5	KPT-5 Nakti-I	High tide	0.717	BDL	48.04
		Low tide	0.493	BDL	33.03
6	KPT-6 Nakti-II	High tide	ND	ND	ND
PATHFINDER CREEK VADINAR					
7	VADINAR-I jetty	High tide	0.968	BDL	64.86
8		Low tide	0.732	BDL	49.04
9	SPM	High tide	0.953	BDL	63.85
10		Low tide	0.598	BDL	

BDL: Below Detectable Limit., ND: Not detected

TABLE:-46. VARIATIONS IN CHLOROPHYLL-a PHEOPHYTIN-a AND ALGAL BIOMASS FROM SAMPLING STATIONS IN DPA HARBOUR AREA, NEAR BY CREEKS AND DPA OOT JETTY IN PATH FINDER CREEK AND SPM NEAR VADINAR DURING NEAP TIDE IN NOVEMBER 2022

Sr.No.	Station	Tide	Chlorophyll-a (mg/m ³)	Pheophytin- a (mg/m ³)	Algal Biomass (Chlorophyll method) mg/m ³
DPA HARBOUR AREA KANDLA CREEK					
1	KPT1	High tide	0.547	BDL	
		Low tide	0.450	BDL	
2	KPT 2	High tide	0.338	BDL	
		Low tide	0.409	BDL	
3	KPT 3	High tide	0.354	BDL	
		Low tide	0.523	BDL	
CREEKS					
4	KPT-4 Khor-I	High tide	0.440	BDL	
		Low tide	0.408	BDL	
5	KPT-5 Nakti-I	High tide	0.205	BDL	
		Low tide	0.426	BDL	
6	KPT-6 Nakti-II	High tide	ND	ND	ND
PATHFINDER CREEK VADINAR					
7	VADINAR-I jetty	High tide	0.799	BDL	
8		Low tide	0.709	BDL	
9	SPM	High tide	0.857	BDL	
10		Low tide	0.987	BDL	

BDL: Below Detectable Limit. ND: Not detected

PHYTOPLANKTON POPULATION:

For the evaluation of the Phytoplankton population in DPA harbour area and within the immediate surroundings of the port, sampling was conducted from 5 sampling locations (3 in harbour area and two in Nakti creek) during high tide period and low tide period of spring tide and neap tide.

The phytoplankton community of the sub surface water in the harbour and nearby creeks was represented by, Diatoms, blue green algae and Dinoflagellates during spring tide period. Diatoms were represented by 26 genera, Blue green algae were represented by 2 genera and Dinoflagellates were represented by 6 genera during the sampling conducted in spring tide in November, 2022. Phytoplankton of the sampling stations at sub surface layer in the harbour area and nearby creeks was varying from 39-243units/ L during high tide period and 115-199 units/L during low tide of Spring Tide. During spring tide sampling phytoplankton communities were dominated by *Skeletonema* sp almost forming a bloom in the Kandla creek and other nearby creek area and abundant population of *Coscinodiscus* sp. and *Chaetoceros* sp.

The phytoplankton community of the sub surface water in the harbour and nearby creeks was represented by Diatoms, Blue green algae and Dinoflagellates during Neap tide period. Diatoms were represented by 24 genera, Blue green algae were represented 2 genera and Dinoflagellates with 5 genera during the sampling conducted in Neap tide in November, 2022. Phytoplankton of the sampling stations at sub surface layer in the harbour area and nearby creeks was varying from 43-299 units/ L during high tide period and 143-193 units/L during low tide of Neap Tide. During Neap tide sampling phytoplankton communities were dominated by, *Ditylum* sp and *Coscinodiscus* sp.

For the evaluation of the Phytoplankton population in DPA OOT jetty area in Path Finder creek sampling was conducted from two sampling locations; Jetty area and SPM area during high tide period and low tide of spring tide and Neap tide period.

The phytoplankton community of the sub surface water in the path finder creeks was represented by Diatoms, Blue green algae and Dinoflagellates during spring tide period. Diatoms were represented by 25 genera, Blue Green algae by 5 genera and Dinoflagellates by 6 genera during the sampling conducted in spring tide in November, 2022. Phytoplankton of the sampling stations at sub surface path finder creek near OOT Jetty area was 209 units/L during high tide period and 177 units/L during low tide of Spring Tide. Phytoplankton of the sampling stations at sub surface layer in the SPM area was varying from 206 units/ L during high tide period and 131 units/ L during low tide of Spring Tide.

The phytoplankton community of the sub surface water in the path finder creeks was represented by Diatoms, Blue green and Dinoflagellates during Neap tide period. Diatoms were represented by 32 genera and Blue green algae by 4 genera and Dinoflagellates by 6 genera during the sampling conducted in Neap tide in November, 2022. Phytoplankton of the sampling stations at sub surface path finder creek near OOT Jetty was varying from 244units/ L during high tide period and 200

units/L during low tide of Neap Tide. Phytoplankton of the sampling stations at sub surface path finder creek near SPM area was varying from 259 units/L during high tide period and 294 units/L during low tide of Neap Tide.

Species Richness Indices and Diversity Indices:

Margalef's diversity index (Species Richness)

Margalef's diversity index (Species Richness) of phytoplankton communities in the Kandla creek and nearby creeks sampling stations was varying from 2.184- 4.688 with an average of 3.346 during the sampling conducted in High tide period of spring tide. While Margalef's diversity index (Species Richness) S of phytoplankton communities in the Kandla creek region and nearby creeks was varying from 1.963- 3.589 with an average of 2.835 during the consecutive low tide period.

Margalef's diversity index (Species Richness) of phytoplankton communities in the stations in Kandla creek and nearby creeks was varying from 2.393-4.279 with an average of 3.586 during the sampling conducted in High tide period of Neap tide. While Margalef's diversity index (Species Richness) of phytoplankton communities in the Kandla creek region and nearby creeks was varying from 2.821- 3.86 with an average of 3.357 during consecutive low tide.

Margalef's diversity index (Species Richness) S of phytoplankton communities in the stations was 4.867 at OOT jetty area and 4.129 at SPM area during the sampling conducted in High tide period of spring tide. While Margalef's diversity index (Species Richness) S of phytoplankton communities in the path finder creek near OOT jetty was 4.443 and 3.692 at SPM during the consecutive low tide period.

Margalef's diversity index (Species Richness) of phytoplankton communities in the stations was 4.73 at OOT jetty area and 4.139 at SPM area during the sampling conducted in High tide period of Neap tide. While Margalef's diversity index (Species Richness) of phytoplankton communities in the path finder creek near OOT jetty was 4.152 and SPM area was 5.454 during the consecutive low tide period.

Shannon-Wiener's index:

Shannon-Wiener's Index (H) of phytoplankton communities in the sampling stations was in the range of 0.786- 1.034 between selected sampling stations with an average value of 0.925 during high tide period of spring tide at Kandla creek and nearby creeks. Shannon-Wiener's Index (H) of phytoplankton communities in the sampling stations was in the range of 0.790-0.915 between selected sampling stations with an average value of 0.855 during consecutive low tide at Kandla creek and nearby creeks.

Shannon-Wiener's Index (H) of phytoplankton communities in the sampling stations was in the range of 0.867–1.022 between selected sampling stations with an average value of 0.932 during high tide period of neap tide at Kandla creek and nearby creeks. Shannon-Wiener's Index (H) of phytoplankton

communities in the sampling stations was in the range of 0.926- 1.001 between selected sampling stations with an average value of 0.951 during consecutive low tide at Kandla creek and nearby creeks. Shannon-Wiener's Index (H) of phytoplankton communities in the stations was 1.037 at OOT jetty area and 0.946 at SPM area during the sampling conducted in High tide period of spring tide. While Shannon-Wiener's Index (H) of phytoplankton communities in the path finder creek near OOT jetty was 1.043 and 0.982 at SPM during the consecutive low tide period of spring tide.

Shannon-Wiener's Index (H) of phytoplankton communities in the stations was 0.998 at OOT jetty area and 1.035 at SPM area during the sampling conducted in High tide period of Neap tide. While Shannon-Wiener's Index (H) of phytoplankton communities in the path finder creek near OOT jetty was 0.942 and at SPM area was 1.036 during the consecutive low tide period.

Typical values are generally between 1.5 and 3.5 in most ecological studies, and the index is rarely greater than 4. The Shannon-Wiener's index increases as both the richness and the evenness of the community increase. This result indicates that diversity of phytoplankton of Kandla Harbour region and nearby creeks is less but with abundant population of few, with relatively few ecological niches and only very few opportunist organisms are really well adapted to this environment and thrive better than other species.

Simpson's diversity index:

Simpson diversity index (1-D) of phytoplankton communities was below 0.9 at all sampling stations in the Kandla Harbour region and nearby creeks, which was varying from 0.778-0.851 between selected sampling stations with an average of 0.823 during high tide period of spring tide. Simpson diversity index (1-D) of phytoplankton communities was below 0.9 at all sampling stations in the Kandla Harbour region and nearby creeks except few, which was varying from 0.787-0.842 between selected sampling stations with an average of 0.814 during consecutive low tide.

Simpson diversity index (1-D) of phytoplankton communities was below 0.9 at all sampling stations except few in Kandla Harbour region and nearby creeks, during high tide period and low tide period during Neap tide also, which was varying from 0.813-0.874 with an average value of 0.847 between selected sampling stations during high tide period and 0.840-0.871 varying from with an average value of 0.858 between selected sampling stations during consecutive low tide period. Low species diversity suggests a relatively few successful species in this habitat.

Simpson diversity index (1-D) of phytoplankton communities in the stations was 0.863 at OOT jetty area and 0.820 at SPM area during the sampling conducted in High tide period of spring tide at Path finder creek. While Simpson diversity index (1-D) of phytoplankton communities in the path finder creek near OOT jetty was 0.876 and 0.867 at SPM during the consecutive low tide period in the path finder creek.

Simpson diversity index (1-D) of phytoplankton communities in the stations was 0.838 at OOT jetty area and 0.881 at SPM area during the sampling conducted in High tide period of Neap tide at Path

finder Creek. While Simpson diversity index (1-D) of phytoplankton communities in the path finder creek near OOT jetty was 0.832 and at SPM area was 0.867 during the consecutive low tide period.

Table:-47 4PHYTOPLANKTON VARIATIONS IN ABUNDANCE AND DIVERSITY IN SUB SURFACE SAMPLING STATIONS IN DPA HARBOUR AREA AT KANDLA CREEK AND , NEAR BY CREEKS DURING SPRING TIDE IN NOVEMBER 2022

Tide	Sampling Station	Abundance In units/L	No of Species observed /total species	% Of diversity	Margalef's diversity index (Species Richness)	Shannon Weiner index H (log ₁₀)	Diversity Index (Simpson's Index) 1-D
HIGH TIDE	1	207	26/34	76.47	4.688	1.034	0.8511
	2	183	22/34	64.71	4.031	1.005	0.8437
	3	193	13/34	38.24	2.28	0.811	0.7778
	4	243	18/34	52.94	3.095	0.9391	0.8192
	5	193	21/34	61.76	3.8	0.9777	0.8281
	6	39	9/34	26.47	2.184	0.786	0.8178
LOW TIDE	1	178	14/34	41.18	2.509	0.8042	0.787
	2	199	20/34	58.82	3.589	0.8982	0.8075
	3	115	14/34	41.18	2.74	0.8696	0.8365
	4	154	18/34	52.94	3.375	0.915	0.8416
	5	163	11/34	32.35	1.963	0.7895	0.7957

Table:-48 PHYTOPLANKTON VARIATIONS IN ABUNDANCE AND DIVERSITY IN SUB SURFACE SAMPLING STATIONS IN DPA HARBOUR AREA AT KANDLA CREEK AND NEAR BY CREEKS DURING NEAP TIDE IN NOVEMBER 2022

Tide	Sampling Station	Abundance In units/L	No of Species observed /total species	% of diversity	Margalef's diversity index (Species Richness)	Shannon Weiner index H (log ₁₀)	Diversity Index (Simpson's Index) 1-D
HIGH TIDE	1	216	24/31	77.42	4.279	0.98	0.8568
	2	229	22/31	70.97	3.865	0.958	0.853
	3	228	22/31	70.97	3.868	1.022	0.8743
	4	299	23/31	74.19	3.859	0.8667	0.8127
	5	254	19/31	61.29	3.251	0.8929	0.8307
	6	43	10/31	32.26	2.393	0.8712	0.8571
LOW TIDE	1	183	18/31	58.06	3.263	0.9504	0.8636
	2	143	15/31	48.39	2.821	0.946	0.8666
	3	178	21/31	67.74	3.86	1.001	0.8708
	4	193	19/31	61.29	3.42	0.931	0.84
	5	193	19/31	61.29	3.42	0.9259	0.8469

Table:-49 ABUNDANCE OF PHYTOPLANKTON SUBSURFACE SAMPLING STATIONS IN DPA HARBOUR AREA AT KANDLA CREEK AND, NEAR BY CREEKS DURING SPRING TIDE IN NOVEMBER2022

Tide	Surface	No of Sampling location	Group of phytoplankton	Phytoplankton Group range Units/L	Genera or species /total Phyto plankton	Species Composition % (Group level)
HIGH TIDE	Sub surface	6	BLUE GREEN ALGAE	0-8	2/34	5.88
			DIATOMS	38-238	26/34	76.47
			DINOFLAGELLATES	0-11	6/34	17.65
			TOTAL PHYTO PLANKTON	39-243	34	
LOW TIDE	Sub surface	5	BLUE GREEN ALGAE	1-6	2/34	5.88
			DIATOMS	110-190	26/34	76.47
			DINOFLAGELLATES	1-7	6/34	17.65
			TOTAL PHYTO PLANKTON	115-199	34	

TABLE:-50 ABUNDANCE OF PHYTOPLANKTON SUBSURFACE SAMPLING STATIONS IN DPA HARBOUR AREA AT KANDLA CREEK AND, NEAR BY CREEKS DURING NEAP TIDE IN NOVEMBER 2022

Tide	Surface	No of Sampling location	Group of phytoplankton	Phytoplankton Group range Units/L	Genera or species /total Phyto plankton	Species Composition % (Group level)
HIGH TIDE	Sub surface	6	BLUE GREEN ALGAE	0-6	2/31	6.45
			DIATOMS	43-293	24/31	77.42
			DINOFLAGELLATES	0-9	5/31	16.13
			TOTAL PHYTO PLANKTON	43-299	31	
LOW TIDE	Sub surface	5	BLUE GREEN ALGAE	2-6	2/31	6.45
			DIATOMS	133-186	24/31	77.42
			DINOFLAGELLATES	3-8	5/31	16.13
			TOTAL PHYTO PLANKTON	143-193	31	

TABLE:-51 PHYTOPLANKTON VARIATIONS IN ABUNDANCE AND DIVERSITY IN SUB SURFACE SAMPLING STATIONS IN DPA OOT AT PATH FINDER CREEK , VADINAR & NEAR BY SPM, DURING SPRING TIDE IN NOVEMBER 2022

Tide	Sampling Station	Abundance In units/L	No of Species observed /total species	% of diversity	Margalef's diversity index (Species Richness S)	Shannon Weiner index H (log ₁₀)	Diversity Index (Simpson's Index) 1-D
HIGH TIDE	Jetty	209	27/36	75.00	4.867	1.037	0.863
	SPM	206	23/36	63.89	4.129	0.946	0.820
LOW TIDE	Jetty	177	24/36	66.67	4.443	1.043	0.876
	SPM	131	19/36	52.78	3.692	0.982	0.867

TABLE:-52 PHYTOPLANKTON VARIATIONS IN ABUNDANCE AND DIVERSITY IN SUB SURFACE SAMPLING STATIONS IN DPA OOT AT PATH FINDER CREEK , VADINAR & NEAR BY SPM, DURING NEAP TIDE IN NOVEMBER 2022

Tide	Sampling Station	Abundance In units/L	No of Species observed /total species	% of diversity	Margalef's diversity index (Species Richness)	Shannon Weiner index H (log ₁₀)	Diversity Index (Simpson's Index) 1-D
HIGH TIDE	Jetty	244	27/42	64.29	4.73	0.998	0.838
	SPM	259	24/42	57.14	4.139	1.035	0.881
LOW TIDE	Jetty	200	23/42	54.76	4.152	0.942	0.832
	SPM	294	32/42	76.19	5.454	1.036	0.867


TABLE:-53 ABUNDANCE OF PHYTOPLANKTON SUBSURFACE SAMPLING STATIONS IN DPAOOT AT PATH FINDER CREEK, VADINAR & NEAR BY SPM, DURING SPRING TIDE IN NOVEMBER 2022

Tide	Surface	No of Sampling location	Group of phytoplankton	Phytoplankton Group range Units/L	Genera or species /total Phyto plankton	Taxon Diversity % (Group level)
HIGH TIDE	Sub surface	2	BLUE GREEN ALGAE	14-20	5/36	13.89
			DIATOMS	180-192	25/36	69.44
			DINOFLAGELLATES	3-6	6/36	16.67
			TOTAL PHYTO PLANKTON	206-209	36	
LOW TIDE	Sub surface	2	BLUE GREEN ALGAE	12-19	5/36	13.89
			DIATOMS	118-156	25/36	69.44
			DINOFLAGELLATES	1-2	6/36	16.67
			TOTAL PHYTO PLANKTON	131-177	36	


Table:- 54 ABUNDANCE OF PHYTOPLANKTON SUBSURFACE SAMPLING STATIONS IN DPA OOT AT PATH FINDER CREEK , VADINAR & NEAR BY SPM, DURING NEAP TIDE IN NOVEMBER 2022

Tide	Surface	No of Sampling location	Group of phytoplankton	Phytoplankton Group range Units/L	Genera or species /total Phyto plankton	Species Composition % (Group level)
HIGH TIDE	Sub surface	2	BLUE GREEN ALGAE	5-7	4/42	9.52
			DIATOMS	238-248	32/42	76.19
			DINOFLAGELLATES	1-4	6/42	14.29
			TOTAL PHYTO PLANKTON	244-259		
LOW TIDE	Sub surface	2	BLUE GREEN ALGAE	4-8	4/42	9.52
			DIATOMS	194-282	32/42	76.19
			DINOFLAGELLATES	2-4	6/42	14.29
			TOTAL PHYTO PLANKTON	200-294		


Species Composition % of Phytoplankton during High tide and Low tide period during spring tide in Kandla creek and nearby creeks




Species Composition % of Phytoplankton during High tide and Low tide period during Neap tide in Kandla creek and nearby creeks



Species Composition % of Phytoplankton during High tide and Low tide period during spring tide in Path Finder Creek, Vadinar



Species Composition % of Phytoplankton during High tide and Low tide period during Neap tide in Path Finder Creek, Vadinar



ZOOPLANKTON POPULATION:

For the evaluation of the Zooplankton population in DPA harbour area and within the immediate surroundings of the port sampling was conducted from 6 sampling locations (3 in harbour area and two in Nakti creek and one in Khoricreek) during high tide period and low tide period of spring tide and Neap tide in November, 2022. The Zooplankton community of the sub surface water in the harbour and nearby creeks during spring tide was represented by mainly six groups; Tintinnids, Copepods, Arrow worms, Mysids, Urochordata, Ciliates and 8 larval forms. The Zooplankton community of the sub surface water in the harbour and nearby creeks during neap tide was represented by mainly six groups; Tintinnids, Copepods, Arrow worms, Mysids, Urochordata, Ciliates and 6 larval forms.

Zooplankton of the sampling stations at sub surface layer in the DPA harbour area and nearby creek was varying from $25-128 \times 10^3 \text{ N/m}^3$ during high tide and $103-144 \times 10^3 \text{ N/m}^3$ during low tide of Spring Tide period. Zooplankton of the sampling stations at sub surface layer in the DPA harbour area and nearby creek was varying from $19-114 \times 10^3 \text{ N/m}^3$ during high tide and $76-106 \times 10^3 \text{ N/m}^3$ during low tide of Neap Tide period.

For the evaluation of the Zooplankton population in DPA OOT jetty area in Path Finder creek and SPM in Vadinar selected 2 sampling locations (1 in jetty area and one near SPM).

During spring tide sampling plankton sample were collected at Jetty area and near SPM during consecutive high tide period and low tide period. During Neap tide sampling Plankton samples were collected from jetty area and SPM during consecutive high tide period and low tide period.

The Zooplankton community of the sub surface water in the path finder creek during spring tide was represented by mainly four groups Tintinnids, Copepods, Urochordata, Ciliates and 4 larval forms. While the Zooplankton community of the sub surface water in the path Finder creeks at Jetty region and SPM during neap tide was represented by four groups, Tintinnids, Copepods, Arrow worms, Urochordata and 5 larval forms.

Zooplankton of the sampling stations at sub surface layer in the DPA OOT Jetty area of path finder creek was $91 \times 10^3 \text{ N/m}^3$ during high tide and $86 \times 10^3 \text{ N/m}^3$ during low tide of Spring Tide period. Zooplankton of the sampling stations at sub surface layer in the DPA SPM area of path finder creek was $101 \times 10^3 \text{ N/m}^3$ during high tide and $70 \times 10^3 \text{ N/m}^3$ during low tide of spring Tide period.

Zooplankton of the sampling stations at sub surface layer in the DPA OOT jetty area in path finder creek was recorded $87 \times 10^3 \text{ N/m}^3$ during high tide and $65 \times 10^3 \text{ N/m}^3$ during consecutive low tide period of Neap tide. Zooplankton of the sampling stations at sub surface layer in the DPASPM area in path finder creek was recorded $64 \times 10^3 \text{ N/m}^3$ during high tide and $87 \times 10^3 \text{ N/m}^3$ during consecutive low tide period of Neap Tide.

Species Richness Indices and Diversity Indices:

Margalef's diversity index (Species Richness)

Margalef's diversity index (Species Richness) of Zooplankton communities in the stations Kandla creek region and nearby creeks was varying from 2.175- 5.186 with an average of 3.450 during the sampling conducted in High tide period. Margalef's diversity index (Species Richness) of Zooplankton communities varying from 2.373-3.823 with an average of 3.261 during the sampling conducted in low tide period during Spring tide.

Margalef's diversity index (Species Richness) of Zooplankton communities in the Kandla creek region and nearby creeks sampling stations were varying from 1.358-3.858 with an average of 2.930 during the sampling conducted in high tide and varying from 2.289- 4.618 with an average of 3.513 during the sampling conducted in low tide during Neap tide period.

Margalef's diversity index (Species Richness) of Zooplankton communities in the sampling station near jetty at Path Finder Creek, Vadinar during the sampling conducted in consecutive high tide period and low tide of spring tide was recorded as 1.995 and 1.796 respectively. Margalef's diversity index (Species Richness) of Zooplankton communities in the sampling station near SPM at Path Finder Creek, Vadinar during the sampling conducted in consecutive high tide period and low tide of spring tide was recorded as 2.600 and 2.118 respectively.

Margalef's diversity index (Species Richness) of Zooplankton communities near Jetty at Path finder creek were varying from 3.807 and 2.396 respectively during the sampling conducted in consecutive high tide period and Low tide period of Neap tide. While Margalef's diversity index (Species Richness) of Zooplankton communities near SPM at Path finder creek were varying from 2.645-3.135 respectively during the consecutive high tide and low tide period.

Shannon-Wiener's index:

Shannon-Wiener's Index (H) of Zooplankton communities in the sampling stations in Kandla Harbour region and nearby creeks was in the range of 0.778-1.164 between selected sampling stations with an average value of 0.939 during high tide period of spring tide. Shannon-Wiener's Index (H) of Zooplankton communities in the sampling stations in Kandla Harbour region and nearby creeks was in the range of 0.795-1.015 between selected sampling stations with an average value of 0.938 during consecutive low tide period.

Shannon-Wiener's Index (H) of Zooplankton communities in the sampling stations in Kandla Harbour region and nearby creeks was in the range of 0.490-0.914 between selected sampling stations with an average value of 0.805 during high tide period of Neap tide. Shannon-Wiener's Index (H) of Zooplankton communities in the sampling stations in Kandla Harbour region and nearby creeks was in the range 0.797-1.041 of between selected sampling stations with an average value of 0.928 during consecutive low tide period.

Shannon-Wiener's Index (H) of Zooplankton communities in the sampling station near jetty at Path Finder Creek, Vadinar during the sampling conducted in consecutive High tide period and low tide of spring tide was recorded as 0.816-0.793 respectively. Shannon-Wiener's Index (H) of Zooplankton communities in the sampling station near SPM at Path Finder Creek, Vadinar during the sampling conducted in consecutive High tide period and low tide of spring tide was recorded as 0.834-0.808 respectively.

Shannon-Wiener's Index (H) of Zooplankton communities near jetty at Path finder creek was varying from 0.956-0.755 respectively during the sampling conducted consecutive high tide period and low tide period of Neap tide. While Shannon-Wiener's Index (H) of Zooplankton communities near SPM at Path finder creek was varying from 0.775-0.751 during the consecutive high tide and low tide period.

Typical values are generally between 1.5 and 3.5 in most ecological studies, and the index is rarely greater than 4. The Shannon-Wiener's index increases as both the richness and the evenness of the community increase. This result indicates that diversity of Zooplankton of Kandla Harbour region and nearby creeks stations is slightly high with very minimum diverse population but very few opportunist organisms are really well adapted to this environment and thrive better than other species.

Simpson's diversity index:

Simpson diversity index (1-D) of Zooplankton communities was below 0.9 most of sampling stations in the Kandla Harbour region and nearby creeks during high tide and low tide of spring tide period except few stations, which was varying from 0.780-0.909 between selected sampling stations with an average of 0.837 during high tide period and was varying from 0.785- 0.864 with an average value of 0.837 between selected sampling stations during low tide.

Simpson diversity index (1-D) of Zooplankton communities was below 0.9 at all sampling stations in the Kandla Harbour region and nearby creeks during high tide and low tide period of Neap tide except few, which was varying from 0.591-0.827 between selected sampling stations with an average of 0.753 during high tide period and was varying from 0.793-0.852 with an average value of 0.820 between selected sampling stations during consecutive low tide. This species diversity suggests a relatively few successful species in this habitat during November, 2022 sampling.

Simpson diversity index (1-D) of Zooplankton communities in the sampling station near jetty at Path Finder Creek, Vadinar during the sampling conducted in consecutive High tide period and low tide of spring tide was recorded as 0.821 and 0.815 respectively. Simpson diversity index (1-D) of Zooplankton communities in the sampling station near SPM at Path Finder Creek, Vadinar during the sampling conducted in consecutive High tide period and low tide of spring tide was recorded as 0.812 and 0.828 respectively.

Simpson diversity index (1-D) of Zooplankton communities in the sampling station near jetty at Path Finder Creek, Vadinar during the sampling conducted in consecutive High tide period and low tide of Neap tide was recorded as 0.836- 0.766 respectively. Simpson diversity index (1-D) of Zooplankton communities in the sampling station near SPM at Path Finder Creek, Vadinar during the sampling conducted in consecutive High tide period and low tide of spring tide was recorded as 0.768 and 0.719 respectively.

TABLE:-55 ZOOPLANKTON VARIATION IN ABUNDANCE AND DIVERSITY IN SUB SURFACE SAMPLING STATIONS IN DPA HARBOUR AREA AT KANDLA CREEK AND NEAR BY CREEKS DURING SPRING TIDE IN NOVEMBER 2022

Tide	Sampling Station	Abundance In $N \times 10^3 / m^3$	No of Species/groups observed /total species/group	% of diversity	Margalef's diversity index (Species Richness S)	Shannon Weiner index H (\log_{10})	Diversity Index (Simpson's Index) 1-D
HIGH TIDE	1	124	26/33	78.79	5.186	1.164	0.9089
	2	114	18/33	54.55	3.589	0.8655	0.7802
	3	102	16/33	48.48	3.243	0.9207	0.8189
	4	128	17/33	51.52	3.298	0.9062	0.8124
	5	107	16/33	48.48	3.21	0.997	0.8686
	6	25	8/33	24.24	2.175	0.7777	0.83
LOW TIDE	1	117	16/33	48.48	3.15	0.9709	0.8609
	2	144	20/33	60.61	3.823	0.9468	0.8238
	3	121	19/33	57.58	3.753	1.015	0.8639
	4	108	16/33	48.48	3.204	0.9609	0.8505
	5	103	12/33	36.36	2.373	0.7949	0.7853

TABLE:-56 ZOOPLANKTON VARIATIONS IN ABUNDANCE AND DIVERSITY IN SUB SURFACE SAMPLING STATIONS IN DPA HARBOUR AREA AT KANDLA CREEK AND NEAR BY CREEKS DURING NEAP TIDE IN NOVEMBER 2022

Tide	Sampling Station	Abundance In $No \times 10^3 / m^3$	No of Species/groups observed /total species/group	% of diversity	Margalef's diversity index (Species Richness S)	Shannon Weiner index H (\log_{10})	Diversity Index (Simpson's Index) 1-D
HIGH TIDE	1	82	18/32	56.25	3.858	0.9017	0.7814
	2	99	16/32	50.00	3.264	0.9138	0.8273
	3	89	13/32	40.63	2.673	0.8264	0.7763
	4	114	18/32	56.25	3.589	0.8478	0.7645
	5	98	14/32	43.75	2.835	0.8503	0.7766
	6	19	5/32	15.63	1.358	0.4901	0.5906
LOW TIDE	1	79	11/32	34.38	2.289	0.797	0.7932
	2	76	21/32	65.63	4.618	1.041	0.8516
	3	106	21/32	65.63	4.289	1.026	0.8446
	4	90	15/32	46.88	3.111	0.9087	0.8177
	5	100	16/32	50.00	3.257	0.865	0.7939

**Table:-57 ABUNDANCE OF ZOOPLANKTON IN SUBSURFACE SAMPLING STATIONS
IN DPA HARBOUR AREAATKANDLA CREEK AND NEAR BY CREEKS DURING
SPRING TIDE IN NOVEMBER 2022**

Tide	Surface	No of Sampling locations	Group of Zooplankton	Abundance of Zooplankton $\times 10^3 / \text{m}^3$ Group Range	Genera or species /total Zooplankton	Taxon Diversity % (Group level)
HIGH TIDE	Sub surface	6	tintinnids	9-26	11/33	33.33
			Copepods	11-51	9/33	27.27
			Arrow worms	0-1	1/33	3.03
			Mysids	0-2	1/33	3.03
			Urochordata	1-6	2/33	6.06
			Ciliates	0-2	1/33	3.03
			Larval forms	4-50	8/33	24.25
			TOTAL ZOOPLANKTON N/ M^3	25-128	33	
LOW TIDE	Sub surface	5	Tintinnids	18-33	11/33	33.33
			Copepods	37-49	9/33	27.27
			Arrow worms	0-4	1/33	3.03
			Mysids	0-2	1/33	3.03
			Urochordata	0-2	2/33	6.06
			Ciliates	0-2	1/33	3.03
			Larval forms	41-65	8/33	24.25
			TOTAL ZOOPLANKTON N/ M^3	103-144	33	

TABLE:-58 ABUNDANCE OF ZOOPLANKTON IN SUBSURFACE SAMPLING STATIONS IN DPA HARBOUR AREA IN KANDLA CREEK AND, NEAR BY CREEKS DURING NEAP TIDE IN NOVEMBER 2022

Tide	Surface	No of Sampling locations	Group of Zooplankton	Abundance of Zooplankton $\times 10^3 / m^3$ Group Range	Genera or species /total Zooplankton	Taxon Diversity % (Group level)
HIGH TIDE	Sub surface	6	Tintinnids	0-14	10/32	31.25
			Copepods	6-49	10/32	31.25
			Arrow worms	0	1/32	3.13
			Mysids	0-6	2/32	6.25
			Urochordata	0-4	2/32	6.25
			Ciliates	0-2	1/32	3.13
			Larval forms	13-50	6/32	18.74
			TOTAL ZOOPLANKTON N/M ³	19-114	32	
LOW TIDE	Sub surface	5	tintinnids	4-17	10/32	31.25
			Copepods	25-45	10/32	31.25
			Arrow worms	0-2	1/32	3.13
			Mysids	0-6	2/32	6.25
			Urochordata	0-5	2/32	6.25
			Ciliates	0-1	1/32	3.13
			Larval forms	27-47	6/32	18.74
			TOTAL ZOOPLANKTON N/M ³	76-106	32	

Table:-59 ZOOPLANKTON VARIATIONS IN ABUNDANCE AND DIVERSITY IN SUBSURFACE SAMPLING STATIONS IN DPA OOT AREA AT PATH FINDER CREEK AND NEAR BY SPM DURING SPRING TIDE IN NOVEMBER 2022

Tide	Sampling Station	Abundance In $\times 10^3 N / m^3$	No of Species/groups observed /total species/group	% of diversity	Margalef's diversity index (Species Richness S)	Shannon Weiner index H (log ₁₀)	Diversity Index (Simpson's Index) 1-D
HIGH TIDE	Jetty	91	10/20	50.00	1.995	0.816	0.821
	SPM	101	13/20	65.00	2.6	0.834	0.812
LOW TIDE	Jetty	86	9/20	45.00	1.796	0.793	0.815
	SPM	70	10/20	50.00	2.118	0.808	0.828

TABLE:-60 ZOOPLANKTON VARIATION IN ABUNDANCE AND DIVERSITY IN SUB SURFACE SAMPLING STATIONS IN DPA OOT AREA AT PATH FINDER CREEK AND NEAR BY SPM DURING NEAP TIDE IN NOVEMBER 2022

Tide	Sampling Station	Abundance In $N \times 10^3 / m^3$	No of Species/groups observed /total species/group	% of diversity	Margalef's diversity index (Species Richness S)	Shannon Weiner index $H (\log_{10})$	Diversity Index (Simpson's Index) 1-D
HIGH TIDE	Jetty	87	18/21	85.71	3.807	0.956	0.836
	SPM	64	12/21	57.14	2.645	0.775	0.768
LOW TIDE	Jetty	65	11/21	52.38	2.396	0.755	0.766
	SPM	87	15/21	71.43	3.135	0.751	0.719


Table:-61 ABUNDANCE OF ZOOPLANKTON IN SUBSURFACE SAMPLING STATIONS IN DPA OOT AREA AND PATH FINDER CREEK AND NEAR BY SPM DURING SPRING TIDE IN NOVEMBER 2022

Tide	Surface	No of Sampling locations	Group of Zooplankton	Abundance of Zooplankton $\times 10^3 / m^3$ Group Range	Genera or species /total Zooplankton	Taxon Diversity % (Group level)
HIGH TIDE	Sub surface	2	Tintinnids	24-32	5/20	25.00
			Copepods	28-38	8/20	40.00
			Urochordata	1-2	2/20	10.00
			Ciliates	0-1	1/20	5.00
			Larval forms	30-36	4/20	20.00
			TOTAL ZOOPLANKTON	91-101	20	
LOW TIDE	Sub surface	2	Tintinnids	17-21	5/20	25.00
			Copepods	30-37	8/20	40.00
			Urochordata	0	2/20	10.00
			Ciliates	0	1/20	5.00
			Larval forms	19-32	4/20	20.00
			TOTAL ZOOPLANKTON	70-86	20	


TABLE:-62 ABUNDANCE OF ZOOPLANKTON IN SUBSURFACE SAMPLING STATIONS IN DPA OOT AREA AT PATH FINDER CREEK AND NEAR BY SPM DURING NEAP TIDE IN NOVEMBER 2022

Tide	Surface	No of Sampling locations	Group of Zooplankton	Abundance of Zooplankton $\times 10^3 / m^3$ Group Range	Genera or species /total Zooplankton	Taxon Diversity % (Group level)
HIGH TIDE	Sub surface	2	tintinnids	9-16	7/21	33.33
			Copepods	23-34	6/21	28.57
			Arrow worms	0	1/21	4.76
			Urochordata	0-2	2/21	9.52
			Larval forms	32-35	5/21	23.82
			TOTAL ZOOPLANKTON	64-87	21	
LOW TIDE	Sub surface	2	tintinnids	6-9	7/21	33.33
			Copepods	29	6/21	28.57
			Arrow worms	0-1	1/21	4.76
			Urochordata	0-3	2/21	9.52
			Larval forms	27-48	5/21	23.82
			TOTAL ZOOPLANKTON	65-87	21	


**Species Composition % of Zooplankton during High tide and Low tide period of spring tide In
Kandla Creek and nearby Creeks**




**Species Composition % of Zooplankton during High tide and Low tide period of Neap tide In
Kandla Creek and nearby Creeks**



Species Composition % of Zooplankton during High tide and Low tide period of Spring tide In Path Finder Creek and near Jetty



Species Composition % of Zooplankton during High tide and Low tide period of Neap tide In Path Finder Creek near jetty and nearby SPM



**TABLE:-63 SYSTEMATIC ACCOUNT OF PHYTOPLANKTON IN THE SAMPLING
LOCATIONS OF DPA HARBOUR AREA AT KANDLA CREEK AND NEARBY CREEKS
DURING NEAP TIDE OF NOVEMBER 2022**

CLASS	ORDER	FAMILY	GENUS/SPECIES	#	Relative Abundance
Cyanophyceae	Nostocales	Oscillatoriaceae	<i>Oscillatoria sp.</i>	B1	Very sparse
	Oscillatoriales	Phormidiaceae	<i>Planktothrix sp.</i>	B2	Very sparse
Coscinodiscophyceae	Biddulphiales	Biddulphiaceae	<i>Biddulphi</i> sp	D1	Abundant
	Chaetocerotales	Chaetocerotaceae	<i>Bacteriastrum sp</i>	D2	Very sparse
			<i>Chaetoceros sp.</i>	D3	Scattered
	Corethrales	Corethraceae	<i>Corethron sp</i>	D4	Very sparse
	Coscinodiscales	Coscinodiscaceae	<i>Coscinodiscus sp.</i>	D5	Dominant
	Hemiaulales	Bellerocheaceae	<i>Bellerochea sp</i>	D6	Very sparse
		Streptothecaceae	<i>Helicotheca sp</i>	D7	Very sparse
	Rhizosoleniales	Rhizosoleniaceae	<i>Rhizosolenia sp.</i>	D8	Sparse
	Lithodesmiales	Lithodesmiaceae	<i>Ditylum sp</i>	D9	Dominant
	Thalassiosirales	Thalassiosiraceae	<i>Planktoniellasp</i>	D10	Very sparse
		Skeletonemataceae	<i>Skeletonemas</i> p	D11	Abundant
	Triceratiales	Triceratiaceae	<i>Odontella sp.</i>	D12	Very sparse
			<i>Triceratium sp.</i>	D13	Very sparse
Bacillariophyceae	Bacillariales	Bacillariaceae	<i>Bacillaria sp.</i>	D14	Very sparse
			<i>Nitzschia sp</i>	D15	Sparse
			<i>Pseudo-nitzschia</i> sp.	D16	Very sparse
	Naviculales	<u>Pleurosigmataceae</u>	<i>Pleurosigma sp.</i>	D17	Very sparse
	Surirellales	Entomoneidaceae	<i>Entomoneis sp.</i>	D18	Very sparse
Fragilariophyceae	Fragilariales	Fragilariaceae	<i>Asterionellopsis sp</i>	D19	Scattered
			<i>Fragilari</i> asp	D20	Very sparse
			<i>Synedr</i> asp	D21	Very sparse

	Striatellales	Striatellaceae	<i>Grammatophora sp</i>	D22	Very sparse
	Thalassionematales	Thalassionemataceae	<i>Thalassionema sp.</i>	D23	Sparse
			<i>Thalassiothrix sp.</i>	D24	Very sparse
Noctiluca / Noctiluciphyceae (Dinokaryota)	Noctilucales	Noctilucaceae	<i>Noctiluca sp.</i>	DF1	Sparse
Dinophyceae	Peridiniales	Protopteridiniaceae	<i>Protopteridinium sp.</i>	DF2	Very sparse
	Gonyaulacales	Pyrophacaceae	<i>Pyrophacus sp.</i>	DF3	Very sparse
		Ceratiaceae	<i>Ceratium furca</i>	DF4	Very sparse
			<i>Ceratium tripos</i>	DF5	Very sparse

TABLE:-64 SYSTEMATIC ACCOUNT OF PHYTOPLANKTON IN THE SAMPLING LOCATIONS IN OF DPA HARBOUR AREA AT KANDLA CREEK AND NEARBY CREEKS DURING SPRING TIDE OF NOVEMBER 2022:

CLASS	ORDER	FAMILY	GENUS/SPECIES	#	Relative Abundance
Cyanophyceae	Nostocales	Oscillatoriaceae	<i>Oscillatoria sp.</i>	B1	Very sparse
	Oscillatoriales	Phormidiaceae	<i>Planktothrix sp.</i>	B2	Very sparse
Coscinodiscophyceae	Biddulphiales	Biddulphiaceae	<i>Biddulphia</i> sp.	D1	Sparse
	Chaetocerotales	Chaetocerotaceae	<i>Chaetoceros sp.</i>	D2	Abundant
	Corethrales	Corethraceae	<i>Corethron sp.</i>	D3	Very sparse
	Coscinodiscales	Coscinodiscaceae	<i>Coscinodiscus sp.</i>	D4	Abundant
	Rhizosoleniales	Rhizosoleniaceae	<i>Rhizosolenia sp.</i>	D5	Sparse
	Leptocylindrales	Leptocylindraceae	<i>Leptocylindrus sp.</i>	D6	Very sparse
	Lithodesmiales	Lithodesmiaceae	<i>Ditylum sp.</i>	D7	Scattered
	Thalassiosirales	Thalassiosiraceae	<i>Planktoniella</i> sp.	D8	Very sparse
		Lauderiaceae	<i>Lauderia sp.</i>	D9	Very sparse
		Skeletonemataceae	<i>Skeletonemas</i> sp.	D10	Dominant
	Triceratiales	Triceratiaceae	<i>Odontella sp.</i>	D11	Very sparse
			<i>Triceratium sp.</i>	D12	Very sparse
Bacillariophyceae	Bacillariales	Bacillariaceae	<i>Bacillaria sp.</i>	D13	Very sparse
			<i>Nitzschia sp.</i>	D14	Very sparse
			<i>Pseudo-nitzschia sp.</i>	D15	Very sparse
	Naviculales	Naviculaceae	<i>Navicula sp.</i>	D16	Very sparse
		Plagiotropidaceae	<i>Plagiotropis sp.</i>	D17	Very sparse
		Pleurosigmataceae	<i>Pleurosigma sp.</i>	D18	Sparse
	Surirellales	Entomoneidaceae	<i>Entomoneis sp.</i>	D19	Very sparse
		Surirellaceae	<i>Surirella sp.</i>	D20	Very sparse
Fragilariophyceae	Fragilariales	Fragilariaceae	<i>Asterionellopsis sp.</i>	D21	Sparse

			<i>Fragilariasp</i>	D22	Very sparse
			<i>Synedrassp</i>	D23	Sparse
	Striatellales	Striatellaceae	<i>Grammatophora sp</i>	D24	Very sparse
	Thalassionematales	Thalassionemataceae	<i>Thalassionema sp.</i>	D25	Scattered
			<i>Thalassiothrix sp.</i>	D26	Sparse
Noctiluca / Noctiluiphyceae (Dinokaryota)	Noctilucales	Noctilucaceae	<i>Noctiluca sp.</i>	DF1	Sparse
Dinophyceae	Peridiniales	Protoperidiniaceae	<i>Protoperidinium sp.</i>	DF2	Very sparse
	Gonyaulacales	Ceratiaceae	<i>Ceratium breve</i>	DF3	Very sparse
			<i>Ceratium furca</i>	DF4	Very sparse
			<i>Ceratium fusus</i>	DF5	Very sparse
			<i>Ceratium tripos</i>	DF6	Very sparse

TABLE:-65 SYSTEMATIC ACCOUNT OF PHYTOPLANKTON IN THE SAMPLING LOCATIONS IN OF DPA OOT AREA AT PATH FINDER CREEK AND NEARBY SPM AT VADINARDURING NEAP TIDE OF NOVEMBER 2022:

CLASS	ORDER	FAMILY	GENUS/SPECIES	#	Relative Abundance
Cyanophyceae	Nostocales	Oscillatoriaceae	<i>Lyngbya sp.</i>	B1	Very sparse
			<i>Oscillatoria sp.</i>	B2	Very sparse
			<i>Spirulina sp.</i>	B3	Very sparse
	Oscillatoriales	Phormidiaceae	<i>Planktothrix sp.</i>	B4	Very sparse
Coscinodiscophyceae	Biddulphiales	Biddulphiaceae	<i>Biddulphia</i> sp	D1	Scattered
	Chaetocerotales	Chaetocerotaceae	<i>Chaetoceros</i> sp	D2	Scattered
	Corethrales	Corethraceae	<i>Corethron</i> sp	D3	Very sparse
	Coscinodiscales	Coscinodiscaceae	<i>Coscinodiscus</i> sp.	D4	Dominant
	Hemiaulales	Bellerocheaceae	<i>Belleroche</i> sp	D5	Very sparse
		Hemiaulaceae	<i>Cerataulina</i> sp.	D6	Very sparse
			<i>Eucampia</i> sp	D7	Very sparse
		Streptothecaceae	<i>Helicotheca</i> sp	D8	Very sparse
	Leptocylindrales	Leptocylindraceae	<i>Leptocylindrus</i> sp	D9	Very sparse
	Lithodesmiales	Lithodesmiaceae	<i>Ditylum</i> sp	D10	Abundant
	Rhizosoleniales	Rhizosoleniaceae	<i>Dactyliosolen</i> sp.	D11	Very sparse
			<i>Rhizosolenia</i> sp.	D12	Sparse
	Thalassiosirales	Skeletonemataceae	<i>Skeletonema</i> sp.	D13	Abundant
		Lauderiaceae	<i>Lauderia</i> sp	D14	Very sparse
		Thalassiosiraceae	<i>Planktoniella</i> sp	D15	Very sparse
	Triceratiales	Triceratiaceae	<i>Odontella</i> sp	D16	Very sparse
			<i>Triceratium</i> sp	D17	Very sparse
Bacillariophyceae	Bacillariales	Bacillariaceae	<i>Bacillaria</i> sp.	D18	Abundant
			<i>Nitzschia</i> sp	D19	Very sparse

			<i>Pseudo-nitzschiasp</i>	D20	Scattered
	Naviculales	Naviculaceae	<i>Meuniera sp.</i>	D21	Very sparse
			<i>Navicula sp</i>	D22	Very sparse
		Pinnulariaceae	<i>Pinnulariasp</i>	D23	Very sparse
		<u>Pleurosigmataceae</u>	<i>Pleurosigma sp</i>	D24	Very sparse
	Surirellales	Entomoneidaceae	<i>Entomoneis sp.</i>	D25	Very sparse
		Surirellaceae	<i>Surirellasp</i>	D26	Very sparse
Fragilariophyceae	Climacospheniales	Climacospheniaceae	<i>Climacosphenia sp.</i>	D27	Very sparse
	Fragilariales	Fragilariaceae	<i>Asterionellopsis sp.</i>	D28	Very sparse
			<i>Synedra sp.</i>	D29	Very sparse
	Striatellales	Striatellaceae	<i>Striatellasp</i>	D30	Very sparse
	Thalassionematales	Thalassionemataceae	<i>Thalassionema sp.</i>	D31	Sparse
			<i>Thalassiothrix sp.</i>	D32	Sparse
Dinophyceae	Peridiniales	Protoperidiniaceae	<i>Protoperidinium sp.</i>	DF1	Very sparse
	Dinophysales	Dinophysaceae	<i>Dinophysis sp.</i>	DF2	Very sparse
	Gonyaulacales	Pyrophacaceae	<i>Pyrophacus sp.</i>	DF3	Very sparse
		Ceratiaceae	<i>Ceratium furca</i>	DF4	Very sparse
			<i>Ceratium fusus</i>	DF5	Very sparse
			<i>Ceratium tripos</i>	DF6	Very sparse

TABLE:-66 SYSTEMATIC ACCOUNT OF PHYTOPLANKTON IN THE SAMPLING LOCATIONS IN OF DPAOOT AREA AT PATH FINDER CREEK AND NEARBY SPM AT VADINAR DURING AND SPRING TIDE OF NOVEMBER 2022:

CLASS	ORDER	FAMILY	GENUS/SPECIES	#	Relative Abundance
Cyanophyceae	Chroococcales	Chroococcaceae	<i>Merismopedia sp.</i>	B1	Very sparse
	Nostocales	Oscillatoriaceae	<i>Lyngbya sp.</i>	B2	Very sparse
			<i>Oscillatoria sp.</i>	B3	Sparse
	Oscillatoriales	Phormidiaceae	<i>Planktothrix sp.</i>	B4	Very sparse
	Stigonematales	Stigonemataceae	<i>Stigonema sp.</i>	B5	Very sparse
Coscinodiscophyceae	Biddulphiales	Biddulphiaceae	<i>Biddulphia</i> sp.	D1	Sparse
	Chaetocerotales	Chaetocerotaceae	<i>Chaetoceros sp.</i>	D2	Dominant
	Corethrales	Corethraceae	<i>Corethron sp.</i>	D3	Very sparse
	Coscinodiscales	Coscinodiscaceae	<i>Coscinodiscus sp.</i>	D4	Abundant
	Hemiaulales	Bellerophyceae	<i>Bellerophyce</i> sp.	D5	Very sparse
		Hemiaulaceae	<i>Cerataulina sp.</i>	D6	Very sparse
		Streptothecaceae	<i>Helicotheca sp.</i>	D7	Very sparse
	Rhizosoleniales	Rhizosoleniaceae	<i>Rhizosolenia sp.</i>	D8	Scattered
	Leptocylindrales	Leptocylindraceae	<i>Leptocylindrus sp.</i>	D9	Very sparse
	Lithodesmiales	Lithodesmiaceae	<i>Ditylum sp.</i>	D10	Abundant
	Thalassiosirales	Thalassiosiraceae	<i>Planktoniella</i> sp.	D11	Very sparse
		Lauderiaceae	<i>Lauderia sp.</i>	D12	Very sparse
	Triceratiales	Triceratiaceae	<i>Odontella sp.</i>	D13	Sparse
			<i>Triceratium sp.</i>	D14	Very sparse
Bacillariophyceae	Bacillariales	Bacillariaceae	<i>Bacillaria sp.</i>	D15	Scattered
			<i>Nitzschia sp.</i>	D16	Very sparse
			<i>Pseudo-nitzschia</i> sp.	D17	Sparse
	Naviculales	Pinnulariaceae	<i>Pinnularia</i> sp.	D18	Very sparse

		<u>Pleurosigmataceae</u>	<i>Pleurosigma</i> sp.	D19	Very sparse
	Surirellales	Entomoneidaceae	<i>Entomoneis</i> sp.	D20	Very sparse
		Surirellaceae	<i>Surirella</i> sp.	D21	Very sparse
Fragilariophyceae	Fragilariales	Fragilariaceae	<i>Asterionellopsis</i> sp	D22	Sparse
			<i>Synedrassp</i>	D23	Very sparse
	Thalassionematales	Thalassionemataceae	<i>Thalassionema</i> sp.	D24	Sparse
			<i>Thalassiothrix</i> sp.	D25	Very sparse
Dinophyceae	Peridiniales	Protoperidiniaceae	<i>Protoperidinium</i> sp.	DF1	Very sparse
	Dinophysales	Dinophysaceae	<i>Dinophysis</i> sp.	DF2	Very sparse
	Gonyaulacales	Pyrophacaceae	<i>Pyrophacus</i> sp.	DF3	Very sparse
		Ceratiaceae	<i>Ceratium furca</i>	DF4	Very sparse
			<i>Ceratium fusus</i>	DF5	Very sparse
			<i>Ceratium tripos</i>	DF6	Very sparse

TABLE:-67 SYSTEMATIC ACCOUNT OF ZOOPLANKTON FROM THE SAMPLING LOCATIONS OF DPA HARBOUR AREA AT KANDLA CREEK AND NEARBY CREEKS DURING NEAP TIDE OF NOVEMBER 2022:

CLASS	ORDER	FAMILY	GENUS/SPECIES	#	RELATIVE ABUNDANCE
Spirotrichea	Tintinnida	Tintinnidiidae	<i>Leprotintinnus</i> sp.	T1	Very sparse
		Codonellidae	<i>Tintinnopsis dadayi</i>	T2	Very sparse
			<i>Tintinnopsis failakkaensis</i>	T3	Very sparse
			<i>Tintinnopsis gracilis</i>	T4	Very sparse
			<i>Tintinnopsis mortensenii</i>	T5	Very sparse
			<i>Tintinnopsis radix</i>	T6	Very sparse
			<i>Tintinnopsis tocaninensis</i>	T7	Very sparse
		Tintinnidae	<i>Amphorellopsis</i> sp.	T8	Very sparse
			<i>Eutintinnus</i> sp.	T9	Very sparse
		Xystonellidae	<i>Favella</i> sp.	T10	Very sparse
Crustacea Subclass: Copepoda	Calanoida	Paracalanidae	<i>Acrocalanus</i> sp.	C1	Sparse
			<i>Parvocalanus</i> sp.	C2	Very sparse
		Acartiidae	<i>Acartia</i> sp.	C3	Very sparse
		Clausocalanidae	<i>Clausocalanus</i> sp.	C4	Very sparse
		Centropagidae	<i>Centropages</i> sp.	C5	Very sparse
		Temoridae	<i>Temora</i> sp.	C6	Very sparse
	Cyclopoida	Oithonidae	<i>Oithona</i> sp.	C7	Abundant
	Harpacticoida	Ectinosomatidae	<i>Microsetella</i> sp.	C8	Scattered
		Euterpinae	<i>Euterpina</i> sp.	C9	Sparse
	Poecilostomatoida	Oncaeidae	<i>Oncaea</i> sp.	C10	Very sparse
Sagittioidea	Aphragmophora	Sagittidae	<i>Sagitta</i> sp.	A1	Very sparse
Malacostraca	Mysida,	Penaeidae	<i>Metapenaeus</i> sp.	M1	Very sparse
	Decapoda	Solenoceridae	<i>Solenocera</i> sp.	M2	Very sparse

Environmental Monitoring Report of Deendayal Port Authority, November - 2022

Appendicularia		Fritillariidae	<i>Fritillaria sp.</i>	U1	Very sparse
		Oikopleuridae	<i>Oikopleura sp.</i>	U2	Very sparse
Oligohymenophorea	Sessilida	Zoothamniidae	<i>Zoothamnium sp.</i>	CI1	Very sparse
Copepoda			Nauplius larvae of copepods	L1	Dominant
Malacostraca Decapoda			Brachyuran zoea	L2	Very sparse
Maxillopoda Thecostraca			Cirripede larvae	L3	Very sparse
			Cyphonautes larvae	L4	Very sparse
			Ophiopluteus larvae	L5	Very sparse
Polychaeta			Trochophore larvae	L6	Very sparse

TABLE:-68 SYSTEMATIC ACCOUNT OF ZOOPLANKTON FROM THE SAMPLING OF DPA HARBOUR AREA AT KANDLA CREEK AND NEARBY CREEKS DURING SPRING TIDE OF NOVEMBER 2022:

CLASS	ORDER	FAMILY	GENUS/SPECIES	#	RELATIVE ABUNDANCE
Spirotrichea	Tintinnida	Tintinnidiidae	<i>Leprotintinnus</i> sp.	T1	Scattered
		Codonellidae	<i>Tintinnopsis dadayi</i>	T2	Very sparse
			<i>Tintinnopsis failakkaensis</i>	T3	Very sparse
			<i>Tintinnopsis gracilis</i>	T4	Very sparse
			<i>Tintinnopsis mortensenii</i>	T5	Very sparse
			<i>Tintinnopsis radix</i>	T6	Sparse
			<i>Tintinnopsis tocaninensis</i>	T7	Very sparse
		Metacyclidiidae	<i>Metacyclis</i> sp.	T8	Very sparse
		Tintinnidae	<i>Amphorellopsis</i> sp.	T9	Very sparse
			<i>Eutintinnus</i> sp.	T10	Very sparse
		Xystonellidae	<i>Favella</i> sp.	T11	Sparse
Crustacea Subclass: Copepoda	Calanoida	Paracalanidae	<i>Acrocalanus</i> sp.	C1	Scattered
			<i>Parvocalanus</i> sp.	C2	Very sparse
		Acartiidae	<i>Acartia</i> sp.	C3	Very sparse
		Clausocalanidae	<i>Clausocalanus</i> sp.	C4	Very sparse
		Centropagidae	<i>Centropages</i> sp.	C5	Very sparse
		Eucalanidae	<i>Subeucalanus</i> sp.	C6	Very sparse
	Cyclopoida	Oithonidae	<i>Oithona</i> sp.	C7	Abundant
	Harpacticoida	Ectinosomatidae	<i>Microsetella</i> sp.	C8	Sparse
		Euterpinae	<i>Euterpina</i> sp.	C9	Sparse
Sagittoidea	Aphragmophora	Sagittidae	<i>Sagitta</i> sp.	A1	Very sparse
Malacostraca	Mysida, Decapoda	Solenoceridae	<i>Solenocera</i> sp.	M1	Very sparse

Appendicularia		Fritillariidae	<i>Fritillaria sp.</i>	U1	Very sparse
		Oikopleuridae	<i>Oikopleura sp.</i>	U2	Very sparse
Oligohymenophorea	Sessilida	Zoothamniidae	<i>Zoothamnium sp.</i>	CI1	Very sparse
Copepoda			Nauplius larvae of copepods	L1	Dominant
Malacostraca			Brachyuran zoea	L2	Sparse
Decapoda					
Maxillopoda			Cirripede larvae	L3	Very sparse
Thecostraca					
			Cyphonautes larvae	L4	Very sparse
			Ophiopluteus larvae	L5	Very sparse
Gastropoda			Opisthobranchia larvae	L6	Very sparse
Streptoneura					
Polychaeta			Trochophore larvae	L7	Sparse
Pelecypoda			Veliger larvae of bivalves	L8	Very sparse

TABLE:-69 SYSTEMATIC ACCOUNT OF ZOOPLANKTON FROM THE SAMPLING LOCATIONS OF DPA OOT AREA AT PATH FINDER CREEK AND NEARBY SPM AT VADINAR DURING NEAP TIDE OF NOVEMBER 2022:

CLASS	ORDER	FAMILY	GENUS/SPECIES	#	RELATIVE ABUNDANCE
Spirotrichea	Tintinnida	Tintinnidiidae	<i>Leptotintinnus</i> sp.	T1	Sparse
		Codonellidae	<i>Tintinnopsis failakkaensis</i>	T2	Very sparse
			<i>Tintinnopsis gracilis</i>	T3	Very sparse
			<i>Tintinnopsis radix</i>	T4	Very sparse
			<i>Tintinnopsis tocanensis</i>	T5	Very sparse
		Tintinnidae	<i>Amphorellopsis</i> sp.	T6	Very sparse
		Xystonellidae	<i>Favella</i> sp.	T7	Very sparse
Crustacea Subclass: Copepoda	Calanoida	Paracalanidae	<i>Acrocalanus</i> sp.	C1	Scattered
			<i>Parvocalanus</i> sp.	C2	Very sparse
	Cyclopoida	Oithonidae	<i>Oithona</i> sp.	C3	Abundant
	Harpacticoida	Euterpinae	<i>Euterpina</i> sp.	C4	Very sparse
		Ectinosomatidae	<i>Microsetella</i> sp.	C5	Very sparse
	Poecilostomatoida	Oncaeidae	<i>Oncaea</i> sp.	C6	Very sparse
Sagittoida	Aphragmophora	Sagittidae	<i>Sagitta</i> sp.	A1	Very sparse
Appendicularia		Fritillariidae	<i>Fritillaria</i> sp.	U1	Very sparse
		Oikopleuridae	<i>Oikopleura</i> sp.	U2	Very sparse
Copepoda			Nauplius larvae of copepods	L1	Dominant
Maxillopoda Thecostraca			Cirripede larvae	L2	Very sparse
Gastropoda Streptoneura			Opisthobranchia larvae	L3	Very sparse
Polychaeta			Trochophore larvae	L4	Very sparse
Pelecypoda			Veliger larvae of bivalves	L5	Very sparse

TABLE:-70 SYSTEMATIC ACCOUNT OF ZOOPLANKTON FROM THE SAMPLING LOCATIONS OF DPA OOT AREA AT PATH FINDER CREEK AND NEARBY SPM AT VADINAR DURING SPRING TIDE OF NOVEMBER 2022:

CLASS	ORDER	FAMILY	GENUS/SPECIES	#	RELATIVE ABUNDANCE
Spirotrichea	Tintinnida	Tintinnidiidae	<i>Leprotintinnus</i> sp.	T1	Abundant
		Codonellidae	<i>Tintinnopsis gracilis</i>	T2	Very sparse
			<i>Tintinnopsis mortensenii</i>	T3	Very sparse
			<i>Tintinnopsis radix</i>	T4	Very sparse
		Xystonellidae	<i>Favella</i> sp.	T5	Scattered
Crustacea Subclass: Copepoda	Calanoida	Paracalanidae	<i>Acrocalanus</i> sp.	C1	Sparse
			<i>Parvocalanus</i> sp.	C2	Very sparse
		Centropagidae	<i>Centropages</i> sp.	C3	Very sparse
		Tortanidae	<i>Tortanus</i> sp.	C4	Very sparse
	Cyclopoida	Oithonidae	<i>Oithona</i> sp.	C5	Abundant
		Euterpinae	<i>Euterpina</i> sp.	C6	Very sparse
	Harpacticoida	Ectinosomatidae	<i>Microsetella</i> sp.	C7	Scattered
	Poecilostomatoida	Corycaidae	<i>Corycaeus</i> sp.	C8	Very sparse
Appendicularia		Fritillariidae	<i>Fritillaria</i> sp.	U1	Very sparse
		Oikopleuridae	<i>Oikopleura</i> sp.	U2	Very sparse
Oligohymenophorea	Sessilida	Zoothamniidae	<i>Zoothamnium</i> sp.	CI1	Very sparse
Copepoda			Nauplius larvae of copepods	L1	Dominant
Malacostraca Decapoda			Brachyuran zoea	L2	Very sparse
Gastropoda Streptoneura			Opisthobranchia larvae	L3	Very sparse
Pelecypoda			Veliger larvae of bivalves	L4	Very sparse

BENTHIC ORGANISMS:

Few Benthic organisms were observed in the collected sediments by using the Van-Veen grabs during the sampling conducted during spring tide period and Neap tide period from DPA harbour region and nearby creek. The Meio-benthic organisms during spring tide were represented by Polychaetes *Tharyx sp.* and *Nereis sp.*, during Neap tide by *Nereis sp.* and few Amphipods. Population of benthic fauna was varying from 10-60- N/m² during spring tide and 0-80 N/m² during Neap tide. The benthic communities at path finder Creek were represented by Polychaetes *Glycera sp.* *Cirratulus sp.* *Nereis sp.* and few Amphipods. Their population was varying as 60 N/m² at OOT jetty premises and 80 N/m² near the SPM area during spring tide and 50 N/m² at OOT jetty premises and 50 N/m² near the SPM area during Neap tide period.

Table:-71 BENTHIC FAUNA IN THE SAMPLING LOCATIONS IN DPA HARBOUR AREA CREEKS DURING SPRING TIDE IN NOVEMBER 2022

ABUNDANCE IN NO/M ² DIFFERENT SAMPLING STATIONS						
REPRESENTATION BY GROUP	DPA HARBOUR			CREEKS		
Benthic fauna						
POLYCHAETES	DPA-1	DPA-2	DPA-3	DPA-4	DPA-5	DPA-6
Family : CIRRATULIDAE <i>Tharyxsp.</i>	20	10	10	0	0	NS
Family :NEREIDAE <i>Nereis sp.</i>	0	0	0	20	40	NS
AMPHIPODA	0	0	0		20	NS
TOTAL Benthic Fauna NUMBER/ M ²	20	10	10	20	60	NS

NS: No sample

Table:-72 BENTHIC FAUNA IN THE SAMPLING LOCATIONS IN DPA HARBOUR AREA CREEKS DURING NEAP TIDE IN NOVEMBER 2022

ABUNDANCE IN NO/M ² DIFFERENT SAMPLING STATIONS						
REPRESENTATION BY GROUP	DPA HARBOUR			CREEKS		
Benthic fauna						
POLYCHAETES	DPA-1	DPA-2	DPA-3	DPA-4	DPA-5	DPA-6
Family :NEREIDAE <i>Nereis sp.</i>	0	0	0	40	60	NS
<i>Amphipoda</i>	0	20	10	10	20	NS
TOTAL Benthic Fauna NUMBER/M ²	0	20	10	50	80	NS

**Table:-73 BENTHIC FAUNA IN THE SAMPLING LOCATIONS IN DPA OOT JETTY AREA,
VADINAR DURING SPRING TIDE IN NOVEMBER 2022**

ABUNDANCE IN NO/M ² DIFFERENT SAMPLING STATIONS		
REPRESENTATION BY GROUP	OOT Jetty Area	SPM area
POLYCHAETES		
Family : Glyceride <i>Glycerasp.</i>	20	40
Family : CIRRATULIDAE <i>Cirratulussp.</i>	0	20
Family: NEREIDAE <i>Nereis sp.</i>	30	10
<i>Amphipoda</i>	10	20
TOTAL Benthic Fauna NUMBER/ M ²	60	80

**Table:-74 BENTHIC FAUNA IN THE SAMPLING LOCATIONS IN DPA OOT JETTY AREA,
VADINAR DURING NEAP TIDE IN NOVEMBER 2022**

ABUNDANCE IN NO/M ² DIFFERENT SAMPLING STATIONS		
REPRESENTATION BY GROUP	OOT Jetty Area	SPM area
POLYCHAETES		
Family : Glyceridase <i>Glycera sp.</i>	20	40
Family: NEREIDAE <i>Nereis sp.</i>	30	10
TOTAL Benthic Fauna NUMBER/ M ²	50	50

CHAPTER-11

CONCLUSIVE SUMMARY & REMEDIAL MEASURES

11.0 Conclusive Summary and Remedial measures Suggested

- The AAQ monitoring of six locations at Deendayal Port Authority indicates that the mean PM_{10} and $PM_{2.5}$ values for four locations viz. Marine Bhavan, Oil Jetty, Estate Office and Coal storage area were found higher than the permissible limit (standards $100 \mu g/m^3$, $60 \mu g/m^3$). The higher concentration of Particulate matter at Marine Bhavan may be due to vehicles emissions during loading-unloading of food grains and timbers; at Estate office due to construction work, vehicles emission produced from trucks, heavy duty vehicles that pass through the road outside Kandla port and Oil jetty area; while at Coal Storage area lifting of coal from grab yard and other coal handling processes. Moreover, the transportation of coal produces pollution from heavy vehicles. At Tuna Port location, concentration of PM_{10} varied from $88-175 \mu g/m^3$ and mean value was observed $145 \mu g/m^3$ which was exceed the prescribed standard limit ($100 \mu g/m^3$), concentration of $PM_{2.5}$ was ranged from $47-87 \mu g/m^3$ and mean was found $71 \mu g/m^3$ which was exceed the standard limit ($60 \mu g/m^3$). At Gopalpuri PM_{10} concentration ranged from $67-168 \mu g/m^3$ and mean was $127 \mu g/m^3$ while $PM_{2.5}$ concentration ranged from $34-94 \mu g/m^3$ and mean was $66 \mu g/m^3$ were found exceed standard limit prescribed by NAAQS.
- At Vadinar, the average concentration of PM_{10} was $114 \mu g/m^3$ and $PM_{2.5}$ was $74 \mu g/m^3$ at Admin Colony which was slightly exceed the standard limit while at Signal building the mean concentration PM_{10} was $100 \mu g/m^3$ and $PM_{2.5}$ was $61 \mu g/m^3$ which were very close to standard limit.
- During winter, the concentration of PM_{10} and $PM_{2.5}$ has been slowly augmented and reached a peak in the evening due to surface inversion of temperature after sunset. Thus, the pollutants are subsequently trapped in the lower layer of the atmosphere due to high atmospheric air pressure.
- Further, precautionary measures and management strategies to minimize the effect of particulate as well as gaseous pollutants have also been suggested for achieving its ambient levels in and around Kandla Port and Vadinar Port, Gujarat, India.
- Drinking water at all the twenty locations was found potable and it was found within in line of BIS standards (IS: 10500-2012).
- Transportation systems are the main source of noise pollution in project areas. Noise sources in port operations include cargo handling, vehicular traffic, and loading / unloading

containers and ships. All sampling location were within the permissible limit day time 75 dB (A) and night time 70 dB (A) for the industrial area.

- The treated sewage water of Kandla STP, Deendayal Port Colony (Gopalpuri) STP and Vadinar were in line with the standards set by the Central Pollution Control Board.
- It was suggested to monitor the STP performance on regular basis to avoid flow of contamination / Polluted water into the sea.
- Good species diversity suggests a relatively successful species in this habitat. A greater number of successful species and a more stable ecosystem. More ecological niches are available and the environment is less likely to be hostile complex food webs environmental change is less likely to be damaging to the ecosystem as a whole.
- The results obtained from the study for biological and ecological parameters in marine water for Arabian Sea at surrounding area of Deendayal Port Authority (DPA) Kandla and Vadinar were not affected by Port activities.
- The mean day time temperature at Deendayal Port was 27.92 °C. The day-time maximum temperature was 32.9°C and minimum was 21.1 °C. The mean night time temperature recorded was 25.47 °C. The night-time maximum temperature was 29.7°C and minimum was 20.0 °C. The mean Solar Radiation in November month was 167.27 w/m². The maximum solar radiation was recorded 759 w/m² in 4th November, 2022 and the minimum solar radiation was recorded 1.80 w/m² in 30th November, 2022. The mean Relative humidity was 69.00 % for the month of November. Maximum Relative humidity was recorded 99.0 % and minimum Relative humidity was recorded 34.0 %. The average wind velocity for the entire month of November was 1.21 m/s. Maximum wind velocity was recorded 10.19 m/s. The wind direction was mostly North-East.
- The results obtained from the study for the month of November 2022 for biological and ecological parameters in marine water for Arabian Sea at surrounding area of Deendayal Port Authority (DPA) Kandla and Vadinar were not affected by Port activities.

Reasons for higher Values of PM₁₀

- The unloading of coal directly in the truck, using grabs cause coal to spread in air as well as coal dust to fall on ground. This settled coal dust again mixes with the air while trucks travel through it.

- Also, the coal loaded trucks were not always covered with tarpaulin sheets and these results in spillage of coal from trucks/dumpers during its transit from vessel to yard or storage site. This also increased PM values around marine Bhavan & Coal storage area.

Remedial Measures

The values of PM₁₀ & PM_{2.5} during the month of November, 2022 were beyond the standard limit at all locations (Coal Storage, Marine Bhavan, Oil Jetty and Estate office, Tuna Port) except Gopalpuri the concentration of particulate matter was slightly exceed. Given below are the remedial measures suggest to minimize the Air pollution.

- During November, 2022 overall ambient air quality of the DPA was within CPCB permissible limits except TSPM, PM₁₀, PM_{2.5} at Coal storage area, Marine Bhavan, Oil Jetty and Estate Office. To improve air quality the port was using number of precautionary measures, such as maintained a wide expanse of Green zone, initiated Inter-Terminal Transfer (ITT) of tractor-trailers, Centralized Parking Plaza, providing shore power supply to tugs and port crafts, the use of LED lights at DPA area helps in lower energy consumption and decreases the carbon foot prints in the environment, time to time cleaning of paved and un paved roads, use of tarpaulin sheets to cover dumpers at project sites etc. are helping to achieve the cleaner and green future at port.

Solution towards the Green port:

Today, it is increasingly recognized that air pollution hurts human health. Consequently, efficient mitigation strategies need to be implementation for substantial environmental and health co-benefits.

The guidelines can be considered a basis for governments for the implementation of a strategic plan focused on the reduction of multi pollutant emission, as well as of the overall air pollution related risk.

- The plantation should be all along the periphery of the port and inside and outside the port along with the road. Trees having high dust trapping efficiency (*Azadirachta indica*, *Cassia fistula*, *Delonix regia*, *Ficus religiosa*, *Pterocarpus marsupium*) are to be grown alongside the roads.
- The water sprinkling should be use at each and every stage of transporting coal up the loading of truck to avoid generation of coal dust.

- The vehicles should be covered during transportation and the vehicle carrying the coal should not be overloaded by raising the height of carriage.
- The water sprinklers should be use during transportation of loaded heavy vehicles on raw road.
- It should be ensure that regular sweeping of coal internal, main road and space a free circulation.
- Practice should be initiated for using mask as preventative measure, to avoid Inhalation of dust particle- Mask advised in sensitive areas.
- Department for use maintenance should have a routine checkup noise level by replacing bearings, tights of all loose parts that can vibrate.
- Speed control is also an effective way to mitigate noise pollution, the lowest sound emission arise from vehicles moving smoothly.
- Use of renewable energy like solar energy should be optimal and ensure to work continuously.
- Keep neat and clean public transport and all basic items at public interaction places as much as possible.
- Technology like Electric cart, Inter-Terminal Transfer (ITT) are worthy selection to reduce Port operation efficiency and fuel cost.
- Conventional RTGCs should be altered as E-RTGCs counting inside the port completely.
- Initiate Natural Gas (CNG) as fuel by all buses and trucks.

Green Ports Initiative

- Deendayal Port is committed to sustainable development and adequate measures are being taken to maintain the Environmental well-being of the Port and its surrounding environs. Weighing in the environmental perspective for sustained growth, the Ministry of Shipping had started “Project Green Ports” which will help in making the Major Ports across India cleaner and greener. “Project Green Ports” will have two verticals - one is “Green Ports Initiatives” related to environmental issues and second is “Swachh Bharat Abhiyaan”.
- The Green Port Initiatives include twelve initiatives such as preparation and monitoring plan, acquiring equipments required for monitoring environmental pollution, acquiring dust suppression system, setting up of waste water treatment plants/ garbage disposal plant, setting up Green Cover area, projects for energy generation from renewable

energy sources, completion of shortfalls of Oil Spill Response (OSR) facilities (Tier-I), prohibition of disposal of almost all kind of garbage at sea, improving the quality of harbour wastes etc.

- Deendayal port has also appointed GEMI as an Advisor for “Making Deendayal Port a Green Port - Intended Sustainable Development under the Green Port Initiatives.
- Deendayal Port has also signed MOU with Gujarat Forest Department in August 2019 for Green Belt Development in an area of 31.942 Ha of land owned by Deendayal Port Trust. The plantation is being carried out by the Social Forestry division of Kachchh.

CHAPTER-12

REFERENCES

12.0 SOURCE OF LITERATURE AND ADDITIONAL REFERENCE

- 1) ALBERT WEST PHAL (1976) Protozoa Blackwell , London
- 2) Arunachalam S, Brantley H, Barzyk TM, Hagler G, Isakov V, Kimbrough E, Naess B, Rice N, Snyder MG, Talgo K, Venkatram A (2015) Assessment of port-related air quality impacts: geographic analysis of population. *Int J Environ Pollut* 58(4):231–250
- 3) Asha, P. S. and Diwakar (2007). Hydrobiology of the inshore waters off Tuticorin in the Gulf. *J. Mar. Biol. Ass. India*, 49: 7-11.
- 4) Astakhov AS, Kalugin IA, Aksentov KI, Daryin AV (2015) Geochemical indicators of paleo typhoons in shelf sediments. *Geochemistry* 4:387–392.
- 5) Bailey D, Solomon G (2004) Pollution prevention at ports: clearing the air. *Environ Impact Assess Rev* 24(7):749–774.
- 6) BANERJEE R.K. (1989) Heavy metals and Benthic foraminiferal distribution along Bombay coast India. *Studies in benthic foraminifera. Tokyo University Press Tokyo* pp 151-157
- 7) Banse K (1995) Zooplankton: Pivotal role in the control of ocean production: I. Biomass and production. *ICES J Mar Sci* 52: 265–277.
- 8) Beaugrand G, and Ibanez F (2004) Monitoring marine plankton ecosystems. II: long-term changes in North Sea calanoid copepods in relation to hydroclimatic variability. *Inter Res Mar Ecol Prog Ser* 284:35-47.
- 9) Benhamed S, Guardiola FA, Martinez S, Sanchez MJM, Sirvent CP, Mars M, Esteban MA (2016) Exposure of the gilthead seabream (*Sparus aurata*) to sediments contaminated with heavy metals down-regulates the gene expression of stress biomarkers. *Toxicol Rep* 3:364–372.
- 10) Best, M. A., Wither, A.W. and Coates, S. (2007). Dissolved oxygen as a physicochemical supporting element in the Water Framework Directive. *Marine Pollution Bulletin* 55(1-6): 53-64.
- 11) Buccolieri R, Cesari R, Dinoi A, Maurizi A, Tampieri F, Di Sabatino S (2016) Impact of ship emissions on local air quality in a Mediterranean City's harbour after the European sulphur directive. *Int J Environ Pollut* 59(1):30–42.
- 12) Chaurasia, S, Singh, S. and Gupta, A.D., 2013. Study on air quality of SKS Ispat and Power Ltd. Raipur (CG), India. *Asian Journal of Science and Technologies*, 4 (4), 48-50.

- 13) Conti ME, Ciasullo R, Tudino MB, Matta EJ (2015) “The industrial emissions trend and the problem of the implementation of the industrial emissions directive (IED)” Air Quality. Atmosph Health 8(2): 151–161.
- 14) CPCB (2013): Guidelines for Manual Sampling & Analyses. (Guidelines for the Measurement of Ambient Air Pollutants, Vol. I), Central Pollution Control Board, National Ambient Air Quality Series: NAAQMS/36/2012-13.
- 15) DAY F. (1889) The fauna of British India Ceylon and Burma- Fishes Vol-1- Vol-2 *Taylor and Francis* London
- 16) DESIKACHARYT.V. (1989) Atlas of diatoms, Madras Science Foundation
- 17) DESIKACHARYT.V.(1959) CyanophytaICAP Monographs on Algae *Indian Council of Agricultural research* New Delhi
- 18) FAIZAYOUSIF AL-YAMANI& MARIA A. SABUROVA(2010) illustrative guide on the flagellates of Intertidal soft sediment *Kuwait Institute for scientific Research* Kuwait
- 19) FAIZAYOUSIF AL-YAMANI, VALERIYSKRYABIN, ALEKSANDRA GUBANOVA, SERGEY KHVOROV AND IRINA PRUSOVA (2011), Marine zooplankton Practical guide from North western Arabian gulf Vol-1 and vol-2 *Kuwait Institute for scientific Research* Kuwait
- 20) FAUVEL P. (1953), The fauna of India Annelida - Polychaeta Indian Press Allahabad
- 21) GajbhiyeSN, Nair VR, and Desai BN (1984). Diurnal variation of zooplankton in Malad creek, Bombay. Indian Journal of Marine Science. 13:75-79.
- 22) Guidelines for the Measurement of Ambient Air Pollutants: National Ambient Air Quality Series/NAAQMS/36/2012-13, Volume-I.
- 23) Gujrat Pollution Control Board Inward No. 143215 dated 06.09.2018.
- 24) HAYWARD P.J AND RYLAND J.S. (1995) Handbook of Marine fauna of north –West Europe oxford University Press London
- 25) Hefmi R., Tanjung1 R., Hamuna B., and Alianto(2019).Assessment of Water Quality and Pollution Index in Coastal Waters of Mimika, Indonesia. *Journal of Ecological Engineering*, 20(2);87-94
- 26) HIGGINS R.P. HAJAMARTHIEL Eds. (1998) Introduction to the study of Meio Fauna
- 27) HORACE G. BARBER AND ELIZABETH Y. HAWORTH (1981) A guide to the Morphology of DIATOMS FRUSTULES.

- 28) Horaginamani, S.M. and Ravichandran, M., 2010. Ambient air quality an urban area and its effects on plants and human beings: A case study of Tiruchirappalli, India. Kathmandu University Journal of Science, Engineering and Technology, 6 (2), 13-19.
- 29) INGRAM HENDEY (1964) An introductory account of smaller Algae of British coastal waters part-V. Bacillariophyceae
- 30) IS1050:2012 Drinking Water-Specification: (second Revision).
- 31) JOHN H. WICKSTEAD(1965) an Introduction to the study of Tropical Plankton .Hutchinson Tropical Monographs
- 32) JOYOTHIBABU,R. MADHU, N.V. MAHESHWARAN, P.A.,NAIRK.K.C., VENUGOPL,P. BALASUBRAMANIAN T.2005) Dominance of Dinoflagellates in micro zooplankton communities in the oceanic region Bay of Bengal and Andaman sea Current science vol.84. 10thNOVEMBER2003
- 33) KASTURIRANGANL.R. (1963) A key for the identification of the Common Planktonic Copepoda of Indian Coastal water
- 34) KusumKK, Vineetha G, Raveendran TV, Nair VR, Muraleedharan KR, Achuthankutty CT and Joseph T (2014) Chaetognath community and their responses to varying environmental factors in the northern Indian ocean. J Plankton Res 36(4): 1146- 1152.
- 35) Lack DA, Cappa CD, Langridge J, Bahreini R, Buffaloe G, Brock C, Cerully K, Coffman D, Hayden K, Holloway J, Lerner B, Massoli P, Li SM, McLaren R, Middlebrook AM, Moore R, Nenes A, Nuaaman I, Onasch TB, Peischl J, Perring A, Quinn PK, Ryerson T, Schwartz JP, Spackman R, Wofsy SC, Worsnop D, Xiang B, Williams E (2011) Impact of fuel quality regulation and speed reductions on shipping emissions: implications for climate and air quality. Environ Sci Technol 45(20):9052–9060.
- 36) Lai KH, Lun VYH, Wong CWY, Cheng TCE (2011) Green shipping practices in the shipping industry: conceptualization, adoption, and implications. Resour Conserv Recycl 55(6):631–638.
- 37) Lalli CM and Parsons TR (1997) Biological Oceanography: An Introduction. DOI<https://doi.org/10.1016/B978-0-7506-3384-0.X5056-7>.
- 38) Lawler, D.M. 2004. Turbidimetry and nephelometry. In:Townshend, A. (Ed.) Encyclopedia of Analytical Science, 2nd edition. London: Academic Press (pp. 343-351).

- 39) Madhu D. Nathani, Mrugesh H. Trivedi (2015). Pre Monsoon Study of Water Quality with reference to Rapid Industrial Development in and around Gandhidham, Kachchh, Gujarat, *International Journal of Scientific & Engineering Research*, 6(9):1851-1862.
- 40) Madhupratap M (1978) Studies on ecology of zooplankton of Cochin backwaters. *Mahasagar Bull Nat Inst Oceanogr* 11: 45-56.
- 41) Madhupratap M (1979) Distribution, community structure and species succession of copepods from Cochin Backwaters. *Indian J Ma Sci* 8: 1-8.
- 42) Madhupratap M (1987) Status and strategy of zooplankton of tropical Indian estuaries: A review. *Bull Plank Soc Jpn* 34: 65-81.
- 43) Madhupratap M (1999). Free living copepods of the Arabian Sea, Distribution and Research Perspectives. *I J Mar Sci* 146-149.
- 44) Madhupratap M and Haridas P (1986) Epipelagic calanoid copepods of the northern Indian Ocean. *Oceanologica Acta* 9(2):105-117.
- 45) MANAL AL-KANDARI, FAIZA Y. AL-YAMANI , KHOLOOD AL-RIFAIE (2009) Marine phytoplankton Atlas of Kuwait's water *Kuwait Institute for scientific Research*
- 46) Meiaraj C. & Jeyapriya S.P. (2019). Marine water quality studies at Tuticorin harbour coastal area. *Indian Journal of Geo Marine Sciences* 48 (06): 943-946.
- 47) MPEDA (1998) Commercial Fishes and shell fishes of India
- 48) Murtini J.T., Ariyani F., Wahyuni I.S., Hak N., Suherman M., Dolaria N., Nurwiyanto. 2001. Inventory research and identification of heavy metal pollution in waters and fish. Technical Report. Pusat Riset Pengolahan Produk dan Bioteknologi Kelautan dan Perikanan. Jakarta. (in Indonesian).
- 49) Narayan R, Saxena K.K. and Chauhan S. (2007). Limnological investigations of Texi Temple pond in district Etawah (U.P.), *Journal of Environmental Biology*, January 28(1):155-157.
- 50) NEWEL G.E. & NEWELL R.C. (1963) Marine plankton a Practical Guide Hutchinson Educational
- 51) NIGAM R.C. AND CHATURVEDIS.K. (2000) Foraminiferal Study from Kharo Creek , Kachchh (Gujarat) North west coast of India. *Indian Journal of marine science* Vol.29 133-189
- 52) OLAV GIERE (1993) Meiobenthology , Microscopic Fauna in Aquatic Sediments m Springer London

- 53) PERRAGALLO (1965) *Diatomees marines de France* A. Asher & Co. Amsterdam
- 54) Robert P. Higgins (Eds.), (1985) *An introduction to the study of Meiofauna* Smithsonian Institution press Washington DC
- 55) Salmin (2005). Dissolved oxygen (DO) and biological oxygen demand (BOD) as indicator to determine water quality. *Oseana*, 30(3):21–26.
- 56) Salvi H., Patel R., Thakur B., Shah K. and Parmar D. (2014). Assessment of Coastal Water Quality Parameters of Selected Areas of Marine National Park & Sanctuary (Okha, Sikka & Khijadiya), *SSRN Electronic Journal*, (pg 1-16)
- 57) STERRER W. STERRER C.S Eds. *Marine fauna and flora of Bermuda A systematic Guide to the identification of Marine Organisms. John Wiley and Sons* New York
- 58) Suresh Gandhi. M. (2009) Distribution of certain ecological parameters and Foraminiferal distribution in the depositional environment of Pak Strait east coast of India. *Indian J. of Marine Science* Vol.33 pp 287-295
- 59) USEPA, 2008. *Air Quality Index: A Guide to Air Quality and Your Health*. Environmental Protection Agency, Office of Air Quality Planning and Standards, Research Triangle Park, USA. 12 pp.
- 60) Vallero (2014): *Fundamentals of Air Pollution* 5th Edition 2014.
- 61) Venkataraman (1993) A systematic account of some south Indian diatoms. *Proceeding of Indian Academy of Science* Vol. X No.6 Sec.B.
- 62) Weiss, R.F., (1970): The solubility of nitrogen, oxygen and argon in water and seawater. *Deep Sea Research and Oceanographic Abstracts* 17(4): 721-735.
- 63) Yap C.K., Chee M.W., Shamarina S., Edward F.B., Chew W. and Tan S.G. (2011). Assessment of Surface Water Quality in the Malaysian Coastal Waters by Using Multivariate Analyses, *Sains Malaysiana* 40 (10)1053–1062

Annexure -F

DEENDAYAL PORT AUTHORITY



Administrative Office Building
Post Box NO. 50
GANDHIDHAM (Kutch).
Gujarat: 370 201.
Fax: (02836) 220050
Ph.: (02836) 220038

www.deendayalport.gov.in

NO.EG/WK/4751/Part (Greenbelt-GUIDE) 196

Dated : 31/5/2022

✓ M/S Gujarat Institute of Desert Ecology,
P.O.Box No. 83,
Opp. Changleshwar Temple, Mundra Road,
Bhuj (Kachchh)- 370 001, Gujarat (India).
Tel.: 02832-329408, 235025.
Tele/Fax: 02832-235027

Email: desert_ecology@yahoo.com

Kind Attn.: Dr.V.Vijay Kumar, Director, M/s GUIDE, Bhuj.

Sub: Greenbelt Development in Deendayal Port Authority and its Surrounding Areas Charcoal site (Phase-I).

Ref.: M/s GUIDE, Bhuj offer vide letter no. M/s GUIDE, Bhuj vide communication no. GUIDE/DPA/GRN/080/2022-23 dated 24/5/2022.

Sir,

Your offer for the subject work submitted vide above referred letter dated 24/5/2022 amounting to Rs. 38,22,900.00 + applicable GST (Rupees Thirty-Eight Lakhs Twenty-Two Thousand and Nine Hundred Only Plus Eighteen Percent GST), with all terms & conditions mentioned in the offer letter, has been accepted **(Copy of offer letter M/s GUIDE attached)**.

2. Scope of work:

Development of Greenbelt in Charcoal site – Kandla, DPA and its surrounding areas. The activities under the Greenbelt Development include; inventory of suitable sites for greenbelt development in DPA, soil & Moisture conservation and management at Plantation sites, selection of suitable species of Plants for plantation, Procurement and plantation of plant saplings and seeds (5000 plants), along with management and monitoring of plantation, including drip/tanker water supply for a period 1 year.

.....Cont.....

3. Obligation of Deendayal Port Authority :

- Assistance regarding the statutory clearance from authorities concerned to be rendered by DPA for field visits/plantation activities.

4. The Terms of Payment:

1. 50% of the project budget to be paid to GUIDE within 15 days from the date of acceptance of Work order by GUIDE.
2. 20% of the project budget to be paid to GUIDE within 15 days from the date of completion of plantation works.
3. 20% of the project budget to be paid to GUIDE within 15 days from the date of submission Progress Report (December 2022).
4. 10% of the project budget to be paid to GUIDE within 15 days from the date of submission of Final Completion Report (May 2023).

5. Time Period : One year (from 5/6/2022 to 4/6/2023).

6. Kindly send the acceptance of this work order & start the work w.e.f. 5/6/2022 .

Thanking you.

Yours faithfully,



Superintending Engineer (PL) & EMC (I/c)
Deendayal Port Authority

Copy To :1) A.O.(W/A) - The proposal has been approved by the Board in its meeting held on 27/5/2022.

The expenditure shall be charged to the scheme
Environmental Services & Clearance thereof
(Allocation: 841/587/9744 WC - 5-13001).

- 2) TPA to CE for kind information of the Chief Engineer, please.
- 3) DA (PL) for further necessary action.
- 4) M/s Precitech Laboratorie ,Vapi, Environmental Management Cell to coordinate with M/s GUIDE,Bhuj.
- 5) RAO, DPA

Annexure -G

Brief Report (Second Season)

Studies on Dredged Materials for the presence of Contaminants and suggesting suitable disposal options

(As per EC & CRZ Clearance accorded by the MoEF & CC, GoI dated 19/12/2016 - Specific Condition No. vii)

DPA Work order No. EG/WK/4751/Part (EC&CRZ-1) / 84. Dt. 18.09.2021.

Submitted by

Gujarat Institute of Desert Ecology
P.B. No. 83, Mundra Road
Opp. Changleshwar temple
Bhuj - Kachchh, Gujarat – 370001, India

Submitted to

Deendayal Port Authority
Administrative Office Building
Post Box NO. 50
Gandhidham (Kachchh)
Gujarat - 370201


October 2022

Project Team

Project Co-Ordinator : Dr. V. Vijay Kumar, Director

S. No	Name & Designation	Role	Background
1.	Dr. K. Karthikeyan Principal Scientist & Head	Principal Investigator	M.Sc., Ph.D. in Environmental Sciences; 15 years of experience in water and sediment studies.
2.	Dr. G. Jayanthi Scientist	Co- Investigator	MSc., MPhil., PhD in Botany; 13 years of Research and teaching experience inclusive of Post-Doctoral experience for 5 years.
3.	Dr. Krushnakant D Baxi Scientific Officer	Co- Investigator	Ph.D in Zoology (Marine Biology) with 5 years of experience
4.	Mr. T. Dhananjayan Sr. Scientific Assistant	Team Member	M.Sc. in Environmental Sciences; 8 years of experience in sediment, water analysis and instrumentation.
5.	Ms. Dipti Parmar Jr. Scientific Assistant	Team member	M.Sc. in Environmental Sciences; 4 years of experience in sediment and water analysis.

Dr. V. Vijay Kumar
Director



Gujarat Institute
of Desert Ecology

Certificate

This is to state that the **Second Season report** of the work entitled, “**Studies on dredged material for the presence of contaminants**” has been prepared in line with the Work order issued by DPA vide No.EG/WK/4751/Part (EC&CRZ-1) / 84. Dt. 18.09.2021 as per the EC & CRZ Clearance accorded by the MoEF & CC, GoI dated 19/12/2016, Specific Condition No. vii. The work order is for a period of Three years from November 2021 – October 2024 for the above-mentioned study.

This Second Season report is for the project period from November 2021 – October 2022.

Authorized Signatory



Institute Seal

P. O. Box No. # 83, Opp. Changleshwar Temple, Mundra Road, Bhuj (Kachchh) - 370 001, Gujarat (India)

Tel : 02832 - 235025 Tele / Fax : 235027

www.gujaratdesertecology.com, E-mail : desert_ecology@yahoo.com

CONTENTS

Chapter No.	Title of the Chapter	Page No.
1	Background	4
2	Sediment Quality (Physico-chemical)	8
3	Sediment Quality (Biological)	18
4	Water Quality (Physico-chemical)	26
5	Water Quality (Biological)	35
6	References	59

One among the twelve major ports of the country, Deendayal Port is located at the tail end of Gulf of Kachchh, which is a largest Creek based Ports in the county which is located in the north-western coast of India in the state of Gujarat. DPA caters the maritime trade requirement of many hinterland states and is well connected by the network of rail and road and serves as a gate way port for export and import of northern and western Indian states of Jammu & Kashmir, Delhi, Punjab, Himachal Pradesh, Haryana, Rajasthan, Gujarat and parts of Madhya Pradesh, Uttaranchal and Uttar Pradesh. About 35% of the country's total export takes place through the ports of Gujarat in which the contribution by Deendayal port is considerable. The port handled a total cargo of 105 MMTPA during 2016-17, 110 MMTPA during 2017-18, 115 MMTPA during 2018-19, 122.5 MMTPA during 2019-2020 and 117.5 MMTPA during 2020-21. DPA is the only major Indian port to handle more than 127 MMT cargo throughput, and it has also registered the highest cargo throughput in its history. The port has handled a total of 3151 vessels during FY 2021-22.

Further, regular expansion of infrastructure and port facilities is under way to cater future logistic requirements. With such capacity, the Port ranks No. 1 among all the major ports in India for 12th Consecutive year. Further, a regular expansion of infrastructure and port facilities is under way to cater future logistic requirements. The port has high commercial importance in the Indian maritime trade as it handled 36.1 million tons (17%) of Cargo out of total Cargo of 213.1 million tons of the maritime Cargo of India during 2015. In addition, regular expansion of infrastructure and port facilities is under way to cater future logistic requirements.

In recent times, Deendayal Port Authority (DPA) has taken up Development of 7 Integrated facilities, and the Ministry of Environment, Forest and Climate Change (MoEF & CC), has put up some conditions while according Environmental and CRZ clearance. One of the conditions is to carry out the *“Study on Dredged Material for presence of contaminants”* as accorded by the MoEF&CC,GoI dated 19/12/2016 - Specific condition no. vii)” which states that *“Dredged materials should be analyzed for presence of contaminants and also to decide the disposal options. Monitoring of dredging activities should be conducted and the findings should be shared with the Gujarat SPCB and Regional Office of the Ministry”*.

1.1 Need of the study

Based on the above condition, DPA has assigned the task of carrying out the study to Gujarat Institute of Desert Ecology (GUIDE), Bhuj. This study will be attempted three times in a year at two specified locations. Further, the study will envisage the evaluation of physico-chemical constituents in the dredged materials in the dumped locations in the study area. GUIDE has received the Work order for this project with project time period being Three years (01.11.2021 – 31.10.2024). In this connection, the study was taken up for evaluation of dredged materials for the presence of contamination was conducted with the methodical investigation of evaluating physical, chemical and biological characteristics of the dredged materials with special reference to pollutants including heavy metal, Petroleum hydrocarbon etc.

1.2. Scope of the study

- a. To monitor the locations where dredged materials are dumped will be conducted.
- b. Dredged materials in the area will be analyzed for the presence of contaminants in two different locations.
- c. Detailed assessment of the dredged materials for physical, chemical and biological characteristics will be studied.
- d. Suggesting suitable disposal options for the dredged material will be made.

1.3. Sampling locations for 2021-22

The study on the presence of contaminants in the dredged materials for the year 2021-22 was designed by considering the location details (Table 1 and Plate 1) as provided to DPA by Hydraulic & Dredging Division regarding location of dumping ground and the details has been shared to GUIDE by DPA in the e-mail dated 24 October 2018. Three seasonal study covering Location 1, Location 2 and Location 3 with the Second season of the study was conducted during 20.04.2022 – 22.04.2022.

Table 1: GPS Co-ordinates of sampling locations

Station	Latitude (N)	Longitude (E)
Location 1 (Offshore)	22° 51' 00" N	70° 10' 00" E
Location 2 (Cargo jetty)	22°56' 31" N	70 13' 00" E
Location 3 (Phang Creek)	23° 04' 28" N	70°13' 28" E

1.4. Details of work done during 3rd Quarter (May – July 2022)

In this quarter, as part of Second season sampling, during April 2022, bottom water and sediment samples were collected from the Offshore and Creek system in three designated locations as earmarked was done. All the samples were subjected for various Physical, Chemical and Biological characteristics both in water (36 Nos.) and sediment samples (18 Nos) following standard methods as prescribed by ICMAM 2012. All the samples were done in triplicates and the data was compared with the limits as prescribed by CPCB for marine waters or with other relevant standards.

Plate 1: Map showing locations of proposed sampling (2021-22)



For the purpose of sediment characterization, the samples from the study area were collected employing standard methodology and the analysis of the samples were also performed as per standard protocol and the data of sediment analysis is presented in this Chapter 1. The sediment samples were collected in pre-fixed stations using a Van-veen type of grab sampler. After collection, the sediment samples were preserved with Rose Bengal and formalin to avoid decomposition of samples and processed for analysis and the samples after collection were brought to the laboratory on the same day of collection and air dried and used for further analysis for the test parameters (Table 2).

Table 2: Physico-chemical and biological characteristics of sediment samples

S. No.	Physico-chemical and Biological parameters
1	pH (1: 10 suspension)
2	Salinity (ppt)
3	Sand (%)
4	Silt (%)
5	Clay (%)
6	Total organic carbon (%)
7	Phosphorus (mg/kg)
8	Sulphur (mg/kg)
9	Petroleum Hydrocarbon ($\mu\text{g/kg}$)
10	Cadmium (mg/kg)
11	Lead (mg/kg)
12	Chromium (mg/kg)
13	Copper (mg/kg)
14	Cobalt (mg/kg)
15	Nickel (mg/kg)
16	Zinc (mg/kg)
17	Magnesium (mg/kg)
18	Macrobenthos

2.1. pH and Salinity (1: 10 suspension)

pH of the sediment is the measure of H^+ ion activity of the sediment water system. It indicates whether the sediment is acidic, neutral or alkaline in nature. Since ions are the carrier of electricity, the electrical conductivity (EC) of the sediment water system rises according to the content of soluble salts. The measurement of EC can be directly related to soluble salts concentration of the sediment at any particular temperature. Ten gram of the finely sieved sediment will be dissolved in 100ml of distilled water to prepare a leachate. This will be subjected to vigorous shaking using a rotator shaker for 1 hour to facilitate proper homogenization of the suspension. The suspension will be allowed to settle for two 2 hours and the supernatant after filtration will be used for the analysis of pH and salinity using the pH and EC meter (Make: Systronics 361) and Refractometer (Make: Atago). Each sample will be analysed in triplicates and the mean values will be taken into consideration.

2.2. Textural analysis (Sand/Silt/Clay)

Sediments will be collected using Van Veen grab whereas intertidal sediments will be collected using a handheld shovel. After collection, the scooped samples will be transferred to polythene bags, labeled and stored under refrigerated conditions. The sediment samples will be thawed, oven dried at 40°C and ground to a fine powder before analyses.

For texture analysis, specified unit of sediment samples will be sieved using sieves of different mesh size as per Unified Sediment Classification System (USCS). Cumulative weight retained in each sieve will be calculated starting from the largest sieve size and adding subsequent sediment weights from the smaller size sieves. The percent retained will be calculated from the weight retained and the total weight of the sample. The cumulative percent will be calculated by sequentially subtracting percent retained from 100%.

2.3.Total organic carbon

Total organic carbon is the carbon stored in sediment organic matter which enters the sediment through the decomposition of plant and animal residues, root exudates, living and dead microorganisms, sediment biota etc. Total Organic carbon in the sediment is oxidized with potassium dichromate in the presence of concentrated sulphuric acid. Potassium dichromate produces nascent oxygen, which combines with the carbon of organic matter to produce CO_2 . The excess volume of $K_2Cr_2O_7$ is titrated against the standard solution of ferrous ammonium Sulphate in presence of H_3PO_4 using Ferroin indicator to detect the first

appearance of unoxidised ferrous iron and thus volume of $K_2Cr_2O_7$ can be found out which is actually required to oxidize organic carbon.

Procedure

Percentage of Total organic carbon in the sediment/sediment will be determined by oxidizing organic matter in the sediment samples by chromic acid and estimating excess chromic acid by titrating it against ferrous ammonium sulphate with ferroin as an indicator. The detailed step-by-step procedure is as follows:

One gm of 0.5 mm sieved sediment will be weighed and put into 500 ml conical flask and to which 10 ml of 1N $K_2Cr_2O_7$ will be added with pipette and swirled. Immediately using a burette, 20 ml Conc. H_2SO_4 will be added and mixed gently until sediment and reagents are mixed. The reaction will be allowed to proceed for 30 min in a marble stone to avoid the damage caused due to release of intense heat due to reaction of sulphuric acid. Further, 200 ml of distilled water will be added slowly and 10 ml of concentrated Orthophosphoric acid and about 0.2 gm NaF will be added and allowed the sample and reagent mixture to stand for 1.5 hrs because the titration end point is better visible in a cooled solution. One ml of ferroin indicator will be added into the conical flask just before the titration and then titrated the excess $K_2Cr_2O_7$ with 0.5 N Ferrous Ammonium Sulphate till the color flashes from yellowish green to greenish and finally brownish red at the end point. Simultaneously a blank test will be also run without sediment sample.

2.4.Total Phosphorus

Phosphorus in sediment is commonly performed by Bray's extraction method and in this method, specific colored compounds are formed with the addition of appropriate reagents in the solution, the intensity of which is proportionate to the concentration of the element being estimated. The color intensity is measured spectrophotometrically. In spectrophotometrically analysis, light of definite wavelength (not exceeding say 0.1 to 1.0 nm in band width) extending to the ultraviolet region of the spectrum constitutes the light source. The photoelectric cells in spectrophotometer measure the light transmitted by the solution.

Fifty ml of the Bray's extractant will be added to 100 ml conical flask containing 5 gm of sediment sample and shaken for 5 minutes and filtered. Exactly 5 ml of the filtered sediment extract will be taken with a bulb pipette in a 25 ml measuring flask and 5 ml of the molybdate reagent with an automatic pipette will be added and diluted to 20 ml with distilled water and shaken well. Further, to this, 1 ml of the dilute Stannous Chloride solution will be added and

volume made upto 25 ml mark and shaken thoroughly. The mixture will be kept for color development and after 10 minutes the readings will be taken in the spectrophotometer at 660 nm wave length after setting the instrument to zero with the blank prepared similarly but without the sediment.

2.5. Total Sulphur

Sulphur in the sediment extract was estimated turbidimetrically using a spectrophotometer. The standards of sulphur were prepared in series such as 2, 4, 6, 8 and 10 ppm working solution from stock solution. In this, 25ml of solution was added in the volumetric flask separately to each flask and 2.5 ml of conditioning reagent solution was also added followed by 5 ml of extraction solution was added. To this mixture, 0.2-0.3 gm of barium chloride was also added and shaken well and made-up to 25 ml with distilled water and the readings were taken at 340nm spectrophotometer.

The sample was analysed by taking 5g of marine sediment into a 100ml conical flask, to which, 25 ml of 0.15 % CaCl_2 solution was added and shaken for 30 minutes. Then this was filtered through Whatman no. 42 filter paper and then 5 ml of sample aliquot was taken in a 25 volumetric flask, to which 2.5 ml of conditioning reagent and 0.2 to 0.3 g of barium chloride powder was added and made up to 25 ml distilled water and shaken well for 2 minutes and the absorbance was read in the same manner as standard solutions.

2.6. Petroleum Hydrocarbons

Sediment after refluxing with KOH-methanol mixture will be extracted with hexane. After removal of excess hexane, the residue will be subjected to clean-up procedure by silica gel column chromatography. The hydrocarbon content will be then estimated by measuring the fluorescence as per standard method.

2.7. Heavy metals

Heavy metals are of concern especially as it relates to the environment are Cadmium (Cd), Lead (Pb), Chromium (Cr), Nickel (Ni), Cobalt (Co), Copper (Cu), Zinc (Zn), Manganese (Mn) etc. For the release of mineral elements from sediment and sediments, wet oxidation of samples are generally performed. Wet oxidation employs oxidizing acids (Tri / Di-acid mixtures).

Sediment sample will be weighed to 1.0 gm and taken in 100ml beaker covered with a watch glass and 12 ml of Aqua regia in (1: 3 HNO_3 : HCl) will be added and the beaker will be

kept in digestion for 3 hours at 100⁰c on a hot plate using sand bath and the samples will be evaporated to near dryness and the samples will be kept cool for 5 mins and then 20 ml of 2% nitric acid will be added and kept for 15 minutes in hot plate for digestion and remove from hot plate and cooled and filtered using Whatman No. 42 mm filter paper and then the final make up to 50 ml with 2 % nitric acid will be made. The extracted sample will be then aspirated to an AAS.

2.8. Results

2.8.1. pH (Hydrogen Ion)

pH values in marine sediments, subatomic concentrations in seawater and deposited in the sediment core. However, these processes are generally depending with cycles of carbon, oxygen, nitrogen, phosphate, silicate, sulphur, iron and manganese and are associated with processes such as heterotrophic respiration, chemoautotrophic activity, photosynthesis, precipitation, and dissolution of calcium carbonate marine water and sediments. In the present investigation pH average values were recorded to be 7.95 ± 0.11 in the offshore, 8.04 ± 0.08 in the cargo jetty and 7.71 ± 0.34 in the Phang creek. Among all the stations, the maximum concentration of pH was recorded to be 8.17 in the cargo jetty station and the minimum concentration of pH was recorded to be 7.02 in the Phang creek station Fig.1




Fig .1 pH (Hydrogen ion) values in the various stations at Deendayal Port

2.8.2. Salinity

In the marine water and sediment, salinity typically varies from 0 to 36 ppt in most estuaries with hyper salinity occurring in many semi-enclosed bays. As well as, salinity concentration is associated with water temperature typically oscillates in diurnal and seasonal cycles in response to atmospheric temperature. In this study, during season two, salinity was observed to the highest concentration of 24.73 ppt in the phang creek station and the lowest concentration of salinity was found to be 7.78 ppt in the offshore station and mean \pm SD salinity of 9.63 ± 2.89 ppt in the offshore station, 21.73 ± 1.30 ppt in the cargo jetty station and 22.36 ± 2.01 ppt in phang creek station. Among all the stations values shown in Fig.2.




Fig.2 Salinity concentration in the various stations at Deendayal Port

2.8.3. Sediment Texture

One of the most important physical characteristics is sediment texture which (Sand, Silt and Clay) to marine benthic groups, in the study was investigated in different stations sediment texture, in which highest sediment texture percentage was observed of sand 54.80 % in the cargo jetty, silt 68.80% in the offshore station and clay 55.30% in the cargo jetty and lowest sediment texture percentage was observed of sand 10.10% in the phang Creek, silt 14.20% in the cargo jetty and Clay 10.60% in the offshore stations and among all the stations and the data shown in the Fig.3




Fig.3. Sediment texture average values in various stations at Deendayal port

2.8.4. Total organic Carbon

The organic carbon in the marine sediment are mainly coming from decomposition from animals, plants and anthropogenic sources such as chemical waste, fertilizers and organic – rich wastes which enrich the marine environment and that organic load settling to the bottom sediments from water column, in the path way that TOC values increasing and it affects the faunal communities. During season two, that TOC mean \pm SD % of $0.41 \pm 0.17\%$ in the offshore station, $0.69 \pm 0.21\%$ in the cargo jetty station and $0.67 \pm 0.09\%$ in phang creek station, among all the station TOC concentration shown in the Fig .4.




Fig.4. Total organic carbon concentration in various stations at Deendayal port

2.8.5. Organic matter

In the marine sediment organic matter is the major reservoir of organic carbon, which is a chemical, physical and biological effect of degradation to produce the organic matter in marine environment. Moreover, composed of material derived from the various planktons and benthic species that comprise the ecology of primary producers and consumers in overlying surface sediment. In the study, during season two, determined the organic matter ranged between 0.41 to 1.50 % among all the stations data shown in the Fig.5.




Fig.5. Organic matter concentration in various stations at Deendayal port

2.8.6. Phosphorus

Phosphorus (P) is an essential nutrient for life that plays a key role in regulating primary productivity in the marine systems. During season two, the maximum concentration of phosphorus was found to be 93.17 mg/kg in the Cargo jetty station and the minimum concentration of phosphorus was found to be 2.70 mg/kg in the offshore station and the average \pm SD being 10.09 ± 4.17 mg/kg in offshore, 30.28 ± 31.16 mg/kg in cargo jetty and 13.82 ± 4.10 mg/kg in phang creek.

2.8.7. Sulphur

Sulphur is a most significant primary source in sediments, the oxidation of sulphur and subsequent processing of oxidation intermediates. However the sulfur cycle of marine sediments is primarily driven by the dissimilatory sulfate reduction to sulfide by anaerobic microorganisms. In the present study, we aimed to examine the sulphur concentration which varies in different seasons, during season two, the maximum concentration of sulphur was recorded to be 28.08mg/kg in the phang creek and the minimum concentration of sulphur was recorded to be 13.0mg/kg in the offshore station, among all stations data shown in Fig.6.




Fig.6. Sulphur concentration in various stations at Deendayal port

2.8.8. Petroleum hydrocarbon

Petroleum hydrocarbon (PHC) contaminating the marine environment which comprises mainly of three classes of groups such as alkanes, olefins, and aromatics. Moreover, the petroleum hydrocarbons has less solubility in marine water and adsorbing by particulate matter showing a long-term persistence on the bottom of sediments and it cause a significant negative impact on benthic aquatic communities in the marine ecosystem. During season two, various stations the PHC ranged between 1.25 to 2.26 μ g/kg and the maximum was observed to be 2.26 μ g/kg in the offshore stations and minimum was observed to be 1.25 μ g/kg in the cargo jetty station Fig.7




Fig. 7. Petroleum hydrocarbon concentration in the various stations at Deendayal port

2.8.9. Heavy metals

The maximum concentration of heavy metals in the sediment samples for the metals such as Nickel, Lead, Cadmium, Chromium, Zinc, Copper, Manganese and Cobalt are in the levels 29.34 mg/kg, 1.44 mg/kg, 0.70 mg/kg, BDL, 21.43 mg/kg, 0.38 mg/kg, 1.50 mg/kg and 8.55 mg/kg respectively. Whereas, the mean concentration of the metals were in the Cargo Jetty location where in the range of 60.46 mg/kg, 9.64 mg/kg, 0.47 mg/kg, BDL, 42.06 mg/kg, 1.02 mg/kg, 3.27 and 33.73 for the metal species Nickel, Lead, Cadmium, Chromium, Zinc, Copper, Manganese and Cobalt respectively. In case of the creek system, the metal concentrations were observed in the range as 24.58 - 32.24 mg/kg for Nickel, 8.56- 13.57 mg/kg for Lead, 0.57-2.15 mg/kg for Cadmium, 0.11 -0.21 mg/kg for Chromium, 28.56-42.80 mg/kg for Zinc, 0.87 - 2.24 mg/kg for Copper, 3.56 - 10.28 mg/kg for Manganese and 9.85 - 18.75 in case of cobalt metal. During the second season, determination of magnesium in the different station in the study area was in the Average \pm SD of 449.53 \pm 155.33 mg/kg (Offshore site), 397.62 \pm 75.65 mg/kg (Cargo Jetty) and 467.60 \pm 29.75 mg/kg (Phang Creek) and the maximum and minimum was 612.48 mg/kg at Control site and 218.56 mg/kg at 1B (Offshore site) respectively.

3.1. Introduction

Earth is unique within the solar system to behold a large amount of water mostly contained in oceans. Life on earth originated in the oceans 3.1 to 3.4 billion years ago, and continuous mixing and dynamic characteristics of the ocean support very high biodiversity mostly unexplored. Indian peninsula surrounded by the Arabian Sea in the West, the Bay of Bengal in the East and the Indian Ocean in the South. The state of Gujarat is the western most in India and having the largest coastline of around 1600 km, along the Arabian sea with both Gulf of Kachchh and Khambhat. Gujarat coasts having different coastal ecosystems like the mangroves, sandy shores, muddy shores, rocky shores, mixed shores, wet sand shores, coral reefs and intertidal mudflats (Brink, 1993; Parasharya and Patel, 2014). Along with the high coastal diversity, there are developmental paradigm also and coastal development was also astonishing with the development of port for easy transportation. Deendayal Port Authority (DPA) is one among the 12 major ports of the country located near Gandhidham of Kachchh district. The port is the largest creek-based port in the country.

The word benthos originated from the Greek word *benthos* meant the depth of the seas. The benthic zone is the substratum zone of any water body mostly begins from the shore and reaches to the bottom of the waterbody and consists of organism living on and attached to or burrowing in the sediments commonly termed as benthos. Benthic community includes diverse group of animals including Gastropod and Bivalve molluscs, corals, sponges, polychaetes and nematode worms, crabs, different crustaceans, echinoderms, etc. Benthos are important predators and scavengers within the food chain and cleans the sea floor or freshwater bodies. Benthic organisms, play an important role as a food source for fish and other higher level of organisms.

The sediments of benthic zone play an important role in providing nutrients for the organisms that live in the benthic zone. The up-down movement of the bottom sediments mainly occurred by these benthic organisms results in a rise of the oxygen concentration of water and hence the overall productivity of the water bodies rich in high level of productivity. Major factors affect which benthic community are depth of water, salinity, temperature, types of substrate, pre-predation ratio and sudden changes in environmental condition. Nowadays, different anthropogenic activities affect aquatic systems including substratum habitat. Most of these animals lack a backbone and are called invertebrate animals.

Based on size, Benthos mainly divided into 3 types namely, Macrobenthos (> 1 mm), Meiobenthos (< 1 mm or > 0.1 mm) and Microbenthos (< 0.1 mm). These animals are further divided into two types Phytobenthos and Zoobenthos and based, on location it is furthermore classified as, Endobenthos, Epibenthos, Hyperbenthos.

The study was conducted summer season at 3 sites of Deendayal Port Authority with the locations namely, Offshore, Cargo Jetty and Phang Creek.

3.2. Methodology

To study the benthic organisms, triplicate samples were collected at each station using Van-veen grab which covered an area of 0.1m². The wet sediment was sieved with varying mesh sizes (0.5 mm-macrofauna) for segregating the organisms. The organisms retained in the sieve were fixed in 5-7% formalin and stained further with Rose Bengal solution for easy spotting at the time of sorting. The number of organisms in each grab sample was expressed as number/ meter square (No/m²). All the species were sorted, enumerated and identified to the advanced taxonomic level possible with the consultation of available literature. The works of Fauvel (1953), Day (1967) were referred for polychaetes; Barnes (1980) and Lyla *et al.* (1999) for crustaceans; Subba Rao *et al.* (1991) and Ramakrishna (2003) for molluscs.

Further, the data were treated with univariate statistical methods in PRIMER (Ver. 6.) statistical software (Clarke and Warwick, 1994)

a) Shannon – Wiener index

In the present study, the data were analyzed for diversity index (H') by following Shannon – Wiener's formula (1949):

$$H' = -\sum^S P_i \log_2 P_i \dots\dots i = 1$$

which can be rewritten as

$$H' = \frac{3.3219 (N \log N - \sum ni - \log ni)}{N}$$

where, H' = species diversity in bits of information per individual

ni = proportion of the samples belonging to the ith species

(number of individuals of the ith species)

N = total number of individuals in the collection and

Σ = sum

b) Species richness(S) was calculated using the following formula given by Margalef (1958)

c) Margalef index (d)

$$d = (S-1) / \log N$$

d) Pielou's evenness index

The equitability (J') was computed using the following formula of Pielou (1966):

$$J' = \frac{H'}{\log_2 S} \text{ or } \frac{H'}{\ln S}$$

Where, J' = evenness; H' = species diversity in bits of information per individual and S = total number of species.

3.3. Results on Species Composition, Population density and Biomass of Macrofauna of selected sites

3.3.1. Location 1 - Offshore site

Data collection was done at six sites (1A, 1B, 1C, 1D, 1E and 1- control). A Total of 4 groups of Benthic community were recorded in all stations at Offshore sites and they are Bivalves, Crustaceans, Gastropods (Mollusca) and Scaphopoda (Mollusca). Data on Density and Biomass expressed in (Nos/m²), (gm/m²) respectively.

Highest population density of benthic organisms was recorded in station 1E-Offshore (2350 nos/m²), whereas lowest in station 1D-Offshore (1425nos/m²). The density range of all stations varied from 1425 nos/m² to 2350 nos/m². Bivalves and Gastropoda were more abundant among all the benthic organisms (Table 3). The highest biomass value (expressed wet weight) of benthic fauna was observed in station 1B-Offshore (8.41 gm/m²) and lowest value was 1E-Offshore (4.14 gm/m²) (Table 3).

3.3.2. Cargo Jetty

In Cargo Jetty, frequently observed Benthic groups were Bivalves-Gastropods than Scaphopoda (Mollusca), and Razor clam (Bivalves). The population density range of 1100 to 4000 nos/m² was recorded between all the stations (Cargo Jetty2A, 2B, 2C, 2D, 2E & 2-Control) during the study period. Highest and Lowest density were recorded in station 2E-Cargo Jetty (4000 nos/m²) and 2B-Cargo Jetty (1100 nos/m²) respectively. The Biomass value indicated a highest value in station 2A- Cargo Jetty (13.86 gm/m²) and lowest in 2B-Cargo Jetty (5.08 gm/m²) (Table 3 and Fig. 8).

3. Phang creek

Six Stations of Phang creek were selected for the study namely 3A, 3B, 3C, 3D, 3E and 3-control-Phang creek. In this Phang creek benthic organisms were represented by Bivalves, Gastropods & *Razor clam* (Mollusca). The population density was highest in station 3Control-Phang creek (3400 nos/m²) and on the other side, lowest density was recorded in

3A-Phang creek (1200 nos/m²). Station 3D-Phang creek comprises highest wet wt (11.81 gm/m²), whereas low at was recorded in 3A-Phang creek (0.87 gm/m²).

Overall result of macrofaunal community showed highest population density in 2E-Cargo Jetty (400 nos/m²) and biomass observed in 2A-Cargo Jetty (13.86 gm/m²). Table 3 showed highest population values of Bivalves in 2E- Cargo Jetty (2800 nos/m²) and lowest value comprised by Scaphopoda 50 nos/m² at 2D and 2-Control (Cargo Jetty). *Optedicerus breviculum* (Common name Mangrove snail-Small Gastropoda shell) was only recorded at 3B and 3-Control site of Phang Creek. The Muddy habitat of Phang creek is preferred for many benthic organisms. This might be due to relatively stable and less polluted environment provided by muddy creek area of Phang creek further added that very low level of predation pressures on benthic community and also might be due to lesser anthropogenic activities in that area. Table 3 showed that average population density and biomass higher in Cargo Jetty area where mostly rocky or covered with coral base providing a unique habitat for gastropod, bivalves and other benthic organisms.

Frequently found species at all sites were *Pirenella cingulata*, *Umbonium vestiarium*, *Optedicerus breviculum*, *Tellina* sp., *Clypeomorus bifasciata*, *Cly Pholas orientalis*, *Dentalium* sp *Dosinia* sp, *Donax* sp, *Anadara* sp, *Turris* sp etc. The percentage of occurrence (Table 3) revealed highest group present was Gastropoda (100%), Bivalves (94.44%) then followed by Razor clam (55.55%), Scaphopoda (38.88%) and others. Lowest percentage of occurrence by *Pirenella cingulata* (5.55%). Compared to three sites, lowest density and biomass was observed at Offshore area (Table 3 and Figure 9) which indicated pollution level or stressful environment, monsoon effect and also might be some chemical and biological changes in water. Detail status of Population density, Group composition and biomass of the benthic community of all selected sites were depicted in (Table 3) and (Figure 9). In all the stations, highest percentage composition recorded by Bivalves (53%) followed by Gastropoda (23%), Razor clam (7%), *Optedicerus breviculum* (5%), Scaphopoda (3%) and 1% comprises by Polychaete, *Pirenella cingulata*(gastropods) and Crustacean (Figure 9). Phytoplankton abundance and their size, zooplankton Body composition, patchy distribution of zooplankton, water currents, ebb and flow tides, and water churning process, changing in structure of muddy, rocky and sandy habitats are the main reasons for biomass and density fluctuation in Benthic communities. In Crustacean most commonly observed species are Crabs and attached Barnacles. Main Gastropods families recorded Trochidae, Cerithidea, Turritellidae, Mitridae and Bucciniae etc. *Nereis* sp, *Capitella* sp, *Nephtys* sp. like polychaete were observed in

samples. More number of the broken bivalves, debris, plat items and broken gastropods are frequently observed in the Microscope.

3.4. Diversity indices of Benthic Community

Table 4 shows various diversity indices calculation, showed that Shannon Diversity Index ranging from (0.444-1.547) indicated very low to near moderate diversity. Highest diversity indices was recorded in Station 3B-Phang creek (1.547) where moderate value of density and biomass of benthos and other side in 1A-Offshore diversity indices value was 0.444 where only two groups were present. The evenness values ranged between (0.634 to 0.960). The highest evenness value is 0.960 observed in station 1C-Offshore and the lowest evenness index value 0.634 was at station 2E-Cargo Jetty and where the population density was recorded highest. Simpson's Index value ranged between 0.273 to 0.776 indicated to lower to moderate diversity.




Figure 8. Population densities of Macro Benthos in various sites




Figure 9. Percentage composition of Macrobenthos in various sites

Table 3. Macrobenthos distribution in different sites of Deendayal Port

Name of Station	Offshore						Cargo Jetty						Phang creek						% of Occurrence
	1A	1B	1C	1D	1E	1-Control	2A	2B	2C	2D	2E	2-Control	3A	3B	3C	3D	3E	3-Control	
Name of Benthic Group																			
Bivalves (Mollusca)	1800	1100	725	1000	1550	1150	1925	625	1900	1100	2800	575	0	525	650	650	600	1000	94.44
Crustacean animals (Crabs, Mysis etc.)	0	0	350	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	5.55
Gastropoda (Mollusca)	350	425	550	425	800	375	1050	350	550	525	550	450	900	225	550	975	800	750	100
Polychaeta worms (Marine Annelids worms)	0	0	0	0	0	0	0	25	0	0	0	0	300	0	0	0	0	0	11.11
Scaphopoda (Mollusca)	0	150	0	0	0	0	225	100	225	50	350	50	0	0	0	0	0	0	38.88
Razor clam (Bivalvia)	0	0	0	0	0	0	325	0	250	225	300	275	0	225	300	125	300	400	55.55
Optediceros breviculum (Gastropoda)	0	0	0	0	0	0	0	0	0	0	0	0	0	500	0	0	0	1250	11.11
Pirenella cingulata (Gastropoda)	0	0	0	0	0	0	0	0	0	0	0	0	0	375	0	0	0	0	5.55
Total Population Density Nos/m²	2150	1675	1625	1425	2350	1525	3525	1100	2925	1900	4000	1350	1200	1850	1500	1750	1700	3400	-
Biomass wet wt gm/m²	6.61	8.41	8.26	5.38	4.14	6.64	13.86	5.08	8.3	7.22	10.45	6.49	0.87	7.68	10.66	11.81	9.51	8.94	-

Table 4: Diversity indices of benthic faunal groups at various station of Deendayal Port (Benthos)

Variables	Offshore						Cargo Jetty						Phang Creek					
	1A	1B	1C	1D	1E	1-Control	2A	2B	2C	2D	2E	2-Control	3A	3B	3C	3D	3E	3 - Control
Taxa_S	2	3	3	2	2	2	4	4	4	4	4	4	2	5	3	3	3	4
Individuals (Nos/m ²)	2150	1675	1625	1425	2350	1525	3525	1100	2925	1900	4000	1350	1200	1850	1500	1750	1700	3400
Dominance_D	0.727	0.504	0.360	0.581	0.551	0.629	0.400	0.433	0.471	0.426	0.522	0.335	0.625	0.224	0.362	0.454	0.377	0.284
Shannon Diversity	0.444	0.840	1.057	0.609	0.641	0.558	1.087	0.990	1.002	1.020	0.930	1.176	0.562	1.547	1.052	0.882	1.028	1.313
Simpson_1-D	0.273	0.496	0.640	0.419	0.449	0.371	0.601	0.567	0.530	0.574	0.478	0.665	0.375	0.776	0.638	0.547	0.623	0.716
Evenness	0.780	0.772	0.960	0.920	0.950	0.873	0.741	0.673	0.681	0.693	0.634	0.810	0.877	0.940	0.955	0.805	0.932	0.929
Menhinick	0.043	0.073	0.074	0.053	0.041	0.051	0.067	0.121	0.074	0.092	0.063	0.109	0.058	0.116	0.077	0.072	0.073	0.069
Margalef	0.130	0.269	0.271	0.138	0.129	0.136	0.367	0.428	0.376	0.397	0.362	0.416	0.141	0.532	0.274	0.268	0.269	0.369

4.1. Introduction

Rapid urbanization and industrial growth showed a significant impact on coastal ecosystems, such as estuaries and the surrounding coastal areas. The presence of a dense human population in their watersheds contaminates the environment (Jha et al., 2015). Coastal environment reference characteristics are necessary to provide a better management solution for the coastal ecosystem (Barbier Edward et al., 2011). Another major activity carried out in industrial port environment in the coastal environment is Dredging which is often carried out to create accesses to oil exploitation, marine/coastal transportation and other waterborne commerce. Dredging in sensitive environments is often accompanied by ecological impacts including damage to flora and fauna, alteration of coastal topography and hydrology, impairment of water quality etc (Adesobande and Associate, 1998). Hence assessing the water for various characteristics will indicate the intensity of pollutants present in such environments.

4.2. Materials and Methods

In the present study, the marine water and marine sediment samples were collected using standard protocol and analysis of the same was done following standard methods for marine water and sediment analysis as prescribed by APHA (2012), NIO manual (1982) and ICMAM Manual (2012). Surface water samples for general analysis were collected using a clean polyethylene bucket while an adequately weighted Niskin sampler was used to collect water samples from the bottom. A glass bottle sampler (1 L) was used for collecting water samples at 1 m below the surface. Parameters such as pH, Temperature, Salinity were recorded on spot using hand held meters and the same was also verified in the Laboratory. The water samples collected were stored in refrigerated conditions until further analysis of other parameters. As per the standard protocol, the fixatives and preservatives were added to the samples in case of parameters such as Dissolved Oxygen using Winkler A&B solution immediately, Chemical Oxygen Demand using concentrated H_2SO_4 to bring the <2 pH and preservation using nitric acid for heavy metals. In case of biological characteristics, the marine water samples for planktonic analysis were added with formalin. In general, all the collected water and sediment samples were stored in a sterile, polythene bottles and ziplock bags in an icebox to maintain suitable conditions till it is brought to the Laboratory. The list

of parameters (Table 5) and the method adopted for the analysis of samples are detailed below.

Table 5: Physico-chemical and biological characteristics of marine water samples

S. No	Physico-chemical and Biological parameters
1	pH
2	Salinity (ppt)
3	Total Dissolved Solids (mg/L)
4	Total Suspended Solids (mg/L)
5	Turbidity (NTU)
6	Dissolved Oxygen (mg/L)
7	Bio-Chemical Oxygen Demand (mg/L)
8	Chemical Oxygen Demand (mg/L)
9	Phenolic compound ($\mu\text{g/L}$)
10	Petroleum Hydrocarbons ($\mu\text{g/L}$)
11	Oil and grease (mg/L)
12	Cadmium (mg/L)
13	Lead (mg/L)
14	Chromium (mg/L)
15	Copper (mg/L)
16	Cobalt (mg/L)
17	Nickel (mg/L)
18	Zinc (mg/L)
19	Magnesium (mg/L)
20	Chlorophyll (mg/m^3)
21	Phaeophytin (mg/m^3)
22	Phytoplankton Phytoplankton cell counts (no/L) Total Genera (no.) Major Genera
23	Zooplankton Biomass ($\text{ml}/100\text{m}^3$) Population ($\text{no}/100\text{m}^3$) Total Group (no.) Major Groups

4.2.1. pH, Temperature and Salinity

A Thermo fisher pH / EC / Temperature meter was used for pH and Temperature measurements. The instrument was calibrated with standard buffers just before use. A suitable volume of the sample was titrated against silver nitrate (20 g/l) with potassium chromate as an indicator. The chlorinity is estimated and from that salinity values were derived using formula.

4.2.2. Total Dissolved Solids (TDS)

The samples were subjected for gravimetric procedure for confirmation of the readings obtained from the hand held meter. About 100 ml of the water sample was taken in a beaker and filtered which was then dried totally in a Hot Air Oven (105°C). TDS values were calculated using the difference in the initial and final weight.

4.2.3. Total Suspended Solids (TSS)

Hundred ml of the sample was filtered through each pre-weighed filter and placed in the Hot air oven at specified temperature as per the protocol for 1 hour. The filter paper was allowed to cool in a desiccator and obtain a constant weight by repeating the drying and desiccation steps.

4.2.4. Turbidity

The sample tube (Nephelometric cuvette) was filled with distilled water and placed in the sample holder. The lid of the sample compartment was closed. By adjusting the '*SET ZERO*' knob, the meter reading was adjusted to read zero. The sample tube with distilled water was removed and the 40 NTU standard solution was filled in the tube and the meter reading was set to read 100. Other standards were also run. The turbidity of the marine water sample was then found out by filling the sample tube with the sample, and the reading was noted.

4.2.5. Dissolved Oxygen (DO) and Biochemical Oxygen Demand (BOD)

DO was determined by Winkler's method. For the determination of BOD, direct unseeded method was employed. The sample was filled in a BOD bottle in the field and incubated in the laboratory for 3 days after which DO was again determined and the difference was calculated.

4.2.6. Chemical Oxygen Demand (COD)

A known quantity of sea water was placed in a 50 ml Erlenmeyer flask and to which 3.0 g of silver sulphate was added and kept in a magnetic stirrer for proper mixing at room temperature to remove the chloride interference in the form of Silver chloride precipitate. The sample with white precipitate turned to a fade lilac mixed coloured precipitate is the indication. At this point, mixing of samples was stopped and the flasks were kept at 40° inclined position. Sedimentation of the coloured precipitate was very quick and 20 ml of the cleared sea water was taken carefully from the upper end of the flask bottom after a rest period of 5-10 min. To the 20ml of sea water sample diluted with 150 ml of distilled water, to

which 10 ml of standard $K_2Cr_2O_7$ was added, to which 30 ml of Sulphuric acid was added. The tubes were connected to condensers and refluxed for 2 hours at $150 \pm 2^\circ C$. After refluxion, the flasks were allowed to cool and titrated against Standard Ferrous Ammonium Sulphate with Ferroin as Indicator. Green blue to wine red is the indication of the end point of the experiment and a blank was run under simultaneous conditions.

4.2.7. Phenolic compounds

Phenols in water (500 ml) were converted to an orange coloured antipyrine complex by adding 4-aminoantipyrine. The complex was extracted in chloroform (25 ml) and the absorbance was measured at 460 nm using phenol as a standard.

4.2.8. Petroleum Hydrocarbons (PHc)

Water sample (1 l) was extracted with hexane and the organic layer was separated, dried over anhydrous sulphate and reduced to 10 ml at $30^\circ C$ under low pressure. Fluorescence of the extract was measured at 360 nm (excitation at 310 nm) with Saudi Arabian crude residue as a standard. The residue was obtained by evaporating lighter fractions of the crude oil at $120^\circ C$.

4.2.9. Oil and Grease

About 500 ml of sample was transferred to the separating funnel and sample bottle was carefully rinsed with 30ml of trichlorotrifluoroethane and add the solvent washings was added to the separating funnel. To this, 5ml of 1:1 HCL was added and shaken vigorously for about 2 minutes If soluble emulsion was formed, then the sample container was shaken for 5 to 10 minutes. Then the layers were allowed to separate and the lower layer (organic layer) was discarded from separating funnel. Then the solvent layer was drained through a funnel containing solvent moistened filter paper into a clean pre weight distillation flask. Then solvent was distilled from distillation flask over a water bath at $70^\circ C$. Then the residue was transferred using minimum quantity of solvent into a clean pre weighed dried beaker and the beaker was placed on water bath for 15 minutes at $70^\circ C$ and evaporate off all the solvent and it was cooled in desiccators for 30 minutes and weight was taken.

4.2.10. Heavy metals

Metals are of great concern especially when it relates to the coastal environment as it has chances of biomagnification from lower organisms to higher organisms through water and sediment. Among common metals are Cadmium (Cd), Lead (Pb), Chromium (Cr), Copper (Cu), Cobalt (Co), Nickel (Ni), Zinc (Zn), Magnesium (Mg) etc. For the release of mineral

elements from sediment and sediments, wet oxidation of samples is generally performed. Wet oxidation employs oxidizing acids (Tri / Di-acid mixtures).

Sediment sample will be weighed to 0.5 gm and taken in 100ml beaker covered with a watch glass and 12 ml of Aqua regia in (1: 3 HNO₃ : HCl) will be added and the beaker will be kept in digestion for 3 hours at 100⁰c on a hot plate using sand bath and the samples will be evaporated to near dryness and the samples will be kept cool for 5 mins and then 20 ml of 2% nitric acid will be added and kept for 15 minutes in hot plate for digestion and remove from hot plate and cooled and filtered using Whatmann No. 42 mm filter paper and then the final make up to 50 ml with 2 % nitric acid will be made. The extracted sample will be then aspirated to an AAS.

4.3 Results

During the current year of study, three locations namely Offshore (Site 1), Cargo Jetty (Site 2) and Phang Creek (Site 3) were monitoring for various Physico-chemical characteristics in the marine water samples and the data is presented in Table 6-8. The description of the values recorded in each station is detailed as below.

4.3.1. Location 1 - Offshore location

The marine water samples in the Offshore locations revealed the pH values ranged between 7.83-8.06 with the average pH being 8.01 which was well within the prescribed limits for Coastal waters. In case of significant parameters like Phenolic compounds, Petroleum hydrocarbon and Oil & Grease, the maximum concentrations observed for the parameters are 19.55 µg/L, 21.61 µg/L and 4.0 mg/L. The data on different heavy metal concentrations observed in the sampling sites are given in Table 6.

4.3.2. Location 2 - Cargo Jetty

The mean pH value among the twelve samples collected in the Cargo Jetty samples are 8.037. Typical Kachchh water salinity concentrations were in the range of 39.33 - 42.79 ppt with the mean salinity of Kandla water was 40.917 ppt which is slightly higher than the salinity of any of the Indian coastal waters. Due to its tail end location, both the Turbidity and Total Suspended Solids concentrations are comparatively high in the waters with the maximum concentrations recorded as 73.59 NTU and 187.91 mg/L. The mean concentrations of Phenolic compounds, Petroleum hydrocarbon and Oil and Grease were recorded to be 13.564 µg/L, 29.290 µg/L and 1.033 mg/L. In addition to this, various toxic heavy metals were recorded which is presented in Table 7.

4.3.3. Location 3 - Phang Creek

In case of the creek system in DPA vicinity, Phang creek was monitored to understand the impact of disposed dredged materials as this is one the pre-designated sites. In this scenario, the pH value of the waters ranged between 8.01 - 8.1 and the mean pH value of this location is 8.055. Further, the possibility of higher load prevailing in the creek systems when compared to Offshore, the maximum concentration of Total Dissolved Solids, Total Suspended Solids and Turbidity concentrations were 43533 mg/L, 302 mg/L and 110.5 mg/L and these characteristics are indicator of a high turbidity nature of this area. Similarly, in case of major polluting parameters are concerned, the concentrations were 15.57 µg/L (Phenolic compounds), 42.38 µg/L (Petroleum hydrocarbon) and 7.2 mg/L in case of Oil and Grease. The highest concentration of Oil and Grease was found from this location. Similar to previous location metal data, the concentrations of metals recorded in the Phang creek is given in Table 8.

Table 6: Physico-chemical characteristics of the marine water from sampling location 1 (Offshore)

S. No	Parameters	1A		1B		1C		1D		1E		Control 1	
		SW	BW	SW	BW	SW	BW	SW	BW	SW	BW	SW	BW
1	Temperature (°C)	28.50	28.00	29.00	28.50	28.80	28.50	28.50	28.00	28.30	28.00	28.00	27.80
2	pH	8.01	8.00	8.01	8.01	7.96	7.83	8.05	8.03	8.03	8.03	8.06	8.04
3	Salinity (ppt)	40.20	36.74	34.15	38.04	36.31	37.17	34.58	36.31	37.60	35.87	38.04	35.44
4	Total Dissolved Solids (mg/L)	42368	42292	41527	42281	41219	41493	40084	40231	40759	41714	42215	42900
5	Total Suspended Solids (mg/L)	226.00	218.00	242.00	228.00	323.00	202.00	256.00	247.00	244.00	221.00	204.00	187.00
6	Turbidity (NTU)	120.10	60.10	153.90	132.90	141.30	139.20	108.20	100.80	146.70	133.60	158.10	104.50
7	Dissolved Oxygen(mg/L)	5.80	5.50	5.70	5.70	5.50	4.90	6.00	5.70	6.40	6.10	5.90	5.90
8	Bio-Chemical Oxygen Demand (mg/L)	1.30	1.20	1.60	1.00	2.30	1.80	1.10	1.10	1.70	1.40	0.90	0.90
9	Chemical Oxygen Demand (mg/L)	52	48	44	42	50	46	48	38	42	40	36	34
10	Phenolic Compounds (µg/L)	14.55	11.91	16.98	12.86	12.05	19.55	12.55	10.80	11.10	15.90	14.48	7.86
11	Petroleum Hydrocarbons (µg/L)	20.515	20.11	18.63	18.64	21.61	21.605	18.93	18.965	17.865	17.91	19.59	19.68
12	Oil and grease (mg/L)	3.20	4.00	2.80	2.80	2.00	2.00	0.80	1.20	2.80	1.60	4.00	2.80
13	Magnesium (mg/L)	1286.52	1187.56	1347.58	1287.98	1187.59	1045.89	1247.89	1148.98	1335.24	1258.47	1542.57	1422.24
14	Nickel (mg/L)	1.84	1.85	2.45	2.22	4.24	3.21	1.80	1.47	2.89	2.41	3.54	2.36
15	Lead (mg/L)	1.21	0.98	1.20	0.98	0.86	0.34	0.28	0.87	1.01	0.58	0.48	1.24
16	Cadmium (mg/L)	0.43	0.22	0.56	0.87	1.45	1.01	1.22	0.89	1.21	1.01	0.48	0.35
17	Chromium (mg/L)	BDL	BDL	BDL	BDL	BDL	BDL	BDL	BDL	BDL	BDL	BDL	BDL
18	Zinc (mg/L)	1.25	0.89	1.47	0.48	2.12	2.01	1.85	1.22	0.58	0.42	0.22	0.18
19	Copper (mg/L)	BDL	BDL	BDL	BDL	BDL	BDL	BDL	BDL	BDL	BDL	BDL	BDL
20	Manganese (mg/L)	1.85	1.48	2.22	2.15	1.48	1.54	0.89	1.22	1.78	1.45	1.62	1.50
21	Cobalt (mg/L)	3.25	2.54	1.28	0.89	2.54	2.48	2.47	1.45	1.58	0.98	1.48	1.22

Note: BDL denotes Below Detection Limit.

Table 7: Physico-chemical characteristics of the marine water from sampling location 2 (Cargo Jetty)

S. No	Parameters	2A		2B		2C		2D		2E		Control 2	
		SW	BW	SW	BW	SW	BW	SW	BW	SW	BW	SW	BW
1	Temperature (°C)	29	28.5	29	28.5	28.9	28	28.7	28.5	29	28.8	29.5	29.2
2	pH	8.09	8.09	7.94	7.92	8.05	7.96	8.08	8.06	8.08	8.05	8.06	8.06
3	Salinity (ppt)	42.79	39.33	41.49	40.63	41.49	40.63	40.63	39.77	41.06	41.06	41.06	41.06
4	Total Dissolved Solids (mg/L)	41146	42035	42887	42285	41929	41658	43796	42232	41308	42098	42762	41774
5	Total Suspended Solids (mg/L)	189	172	190	174	201	189	187	174	212	192	195	180
6	Turbidity (NTU)	67	65.7	74.1	73.9	99.8	96.7	54.2	53.6	98.5	64.1	55.6	79.9
7	Dissolved Oxygen(mg/L)	6.42	5.61	5.81	5.94	5.72	5.14	5.52	5.34	5.35	5.51	5.81	5.34
8	Bio-Chemical Oxygen Demand (mg/L)	1.2	0.5	1.2	0.8	0.9	0.7	0.5	0.42	1.02	0.72	1	0.3
9	Chemical Oxygen Demand (mg/L)	42	38	44	40	52	38	34	32	44	42	38	32
10	Phenolic Compounds (µg/L)	12.77	9.7	6.74	7.82	11.98	24.19	20.6	6.24	20.74	12.99	8.11	20.89
11	Petroleum Hydrocarbons (µg/L)	30.865	30.975	29.425	29.335	27.875	27.49	32.925	33.235	26.18	26.08	29.205	27.895
12	Oil and grease (mg/L)	0.8	1.2	0.4	0.8	0.8	1.2	0.4	1.6	1.2	0.4	2.8	0.8
13	Magnesium (mg/L)	1548.25	1347.23	1258.59	11875.69	1358.47	1258.47	1547.38	1482.36	1542.82	1462	1358.68	1284.49
14	Nickel (mg/L)	0.32	0.28	0.58	0.45	1.25	0.89	1.14	0.98	1.25	0.87	0.98	0.87
15	Lead (mg/L)	0.35	BDL	BDL	0.25	0.18	BDL	0.21	0.15	BDL	0.98	BDL	BDL
16	Cadmium (mg/L)	0.02	0.03	0.07	0.05	0.18	0.09	0.54	0.24	0.05	BDL	BDL	BDL
17	Chromium (mg/L)	BDL	BDL	BDL	BDL	BDL	BDL	BDL	BDL	BDL	BDL	BDL	BDL
18	Zinc (mg/L)	0.89	0.75	1.25	1.01	1.54	1.21	0.98	0.75	1.1	0.58	1.48	1.3
19	Copper (mg/L)	0.25	BDL	0.18	0.16	BDL	0.25	0.2	BDL	0.21	BDL	0.08	BDL
20	Manganese (mg/L)	3.21	2.58	3.11	3.18	2.45	2.78	1.58	1.48	2.01	BDL	BDL	0.28
21	Cobalt (mg/L)	1.22	BDL	1.22	0.89	0.45	1.32	0.89	BDL	0.21	BDL	0.22	0.67

Note: BDL denotes Below Detection Limit

Table 8: Physico-chemical characteristics of the marine water from sampling location 3 (Phang Creek)

S. No	Parameters	3A		3B		3C		3D		3E		Control 3	
		SW	BW	SW	BW	SW	BW	SW	BW	SW	BW	SW	BW
1	Temperature (°C)	29.2	29	28.8	28.7	28.5	28.3	29	28.8	30	29.5	28.9	28.7
2	pH	8.01	8.01	8.08	8.06	8.03	8.02	8.1	8.07	8.03	8.07	8.09	8.09
3	Salinity (ppt)	42..36	38.04	37.17	38.47	43.22	40.36	39.33	39.33	43.22	44.09	40.63	40.63
4	Total Dissolved Solids (mg/L)	40235	42246	43315	40769	42393	40806	41903	42599	41307	42128	43533	41175
5	Total Suspended Solids (mg/L)	245	221	287	262	302	274	287	268	301	289	301	278
6	Turbidity (NTU)	90	104.1	89.4	93.3	71.1	68.7	110.5	108.1	102.5	73.2	94.7	95.3
7	Dissolved Oxygen(mg/L)	5.54	5.31	5.72	5.51	5.32	5.1	5.37	5.24	5.38	5.11	5.47	5.26
8	Bio-Chemical Oxygen Demand (mg/L)	1.4	1.2	0.9	0.7	0.8	0.7	0.82	0.46	0.92	0.52	0.74	0.42
9	Chemical Oxygen Demand (mg/L)	38	32	40	34	42	36	48	40	34	32	40	36
10	Phenolic Compounds (µg/L)	12.27	13.35	15.57	14.71	11.98	15.14	13.35	15.49	5.38	10.26	12.77	10.19
11	Petroleum Hydrocarbons (µg/L)	24.93	25.07	35.14	35.325	42.285	42.38	25.38	25.44	21.875	21.85	26.005	27.325
12	Oil and grease (mg/L)	4	4.4	3.2	1.6	6.4	6	3.6	4	7.2	4.4	6.8	7.2
13	Magnesium (mg/L)	1536.65	1487.59	1325.25	1258.45	1456.25	1352.56	1478.59	1254.69	1458.87	1602.25	1458.56	1324.87
14	Nickel (mg/L)	BDL	BDL	0.85	0.48	1.22	0.25	1.02	0.89	1.21	0.22	0.45	1.12
15	Lead (mg/L)	0.03	BDL	0.52	0.42	BDL	BDL	BDL	BDL	0.24	0.32	BDL	BDL
16	Cadmium (mg/L)	0.18	0.11	0.25	0.45	0.36	0.34	0.48	0.35	0.48	0.35	BDL	0.25
17	Chromium (mg/L)	BDL	BDL	BDL	BDL	BDL	BDL	BDL	BDL	BDL	BDL	BDL	BDL
18	Zinc (mg/L)	0.35	0.28	0.48	0.32	1.28	0.65	1.04	0.75	0.46	0.57	0.32	0.25
19	Copper (mg/L)	0.32	0.18	BDL	0.24	0.62	0.21	BDL	0.58	0.4	0.08	0.2	0.34
20	Manganese (mg/L)	3.24	1.18	2.25	1.14	3.78	1.25	1.54	2.54	2.35	3.58	2.78	1.58
21	Cobalt (mg/L)	1.32	1.21	1.65	2.58	2.58	1.36	1.36	1.54	2.14	2.87	1.56	0.89

Note: BDL denotes Below Detection Limit

5.1. Introduction for Plankton

Planktons denotes a group of organisms either animal (zooplankton) or plants (phytoplanktons) origin. Major phytoplankton in sea water are Diatoms (Tiwari and Nair, 1998; Thakur et al, 2015), Cocolithophores, Silicoflagellates, Blue green algae (Cyanobacteria) and Dinoflagellates. Diatoms constitute the major part of the phytoplankton in sea water. Zooplankton comprises the second level in the food chain and includes Tintinnids, Foramoniferan, Radiolarians, Amphipoda, Copepoda, Calanoida, Chaetognaths, larvae of benthic invertebrates and fish larvae etc. (Gajbhiye and Abidi, 1993; Thirunavukkarasu, 2013; Chakrabarty et al. 2017). Many species spend their entire lifecycle as zooplankton, whereas, barnacles, Copepoda and other Crustacean includes different Nauplius stages (larval stages) of zooplankton within their lifecycle also known as meroplankton. The planktonic stages of invertebrates are economically important as a food for pelagic fishes. Zooplankton require a constant supply of oxygen (Dodson, 1992).

The zooplankton may be classified according to their habitat and depth, distribution, size and duration of planktonic life period (Omori and Ikeda, 1984). There are the two main classification on the bases of habitat which are Marine plankton or Haloplankton and Freshwater plankton or Limnoplankton. Marine plankton is further divided in to 3 types; Oceanic plankton, Neritic plankton and Brackish water plankton. Oceanic plankton or Off-shore plankton generally found in surface water and continental shelf region water whereas neritic zooplankton means occurring to continental zone to neritic or deep sea (Besiktepe et al, 2015). Brackish water plankton generally inhabiting brackish water like mangrove, estuaries and sea vegetation area.

Size is very important to understanding about the classification of both zooplankton and phytoplankton. Based on size, various categories of plankton are smallest one Picoplankton (0.2-2 μm), Nanoplankton (2-20 μm), Microplankton (20-200 μm), Mesoplankton (200 μm -2 mm), Macroplankton (2-20 mm) and Megaplakton(> 20 mm) .

Phytoplankton are primary producers of sea whereas as a primary consumer are zooplankton which play precious role to control the primary producers in sea. Benthic organisms and higher vertebrate animals uses plankton as a food material in Ocean life. Zooplankton and Phytoplankton are main prey food sources for different Fishes. The main food items of mesopelagic fishes are zooplankton larvae, juvenile fish and many small invertebrate animals.

Population of plankton and other marine living organisms on which the whole aquatic life depends directly or indirectly is largely governed by the interaction of a number of biological, chemical and physical processes and tolerance to one or more of these conditions (Reid and Wood 1976). Understanding of diversity and distribution of marine organisms would not be complete without

consideration of abiotic and biotic factors of marine environment. Amongst the various abiotic factors affecting the survival of marine invertebrates in coastal and estuarine regions, salinity and temperature are of primary importance (Rao and Balasubramanian, 1996; Sreenivasulu et al, 2017). Planktons are affected by changes in biotic and abiotic factors of environment and can rapidly respond to climatic changes. The population of plankton diversity is largely related to Seasonal and Monthly variability in Physical, Chemical and Biological parameters; Interspecific competition among the Zooplankton; Inter-relationship for prey and predator between zooplankton and their mostly predator animals; Grazing ratio of Zooplankton; Suspension of sediment; Fluctuation in Phytoplankton abundance; Waves, Currents and Tidal turbulence effect; Fluctuation in Chlorophyll a and Nutrients; Input of Organic and other Pollution creating sources; Fish potential ratio; Monsoon effect; Suddenly changes in atmosphere; Peak time of every seasons and its effect; Vertical migration of Zooplankton; Food selection pattern of predator; Collection time and number of collected samples, mixing of water column, high surface action, Seasonal upwelling and down welling process in water column.

5.2. Methodology

5.2.1 Estimation of Chlorophyll and Phaeophytin

Estimating Chlorophyll and Phaeophytin was done using known volume of water (500 ml) was filtered through a 0.45µm Millipore membrane filter paper and the pigments retained on the filter paper were extracted in 90% acetone overnight at 50°C. The extinction of the acetone extract was measured using fluorimeter before and after treatment with dilute acid (0.1N HCl).

5.2.2. Phytoplankton sampling and analysis

Phytoplankton samples were collected in the ten prefixed sampling sites using a standard plankton net with a mesh size of 51 µm. Plankton nets are with a square mouth covering an area of 0.900 cm² (30cm square mouth) fitted with a flow meter (Hydrobios). Nets were towed from a moving boat for 10 minutes and the plankton adhering to the net was concentrated in the net bucket. Plankton soup from the net bucket was transferred to a pre-cleaned and rinsed container and preserved with 5% neutralized formaldehyde. The containers were appropriately labelled. The initial and final flow meter reading was noted down for calculating the amount of water filtered to estimate plankton density. As per flow meter reading, a total amount of 165m³ of water was filtered by the net. One liter of water was separately collected for density estimation to counter check density estimation obtained by the flow meter reading. Quantitative analysis of phytoplankton (cell count) was carried out using a sedge wick-Rafter counting chamber. One ml of soup added to a Sedgwick

counting chamber was observed under an inverted compound microscope. The number of cells present in individual cells of the counting chambers (1/1000) was noted and identified up to a generic level. Several observations were fixed to represent the entire quantity of the soup (generally more than 30 times) and the recorded data were used to calculate the density (No/l) using the formula, $N = n \times v / V$ (where N is the total no/l; n is an average number of cells in 1 ml; v is the volume of concentrate; V is the total volume of water filtered). The phytoplankton diversity richness and evenness were past software.

5.3. Phytopigments

The concentration of phytopigments are directly proportional to the turbidity of the waters and in general, Kandla waters owing to the high turbidity restricts sunlight penetration essential for nutrient uptake by phytoplankton and thus inhibiting primary production. The concentration of chlorophyll pigment in the water samples ranged from 0.31-1.31 mg/m³ with a mean \pm SD being 0.60 \pm 0.28 mg/m³ in the Offshore (Table 9), 0.17 to 0.52 mg/m³ with mean \pm SD of 0.356 \pm 0.098 mg/m³ in the Cargo Jetty (Table 10) and 0.21 to 0.75 mg/m³ with mean \pm SD being 0.391 \pm 0.149 mg/m³ in the Phang creek location (Table 11).

The another phytopigment estimated was Phaeophytin, which is one of the breakdown products of Chlorophyll was also estimated in the water samples collected from all the three locations and the concentration of Phaeophytin in the marine water samples were in the concentrations such as 0.19 – 0.73 mg/m³ with a Mean \pm SD of 0.35 \pm 0.16 mg/m³ in the Offshore location. In case of Cargo Jetty location, the concentration of the secondary pigment was in the range of 0.11 – 0.41 mg/m³ with a Mean \pm SD of 0.256 \pm 0.082 mg/m³ and in case of the creek location, the concentration of phaeophytin was almost similar when compared to the other two locations and was ranging between 0.18 – 0.51 mg/m³ with a Mean \pm SD of 0.306 \pm 0.111 mg/m³ (Table 11). An optimum ration of Chlorophyll to Phaeophytin of above 1.5 as expected for natural estuarine and coastal waters.

Table 9: Chlorophyll and Phaeophytin concentration observed in the Offshore site

Parameters	1A		1B		1C		1D		1E		1 Control	
	SW	BW	SW	BW	SW	BW	SW	BW	SW	BW	SW	BW
Chlorophyll	1.31	0.67	0.81	0.61	0.66	0.36	0.66	0.4	0.31	0.35	0.35	0.67
Phaeophytin	0.41	0.28	0.73	0.56	0.21	0.31	0.19	0.21	0.28	0.27	0.29	0.47

Table 10: Chlorophyll and Phaeophytin concentration observed in the Cargo Jetty site

Parameters	2A		2B		2C		2D		2E		2 Control	
	SW	BW	SW	BW	SW	BW	SW	BW	SW	BW	SW	BW
Chlorophyll	0.45	0.17	0.33	0.35	0.35	0.27	0.52	0.35	0.32	0.51	0.35	0.3
Phaeophytin	0.34	0.11	0.27	0.27	0.27	0.18	0.41	0.22	0.19	0.32	0.3	0.19

Table 11: Chlorophyll and Phaeophytin concentration observed in the Phang Creek site

Parameters	3A		3B		3C		3D		3E		3 Control	
	SW	BW	SW	BW	SW	BW	SW	BW	SW	BW	SW	BW
Chlorophyll	0.36	0.22	0.37	0.42	0.58	0.36	0.3	0.21	0.31	0.39	0.75	0.42
Phaeophytin	0.21	0.2	0.21	0.34	0.5	0.32	0.27	0.18	0.28	0.27	0.51	0.38

5.4. Phytoplankton

The study was conducted at 3 sites (or regions) at Deendayal Port and near area where dredging activities is going on Creek and the stations are Offshore, Cargo Jetty and Phang Creek.

Offshore

In this site, frequently observed species were *Coscinodiscus oculus-iridis*, *Coscinodiscus radiatus*, *Coscinodiscus granii*, *Gyrosigma* sp, *Synedra ulna*, & *Thalassiosira fraunfeldii* colony, *Thalassiosira nitzschoides* colony, *Triceratium broeckii*. whereas less observed species were *Ceratium furca*, *Ceratium tripos*, *Entomoneis* sp, *Pinnularia* sp, *Protoperidinium* sp, *Pyrophacus* sp, *Triceratium favus*. Highest population density was recorded at site 1C-Offshore (896000 NoS/l) and low density recorded at site 1control-Offshore (33120 no/l). The maximum number of species observed in site 1a-Offshore (21 nos.) followed by 1B-Offshore (19 nos.), 1C-Offshore (11 nos), 1E-Offshore (10 nos) and 1D-1Control-Offshore (8 nos). The population density greatly varied (33120 nos/l to 896000nos/l). Among all recorded Phytoplankton Centric diatoms were 18, Pennate diatom- 9, Dinoflagellated -4 and Unidentified -1. Dinoflagellats like *Ceratium furca*, *Ceratium tripos*, *Protoperidinium* sp and *Pyrophacus* sp were recorded which are sometimes responsible for Algal Blooms in water.

Cargo jetty

The population density greatly varied between 34240 Nos/l to 62080 Nos/l. Highest density value recorded at 2B-Cargo Jetty (62080 No/l) and lowest value was at 2D-Cargo Jetty (34240). The highest number of species noticed in the site 2B- Cargojetty (17 nos.) where as density was also higher and lowest number of species noticed at 2C and 2E-Cargo Jetty (12 nos.). In this Cargo Jetty station commonly or frequently observed species were *Coscinodiscus granii*, *Coscinodiscus oculus-iridis*, *Coscinodiscus radiatus*, *Navicula* sp, *Pleurosigma* sp, *Thalassionema frauenfeldii* colony, *Thalassionema nitzschoides* colony, *Thalassiosira* sp. The rarely found species were *Biddulphia*, *Cyclotella* sp, *Odontella* sp,, *Surirella* sp, *Tripos azoricus*, *Coccolithoohores* etc. Among all Phytoplankton 18 Centric Diatoms, 2 Dinoflagellated cysts, 1 Coccolithophore, 1 Green algae, 9 Pennate Diatoms and 1 unidentified phytoplankton recorded.

Phang Creek

The population density of phytoplankton ranged from 26240 No/l to 71040 No/l same way species availability ranged from 12 to 25 nos. Maximum and minimum value of population density were recorded in site 3A-Phang Creek (71040 No/l) to 3E-Phang Creek (26240 No/l). Highest number of species recorded in site 3D-Phang Creek (25 nos) and lowest in site 3A-Phang Creek (12 nos).

Coscinodiscus centralis, *Coscinodiscus oculus-iridis*, *Coscinodiscus radiatus*, *Coscinodiscus granii* *Coscinodiscus wailesii*, *Euglena sp*, *Planktoniella blanda*, *Synedra sp*, *Synedra ulna* *Thalassiosira leptopus* were frequently noticed in samples whereas less observed species were *Planktoniella sol*, *Thalassiosira ecenntrica*, *Triceratium favus*, *Oscillatoria sp*, *Ditylum brightwellii* in this site.

Overall view of Phytoplankton showed that a total 54 species of Marine phytoplankton were identified during winter season of the year 2022. Among them, 25 were Centric diatoms, 14 were Pennate diatoms, 6 were Dinoflagellates, 1 was a Blue Green Algae, 1 was a Coccolithophores, 1 belong to Silicoflagellata, 2 were Green algae, 1 species was Unidentified. Plankton identification, both zooplankton and phytoplankton, was done by using relevant identification and taxonomic keys and with standard literatures, monographs and research articles. Some species like *Biddulphia sp*, *Thalassiosira leptopus*, *Climacosphaenica sp*, *Tripes azoricus*, *Pediastrum sp*, *Ditylum brightwellii*, *Protoperidinium sp*, *Scenedesmus sp*. were rarely recorded during sample analysis. Input of the fresh water indicated by the presence of some common fresh water species like *Euglena sp*, *Green algae*, *Oscillatoria sp*, *Pediastrum sp*, *Scenedesmus sp*. Highest phytoplankton density was observed at the site 1C-Offshore (89600 No/l) and lowest was observed at site 3E-Phang creek (26240 No/l) (Table 12). Total number of highest species observed at site 1A-Offshore (21 nos) and lowest in site 1D-Offshore and also 1-control-Offshore (8 nos). **During** laboratory analysis some Dinoflagellate species were also recorded like *Ceratium tripes*, *Protoperidinium sp*, *Pyrophacus sp*, *Tripes azoricus*. **Some** Blue green algae represented by *Oscillatoria sp* and *Scenedesmus sp*. The high population density composed by species like *Coscinodiscus granii*, *Coscinodiscus radiatus*, *Coscinodiscus granii*, *Planktoniella blanda*, *Thalassiosira sp*, *Thalassionema frauenfeldii colony*, *Thalassionema nitzschioides colony* and *Synedra ulna*. (Table 12). This result indicated that genus *Coscinodiscus sp*. was very common with good numbers in all sites. In some sites, least number of species and low density of phytoplankton might be responsible due to by the high Pre-Predation ratio,

Pollution, High turbidity, Total suspended solids, Water current of water and suddenly changes in favourable environment conditions. The individual density of species of sites viz. has been depicted in Table 12. All values of zooplankton density, list of zooplankton and others shown in Table 12.

5.4.4. Diversity Indices of Phytoplankton

The Table 13 shows diversity indices calculation for phytoplankton showed that the Shannon Index ranged from (1.784 to 3.004) indicated moderate to slightly higher level of diversity status. High Shannon Index was recorded at 3D-Phang Creek (3.004) and low at 1control-Offshore (1.784). Lowest evenness recorded at site 1A-Offshore (0.486) where highest phytoplankton numbers (21 Nos) were noticed, whereas highest was in at 3E-Phang Creek (0.925) where density was low recorded (26240 nos/l). Simpson dominance index 1-D-Offshore was showed the range from 0.782 to 0.940 whereas higher value in 3D-Phang Creek (0.940) and lowest was at in 1-Control-Offshore (0.782) (Table 13)

Table 12. Density of Phytoplankton at different sites of Deendayal Port

Name of Sites	Offshore						Cargo Jetty						Phang Creek					
	1A	1B	1C	1D	1E	1 control	2A	2B	2C	2D	2E	2 control	3A	3B	3C	3D	3E	3 control
Genus of Phytoplankton																		
<i>Actinocyclus sp</i>	0	960	8320	0	3040	3520	2720	0	0	1920	0	0	0	2240	0	4480	0	0
<i>Biddulphia sp.</i>	1760	0	0	0	0	0	0	0	0	640	0	0	0	0	0	0	0	0
<i>Campylodiscus sp.</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Ceratium furca</i>	640	1440	0	0	0	0	0	0	0	0	0	0	0	0	0	1280	0	0
<i>Ceratium sp.</i>	0	0	0	0	0	0	0	0	0	0	0	800	0	0	0	0	0	0
<i>Ceratium tripos</i>	0	960	0	0	0	0	0	0	0	0	0	0	0	0	0	2080	0	0
<i>Climacosphaenia sp</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	960	0	0	0	0
<i>Coccolithophores</i>	0	0	0	0	0	0	0	0	0	800	0	0	0	0	0	1760	0	0
<i>Coscindiscus centralis</i>	0	0	0	0	0	0	0	0	0	1440	0	2560	0	960	1600	1280	1440	2080
<i>Coscindiscus oculus-iridis</i>	3040	4320	10560	9920	4640	3360	4160	4160	2560	1440	960	0	11840	2400	3200	0	800	0
<i>Coscindiscus radiatus</i>	1600	4480	4160	0	2560	3040	2560	4480	1920	0	3200	3360	800	960	1920	5760	2080	4160
<i>Coscinodiscus granii</i>	16480	5440	29920	10080	11360	13120	12000	13280	13760	0	13920	9920	11680	9600	3360	2720	2080	3200
<i>Coscinodiscus sp</i>	0	2720	0	0	0	0	3200	0	0	0	0	4320	0	0	0	1760	0	0
<i>Coscinodiscus wailesii</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	8480	1600	3200	2560	1920
<i>Cyclotella sp.</i>	0	0	0	0	0	0	0	0	0	1440	0	2560	0	0	1760	4320	0	1600
<i>Dictyocha sp</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1120	0	0
<i>Ditylum brightwelli</i>	0	0	0	0	0	0	0	0	0	0	0	0	480	0	0	0	0	0
<i>Entomoneis sp</i>	640	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Euglena sp.</i>	0	0	0	0	0	0	0	0	0	0	0	0	2560	3040	0	1760	2400	0
<i>Green algae</i>	0	0	0	0	0	0	0	0	0	0	0	800	0	800	0	1120	0	0
<i>Guinardia sp</i>	0	480	0	0	0	0	0	0	0	0	0	640	0	0	640	0	0	0
<i>Gyrosigma sp.</i>	1120	0	0	2080	0	0	1440	2880	0	0	0	0	0	0	0	0	0	0
<i>Navicula lyra</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	0	800	960	0	0
<i>Navicula sp.</i>	640	1440	0	0	0	0	800	1440	1280	1440	0	0	0	0	2240	0	0	0
<i>Nitzschia sp.</i>	320	0	0	0	0	0	0	0	0	0	0	800	0	0	960	0	0	0
<i>Odontella sinensis</i>	0	0	0	0	0	0	0	0	0	0	0	1440	0	0	0	0	320	0
<i>Odontella mobiliensis</i>	0	0	0	0	0	0	1920	0	0	0	2560	0	0	0	0	0	1120	1600
<i>Odontella sp.</i>	0	0	0	0	0	0	0	0	0	0	960	0	0	0	0	0	0	0
<i>Oscillatoria sp.</i>	0	0	0	0	0	0	0	0	0	960	1280	0	0	1120	0	0	0	0
<i>Pediastrum sp.</i>	0	0	0	0	0	0	0	800	0	0	0	0	0	0	0	0	0	0
<i>Pinnularia sp.</i>	800	0	0	0	0	640	640	0	0	0	0	0	0	0	0	0	0	0
<i>Planktoniella blanda</i>	1440	0	8160	2560	4320	3840	0	3360	0	4480	0	0	6400	3840	3360	4320	1760	5920
<i>Planktoniella schutt</i>	1920	0	3360	0	0	0	0	0	0	0	0	2080	3520	0	0	0	0	0

<i>Planktoniella sol</i>	0	0	0	0	1920	0	0	1920	0	1760	0	0	0	800	0	0	0	0
<i>Pleurosigma sp.</i>	0	3040	0	0	320	0	1600	2080	1440	2240	0	0	3040	0	0	1920	1280	0
<i>Protopteridinium sp.</i>	0	960	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Pyrophacus sp.</i>	640	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Scenedesmus sp.</i>	0	0	0	0	0	0	0	3840	0	0	0	0	0	0	0	0	0	0
<i>Stellate trichome microplant parts</i>	0	0	0	0	0	0	1280	0	0	0	0	0	0	0	0	0	0	0
<i>Surirella sp.</i>	0	1920	0	0	1120	0	0	1280	0	1440	0	0	0	0	0	0	1920	0
<i>Synedra sp.</i>	1440	0	0	1120	0	0	0	0	960	0	2720	0	0	3040	800	1280	1600	1120
<i>Synedra ulna</i>	0	2880	1120	0	2880	1600	3840	0	1600	0	2720	2240	0	0	480	1440	0	960
<i>Thalassionema frauenfeldii colony</i>	0	8480	4960	4160	0	0	1120	6720	1760	3360	7520	0	5920	5920	3040	4320	2240	3200
<i>Thalassionema nitzschioides colony</i>	5120	12000	6560	5920	0	0	960	2560	4160	2720	3360	5120	0	5280	0	7520	2560	4160
<i>Thalassiosira ecentrica</i>	1760	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1760	0	0
<i>Thalassiosira ferelineata</i>	0	4960	0	0	0	0	0	0	0	0	0	0	8960	0	0	0	0	0
<i>Thalassiosira leptopus</i>	1440	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Thalassiosira sp</i>	160	14400	11040	6560	6720	4000	3840	8800	4160	6560	3840	1920	13120	8480	5120	2240	2080	5440
<i>Triceratium broeckii</i>	800	2080	1440	0	0	0	0	960	1600	1600	0	0	2720	1600	0	1600	0	1280
<i>Triceratium favus</i>	480	0	0	0	0	0	0	1600	480	0	0	0	0	320	0	480	0	0
<i>Triceratium sp.</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Tripes azoricus</i>	0	0	0	0	0	0	0	1920	0	0	0	0	0	0	0	0	0	0
<i>Unidentified sp.</i>	320	8000	0	0	0	0	0	0	0	0	3200	0	0	640	640	640	0	4480
Density of Phytoplankton (diff. sites wise.) (no/lit)	42560	80960	89600	42400	38880	33120	42080	62080	35680	34240	46240	38560	71040	60480	31520	61120	26240	41120
Total=877920 No/l																		
Total No Of Genus/Species= 53																		

Table 13. Diversity Indices of Phytoplankton at different selected sites of Deendayal Port

	Offshore						Cargo jetty						Phang Creek					
	1A	1B	1C	1D	1E	1-control	2A	2B	2C	2D	2E	2-control	3A	3B	3C	3D	3E	3-control
Taxa_S	21	19	11	8	10	8	15	17	12	16	12	14	12	19	16	25	15	14
Individuals (Nos/m²)	42560	80960	89600	42400	38880	33120	42080	62080	35680	34240	46240	38560	71040	60480	31520	61120	26240	41120
Dominance_D	0.182	0.095	0.170	0.171	0.161	0.218	0.129	0.103	0.194	0.093	0.150	0.125	0.127	0.096	0.089	0.060	0.075	0.093
Shannon Diversity	2.324	2.594	2.054	1.886	2.022	1.784	2.375	2.533	2.046	2.576	2.177	2.34	2.205	2.567	2.567	3.004	2.63	2.49
Simpson_1-D	0.818	0.905	0.830	0.829	0.839	0.782	0.871	0.897	0.806	0.908	0.850	0.876	0.873	0.904	0.911	0.940	0.925	0.907
Evenness	0.486	0.705	0.709	0.824	0.756	0.744	0.717	0.741	0.645	0.822	0.735	0.741	0.756	0.686	0.814	0.807	0.925	0.861
Menhinick	0.102	0.067	0.037	0.039	0.051	0.044	0.073	0.068	0.064	0.086	0.056	0.071	0.045	0.077	0.090	0.101	0.093	0.069
Margalef	1.88	1.59	0.88	0.66	0.85	0.67	1.32	1.45	1.05	1.44	1.02	1.23	0.98	1.64	1.45	2.18	1.38	1.22

5.5. Zooplankton

The study was conducted at 3 sites in Deendayal Port area and nearby areas where dredging activities are going on. The three selected study stations are Offshore, Cargo Jetty and Phang Creek.

Offshore

The *Ostracoda*, *Sponge spicules*, *Eggs capsules of Littorinids*, *Euterpina sp (Harpacticoida)*, *Globigerina sp (Foraminifera)*, *Nauplius larva of Copepoda*, *Nauplius larva of Barnacles*, *Tintinnopsis radix (Tintinnida)*, *Copepoda egg sacs(pouch)* were the mostly common zooplankton throughout observed in all sites of Offshore points. Highest population density was recorded at site 1D-Offshore (128800 No/100m³) and lowest in 1-control-Offshore (44000 No/100m³). **Site 1B-Offshore** has maximum number of species (28 nos) whereas minimum was found in the site 1-control-Offshore (15 nos). High biomass was observed in the site 1Control-Offshore (55.97 ml/100m³) and low biomass was in site 1E-Offshore (10.17 ml/100m³). The range of the population density, biomass and number of species were (44000 to 128800 no/100m³), (10.17 to 55.97 ml/100m³) and (12 to 33 nos) respectively in all sites. Less observed species are *Ammonia sp* (Foraminifera), *Arcella sp* (Protozoa) *Sagitta sp*, *Dentilium*, *Calcarina sp* (Foraminifera), *Spirulina sp* (Foraminifera), *Centropages sp* (Calanoida) etc. rarely recorded in this station. Total 52 zooplankton was recorded in Offshore station adding that more composition of zooplankton by phylum Crustacea and Foraminifera as shown in Table 14.

Cargo Jetty

The population density of zooplankton varied from 47320 No/100m³ to 96140 No/100m³. Maximum density was noticed in site 2C-Cargo Jetty (96140 no/100m³) and minimum was at site 2Control-Cargo Jetty (47320 no/100m³) as given in Table 15. The site 2C-Cargo Jetty comprises highest number of species (33 nos) and minimum number of species was observed in site 2B-Cargo Jetty (15 nos). Biomass ranged between 15 to 57.14 ml/100m³ where highest biomass noted in site 2B-Cargo Jetty and lowest in 2A-Cargo Jetty. Frequently observed species were *Centropages sp* (Calanoida), *Clausocalanus sp* (Calanoida) *Zoea larva of Crustacean*, *Oithona sp* (Cyclopoida), *Subeucalanus sp* (Calanoida), *Tintinnopsis beroidea* (Tintinnida), *Tintinnopsis radix* (Tintinnida), and *Egg capsules of Littorinids* whereas less observed species were *Nodosaria sp* (Foraminifera), *Copepoda egg sacs(pouch)*, *Euchaeta sp*

(Calanoida), *Diacyclops sp* (Cyclopoida), *Leprotintinnus nordqvistii* (Tintinnida), *Leprotintinnus simplex* (Tintinnida). Total recorded zooplankton was 60 nos. in Cargo Jetty.

Phang Creek

This Creek area was represented by the zooplankton fauna majority of Egg capsule of Littorinids, Nauplius larva of Copepoda, Sponge spicules, *Clausocalanus sp* (Calanoida), *Oithona sp* (Cyclopoida), *Tintinnopsis beroidea* (Tintinnida). **Very** less time or rarely recorded species were *Amphipoda*, *Cibicides sp* (Foraminifera), Coccolithophores, *Eponidis sp* (Foraminifera) The range of zooplankton Biomass was between 0.50 to 30.49 ml/100m³. Highest Biomass was recorded in site 3D-Phang creek (30.49 ml/100m³) and lowest in site 3B-Phang creek (0.50 ml/100m³). Maximun and Minumum species count was at in site 3A,3C and 3D-Phang creek (25 nos) and 3E-Phang creek (12 nos) respectively. Population density was maximum recorded in site 3C-Phang creek (101600 No/100m³) and minimum in site 3E-Phang creek (36360 No/100m³).

Overall assessment of zooplankton showed that the total number of 38 Zooplankton recorded during monsoon season. Out of these (86 nos) zooplankton, 52 zooplankton recorded in Offshore region, 60 zooplankton at Cargo Jetty and 55 zooplankton in Phang Crek region. The recorded zooplankton of all 3 stations mainly representing Phylum Arthropoda (Crustacea) as presented in Table 16. Protozoa (mainly foraminifera and tintinnids), Porifera (Sponge spicules) Generally zooplankton population dynamics and studies emphasize is given up to group level rather than to species level because of microscopic size of zooplankton so owing to the difficulty in identifying the zooplankton as some species are considered as a group or genus level. The most dominant or frequently observed species were *Clausocalanus sp* (Calanoida), *Egg capsules of Littorinids*, *Ostracoda*, *Tintinnopsis radix*, *Oithona sp* (Cyclopoida), *Zoea larva of Crustacea*, *Sponge spicules*, *Globigerina sp* (Foraminifera) and other Foraminifera. The range of Population density, Biomass and Number of Species were (36360 to 128800 no/100 m³), (0.50 to 57.14 ml/100m³) and (12 to 33 nos) respectively. **Average** high biomass noted at Cargo Jetty followed by Offshore and Phang creek (Table 14,15,16). Highest population density was recorded in site 1D-Offshore (128800 no/100m³) and lowest was recorded in site 3E-Phang Creek (36360 No/100m³). Among all recorded zooplankton, majority dominance occurrence was by the Copepoda, Crustacean larvae, Spong Spicules, Foraminifera (Protozoa), Tintinnids (Protozoa), Egg

capsules of Littorinids (Mollusca). Maximum zooplankton faunal composition was dominated by the Phylum Arthropoda, Mollusca, Protozoa and Porifera. The Chaetognatha and tunicata groups were only represented by the one species namely Sponge spicules, *Sagitta sp* and *Oikopleura sp* respectively. In Offshore, maximum Occurrence (%) was by the Egg Capsules of Littorinids (18.33%) and minimum by the Radiolarian (0.10%). In Cargo Jetty, maximum Percentage of Occurrence (%) by the Eggs of Littorinids (14.31%) and minimum by the Nodosaria sp (0.07%) (Foraminifera).. In Phang Creek maximum Occurrence (%) was by the Egg capsules of Littorinids (12.42%) and minimum (0.08%) by the *Cibicides sp* (Foraminifera).

During analysis, some Species of Foraminifera and Spicules of sponge were frequently observed. These both are very important for paleontological study aspects and also for evolutionary, ecological and environmental rebuilding. Some species of Ostracoda, Foraminifera and Sponge spicules are considered in microfossils materials. Some deep sea species also recorded that is indication of water circulation pattern. Data on zooplankton density, list of zooplankton is shown in Table (14, 15 & 16).

Diversity Indices of Zooplankton

Table 17 shows *diversity zooplankton*. The Shannon-wiener diversity index (H') fluctuated between 2.42 to 3.22 indicated moderate to quite high range of diversity with a maximum value in site 2C-Cargo Jetty (3.22) and minimum value in site 3E-Phang creek (2.42). Range of the evenness was 0.514 to 0.938 where highest and lowest recorded in site 3E-Phang Creek (0.938) where lowest density was recorded and 1D-Offshore (0.514) respectively. Highest Simpson index 0.95 noted at site 2C-Cargo Jetty whereas lowest in site 1A (0.88).

Table 14. Density of Zooplankton at Offshore site of Deendayal Port

Name of Genera/Group	1A	1B	1C	1D	1E	1 Control	Individual total density (no/100m ³)	% of Occurrence
Acartia sp (Calanoida)	0	0	5120	0	0	0	5120	1.06
Ammonia sp. (Foraminifera)	0	0	0	0	3360	0	3360	0.70
Arcella sp (Amoebozoa)	0	0	1280	0	0	0	1280	0.27
Bolivina sp.(Foraminifera)	0	0	7040	3360	1440	0	11840	2.46
Calcarina sp. (Foraminifera)	0	0	0	0	800	0	800	0.17
Centropages sp. (Calanoida)	0	1440	0	0	0	0	1440	0.30
Clausocalanus sp (Calanoida)	2560	5120	0	3520	0	0	11200	2.32
Copepoda egg sacs (egg pouch)	0	4320	3840	0	0	6880	15040	3.12
Cyclops sp (Cyclopoida)	0	0	0	6560	0	0	6560	1.36
Cyphonautes larva of bryozoans	640	0	0	0	1920	0	2560	0.53
Dentalium	0	0	0	0	0	640	640	0.13
Diacyclops sp. (Cyclopoida)	5440	1920	0	0	0	0	7360	1.53
Egg Capsules of Littorinids	12480	13120	13920	43040	0	5760	88320	18.33
Eucalanus sp. (Calanoida)	0	0	0	3200	0	0	3200	0.66
Euchaeta sp (Calanoida)	0	0	0	1600	0	0	1600	0.33
Euterpina sp (Harpacticoida)	960	3520	0	0	480	2560	7520	1.56
Eutintinnus apertus (Tintinnida)	0	0	0	2240	1920	0	4160	0.86

Globigerina sp. (Foraminifera)	2720	3520	12320	4640	4160	0	27360	5.68
Labidocera sp. (Calanoida)	0	1120	0	0	0	0	1120	0.23
Larva of Hydrozoa (Phylum: Cnidaria)	0	2880	0	0	3040	1440	7360	1.53
Leprotintinnus nordqvistii (Tintinnida)	0	2080	0	0	2720	0	4800	1.00
Leprotintinnus simplex (Tintinnida)	0	0	0	0	1760	0	1760	0.37
Nauplius larva of Copepoda	1920	3360	1600	0	7360	2720	16960	3.52
Nauplius larva of Harpacticoida	0	0	0	0	0	1600	1600	0.33
Nauplius larvae of Barnacles	2720	2720	0	0	1920	1120	8480	1.76
Nauplius larvae of Crustacea	0	0	0	0	0	4960	4960	1.03
Nonion sp. (Foraminifera)	0	0	0	2240	960	0	3200	0.66
Oithona sp. (Cyclopoida)	0	5120	9120	3520	0	0	17760	3.69
Ophiopluteus larva of (Echinodermata)	1440	0	0	0	1440	0	2880	0.60
Ostracoda	1120	320	2720	0	4640	4640	13440	2.79
Other Calanoida	0	0	0	16960	0	1280	18240	3.78
Other Cyclopoida	0	1440	0	7040	0	0	8480	1.76
Parvocalanus sp (Calanoida)	1760	1920	0	0	0	0	3680	0.76
Quinqueloculina sp.(Foraminifera)	0	10240	0	2720	0	0	12960	2.69
Radiolaria skeleton	320	800	0	320	160	0	1600	0.33
Radiolaria sp (Protozoa)	0	320	0	160	0	0	480	0.10

Rosalina sp. (Foraminifera)	800	1440	3200	4640	4960	0	15040	3.12
Sagitta sp (arrow worm)	1120	0	0	0	0	0	1120	0.23
Small Gastropoda	0	0	0	1280	0	0	1280	0.27
Spirillina sp. (Foraminifera)	0	0	0	0	3360	0	3360	0.70
Spiroloculina sp (Foraminifera)	1440	1920	1760	0	0	0	5120	1.06
Sponge spicules	10880	8480	10240	9920	5760	5920	51200	10.62
Temora sp (Calanoida)	0	0	3040	0	5280	2720	11040	2.29
Thermocyclops sp. (Cyclopoida)	0	0	0	0	3360	0	3360	0.70
Tintinnopsis beroidea (Tintinnida)	3680	0	0	960	0	0	4640	0.96
Tintinnopsis cylindrica (Tintinnida)	1280	4480	4000	2720	0	0	12480	2.59
Tintinnopsis lobiancoi (Tintinnida)	0	2240	0	3680	0	0	5920	1.23
Tintinnopsis orientalis (Tintinnida)	0	2720	5760	0	1280	0	9760	2.03
Tintinnopsis radix (Tintinnida)	1920	1120	5600	2720	1920	0	13280	2.76
Veliger larvae of Bivalve	0	640	3520	1760	0	1760	7680	1.59
Zoea larva of Crustaceans	0	4000	0	0	2880	0	6880	1.43
Unidentified sp.	0	0	640	0	0	0	640	0.13
Total No. Of Genera/Groups =52								
Site-wise Total Density (no/100m³)	55200	92320	94720	128800	66880	44000	Total Density =481920	100%
Biomass (ml/100m³)	11.24	13.38	15.67	16.91	10.17	55.97		

Table 15. Density of Zooplankton at Cargo Jetty site of Deendayal Port

Name of Genera/Group	2A	2B	2C	2D	2E	2 Control	Individual total density (no/100m³)	% of Occurrence
Acartia sp (Calanoida)	0	1600	960	800	2080	0	5440	1.27
Acrocalanus sp. (Calanoida)	1920	0	0	1280	0	0	3200	0.75
Bolivina sp.(Foraminifera)	0	0	1920	2240	0	0	4160	0.97
Calcarina sp. (Foraminifera)	0	0	960	960	0	0	1920	0.45
Centropages sp. (Calanoida)	640	2240	1760	960	1760	0	7360	1.72
Clausocalanus sp (Calanoida)	1920	1760	2560	1920	2880	0	11040	2.58
Copepoda egg sacs (egg pouch)	1280	0	0	0	0	0	1280	0.30
Corycaeus sp (Calanoida)	0	0	0	1440	0	0	1440	0.34
Cyphonautes larva of bryozoans	2720	0	0	0	1440	1600	5760	1.35
Diacyclops sp. (Cyclopoida)	0	0	0	1760	0	0	1760	0.41
Egg Capsules of Littorinids	11680	8640	9920	14880	5600	10400	61120	14.31
Euchaeta sp (Calanoida)	0	0	0	0	1440	0	1440	0.34
Euterpina sp (Harpacticoida)	7040	3520	0	2080	2880	0	15520	3.63
Eutintinnus sp. (Tintinnida)	0	0	0	0	1920	0	1920	0.45
Fish larva	0	0	0	0	1120	0	1120	0.26
Globigerina sp. (Foraminifera)	0	0	6400	12480	3360	2400	24640	5.77

Heterolaophonte (Harpacticoida)	0	0	0	0	1760	0	1760	0.41
Larva of Crustacea	0	0	0	0	640	0	640	0.15
Larva of Hydrozoa (Phylum: Cnidaria)	1920	0	300	1440	0	0	3660	0.86
Leprotintinnus nordqvistii (Tintinnida)	0	0	0	1760	0	0	1760	0.41
Leprotintinnus pellucidus (Tintinnida)	0	0	0	1920	1120	0	3040	0.71
Leprotintinnus simplex (Tintinnida)	0	0	0	0	3360	0	3360	0.79
Microsetella sp (Harpacticoida)	0	2520	0	0	0	1760	4280	1.0
Nauplius larva of Calanoida	0	0	3040	0	0	0	3040	0.71
Nauplius larva of Copepoda	0	0	4320	2720	0	0	7040	1.65
Nauplius larvae of Barnacles	4160	0	1760	2240	1760	0	9920	2.32
Nauplius larvae of Cyclopoida	0	0	4000	0	0	0	4000	0.94
Nodosaria sp (Foraminifera)	0	0	320	0	0	0	320	0.07
Oithona brevicornis	0	0	1440	0	0	0	1440	0.34
Oithona sp. (Cyclopoida)	3360	3520	4320	2400	2240	2400	18240	4.27
Ophiopluteus larva of (Echinodermata)	0	0	1440	960	0	0	2400	0.56
Ostracoda	3840	3840	1440	0	0	1720	10840	2.54
Other Calanoida	3040	2720	0	0	0	0	5760	1.35
Other Cyclopoida	1760	0	1280	0	0	0	3040	0.71
Paracalanus sp. (Calanoida)	2240	0	3200	0	0	0	5440	1.27
Parvocalanus sp (Calanoida)	1920	1280	0	1760	0	1920	6880	1.61
Polychaeta larvae (Annelida)	0	2560	0	0	0	0	2560	0.60

Pseudodiaptomus sp (Calanoida)	0	0	0	0	2080	0	2080	0.49
Quinqueloculina sp.(Foraminifera)	0	0	1760	3680	3840	0	9280	2.17
Radiolaria skeleton	0	0	0	0	320	480	800	0.19
Radiolaria sp (Protozoa)	0	0	320	0	160	0	480	0.11
Rosalina sp. (Foraminifera)	0	0	1920	3520	0	800	6240	1.46
Sagitta sp (arrow worm)	0	1600	0	0	0	0	1600	0.37
Small Gastropoda	0	0	1600	0	0	0	1600	0.37
Spirillina sp. (Foraminifera)	0	0	0	320	0	640	960	0.22
Spiroloculina sp (Foraminifera)	0	0	1920	640	2720	1920	7200	1.69
Sponge spicules	16320	8320	7040	0	0	8320	40000	9.36
Subeucalanus (Calanoida)	3360	0	0	1920	1280	1920	8480	1.98
Temora sp (Calanoida)	4640	0	0	0	0	1760	6400	1.75
Tintinnopsis beroidea (Tintinnida)	1920	2560	5280	6240	2720	1600	20320	4.76
Tintinnopsis cylindrica (Tintinnida)	0	0	0	0	0	1440	1440	0.34
Tintinnopsis lobiancoi (Tintinnida)	0	0	0	0	1440	0	1440	0.34
Tintinnopsis mortenseni (Tintinnida)	0	0	800	0	0	0	800	0.19
Tintinnopsis radix (Tintinnida)	2560	0	6400	1760	1920	2240	14880	3.48
Tintinnopsis sp (Tintinnida)	0	0	4960	0	0	1600	6560	1.54
Tintinnopsis tubulosa (Tintinnida)	0	0	2400	3200	0	1920	7520	1.56

Triloculina sp (Foraminifera)	0	0	1280	0	0	0	1280	0.30
Veliger larvae of Bivalve	0	0	2560	1760	3040	0	7360	1.72
Zoea larva of Crustaceans	2080	8480	6560	5120	8960	0	31200	7.30
Unidentified sp.	0	0	0	320	0	480	800	0.19
Total No. Of Genera/Groups =60								
Site-wise Total Density (no/100m³)	80320	55160	96140	84480	63840	47320	Total Density =427260	100%
Biomass (ml/100m³)	15	57.14	20	27.27	20.55	41.03		

Table 16. Density of Zooplankton at Phang Creek site of Deendayal Port

Name of Genera/Group	3A	3B	3C	3D	3E	3 Control	Total density (no/100m3)	% of Occurrence
Acartia sp (Calanoida)	0	0	1440	1440	0	0	2880	0.74
Acrocalanus sp. (Calanoida)	1280	0	2880	0	0	2080	6240	1.60
Amphipoda	0	0	0	800	0	0	800	0.21
Arcella sp (Amoebozoa)	1120	0	0	0	0	0	1120	0.29
Centropages sp. (Calanoida)	2560	0	0	0	0	0	2560	0.66
Cibicides sp (Foraminifera)	0	320	0	0	0	0	320	0.08
Clausocalanus sp (Calanoida)	7360	7200	9920	0	0	4960	29440	7.57
Clytemnestra sp (Harpacticoida)	0	0	0	1120	0	1280	2400	0.62
Cyclops sp (Cyclopoida)	0	960	2240	0	0	0	3200	0.82
Cyphonautes larva of bryozoans	800	0	0	1600	0	0	2400	0.62
Diacyclops sp. (Cyclopoida)	0	0	1440	0	0	1280	2720	0.70
Egg Capsules of Littorinids	9920	7520	12160	3040	4960	10720	48320	12.42
Eponides sp (Foramonifera)	0	800	0	0	0	0	800	0.21
Eucalanus sp. (Calanoida)	1280	2560	0	2080	0	0	5920	1.52
Euterpina sp (Harpacticoida)	0	0	2560	0	3520	3360	9440	2.43
Eutintinnus apertus (Tintinnida)	2400	0	0	800	0	0	3200	0.82
Eutintinnus lususundae (Tintinnida)	0	0	0	0	0	2080	2080	0.53
Eutintinnus sp. (Tintinnida)	1600	0	0	0	0	0	1600	0.41
Gastrula embryo of Seastar	0	0	0	800	0	0	800	0.21

Globigerina sp. (Foraminifera)	0	2560	0	7040	2080	0	11680	3.00
Labidocera sp. (Calanoida)	0	0	800	960	0	800	2560	0.66
Larva of Hydrozoa (Phylum: Cnidaria)	0	800	0	0	0	0	800	0.21
Leprotintinnus nordqvistii (Tintinnida)	1280	1440	0	0	0	0	2720	0.70
Microsetella sp (Harpacticoida)	2080	5440	0	2880	0	0	10400	2.67
Mysis larva	0	1120	0	960	0	0	2080	0.53
Nauplius larva of Copepoda	4800	5120	2080	10080	3360	5760	31200	8.02
Nauplius larvae of Barnacles	2400	3360	5120	3520	0	1760	16160	4.15
Nauplius larvae of Cyclopoida	0	0	0	1760	0	0	1760	0.45
Nonion sp. (Foraminifera)	0	0	0	640	0	0	640	0.16
Oithona sp. (Cyclopoida)	1920	3040	9440	1600	0	4000	20000	5.74
Ophiopluteus larva of (Echinodermata)	0	0	2080	2720	0	1920	6720	1.73
Ostracoda	1440	1280	2880	0	3840	0	9440	2.43
Other Calanoida	0	0	0	1280	0	3200	4480	1.15
Other Cyclopoida	1280	0	2240	0	0	4160	7680	1.97
Paracalanus sp. (Calanoida)	0	0	3680	0	0	0	3680	0.95
Parvocalanus sp (Calanoida)	0	2400	0	0	0	1440	3840	0.99
Planispirinella sp (Foraminifera)	0	0	480	0	0	0	480	0.12
Polychaeta larvae (Annelida)	0	0	2240	0	0	0	2240	0.58

Pontellopsis sp. (Calanoida)	480	0	0	0	0	0	480	0.12
Quinqueloculina sp.(Foraminifera)	5920	1280	0	0	0	2080	9280	2.39
Sagitta sp (arrow worm)	0	0	0	0	1600	0	1600	0.41
Sponge spicules	10880	5280	0	3040	3200	2080	24480	6.29
Subeucalanus (Calanoida)	0	0	0	0	0	2720	2720	0.70
Temora sp (Calanoida)	2720	1280	7520	1120	3520	0	16160	4.15
Textularia sp. (Foraminifera)	0	0	0	1760	0	0	1760	0.45
Tintinnopsis beroidea (Tintinnida)	1440	1440	0	1760	3200	0	7840	2.02
Tintinnopsis cylindrica (Tintinnida)	0	0	6880	0	0	1440	8320	2.14
Tintinnopsis karajacensis (Tintinnida)	0	0	0	0	0	800	800	0.21
Tintinnopsis lobiancoi (Tintinnida)	0	0	2720	0	0	0	2720	0.70
Tintinnopsis orientalis (Tintinnida)	3840	0	8480	0	0	0	12320	3.17
Tintinnopsis radix (Tintinnida)	0	0	0	0	2560	0	2560	0.66
Tintinnopsis tubulosa (Tintinnida)	1920	2400	2080	3360	960	0	10720	2.76
Veliger larvae of Bivalve	0	0	7520	0	0	0	7520	1.93
Zoea larva of Crustaceans	1760	2560	2560	2400	3560	0	12840	3.30
Unidentified sp.	320	0	160	0	0	0	480	0.12
Total No of Genera/ Groups =55								
Site-wise Total Density (no/100m³)	72800	60160	101600	58560	36360	57920	Total density =387400	100%
Biomass (ml/100m³)	2.50	0.50	13.57	30.49	19.09	11.59		

Table 17. Diversity indices of Zooplankton at different sites of Deendayal Port

Variables	Offshore						Cargo jetty						Phang Creek					
	1A	1B	1C	1D	1E	1-control	2A	2B	2C	2D	2E	2-contrl	3A	3B	3C	3D	3E	3-control
Taxa_S	19	28	18	23	24	15	21	15	33	30	27	20	25	22	25	25	12	20
Individuals (Nos/m²)	55200	92320	94720	128800	66880	44640	80320	55160	96140	84480	63840	47320	72800	60160	101600	58560	36360	59520
Dominance_D	0.12	0.06	0.08	0.15	0.06	0.10	0.09	0.10	0.05	0.08	0.06	0.10	0.08	0.07	0.07	0.07	0.09	0.08
Shannon Diversity	2.50	3.01	2.65	2.47	2.97	2.48	2.72	2.51	3.22	2.98	3.08	2.64	2.87	2.83	2.92	2.94	2.42	2.81
Simpson_1-D	0.88	0.94	0.92	0.85	0.94	0.90	0.91	0.90	0.95	0.92	0.94	0.90	0.92	0.93	0.93	0.93	0.91	0.92
Evenness	0.639	0.722	0.786	0.514	0.813	0.795	0.726	0.818	0.755	0.655	0.802	0.701	0.705	0.773	0.740	0.754	0.938	0.792
Menhinick	0.081	0.092	0.058	0.064	0.093	0.071	0.074	0.064	0.106	0.103	0.107	0.092	0.093	0.090	0.078	0.103	0.063	0.086
Margalef	1.649	2.362	1.484	1.87	2.07	1.308	1.771	1.282	2.789	2.556	2.35	1.765	2.144	1.908	2.082	2.186	1.047	1.819

6.0. References

- Airoidi, L., & Beck, M. (2007). Loss, status and trends for coastal marine habitats of Europe. *Oceanography and Marine Biology: An Annual Review*, 45, 345–405.
- Barbier Edward, B., Hacker Sally, D., Kennedy, C., Koch, E. W., Stier, A. C., and Silliman, B. R. (2011). The value of estuarine and coastal ecosystem services. *Ecol. Monogr.* 81, 169–193. doi: 10.1890/10-1510.1
- Barnes, R D, 1980. Invertebrate Zoology Saunders College, Philadelphia 108pp
- Besiktepe, S., Tang, K. W., Mantha, G. (2015). Seasonal variation of abundance and live/dead compositions of copepods in Mersin Bay, northeastern Levantine Sea (eastern Mediterranean). *Turk Zool.* 39: 494-506. doi:10.3906/zoo-1405-23.
- Bhaskar P. V., Roy R., Gauns M., Shenoy D. M., Rao V. D., Mochamadkar S., 2011. “Identification of non-indigenous phytoplankton species dominated bloom off Goa using inverted microscopy and pigment (HPLC) analysis”, *J Earth Syst Sci*, pp.1145–1154.
- Bhunja, A.B. and Choudhury, A., 1998. Studies on the seasonal abundance and biomass of Crustacean Zooplankton and Chaetognaths in relation to ecological parameter of a tidal Creek (Mooriganga), of Sagar Island (north), Sunderbans, West Bengal. *Indian Journal of Marine Science* 28, 93-198.
- Boyd, S. E., Limpenny, D. S., Rees, H. L., & Cooper, K. M. (2005). The effects of marine sand and gravel extraction on the macrobenthos at a commercial dredging site (results 6 years post-dredging). *ICES Journal of Marine Science*, 62(2), 145–162.
- Brink, K. H. (1993). The coastal ocean progresses effort. *Oceanus*, 36, pp. 47-49.
- Chakrabarty, M., Banerjee, A., Mukherjee, J., Rakshit, N., Ray, S. (2017) Spatial pattern analysis of zooplankton community of Bakreswar reservoir, India. *Energ. Ecol. Environ.* 2(3): 193-206. doi: 10.1007/s40974-017-0057-8.
- Chattopadhyay, J., R.R. Sarkar & S. Pal 2003. Dynamics of nutrient–phytoplankton interaction in the presence of viral infection. *BioSystems*, 68: 5–7.
- Clark, K R and Warwick, R M 1994. Change in Marine Communities, An Approach to Statistical Analysis and Interpretation Natural Environment Research Council, Plymouth Marine Laboratory, Plymouth, pp144.
- Cloern J. E., “Phytoplankton bloom dynamics in coastal ecosystems: A review with some general lessons from sustained investigation of San Francisco Bay”, California. 1996. *Rev Geophys*, 34(2), pp.127–168, 1996.
- Davies, O.A., C. C. Tawari and J. F. N. Abowei, 2008. Zooplankton of Elechi Creek, Niger Delta, Nigeria. *Environ. Ecol.*, 26 (4c) : 2346 - 2441.
- Davis, B. J. 1977. Distribution and temperature adaptation in the teleost fish genus *Gibbonsia*. *Mar. Biol.* 42: 315-320.
- Day, J H 1967. A monograph on the Polychaeta of Southern Africa Pts I and II, *Brit Mus Nat Hist*, 656, 1-878.
- Dekker, R. 1989. The macrozoobenthos of the subtidal western Dutch Wadden Sea. I. 468 Biomass and species richness. *Netherlands Journal of Sea Research* 23: 57–68.
- Desai, S. R, 2008. Subashchandran, MD, Ramachandra TV; Phytoplankton Diversity in Sharavati River Basin, Central Western Ghats. *Journal of Sediment and Water Sciences*, 2008; 1(1):7- 66.
- Descy, J. P, 1993. Ecology of the phytoplankton of river Moselle: Effect of disturbance on community structure and diversity. *Hydrobiologia*, 249(1-3): 111-116.
- Dodson, S. (1992). Predicting crustacean zooplankton species richness. *Limnol Oceanogr.* 37(4): 848-856.

- Dodson, S.I. and Frey, D. G. (2001). Cladocera and other branchiopoda. Ecology and classification of North American Freshwater Invertebrates. Academic Press. London. 850 – 914.
- Fauvel, P, 1953 The Fauna of India Including Pakistan, Ceylon, Burma and Malaya Annelida, Polychaeta, Allahabad pp507.
- Figueredo, C. C. & A. Giani 2001. Seasonal variation in the diversity and species richness of phytoplankton in a tropical eutrophic reservoir. *Hydrobiologia*, 445: 165-174.
- Gajbhiye, S.N. & Abidi, S.A.H. (1993). Zooplankton distribution in the polluted environment around Bombay. *Environment Impact on Aquatic & Terrestrial Habitats*. pp. 127-142.
- Gao, X. & J. Song 2005. Phytoplankton distributions and their relationship with the environment in the Changjiang Estuary, China. *Marine Poll. Bull.*, 50: 327-335.
- Garzke, J., Sommer, U., Ismar, S.M.H. (2017). Is the chemical composition of biomass the agent by which ocean acidification influences on zooplankton ecology. *Aquat Sci.* 79(3): 733-748. doi: 10.1007/s00027-017-0532-5.
- Goswami, S.C & Padmavathi, G. (1996). Zooplankton production, composition and diversity in the coastal water of Goa. *Indian Journal of. Marine Science* 25, 91-97.
- Gray, J. S. (1997). Marine biodiversity: Patterns, threats and conservation needs. *Biodiversity and Conservation*, 6, 153–175.
- GUIDE, (2011). Comprehensive Terrestrial EIA (including Mangroves) for the Proposed Multi-Project SEZ at Kandla. EIA report submitted to Mumbai Regional Centre of National Institute of Oceanography, Dona Paula, Goa.
- Hambler, C and Speight, M R 1995. Biodiversity conservation in Britain, science replacing tradition *British Wildlife*, 6, 137-147 yla, P S, S Velvizhi and S Ajmal Khan 1999 A Monograph on the amphipods of Parangipettai coast Annamalai University, India 78.
- Harkantra, S. N., A. Nair, Z. A. Ansari and A. H. Parulekar, 1980. Benthos of the shelf region along the West coast of India. *Indian J. Mar. Sci.*, 9: 106-110.
- Ingle B, Sivadas S, Goltekar R, Clemente S, Nanajkar M, Sawant R, D'Silva C, Sarkar A, Ansari Z (2006) Ecotoxicological effect of grounded MV River Princess on the intertidal benthic organisms of Goa. *Environ /ntemat.* 32:284-289.
- Jagadeesan, L., Jyothibabu, R., Anjusha, A., Arya, P. M., Madhu, N. V., Muraleedharan, K. R., and Sudheesh, K., 2013. Ocean currents structuring the mesozooplankton in the Gulf of Manner and the Palk Bay, southeast coast of India. *Progress in Oceanography*, 110: 27-48.
- Jegadeesan, P., 1986. Studies on environmental inventory of the marine zone of Coleroon estuary and inshore waters of Pazhayar, Southeast coast of India. *Ph. D., Thesis, Annamalai University, India.*
- Jha, D. K., Devi, M. P., Vidyalakshmi, R., Brindha, B., Vinithkumar, N. V., and Kirubakaran, R. (2015). Water quality assessment using water quality index and geographical information system methods in the coastal waters of Andaman Sea, India. *Mar. Pollut. Bull.* 100, 555–561. doi: 10.1016/j.marpolbul.2015.08.032.
- Jickells, T. D. (1998). Nutrient biogeochemistry of the coastal zone. *Science* 281, 217–222. doi: 10.1126/science.281.5374.217.
- Jones, G., and Candy, S. (1981). Effects of dredging on the macrobenthic infauna of Botany Bay. *Marine and Freshwater Research*, 32(3), 379–398.
- Kadam S.S. and L. R. Tiwari, 2012. Zooplankton Composition in Dahanu Creek-West Coast of Ind. Res. J. Rec. Sci., 1(5): 62-65.
- Karr, J .R., J D. Allen, and A. C. Benke 2000 River conservation in the United States and Canada. In P. J. Boon, Davies and B .R. Petts, G E (Ed.), *Global perspectives on River conservation*, pp 3–39 Science, Policy, and Practice. Wiley, New York.

- Krishnamurthy, K. & Santhanam, R. (1975). Ecology of Tintinids (Protozoa: Ciliata) in Porto Novo region. *Indian Journal of Marine Science* 4, 181-184.
- Kumar, A., 1995 Studies of pollution in river Mayurakshi in south Bihar. *Indian Journal of Environmental Pollution*, 2(1): 21-26.
- Levandowsky, M., 1972. An ordination of phytoplankton population in ponds of varying salinity and temperature. *Ecology*, 53(3): 398-407.
- Lyla, P S., Velvizhi, S and Ajmal Khan, S 1999. A Monograph on the amphipods of Parangipettai coast Annamalai University, India pp78.
- Magurran, A 1991. Ecological Diversity and Its Measurement Princeton University Press, Princeton, pp178.
- Mahapatro, D R C., Panigrahy, K and Samal, R N 2011. Macrobenthos of shelf zone off Dhamara estuary, Bay of Bengal. *J Oceanog Mar Sci* 22, pp 32-42.
- Margalef, R 1958. Information theory in ecology. *Gen Syst*, 3, 36-71.
- Marine Biology Organization (MBO), 2007. Zooplankton Retrieved from: <http://www.marinebio.com/oceans/zooplankton.Askp>. 62k, (Accessed on: September 29, 2006).
- Maurer, D., Watling, L., Kinner, P., Leathem, W and Wethe, C 1978. Benthic invertebrate assemblages of Delaware Bay. *Mar Biol*, 45, 65-78.
- Maya, M. V., M. A. Soares, R. Agnihotri, A. K. Pratihary, S. Karapurkar, H. Naik & S. W. A. Naqvi 2011. Variations in some environmental characteristics including C and N stable isotopic composition of suspended organic matter in the Mandovi estuary. *Environ. Monit. Assess.*, 175: 501-517.
- Mishra, S., and Panigrahy, R. C. (1999). Zooplankton ecology of the Bahuda estuary (Orissa), east coast of India. *Indian Journal of Marine Science* 28, 297-301.
- Mitra, A., Zaman, S., Sett, S., Raha, AK and Banerjee, AK 2014. Phytoplankton cell volume and diversity in Indian sundarban. *Ind J Mar Sci* 43, 2 208-215.
- Moura, A. N., Bittencourt-Oliveira, M. C & Nascimento, E. C. (2007). Benthic Bacillariophyta of the Paripe River estuary in Pernambuco state, Brazil. *Braz. J. Biol.* 67(3): 393-401.
- Murugan, A., 1989. Ecobiology of Cuddalore, Uppanar backwaters, Southeast coast of India. *Ph.D., Thesis, Annamali University, India*.
- Murugesan, P., 2002. Benthic biodiversity in the marine zone of Vellar estuary (Southeast Coast of India). *Ph. D., Thesis Annamalai University, India*, 359pp.
- Nair, VR 2002. Status of flora and fauna of Gulf of Kachchh, India NIO, Goa, pp 1-258.
- Newell, R. C., Seiderer, L. J., & Hitchcock, D. R. (1998). The impact of dredging works in coastal waters: a review of the sensitivity to disturbance and subsequent recovery of biological resources on the sea bed. *Oceanography and Marine Biology: An Annual Review*, 36, 127-178.
- NIO National Institute of Oceanography 1998. Environmental studies for proposed BPCL jetty and associated facilities at Kandla Part-I Rapid marine EIA, May 1998 NIO Mumbai.
- Omori, M. and Ikeda, T. (1984). Methods in Marine Zooplankton ecology. John Wiley & Sons, New York.
- Parasharya D and Patel B. (2014). Spawning aggregation of *Melibe viridis* Kellart (1858) from Gulf of Kachchh-Western India. *International Journal of Scientific and Research Publication*, 4(3), ISSN 2250-3153.
- Parulekar, A. H., Dhargalkar, V. K., & Singbal, S. Y. S. (1980). Benthic studies in Goa estuaries. Part 3. Annual cycle of macrofaunal distribution, production and trophic relations. *Indian J Mar Sci*.

- Perumal P, Sampathkumar P, Karuppasamy PK (1999) Studies on the bloom-forming species of phytoplankton in the Vellar estuary, southeast coast of India. *Ind J Mar Sci* 28: 400-403.
- Pielou, E C 1966. The measurement of diversity in different types of biological collections. *J Theoret Biol* 13, 131-144.
- Plafkin, J. L., Barber, M. T., Poter, K. D., Gross, S. K. and Highes, R. M. 1989. *Rapid bioassessment protocol for use in streams and rivers for benthic macro invertebrates and fish*. EPA/444/ 4-89/001. Office of water regulation and standards. U.S. Environmental Protection Agency, Washington DC, USA.
- Prabhakar. C., K. Saleshrani & Enbarasan 2011. Studies on the ecology and distribution of phytoplankton biomass in Kadalur coastal zone Tamil Nadu, India. *Curr. Bot.*, **2(3)**: 26-30.
- Ramakrishna, D A 2003. Manual on identification of schedule molluscs from India 40pp
- Ramakrishna, T C R., Sreeraj, C., Raghunathan, R., Raghuraman, P and Yogesh Kumar, J S 2011. An account of additions to the Ichthyofauna of Andaman and Nicobar Islands Records of the Zoological Survey of India, Occasional Paper no 326, 1-140 Published-Director, Zool Surv India.
- Rao, K.K and Balasubramanian, T. (1996). Distribution of Foraminifera in the Cochin Estuary. *J.mar.biol. Ass. India*. 38(1 and 2): 50-57.
- Reid, G. K, 1961. Ecology in inland waters and estuaries. New York.375
- Reid, G. K., Wood, R. D. (1976). Ecology of inland waters and estuaries. Toronto, Ontario, D. Van Nostrand Co., pp. 138–146.
- Reid, G. K., Wood, R. D. (1976). Ecology of inland waters and estuaries. Toronto, Ontario, D. Van Nostrand Co., pp. 138–146. Thakur, B., Chavda, C., & Salvi, H. (2015). Phytoplankton diversity at some selected sites of the Gulf of Kachchh, Gujarat, India. *Bulletin of Environmental and Scientific Research*. 4(4): 7-12. ISSN. 2278-5205.
- Saravanan, K. R., Sivakumar, K., and Choudhury, B. C. (2013). “Important coastal and marine biodiversity areas of India,” in *Coastal and Marine Protected Areas in India: Challenges and Way Forward*, ENVIS Bulletin: Wildlife & Protected Areas, Vol. 15, ed. K. Sivakumar (Dehradun: Wildlife Institute of India), 134–188. doi: 10.1007/978-3-642-38200-0_30
- Shannon, C E and Wiener, W 1949. The Mathematical theory of Communication Univ of Illinois Press, Urbana
- Sinha B. and M. R. Islam, 2002. Seasonal variation in zooplankton population of two lentic bodies and Assam State Zoo cum Botanical garden, Guwahati, Assam, *Eco.EnvIRON. Cons.*, 8: 273-278.
- Sivasamy, S.N., 1990. Plankton in relation to coastal pollution at Ennore, Madras coast. *Indian J. Marine Sci.*, 19: 115-119.
- Sreenivasulu, G., Jayaraju, N., Raja Reddy, B.C.S., Prasad, T. L., Nagalakshmi, K., Lakshmana, B. (2017). Foraminiferal research in coastal ecosystems of India during the past decade: A review. *Geo ResJ*. 13: 38–48.
- SubbaRao, N V., Surya Rao, K V and Maitra, S 1991. Marine molluscs State Fauna Series 1, Part 3 Fauna of Orissa. Zool Surv India, Kolkata, 1–175.
- Tabassum, A. and Saifullah, S. (2012). *Centric Diatoms from the North Arabian Sea Shelf of Pakistan*. LAP. BOOK Lambert Academic Publishing. ISBN: 978-3-659-28532-5.
- Takai N, Mishima Y, Yorozu A, Hoshika A (2002) Carbon sources for demersal fishes in the western Seto Inlan Sea, Japan, examined by delta 13C and delta 15N analyses. *Limnol Oceanogr*. 47(3):730-741.

- Taylor, B. E. (1998). Analyzing population dynamics of zooplankton. Published by the American Society of Limnology and Oceanography, Inc. *Limnol.Oceanogr* 33(6, part 1): 1266-1273.
- Thakur, B., Chavda, C., & Salvi, H. (2015). Phytoplankton diversity at some selected sites of the Gulf of Kachchh, Gujarat, India. *Bulletin of Environmental and Scientific Research*. 4(4): 7-12. ISSN. 2278-5205.
- Thangaraja, G. S., 1984. Ecobiology of the marine zone of the Vellar estuary. *Ph. D. Thesis, Annamalai University, India.*
- Thirunavukkarasu, K., Soundarapandian, P., Varadharajan, D., Gunalan, B.(2013).Zooplankton Composition Structure and Community Structure of Kottakudi and Nari Backwaters, South East of Tamilnadu. *J. Environ Anal Toxicol.* 4(1): 200. doi:.10.4172/2161-0525.1000200
- Thivakaran, GA and Kundu, S 2012. Ecology of Macrobenthos in the Coastal Waters of Gulf of Kachchh, Gujarat In, *Marine Biodiversity, Present Status and Prospects* Eds, P Santhanam and P Perumal Narendra Publishing House, New Delhi 207-225.
- Tiwari, L.R and Nair, V.R. (1998). Ecology of Phytoplankton from Dharmatar creek, West Coast of India. *Indian. J. MarineScience.* 27 (3 & 4).
- Uptake, A. (1999). Primary production by phytoplankton and microphytobenthos in estuaries. *Estuaries*, 29, 93.

Annexure -H

CSR Activities at Decendayal Port Trust

Details of CSR

Sr. No	Year	Board Resolution For Budget Provision	Board Approved Budget Provision	Board Resolution for approval of the CSR activities	Board Approved Amount For CSR Activities	Actual exp. upto Nov'20 (Rs. In Lakhs)	Net balance (Rs. In Lakhs)	Remarks
1	2	3	4	5	6	7	6-7	
1	2011-2012	369 of 28.03.2012	3.00 Cr					
2	2012-2013	17 of 31.05.2012	4.00 Cr					
3	2013-2014	99 of 30.09.2013	6.43 Cr	64 of 30.08.2012	564.00 Lakh	564.00	Nil	Works completed
4	2014-2015	322 of 21.11.2014	1.07 Cr	20 of 16.04.2015	236.22 Lakh	188.18	8.04	Works in progress
5	2015-2016	151 of 12.02.2016	1.50 Cr	48 of 12.08.2016	28.00 Lakh	5.00	23.00	Works in progress
6	2016-2017	138 of 06.01.2017	2.60 Cr	52 of 2.8.2017	140.30 Lakh	146.00	-5.70	Works completed
7	2017-2018	41 of 2.08.2017	7.02 Cr	15 of 04.05.2018	155.10 Lakh	115.37	39.73	Works in progress
8	2018-19	51 of 07.08.2019	6.70 Cr	111 of 4.12.2018	154.90 Lakh	50.50	104.40	Works in progress
					1278.52 Lakh	1069.05	209.47	
9	2019-20	58 of 10.10.2019	5.49 Cr	92 of 06.12.2019	1838.57 Lakh	Nil		MoS approval is awaited
		Total	37.81 Cr		3117.09 Lakh	Spent in PM Fund for COVID-19-800 Lakhs		

Year-wise details of CSR works undertaken by DPT during 2012 – 13 to 2019 – 20 are given in **Tables 7.3a, 7.3b, 7.3c, 7.3d, 7.3e, 7.3f and 7.3g.**

Table 7.3a: CSR Works Undertaken by DPT during 2011-12 and 2012 – 13

Sl. No.	Name of Work	Cost (Rs. In lakhs)
1	Repair of road from Dr. Baba Saheb Ambedkar Circle to NH 8A (via Ganesh Nagar)	518
2	Repair of road from S.T. Bus Stand to Sunderpuri Cross Road via Collector Road	
3	Repair of road from NH 8A Railway Crossing to Maninagar (along railway track)	
4	Repair of road from Khanna Market Road (Collector Road) to Green Palace Hotel	
5	Construction of internal roads at “Shri Ram” Harijan Co-operative Housing Society (near Kidana)	
6	Construction of cremation ground and graveyard with other facilities at Vadinar	19.44
7	Providing cement concrete internal roads in Village Vadinar Stage - I	16.16
8	Approach road provided for developing tourism at Village Veera near Harsidhi Mata Temple	4.65
9	Water tank along with R.O. provided near developing tourism area	0.30
10	Creating facilities of flooring and steps surrounding lake to stop soil erosion and attract tourists at Village Veera.	4.80
	TOTAL	563.35

Table 7.3b: CSR Works Undertaken by DPT during 2014-15

Sl. No.	Name of Work	Cost (Rs. In lakhs)
1	Construction of community hall – cum – school at Maheshwari Nagar, Gandhidham	51.90
2	Renovation of “Muktidham” (cremation ground) at Kandla	10.65
3	Sunderpuri – 1 Valmiki Community Hall	5.00
4	Sunderpuri – 2 Valmiki Community Hall	5.00
5	Ganeshnagar Community Hall	10.00
6	Jagjivan Maheshwari Community Hall	10.00
7	Various works of road at Sapnanagar	99.19
8	Construction of compound wall in the dam of Jogninar Village	14.48
	TOTAL	206.22

Table 7.3c: CSR Works Undertaken by DPT during 2015-16

Sl. No.	Name of Work	Cost (Rs. In lakhs)
1	Construction of Bus Stand at Vadinar Village	10.00
2	Providing drainage system at Vadinar Village	6.00
3	Providing and laying of water supply lines in Vadinar Village	6.00
4	Road from Gandhidham Post Office to Merchantile Marine Department Office along with toilet facilities	60.00
5	Construction of toilets for girls / women at Khari Rohar, Village	3.00
6	Construction of toilets for girls at Mathak Primary School, Mathak, Village	3.00
	TOTAL	88.00

Table 7.3d: CSR Works Approved by DPT Board for 2016-17

Sl. No.	Name of Work	Cost (Rs. In lakhs)
1	RCC community hall at Harsidhi Mata Temple, Village Veera, Anjar Taluka	19.00
2	Fabricated Community Hall at Sanghad Village, Anjar Taluka	21.00
3	CSR Works for Shri Maheshwari Meghavad Samaj, Gandhidham at graveyard behind Redison Hotel	8.00
4	CSR Works for Shri Dhanraj Matiyadev Mukti Dham, Sector 14, Rotary Nagar, Gandhidham	30.50
5	CSR Works for Nirvasit Harijan Co-operative Housing Society, Gandhidham Health Cum Education Centre	41.00
6	CSR Works for Shri Rotary Nagar Primary School, Gandhidham	2.80
7	CSR Works at NU-4, NU-10(B) Sapnanagar & Saktinagar, Golden Jubilee Park at Gandhidham	18.00
	TOTAL	140.30

Table 7.3e: CSR Works Approved for 2017-18

Sl. No.	Name of Work	Proposal Received from / / Name of Organization / N.G.O	Cost (Rs. In lakhs)
1	CSR Works at Shri Ganesh Nagar High School, Gandhidham	Principal, Shri Ganesh Nagar Govt High School, Gandhidham	38.30 Lakhs
2	CSR Works for MOLANA AZAD Primary School, Kandla	Shri M L Bellani, Trustee, DPT, Shri Kandla Port Education Society, New Kandla	7.00 Lakhs
3	Grant financial contribution for facility of Army Cantonment for 50 nos. air coolers at Kutch Border Area	Shri Vinod L Chavda, MP	15 Lakhs
4	40% of the estimated cost of providing drainage lines at Tuna and Vandi villages under Swachh Bharat Abhiyan.	Shri Sarpanch, Tuna Village & Vandi village & Shri M L Bellani, Trustee, DPT	Rs. 39.80 Lakhs <i>Approx. estimated Cost Rs.99.50 Lakhs, of which 40% to be contributed by DPT.</i>
5	CSR works for S.H.N. Academy English School (managed by Indian Inst. Of Sindhology – Bharati Sindhu Vidyapeeth), Adipur	Director, S.H.N Academy English School	40 Lakhs
6	Construction of internal roads at Bhaktinagar Society, Kidana	Smt Maltiben Maheshwari, MLA	15 Lakh
	TOTAL		155.10

Table 7.3f: CSR Works Approved for 2018-19

Sl. No.	Name of Work	Proposal Received from / / Name of Organization / N.G.O	Cost (Rs. In lakhs)
1	CSR work to Donate 100 Nos of Computers to Daughters of Martyred Soldiers in the country under the "BETI BACHAO BETI PADHAO" program by Atharva Foundation, Mumbai	Chairman, Atharva Foundation, Mumbai	24.00
2	CSR work to Donate ONE (40 Seater) School Bus for Deaf Children Students for the Institute of Mata Lachmi Rotary Society, Adipur	Mata Lachmi Rotary Society, Adipur	18.00
3	CSR work to Providing One R.O Plant with Cooler at PanchyatPrathmikSala, Gadpadar Village for the ANARDE Foundation, Kandla&Gandhidham Center.	Dist. Rural Development Officer, Annarde Foundation-Kandla & Gandhidham	1.50
4	CSR work for Providing Drainage Line at MeghparBorichi village, AnjarTaluka	Shri Vasanbhai Ahir, MLA, Gujarat Govt	25.00
5	CSR work for Construction of Health Centre at Kidana Village	Shri Vinod L Chavda, MP	13.00
6	CSR work to provide 4 Nos. of Big Dust Bin for MithiRoharJuth Gram Panchayat.	Shri Sarapanch, Mithi RoharJuth Gram Panchayat	3.40

Sl. No.	Name of Work	Proposal Received from / / Name of Organization / N.G.O	Cost (Rs. In lakhs)
7	CSR work for Renovation & construction of shed at CharanSamaj, Gandhidham –Adipur.	Shri Vinod L Chavda, MP	10.00
8	CSR Work for Renovation/Repairing of Ceiling of School Building at A. P Vidhyalay, Kandla.	Smt Maltiben K. Maheshwary, MP, Gandhidham.	10.00
9	CSR work for Construction of Over Head Tank & Providing 10 Nos of Computers (for students) of NavjivanViklangSevashray, Bhachau, Kutch	Shri Jitendra Joshi, Founder Secretary, Shri Navjivan Viklang Sevashray, Bhachau, Kutch	9.50
10	CSR work to Provide Books & Tuition fees for Educational facilities to weaker section children of ValmikiSamaj, Kutch.	Shri Manohar Jala, Chairman of "National Commission of Safai Karamcharis"	2.00
11	CSR work to provide Water Purifier & Cooler for the ST. Joseph's Hospital, Gandhidham	Smt. Maltiben K Mahewari, MLA ,Gandhidham	1.50
12	CSR work for Construction of Second Floor (Phase – I) for Training Centre of "GarbhSanskran Kendra" "Samarth Bharat Abhiyan" of Kutch Kalyan Sangh, Gandhidham	Shri Vinod L Chavda, MP, Kutch	37.00
TOTAL			154.90

Table 7.3g: CSR works approved for the year 2019-20 (approval from Ministry of Shipping still awaited)

Sl. No.	Name of Work	Proposal Received from / / Name of Organization / N.G.O	Cost (Rs. In lakhs)
1	CSR activities for Providing Drainage line at Nani Nagalpar village.	Sarpanch of Village:-Nani Nagalpar, Taluk: Anjar.	3.00
2	CSR activities for Development of ANGANWADI Building at School no- 12 at Ward no 3 & 6 at Anjar.	Shri Vasanhbai Ahir, MLA	7.00
3	CSR activities for Improving the facilities of Garden at Sapna Nagar(NU-4)& (NU-10 B),Gandhidham.	Shri K P Maheshwari, Resident Sapnanagar, Gandhidham	18.00
4	CSR activities for Providing of Plastic Shredding Machine to Mirror Charitable Trust, Gandhidham.	Mirror Charitable Trust ,Gandhidham	4.75
5	CSR activities for development of School premises of Shri Guru Nanak Edu. Society, Gim.	Shri Guru Nanak Education Society, Gandhidham.	30.00
6	CSR activities for the improvement of the facilities at St. Joseph Hospital & Shantisadan at Gandhidham	St. Joseph Hospital Trust, Gandhidham	20.00
7	CSR activities for the improvement of the facilities at SVP (SardarValabhbbhai Patel) Multipurpose Hall at Gandhidham	Request from MarwadiYuva Munch & UNION Gandhidham	500.00
8	Consideration of Expenditure for running of St Ann's High School at Vadinar of last 5 years 2014 to 2019 under CSR.	Proposal from COM, OOT Vadinar, DPT	825.00
9	CSR activities for development of school premises of Shri Adipur Group Kanya Sala no-1 at Adipur	Principal, Shri Adipur Group KanyaSala, Adipur	6.50
10	CSR activities for development of school premises of Shri Jagjivan Nagar Panchyat Prathmiksala, Gandhidham.	Principal, Shri Jagjivan Nagar Panchyat Prathmiksala, Gandhidham.	16.50
11	CSR activities for development of school premises of Ganeshnagar Government high school, Gandhidham.	Shri Vinod L Chavda, MP, Kutch	9.00
12	CSR activities for improving greenery, increase carbon sequestration and beat Pollution at Kandla, DPT reg.	Work awarded to Forest Department , Bhuj	352.32
13	CSR activities for providing infrastructures facilities at "Bhiratna Sarmas Kanya Chhatralaya" under the Trust of Samaj Nav- Nirman at Mirjapur highway, Ta Bhuj.	SamajNav- Nirman at Mirjapur highway, Ta Bhuj.	46.50
TOTAL			1838.57

List of CSR applications received from various NGOs , Organizations , Village Sharpanchs etc for the FY 2021-22 .

Sr.No	Name of Scheme	Proposal Received from / Name of Organization / N.G.O	Brief Details
1	CSR activities for the development of gardening at Sector -5 , Gim	Shri Sarvodaya Co-Operative Housing Society Ltd	Appx Cost – Rs 25.00 Lakhs Cost for – Comp wall, Benches, Plantation, walkway, other facilities (Land is reserved for Garden development only since from 50 years)
2	CSR activities for providing various facilities in SHRI GANESHNAGAR GOVT HIGHSCHOOL, GANDHIDHAM	Principal of School	Appx cost –Rs 20.00 Lakhs (Two times CSR works carried out at school by DPT)
3	CSR activities for the VadhiyarVankarSamajvaadi, NaviSunderpuriGim	SmtMaltiben K Maheswari, MLA	Appx Cost Rs 6.00 Lakhs Cost for Const. of Comp Wall
4	CSR activities for Construction work of Cabin at Oslo Area- Gim	SmtMaltiben& Shri VinadChavda	Cost not mentioned.
5	CSR activities & Land requirement forAkhil Kutch SamastaMeghvanshiGurjarmeghwal Charitable Trust ,Gim.	Shri Akhil Kutch SamastaMeghvanshiGurjarmeghwal Charitable Trust. Shri Dharmendra R Gohil	Cost Not mentioned. (demand of Land for development of SAMAJ VADI in Gandhidham)
6	CSR Activities for providing Water supply pipe line, Play ground and sports equipment, electric facilities, drinking water facilities for poor people & Fishermen at VANDI Village.	Shri R RKhambhra, PRO , Collector Office, Bhuj.	Appx Cost Rs 51.00 Lakhs (Last year also applied by village Sarpanch) & Recommended by Shri VASANBHAI AHIR, MLA, Shri V L Chavda, MP)
7	CSR activities for the Tuna village,	Sarpanch, Tuna village	Appx Cost Rs. 25 Lakhs Cost for :-

List of CSR applications received from various NGOs , Organizations , Village Sharpnchs etc for the FY 2021-22 .

Sr.No	Name of Scheme	Proposal Received from / Name of Organization / N.G.O	Brief Details
	Ta -Gim		2 No Fab shed 20'x20'x1250= 10 Lakh 2 Nos of Agnawadi =10 Lakh Fab shed for school=5 Lakh
8	CSR activities for the Global Vision India Foundation, Gim	Global vision India Foundation, G'dham	Requirement of Land –OR- Old building at Gandhidham for foundation of welfare activities.
9	CSR activities for the UNITED ORPHANAGE FOR THE DISABLED, TAMIL NADU	UNITED ORPHANAGE FOR THE DISABLED, TAMIL NADU	Cost Rs 25,000.00 (Winter sweaters for children)
10	CSR activities for the Garden Development on already bounded area with Compound wall near Plot no 448 Sector-1/A, Gandhidham.	Residents, near Plot no 448, Sector-1/A, Gim.	AppxCost Rs 20.00 Lakhs (Requirement to provide benches, drinking water facility, plantation, lightings & walkways in side bounded area)
11	CSR activities for donation of Land for the Shri SUNDARPUI Govt Primary School, Gim	SmtMalti ben Maheshwari, MLA	(request for Land Requirement)
12	CSR activities for Extension of Adarsh Primary School building, Adipur	GandhidhamMatri Mandal, English Medium School, Adipur	Appx Cost Rs. 40.00 Lakhs (Construction for 4 Rooms extension) (Trust registered under Societies Registration Act XXI -1860, Reg No F-42 dtd 23.9.1965. Land belong to Trust)
13	CSR Activities for providing HD projector for KANYA MAHA VIDYALAYA, Adipur	Principal, KANYA MAHA VIDYALAYA, Adipur	Cost Rs 1.50 Lakhs (School Managed by G'dhamMaitry Mandal, Adipur)

List of CSR applications received from various NGOs , Organizations , Village Sharpnchs etc for the FY 2021-22 .

Sr.No	Name of Scheme	Proposal Received from / Name of Organization / N.G.O	Brief Details
14	CSR activities for DONATION various Medical Equipment for the Hospital of Gandhidham Jain SevaSamiti, Adipur	Gandhidham Jain SevaSamiti, Adipur	Cost for :- 1) Fresenius Haemodialysis Machine Rs 38.00 Lakh 2) Maltislice Helical CT Scanner- Rs 52.00 Lakhs 3) Others Rs 54.00 Lakhs (Total Appx Cost Rs 144 Lakhs)
15	CSR activities for SHRI VIDI JUTH GRAM PANCHAYAT, Vidi, Anjar	Sarpanch, Vidi Gram	Appx Cost Rs 30.00 Lakhs Cost for- Drainage , Garbage vehicle, and Cattle shed (Already applied earlier at Sr-5/12)
16	CSR activities for SOS CHILDREN'S VILLAGES INDIA, Madhapar, Bhuj	Director, SOS Children's Village of India-Bhuj	Appx Cost Rs 31.00 Lakhs (request for Financial support towards parentless and abandoned Children Education support located at Bhuj) & support to women working in SOS.
17	Gujarat Biodiversity Board, Gandhinagar invites to involved National & Global endeavour of conservation of biodiversity by creating financial partnership with GBB under CSR programme of expenditure to be incurred 187 Lakh.	GUJARAT BIODIVERSITY BOARD, GANDHINAGAR	Requirement- Financial Support from DPT for AppxRs 1.88 Cr. (Cost for various meetings, collection of primary data from villagers , processing of documentation, printing , TA DA of Technical support & Miscexp for 150 Peoples Biodiversity Register (PBR).

List of CSR applications received from various NGOs , Organizations , Village Sharpnachs etc for the FY 2021-22 .

Sr.No	Name of Scheme	Proposal Received from / Name of Organization / N.G.O	Brief Details
18	CSR activities for providing furniture & Home appliances for ROJAVANAM TRUST at Madurai.	Shri Arul Kannan, Director	Appx Cost Rs 30 Lakhs (seeking help to provide facilities to Aged & Homeless people living in Trust and Purchasing of New Ambulance)
19	CSR activities for providing Dialysis Machine for treatment of Kidney patients at "ST JOSEPH'S HOSPITAL TRUST" at Gandhidham.	Sr. Franciline, Administrator of Hospital.	Appx Cost Rs 31.36 Lakhs (Cost of 5 Nos of Dialysis Machines for treatment of kidney patients)
20	CSR activities for providing facilities in Girls Hostel of Gasturba Gandhi BalikaVidhyalay, Gandhidham.	Shri Vinod L Chavda, MP	Appx cost Rs 30 Lakhs. (Cost of Comp Wall, Entrance gate, Girls toilets etc)
21	CSR works for providing Oxygen Generator Plant and 45 KV Silent Generator for COVID HOSPITAL at Swami LilashahKutia, Adipur.	Secretary, BHARAT VIKAS PARISHAD, Gandhidham	Appx Cost Rs 80.00 Lakhs (Facilities for 100 Beds of COVID patient which it to be extend upto 240 Beds)
22	CSR works for providing Two Numbers of Oxygen Concentrator and others medical equipment for the Trust ,Antarjal, Gim.	President SHRI SARV JEEV KALYAN TRUST, ANTARJAL, Gandhidham	Appx Cost Rs21.50 Lakhs (Facilities to be provided for the treatment of CORONA PATIENTS at their trust.)
23	CSR works for providing Fabricated Shed , Construction of Compound Wall and Land levelling for the Cattle of GauSevaSamiti-Tappar at Gram-Tappar, Ta Anjar.	Shri Vinod Chavda, MP & Presedent , GauSevaSamiti, village Tappar, Ta-Anjar	Appx Cost Rs84 Lakhs (Facilities to be provided for Cattle shelters at Village.) (Land belongs to Gram-panchayat)
24	CSR works for Construction of Auditorium Hall at RSETI (Rural Self Employment Training Institute) at	Shri Vinod Chavda, MP & Director of RSETI, Bhuj	Cost not mentioned. (Facilities to be provided

List of CSR applications received from various NGOs , Organizations , Village Sharpnchs etc for the FY 2021-22 .

Sr.No	Name of Scheme	Proposal Received from / Name of Organization / N.G.O	Brief Details
	Bhujodi-Bhuj.		for the people needs Self-employment activities.)
25	CSR works for Providing of Furniture for the School "SHRI GALPADAR PANCHAYAT PRATHMIC KUMAR GROUP SALA " atGalpadar Village Ta Gim.	Principal, SHRI GALPADAR PANCHAYAT PRATHMIC KUMAR GROUP SALA " atGalpadar Village Ta Gim.	Cost not mentioned. (Facilities to be provided for the Students of Workers & poor village people who study in the school.)
26	Construction of Shed, hall and Gate for the DADA Bhagwandas Charitable Trust, Adipur. (Sr no -4)	Shri Vinod Chavda, MP & DADA BHAGWANDAS CharitableTrust, Gandhidham	<u>As per CSR Guideline-</u> ➤ Promoting gender equality and empowering women ➤ Eradicating extreme hunger and poverty (Considered shed and hall) Fab Shelter Shed - 30'x100' x 1250=37.00 Lakh & RCC Hall – 20'x100'x1500=30.00 Lakh (Appx Cost Rs67.00 Lakhs) Land authority belongs to Trust given by GDA and NOC given by SRC.Doc submitted.
27	CSR work for reconstruction of the Internal Roads of the Sector-9B-C and Sector-10 area in Gandhidham.	President, Shri TejaKangad, The Gandhidham Chamber of Commerce and Industry, Gandhidham.	Cost not mentioned.

List of CSR applications received from various NGOs , Organizations , Village Sharpanchsetc for the FY 2021-22 .

Sr.No	Name of Scheme	Proposal Received from / Name of Organization / N.G.O	Brief Details
	<u>CSR Applications kept pending in last year Agenda:-</u>		
27	CSR Activities for providing Water supply pipe line, Play ground and sports equipment, electric facilities, drinking water facilities for poor people & Fishermen at VANDI Village. (Sr no-3)	Sarpanch ,Village-VANDI , Ta- Anjar (Recommd. By Shri VASANBHAI AHIR, MLA, Shri V L Chavda, MP)	<u>As per CSR Guideline-</u> ➤ Env Sustainability ➤ Eradicating extreme hunger and poverty (to be Consider for health Center ,Drainage line, Water sump etc activities) (Appx Cost - 51.00 Lakhs) (Land authorization of Gram Panchayat)
28	Construction of Shed, hall and Gate for the DADA Bhagwandas Charitable Trust, Adipur. (Sr no -4)	DADA BHAGWANDAS CharitableTrust, Gandhidham (Recommd. By Shri V L Chavda, MP)	<u>As per CSR Guideline-</u> ➤ Promoting gender equality and empowering women ➤ Eradicating extreme hunger and poverty (Considered shed and hall) Fab Shed - 30'x100' x 1250=37.00 Lakh & RCC Hall – 20'x100'x1500=30.00 Lakh (Appx Cost Rs 67.00 Lakhs) Land authority belongs to Trust given by GDA and NOC given by SRC. Doc submitted.
29	10 Nos of Computers required for ShirMaheswarinagar Panchayat Girls Primary School, Gandhidham& Boys Group School, Gandhidham. (Sr no-8)	Maheswarinagar Panchayat Primary Kanya Sala, Gandhidham (Contact no 9913903686)	AppxRs 5.00 Lakhs <u>As per CSR Guideline-</u> ➤ Promotion of Education (to be consider for 20 Computers)

List of CSR applications received from various NGOs , Organizations , Village Sharpnchsetc for the FY 2021-22 .


Sr.No	Name of Scheme	Proposal Received from / Name of Organization / N.G.O	Brief Details
			Visited the site. Land belongs to MahewariMeghwadSamaj given by SRC for school purpose, doc are awaited.
30	Construction of Shed and Roof at JeparMatiyadev, shamsanbhumi at Kidana village & Maheswari Community Hall at JuniSundarpuri, Gandhidham. (Sr no-10)	Shri VINOD CHAVDA, MP	AppxRs 15.00 Lakhs (Land authorization not mentioned)
31	Drainage, road, Dust bins, & shed for Cattle shelters at VIDI Village, Ta –Anjar. (Sr no- 12)	Village- VIDI, Ta: Anjar	AppxRs 30.00 Lakhs <u>As per CSR Guideline-</u> ➤ Env Sustainability ➤ Eradicating extreme hunger and poverty (Consider for Garbage vehicle & Drainage Cost)
32	Education, Women empowerment and Primary health care services at Kutch area. (Sr no-13)	Light of Life Trust, Mumbai.	Cost not mentioned.
33	Request for Help Divyang persons to employment by providing machineries. (Sr no-14)	Kutch DivyangSangthan, Gandhidham.	Cost not mentioned
34	Construction of 2 nd Floor of Shri MaheswariMeghwadSamaj, Gandhidham. (Sr no-20)	Shri MaheswariMeghwadSamaj, Gandhidham	AppxRs. 15.00 Lakhs (Visited the site and Land ownership documents awaited) (Name plate of DPT fixed at the Asset)

List of CSR applications received from various NGOs , Organizations , Village Sharpanchsetc for the FY 2021-22 .

Sr.No	Name of Scheme	Proposal Received from / Name of Organization / N.G.O	Brief Details
35	Installation of Mini Science Center at Anjar and Gandhidham. (Sr no-21)	STEM Learning Pvt Ltd, Mumbai.	Cost not mentioned.
36	CSR work for Shri Rampar Gram Panchayat. ➤ Wall Plastering for Cattles -7 Lakhs ➤ Shed for Cattel's-15 Lakhs (Sr no-25)	Shri Sarpanch, Rampar Village.	AppxRs 22.00 Lakhs (Land authorization of Gram Panchayat and under taking submitted by applicant)
37	CSR activities for the 45,000 Patients over the period of 3 years by "SMILE FOUNDATION", Mumbai. 1. Concept for Nutrition covering 3 years 2. Concept for Mobile Health Unit reaching beneficiaries for 3 years 3. Concept for Vocational Training with NGO (Sr no-29)	Proposal from "SMILE FOUNDATION " Mumbai.	Appx Cost- Rs 539 Lakhs for 3 years
38	Development of Park in Public utility plot in between Block "C" & "D" of Sapna Nagar (NU-4) , Gandhidham (Sr no -31)	Shri RAVI MAHESHWARI, DPT	Land belongs to DPT earmarked for recreational purpose. (Total Cost – Rs88.75 Lakhs)
39	CSR works for NariJanshsktiVikas Foundation at Gandhidham near Shakti Nagar. (Sr no-33)	NariJanshsktiVikas Foundation, Ahmedabad	➤ Promoting gender equality and empowering women ➤ Env Sustainability ➤ Under promotion of education (Consider for Computers with printers, Sewing machine & RO plant Cost Rs 48 Lakhs)

Annexure -I

DEENDAYAL PORT TRUST



Administrative Office Building
Post Box NO. 50
GANDHIDHAM (Kutch).
Gujarat: 370 201.
Fax: (02836) 220050
Ph.: (02836) 220038

www.deendayalport.gov.in

NO.EG/WK/4783/V/131

Dated : 05/02/2021

To,
M/s Precitech Laboratories Pvt Ltd,
1st Floor, Bhanujyot Complex,
Plot No C5/27, B/h Panchratna Complex,
Nr. GIDC Char Rasta,
VAPI-396195.

Sub: **Work order** for "STRENGTHENING OF EXISTING ENVIRONMENTAL MANAGEMENT CELL AT DEENDAYAL PORT TRUST: Appointment of environment experts for two years further extendable for one year"-**reg.**

Ref: 1) Tender dated 21.06.2019 submitted by M/s Precitech Laboratories Pvt.Ltd, Vapi.
2) Letter of Acceptance vide no-EG/WK/4783/V/100 dtd 01(04).01.2021
3) Letter from DPT no E/WK/4783/V/103 dtd 06.01.2021
4) Performance Guarantee submitted by M/s Precitech Laboratories Pvt Ltd in the form of Bank Guarantee of Rs. 3,60,000.00 vide Bank Guarantee no. 1102921BG0000016 dated 19.01.2021 issued by State Bank of India, Vapi.

Sir,

Kindly refer above cited Letter of Acceptance dtd 01(04).01.2021.

- 2) You shall have to provide Key Experts as per tender requirement during the entire contract period. Accordingly, you shall have to submit the qualification and experience certificates of the Key experts to be appointed at DPT, as per tender conditions for verification & approval.
- 3) Please submit the Agreement of contract as per tender conditions no 1.29.
- 4) Kindly commence the work on or before 15.02.2021.


.....Cont.....

- 2 -

Please note that the time period for providing Consultancy services for the subject work will be initially for two years and further extendable for one year on mutual consent as per tender conditions.

Thanking you.

Yours faithfully,


Superintending Engineer (Design & EMC (i/c))
Deendayal Port Trust

Annexure -J

DEENDAYAL PORT TRUST

ISO 9001 : 2008 : ISO 14001 : 2004

Ph. : 02836-220167

Fax: 02836-233172

website: deendayalport.gov.in

e-mail : secretary@deendayalportgov.in



General Administration Deptt.
Administrative Office Building,
Post Box No. 50,
Gandhidham (Kutch) 370 201

By Speed Post / E-mail

No. GA/PS/4292/HE(PF)/2017/ 304

Dated, 17 January, 2022

OFFER OF CONTRACTUAL ENGAGEMENT AS MANAGER(ENVIRONMENT), IN DEENDAYAL PORT TRUST.

With Reference to your application for contractual engagement as Manager – Environment, in response to the advertisement, inviting applications for the subject position, on assessment and interview before the Services Selection Committee on 06.01.2022, the Competent authority has been pleased to offer the contractual engagement as Manager (Environment) in Deendayal Port Trust, purely on contractual basis, subject to the following terms and conditions :

a) Roles & Responsibilities

- Develop, implement and manage long term port environmental programmes such as the Green Marine Programme, sustainability plan, air strategies, tenant environment plan and tenant lease management.
- Represent the Port in local, state and federal agency meetings.
- Assist in the development and updating of the Port's comprehensive scheme of Harbour improvements and strategic plan.
- Monitor and conduct regular mock drills to train the employees at different levels.

b) Remuneration :-

Your consolidated remuneration per month will be Rs.1,00,000/- (Rupees One Lakh Only). Suitable increase depending upon the performance and variation in the AICP index may be given after successful completion of yearly service. Applicable taxes will be deducted at the time of payment.

c) Period of Contract :

The contract will be for a period of 3 years, extendable by another two years, subject to satisfactory performance.

d) Duty Hours :

You may be posted at/under any department/authority of Deendayal Port Trust, as per requirement, Duty Hours are from 10.00 AM to 06.00 PM or as may be decided by the Administration from time to time. In case of requirement, you may have to work beyond the normal duty hours, for which no other compensation, monetary or otherwise will be considered.

.....
(Mukkannawar Utkarsh Suresh)

Contd....

You will normally be entitled to a weekly off on Sunday. If situation warrants, the weekly day of rest may be changed with prior intimation. For work on any weekly day off / declared national holiday in exigencies of work, a compensatory day of rest as per the convenience of the Administration, in lieu thereof, will be granted and for which no other compensation, monetary or otherwise will be considered.

Failure to report for duty will entail deduction of wages on pro-rata basis.

- e) Medical facility : Only Outdoor Medical treatment facility for self and your spouse will be provided in the Port Trust Hospital. No other medical facilities will be provided to you/ your family.
- f) Leave entitlement : 10 days leave in a year and National Holidays will be given. No other leave will be admissible and for any absence beyond the said leave, pro-rata deduction will be made from the consolidated remuneration.
- g) Accommodation : Suitable accommodation, if available, may be provided, subject to recovery of charges under FR-45A, and the element of HRA excluded from the lumpsum remuneration.
- h) Your engagement on contractual basis is subject to strict adherence to the norms and conduct.
- i) The engagement can be terminated by giving one month's notice in writing from either side. However, in case of unsatisfactory performance or for any act considered derogatory/ detrimental to the interest of Deendayal Port Trust, this contractual engagement will be terminated forthwith.
- j) If you leave without notice or without acceptance of notice of termination, the amount due i.e., consolidated remuneration payable will be forfeited.
- k) You shall not claim any right/title/interest on par with the regular employees of the Port or otherwise.
- l) You shall not have any claim/right whatsoever for regular appointment / absorption in Deendayal Port Trust under any circumstances.
- m) Your contractual engagement is subject to verification of antecedents by the police. If any adverse report is received from the Police, your contractual services are liable to be terminated forthwith.
- n) You will not be permitted to take any other assignment during the period of contract with Deendayal Port Trust.

.....
(Mukkannawar Utkarsh Suresh)

Contd....

- l) On official tour outside Head Quarters, you will be entitled to TA/DA as admissible under the rules.
- m) The terms and conditions shall be amended / modified depending upon the requirement of the Port. Any dispute(s)/difference(s) shall be decided solely by the Chairman, Deendayal Port Trust, which shall be final and binding.
- n) You are required to submit discharge letter / relieving letter from your present employer at the time of joining Deendayal Port Trust, without you may not be allowed to join.
- o) The contractual engagement is subject to your being found medically fit as per the requirements of Deendayal Port Trust.

2. You have to report for medical examination before the Medical Board of DPT at Gopalpuri Hospital on any working day between 10.00 hrs to 12.00 hrs.

3. If you agree to the above terms and conditions, you may convey acceptance by signing the duplicate of the letter in token of your acceptance and submit the same to this office and call at this office with all certificates and two copies of passport size photographs latest by 27th January, 2022 failing which the offer of contractual engagement stands automatically cancelled.


Secretary
Deendayal Port Trust

To
Shri. Mukkanawar Utkarsh Suresh,
21/1, Madhukunj Housing Society,
Near Canara Bank, Panchavati,
Pashan, Pune, Maharashtra - 411008.
Email : utkaish@gmail.com

I accept the above terms and conditions and will report for duty on _____.

Name :

Date :

Copy to: CMO - for conducting Medical Examination.